



OPEN ACCESS

EDITED BY

Giulia Marrone,
University of Rome Tor Vergata, Italy

REVIEWED BY

Alicia Ruiz,
University of Granada, Spain
Giulia Cafiero,
Bambino Gesù Children's Hospital (IRCCS), Italy

*CORRESPONDENCE

Pasqualina Buono
✉ pasqualina.buono@uniparthenope.it

RECEIVED 26 May 2023

ACCEPTED 28 July 2023

PUBLISHED 10 August 2023

CITATION

Mancini A, Cerulli C, Vitucci D, Lasorsa VA, Parente D, Di Credico A, Orrù S, Brustio PR, Lupo C, Rainoldi A, Schena F, Capasso M and Buono P (2023) Impact of active lifestyle on the primary school children saliva microbiota composition.

Front. Nutr. 10:1226891.

doi: 10.3389/fnut.2023.1226891

COPYRIGHT

© 2023 Mancini, Cerulli, Vitucci, Lasorsa, Parente, Di Credico, Orrù, Brustio, Lupo, Rainoldi, Schena, Capasso and Buono. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Impact of active lifestyle on the primary school children saliva microbiota composition

Annamaria Mancini^{1,2}, Claudia Cerulli³, Daniela Vitucci^{1,2}, Vito Alessandro Lasorsa², Daniela Parente^{2,4}, Andrea Di Credico^{5,6}, Stefania Orrù^{1,2}, Paolo Riccardo Brustio⁷, Corrado Lupo⁸, Alberto Rainoldi⁸, Federico Schena⁹, Mario Capasso^{2,4} and Pasqualina Buono^{1,2*}

¹Department of Movement Sciences and Wellness, University Parthenope, Naples, Italy, ²CEINGE-Biotecnologie Avanzate "Franco Salvatore", Napoli, Italy, ³Department of Movement, Human and Health Sciences, University of Rome "Foro Italico", Rome, Italy, ⁴Department of Molecular Medicine and Medical Biotechnologies, University of Naples Federico II, Naples, Italy, ⁵Reprogramming and Cell Differentiation Lab, Center for Advanced Studies and Technology (CAST), Chieti, Italy, ⁶Department of Medicine and Aging Sciences, University "G. D'Annunzio" of Chieti-Pescara, Chieti, Italy, ⁷Department of Clinical and Biological Sciences, University of Torino, Turin, Italy, ⁸Department of Medical Sciences, University of Torino, Turin, Italy, ⁹Department of Neuroscience, Biomedicine and Movement, University of Verona, Verona, Italy

The aim of the study was to evaluate the effects of Active or Sedentary lifestyle on saliva microbiota composition in Italian schoolchildren.

Methods: Male (114) and female children (8–10 years) belonging to five primary schools in the neighborhoods of Turin were classified as active (A) or sedentary (S) based on PAQ-C-It questionnaire. PCR amplification of salivary DNA targeted the hypervariable V3–V4 regions of the 16S rRNA bacterial genes. DADA2 workflow was used to infer the Amplicon Sequence Variants and the taxonomic assignments; the beta-diversity was obtained by PCoA with the UniFrac method; LEfSe algorithm, threshold at 5%, and Log LDA cutoff at ± 0.5 were used to identify differently abundant species in A compared to S saliva sample. Daily food intake was assessed by 3-Days food record. The metabolic potential of microbial communities was assessed by PICRUSt.

Results: No significant differences were found in individual's gender distribution ($p = 0.411$), anthropometry, BMI ($p > 0.05$), and all diet composition between A and S groups ($p > 0.05$). Eight species were differently abundant: *Prevotella nigrescens* (LDA score = -3.76 ; FDR = 1.5×10^{-03}), *Collinsella aerofaciens* (LDA score = -3.17 ; FDR = 7.45×10^{-03}), *Simonsiella muelleri* (LDA score = -2.96 ; FDR = 2.76×10^{-05}), *Parabacteroides merdae* (LDA score = -2.43 ; FDR = 1.3×10^{-02}) are enriched in the A group; *Gemella parahaemolysans*, *Prevotella aurantiaca* (LDA score = -3.9 ; FDR = 5.27×10^{-04}), *Prevotella pallens* (LDA score = 4.23 ; FDR = 1.93×10^{-02}), *Neisseria mucosa* (LDA score = 4.43 ; FDR = 1.31×10^{-02} ; LDA score = 2.94 ; FDR = 7.45×10^{-03}) are enriched in the S group. A prevalence of superpathway of fatty acid biosynthesis initiation (*E. coli*) and catechol degradation II (meta-cleavage pathway) was found in saliva from A compared to S children.

Conclusion: Our results showed that active children had an enrichment of species and genera mainly associated with a healthier profile. By contrast, the genera and the species enriched in the sedentary group could be linked to human diseases.

KEYWORDS

lifestyle, exercise, saliva, microbiota, children

1. Introduction

The human gut microbiota is intimately associated with different aspects of human health and disease. Its characterization could help diagnosis, prognosis, and therapy settings by giving over 150 times more genetic information than the human genome alone (1). The microbiota composition depends on spatial distribution and age; in general, the microbiota diversity increases over the time and decreases in elderly (2, 3). In children of about 3 years, gut microbiota becomes similar to that of adults, with five predominant bacterial phyla: Firmicutes, Bacteroidetes, Actinobacteria, Proteobacteria and Verrucomicrobia (4). Recent findings established the role of environmental factors on microbiota composition suggesting a close cross-talk between the lifestyle and the diversity of microorganisms populating the intestine (5). To date, the effects of exercise on human health have been thoroughly studied. In the last decade, many evidences supported a dynamic relationship between the composition of gut microbiota (GM) and physical activity levels in animal models (6–10) and in human (11–14). In particular, the theory that exercise is able to enrich the diversity of the human gut microbiota depending on the volume of training resulting in improved health status of the host, has been supported (12, 15). In particular, GM diversity has been associated to cardiorespiratory fitness (11, 16, 17) and to VO₂max in adults (18). Moreover, Barton et al. (18) highlighted in GM, by a metagenomic and metabolomic approach, relative increases in fecal pathways and metabolites, e.g., short-chain fatty acids (SCFAs) produced by microbes, associated with higher muscle turnover and overall health in professional athletes compared with sedentary controls.

While in adults there are some evidences of the influence of physical activity on GM composition, in children or pre-adolescents, very poor results have been provided. Recent reports indicated that the GM profile is associated with the body mass index and could be modulated by exercise training and lifestyle habits in obese children (19–22). Furthermore, several evidences show that the salivary microbiota mirrors the gut microbiota and that some oral bacteria colonize the gut and have been associated both to oral and systemic health. (23–26).

Despite these recent advances, the complete landscape of the association between the saliva profile and lifestyle habits in children is still to be clarified. Further, no data on saliva microbiota composition have been provided in Italian schoolchildren associated to Active compared to Sedentary status, to date. Thus, the principal aim of this study was to analyse the possible association between saliva microbiota compositions and lifestyle in Active compared to Sedentary cohort of 8–10-year-old Italian school-aged children living in the neighborhoods of Turin (northwest Italy).

We conducted this study by hypothesizing that active lifestyle could be associated with saliva microbiota profiles contributing to host health promotion. Indeed, the main aim of our work was to identify the differences in the saliva of Active compared to Sedentary schoolchildren. In order to study the microbiota composition, we sequenced the bacterial 16S rRNA of saliva biospecimens and assessed their differential abundance.

2. Materials and methods

2.1. Participants

One hundred and thirty children (8–10 years) belonging to five primary schools in the neighborhoods of Turin (northwest Italy) were

enrolled. All information on the aim of the study has been provided to children's parents/guardians and teachers as previously described (27).

Children meeting any of the following criteria were excluded from the study: (i) recent infections (1 month prior to sample collection), (ii) having disorders affecting diet or physical activity, and (iii) recent usage of either antibiotic, prebiotics and probiotics supplements (1 month prior to data and sample collection). The enrolled children were classified in two groups: active (A) and sedentary (S) on the basis of Physical Activity Questionnaire for Older Children (PAQ-C-It cut-off score of 2.75), the related procedures are detailed in Lupo et al. (27). Parents/guardians and teachers provided written informed consent for participation to the study, according to the ethical standards provided in the 1964 Declaration of Helsinki. Ethics committee on human research of the University of Turin (9 March 2020: Protocol #134691) and Naples (17 January 2020: Protocol #376/19) approved the study. The procedures used to take anthropometric measures were described in Lupo et al. (27); briefly, stature was measured by a portable stadiometer (Model 214; Seca, Hamburg, Germany), body mass was measured by an electronic scale (Model 876; Seca, Hamburg, Germany), participants' waist circumference was measured in the standing position, midway between the lowest rib and the iliac crest by Ana elastic meter. The Body Mass Index (BMI) was calculated as body mass divided by height squared (kg/m²).

To estimate the daily food intake, all participants filled the questionnaire (3-Days food records). Records were processed using Winfood software (Medimatica S.u.r.l., Colonnella, TE, Italy). Statistical analysis was performed through a one-way ANOVA (Statview software).

2.2. Saliva sample collection and genomic DNA extraction

The donor was asked not to eat and not to use oral hygiene products 1 h before saliva collection. At least 2 mL of unstimulated saliva was collected, put on ice and stored at –80°C until the analysis. DNA was extracted from saliva samples using the MagPurix Bacterial DNA Extraction Kit (ZP02006; Zinexts Life Science Corp.) according to the manufacturer's instructions. DNA was quantified using the Qubit dsDNA BR and HS assay kit (Life Technologies, CA, United States).

2.3. Preparation of the 16S metagenomic sequencing library

PCR amplification was conducted to target the hypervariable V3–V4 regions of the 16S rRNA bacterial genes. Specific primers with barcodes and high-efficiency enzymes were used to perform PCR. The PCR primers were: forward 341F: CCTAYGGGRBGCASCAG; reverse 806R: GGACTACNNGGTATCTAAT. The PCR products of 450–500 bp were collected with 2% agarose gel electrophoresis. To build library, same amount of PCR products from each sample is pooled, end-repaired, A-tailed and further ligated with Illumina adapters. The library QC was performed with Qubit and real-time PCR for quantification and with bioanalyzer to check the insert size distribution. Libraries were sequenced on a paired-end Illumina

platform to generate 250 bp paired-end raw reads. The raw sequencing data are available in Zenodo (<https://doi.org/10.5281/zenodo.7920752>; Publication date: May 10, 2023).

2.4. Bioinformatic analysis and statistics

We used the R platform for statistical analysis and for the data processing. We applied the DADA2 workflow (28) to infer the Amplicon Sequence Variants (ASVs) and for the taxonomic assignments.

In brief, we first filtered and trimmed raw sequencing reads in order to remove low quality bases and adapter contamination. Then, we removed identical reads. Moreover, the reads were denoised, merged filtered to remove artifacts (PCR, and PhiX related chimeras). We obtained the ASVs quantifications and assigned taxonomy annotations (including the Species level) using the SILVA database of non-redundant sequences (version: v138, nr99) (29). The data were structured in objects including the ASVs quantifications, the taxonomy annotations, the sample group data and the phylogenetic tree using the phyloseq and the APE packages (30, 31). Finally, based on the initial DNA concentration, we removed possible contaminant ASVs by using the “prevalence” method of the decontam package (32).

Downstream analyses were performed using the MicrobiomeAnalystR package (33, 34) and included data normalization, measures of diversity and differential abundance estimation.

Briefly, we normalized the ASV counts based on their abundance (low count filter: for any ASV to be retained, at least 20% of its values should contain at least 4 counts) and variance (low variance filter: based on Inter-quantile range $\pm 10\%$). This, because ASVs with small counts (in few samples) could represent sequencing errors. Moreover, ASVs that are closely constant in all samples could be excluded from the comparative analyses. Finally, we used the total sum scaling in order to bring all the samples to the same scale.

We evaluated the alpha-diversity by calculating the Abundance-based coverage estimator (ACE) and a nonparametric estimator of species richness (Chao1) indices and by the Fisher metrics (to consider both richness and evenness). The degree to which the species composition changes between the two groups (the beta-diversity) was obtained by PCoA (Principal Coordinates Analysis) of the distances calculated with the un-weighted UniFrac method and the statistical significance assessed by the PERMANOVA test.

We also used the rarefaction curves to evaluate whether the samples were sufficiently sampled and sequenced to represent their species richness. We assessed the statistical significance of comparisons between the two groups of samples under study by using the Mann-Whitney test.

The differential abundance was assessed by the LefSe (Linear Discriminant Analysis Effect Size) algorithm (35) for biomarker discovery and interpretation of metagenomics data. It involves the Kruskal-Wallis rank sum test to identify features (e.g., Species or Genera) with significant differential abundance in the two groups, followed by linear discriminant analysis (LDA) to evaluate the relevance (the effect size) of the selected features. Different abundant features were considered if the FDR adjusted *value of p* was less than or equal to 0.05 and if the Log LDA was greater than or less than 0.5. We used Phylogenetic Investigation of Communities by Reconstruction of Unobserved States (PICRUST) to assess the

metabolic potential of microbial communities (KEGG pathways). In this analysis, we started from the ASVs belonging to the significant genera obtained by LefSe algorithm.

3. Results

3.1. Cohort characteristics

Anthropometric characteristics and eating habits of the children enrolled in this study are shown in Table 1. No significant differences in individuals' gender (Chi-square 0.6748; $p=0.411399$), anthropometric characteristics such as height, weight, BMI and the waist/height ratio were observed ($p>0.05$). Similarly, no significant differences in all diet components analyzed in A and S groups were found (Table 1).

3.2. Sequencing reads processing and taxonomic assignments

The Illumina sequencing of the hypervariable V3–V4 regions of the 16S rRNA bacterial genes generated 2×250 bp paired-end reads. On average, we obtained 169.124 reads per-sample (Supplementary Figure S1A). Overall, the percentage of bases with quality scores above 20 and 30 (Q20 and Q30, respectively) was of 96.45 and 91.22, respectively (Supplementary Figures S1B,C). The percentage of GC nucleotides was of 52.03 (Supplementary Figure S1D). The set of reads was used to run the DADA2 workflow including the filtering and trimming (median = 169.02), denoising of forward (median = 165.00) and reverse (median = 164.80) reads; merging (median = 149.27) and chimeric reads removal (median = 114.43; Supplementary Figure S1E). After merging, the median length of reads was of 424 bp. Overall, starting from filtered reads, we obtained a merging rate of 88.38% and a final rate of read processing (non-chimera over merged reads) of 67.62% (Supplementary Figure S1F). Details on read processing are reported in Supplementary Table S1.

For taxonomic assignments we used the SILVA database of non-redundant sequences (version: v138, nr99). Overall, we could identify a total of 14.197 taxa (ASVs) that were annotated to the seven taxonomic ranks as follows. All the ASVs were taxonomically assigned to the kingdom of bacteria. The 98.06% of the ASVs was annotated at the phylum level (43 phyla), the 96.97% at the class level (94 classes), the 94.93% at the order level (195 orders), the 88.43% at the family level (231 families), the 81.04% at the genus level (404 genera) and the 4.86% was annotated up to the species level (257 species; Supplementary Figure S2A). From the initial set of annotated ASVs, we discarded a total of 68 taxa as possible contaminants. Moreover, as described in Methods we removed low abundant and low variable ASVs to obtain the final set of 472 ASVs that was normalized and used for downstream analyses.

3.3. Diversity estimates

At the genus level, alpha-diversity estimates were significantly different between the two groups, A and S. Indeed, ACE (Figure 1A),

TABLE 1 Anthropometric characteristics and eating habits of Active (A) and Sedentary (S) children.

	Total	Active (A)	Sedentary (S)
Gender M/F	69/45	33/18	36/27
Age (years)	8–10	8–10	8–10
<i>Anthropometric data</i>			
Height (cm)	142.8 ± 7.5	143.5 ± 8.0	142.2 ± 7.0
Weight (kg)	38.7 ± 9.8	38.2 ± 9.3	39.2 ± 10.2
BMI (kg/m ²)	18.9 ± 3.8	18.5 ± 3.7	19.2 ± 3.9
Waist/Height (cm)	0.5 ± 0.06	0.45 ± 0.05	0.5 ± 0.06

Eating habits (Average daily intake)	Active	Sedentary
Calories (kcal)	1464.7 ± 284.2	1491.1 ± 281.1
Carbohydrates (%)	48.1 ± 5.8	48.9 ± 6.2
Carbohydrates (g)	188.2 ± 43.8	194.0 ± 43.6
Starch (g)	75.8 ± 27.3	70.8 ± 27.0
Oligosaccharides (g)	50.7 ± 19.6	55.9 ± 18.6
Oligosaccharides/Carbohydrates (%)	26.7 ± 7.1	29.1 ± 8.7
Lipids (%)	36.1 ± 5.8	36.1 ± 5.5
Lipids (g)	58.5 ± 13.5	59.7 ± 13.7
Saturated fatty acids (g)	14.2 ± 4.6	15.3 ± 4.7
Monosaturated fatty acids (g)	18.7 ± 5.6	18.7 ± 6.0
Polysaturated fatty acids (g)	4.8 ± 1.4	5.0 ± 1.9
Saturated/ fatty acids (%)	37.4 ± 6.5	39.5 ± 6.3
Cholesterol (mg)	135.6 ± 64.7	144.9 ± 78.9
Proteins (%)	15.5 ± 2.0	15.0 ± 2.2
Proteins (g)	56.9 ± 13.8	55.9 ± 13.7
Animal proteins/Proteins (%)	70.6 ± 9.3	72.6 ± 14.1
Vegetal proteins/Proteins (%)	29.4 ± 9.3	27.4 ± 14.2
Total fiber/1,000 (kcal)	7.8 ± 3.2	6.8 ± 2.2
Total fiber (g)	11.7 ± 6.2	10.0 ± 3.3

Chao1 (Figure 1B) and Fisher (Figure 1C) indices showed $p < 0.01$ (Mann–Whitney test). The beta-diversity analysis, as measured by unweighted UniFrac distances, showed a significantly different microbial composition between the two groups ($r^2 = 0.026$, $p = 0.001$; Figure 1D), the diversity was also confirmed by using the weighted UniFrac distance metric ($r^2 = 0.018$, $p = 0.001$; Figure 1E). As previously reported (36), unweighted and weighted UniFrac distance measures can be considered as quality-based and quantity-based indexes, respectively. Indeed, we can assess that the observed variation between the two groups was due to the different taxa abundances and to the types of taxa in their microbiome. Moreover, the rarefaction analysis clearly evidenced the capacity to capture the species richness from the results of sampling and sequencing in both

groups without any statistically significant difference ($p = 0.084$; Supplementary Figure S2B).

3.4. Abundance estimates

We evaluated and compared the taxa abundance in the final set of 472 filtered and normalized ASVs. Overall, we identified 8 phyla, 12 classes, 30 orders, 46 families, 84 genera and 96 species.

At the phylum level, on average, the most abundant bacteria were Firmicutes, Bacteroides and Proteobacteria accounting for the 32.08, 26.97% and the 25.58% of the taxa, respectively (Supplementary Figure S3A). The most represented classes were Bacteroides (26.97%), Gammaproteobacteria (25.58%) and Bacilli (21.36%; Supplementary Figure S3B). The most prevalent orders were Bacteroides (26.63%), Lactobacillales (19.58%), Pasteurellales (12.88%) and Burkholderiales (12.12%; Supplementary Figure S3C). Among the most abundant families, we found Prevotellaceae (23.07%), Streptococcaceae (17.66%), Pasteurellaceae (12.88%) and Neisseriaceae (11.77%; Supplementary Figure S3D). At the genus level, we found the *Prevotella* (19.74%), *Streptococcus* (17.66%), *Haemophilus* (11.87%), *Neisseria* (11.57%) and *Veillonella* (6.06%; Figure 2A). Finally, the top abundant species that we were able to classify were *Prevotella melaninogenica* (10.86%), *Fusobacterium periodonticum* (4.44%), *Haemophilus parainfluenzae* (2.27%), *Rothia mucilaginosa* (2.10%) and *Veillonella dispar* (1.55%; Figure 2B).

3.5. Differential abundance estimates

As described in Methods, we used the LefSe algorithm to perform the differential abundance analysis and to identify the taxa that could explain the differences between the two groups A and S. We set the threshold at 5% and the Log LDA cutoff at ± 0.5 . Interestingly, using these very stringent criteria, we found *Coriobacteriaceae* family as more abundant in the A compared to the S group (LDA score = -3.17 ; FDR = 0.021).

Further, at the genus level, we found that ten genera were responsible for the differences between the two groups. In particular, *Agathobacter* (LDA score = -3.40 ; FDR = 0.015), *Escherichia-Shigella* (LDA score = -3.37 ; FDR = 7.68×10^{-4}), *Collinsella* (LDA score = -3.17 ; FDR = 0.012), *Simonsiella* (LDA score = -2.95 ; FDR = 0.044), *Eubacterium-yurii* group (LDA score = -2.79 ; FDR = 0.041) and *Parabacteroides* (LDA score = -2.43 ; FDR = 0.015) were more abundant in the A group. On the contrary, *Mogibacterium* (LDA score = 2.71; FDR = 9.21×10^{-4}), *Stomatobaculum* (LDA score = 3.24; FDR = 0.44), TM7x (also known as *Nanosynbacter lyticus*, LDA score = 3.90; FDR = 0.045) and *Granulicatella* (LDA score = 4.14; FDR = 0.045) were more abundant genera in the S group (Figures 3A,B; Supplementary Table S2). Eight species showed significant differences in the LefSe analysis. Indeed, *Prevotella nigrescens* (LDA score = -3.76 ; FDR = 1.5×10^{-3}), *Collinsella aerofaciens* (LDA score = -3.17 ; FDR = 7.45×10^{-3}), *Simonsiella muelleri* (LDA score = -2.96 ; FDR = 2.76×10^{-5}), *Parabacteroides merdae* (LDA score = -2.43 ; FDR = 1.3×10^{-2}), were the most represented species in the A group. Conversely, *Gemella parahaemolysans*, *Prevotella aurantiaca* (LDA score = -3.9 ; FDR = 5.27×10^{-4}), *Prevotella pallens* (LDA score = 4.23; FDR = 1.93×10^{-2}), *Neisseria mucosa* (LDA score = 4.43;




FIGURE 1

Diversity and distance measures between active (A) and sedentary (S). (A) Alpha-diversity measured by ACE index ($p = 0.0099$). (B) Alpha-diversity measured by Chao1 index ($p = 0.0097$). (C) Alpha-diversity measured by Fisher index ($p = 0.0039$). (D) Principal Coordinates Analysis plot of beta-diversity index measured by unweighted UniFrac distances ($p = 0.001$). (E) Principal Coordinates Analysis plot of beta-diversity index measured by weighted UniFrac distances ($p = 0.001$). (A), B, C, F: Mann–Whitney test. (D), E: PERMANOVA test.

$FDR = 1.31 \times 10^{-02}$) were more abundant species in the S group (LDA score = 2.94; $FDR = 7.45 \times 10^{-03}$; Supplementary Table S3; Figures 3C,D).

3.6. Metabolic pathways reconstruction

PICRUSt analysis highlighted the predominance of super pathway of hexitol degradation, L-glutamate degradation VII (to propionate), 2-methylcitrate cycle II, tetrapyrrole biosynthesis I, L-histidine degradation II, superpathway of beta-D-gluconide and D-gluconate

degradation, biotin biosynthesis I, and L-arginine biosynthesis pathways activation in saliva from S compared to A children. Conversely, we found a prevalence of superpathway of fatty acid biosynthesis initiation (*E. coli*) and catechol degradation II (meta-cleavage pathway) in saliva from A respect to S children (Figure 4).

4. Discussion

The aim of the study was to evaluate the effects of Active or Sedentary lifestyle on saliva microbiota composition in Italian




FIGURE 2 Overall taxonomic distribution. The Figure reports the overall abundance of the identified taxa. (A) Genus level. (B) Species level. Each plot shows the top ten abundant taxa.

schoolchildren living in the outskirts of Turin. The participants were classified as Active (A) or Sedentary (S) according to the cut-off score of 2.75 for PAQ-C-It (27). We evidenced an enrichment of several genera, such as *Agathobacter*, *Collinsella*, *Simonsiella*, and *Parabacteroides* in the children’s saliva from A compared to S group.

Among these, four species were differentially represented: *Prevotella nigrescens*, *Collinsella aerofaciens*, *Simonsiella muelleri*, and *Parabacteroides merdae*. Increased abundance of *Agathobacter* and *Prevotella* at both genus and species levels, was reported in GM of cross-country and marathon athletes; although an inverse correlation





FIGURE 3 Differently abundant taxa. **(A)** Dot plot showing the differently abundant genera. **(B)** Box plots showing the Normalized abundance levels of genera reported in panel A. **(C)** Dot plot reporting the differently abundant species. **(D)** Box plots showing the normalized abundance levels of species reported in panel C. In panels A and C, the dot size is proportional to the score of the LDA algorithm. The dot graduation color is proportional to the significance level as determined by FDR adjustment of Kruskal-Wallis rank sum test *p* values. A: Active; S: Sedentary.



was found for *Prevotella* and sucrose intake and a positive correlation for *Agathobacter* and dietary fiber content (37), we did not find statistically significant differences in all dietary components, including fiber content in A compared to S children. *Parabacteroides* are involved in host health promotion by regulating different pathways including inflammation, obesity and cancer prevention (38). Moreover, recent data suggest an anti-seizure and anti-cancer functions for *Parabacteroides merdae* (39, 40) and increased abundance have been also found in the GM of centenarians living in East China (41). As no differences in BMI were found between A or S children belonging to our cohort, we speculate that the prevalence of *Parabacteroides merdae* in saliva of group A children could be associated to the higher level of daily Physical Activity Amounts (PAA) when compared to group S. Furthermore, the increased abundance of *Parabacteroides* in group A resulted in the enrichment of the superpathway of fatty acid biosynthesis initiation that we found by metabolic pathway reconstruction. Of note, *Parabacteroides* are also involved in regulating different processes as carbohydrates metabolism and metabolites secretion, including Short Chain Fatty Acids (SCFAs) (42, 43). Among them, acetate, propionate, and butyrate are the main metabolites produced by several anaerobic bacteria from the fermentation of complex starch and dietary fibers. The available mechanistic data strongly suggest that SCFAs exert their powerful anti-inflammatory, antitumorigenic and even antimicrobial effects in the preventing gastro-intestinal dysfunction, obesity and type 2 diabetes mellitus (44, 45). In line with these evidences, several studies conducting in patients with type 1 and type 2 diabetes, liver cirrhosis, inflammatory bowel disorders (IBD) and atherosclerosis have shown a reduction in the abundance of SCFA-producing bacteria gut (46, 47). Gut microbiota of athletes have an enriched profile of SCFAs, previously associated to a healthier status and a lean phenotype (44, 48). In skeletal muscle, SCFAs can be oxidized, incorporated into glucose *via* gluconeogenesis or increase

the bioavailability of glucose, glycogen and fatty acids during exercise (49). Similarly, increased abundance in GM of taxa as *Firmicutes* and *Feacalibacterium prausnitzii* together with *Akkermansia*, producing butyrate, have been associated to exercise in athletes and non-athletes' controls with improvement in lipid oxidation, healthier profile and reduced risk for obesity and metabolic diseases, independently from body composition and diet (50–53). Further, similarly to our results, the association of a healthier profile with a reduction in *Bacteroides* species together with an increase in *R. hominis*, *A. Muciniphyla* and *F. prausnitzii* species have been described in GM from Active compared to Sedentary adults (54–56).

In group S we found an increased abundance of *Gemella parahemolyan*, *Prevotella aurantiaca*, *Prevotella pallens* and *Neisseria mucosa* species and of the *TM7x* genus as compared to group A. Notably, previous studies reported the abundance of *Neisseria mucosa* as sixfold higher in obese adolescents compared to normal-weight controls (57). Suggesting that although the Sedentary children are normal-weight, they present a predictive marker linked to obesity. Moreover, *Prevotella* species, habitually present in the oral microbiome, have constant and direct access to the gastrointestinal tract *via* saliva swallowing. Here, they could act as commensals but also as potentially harmful agents (58). Furthermore, the group S showed an increased abundance of the genus *TM7x* (also known as *Nanosynbacter lyticus*) which is an obligate epibiont parasite of the bacteria *Actinomyces odontolyticus* (not significantly enriched in our data) (59, 60). *TM7x* have been previously associated to different human inflammatory mucosal diseases such as the periodontitis (61). Moreover, *TM7x* have been considered as biomarker of active disease in patients with ulcerative colitis (62).

In group S, the metabolic pathway reconstruction highlighted the enrichment of L-glutamate degradation and L-arginine biosynthesis pathways. Interestingly, the dysregulation of L-glutamate and L-glutamine pathways have been associated with poor survival in

colon cancer patients (63, 64). L-glutamate signaling triggers oxidative and nitrosative stress pathways which are essential for the production of ROS that can induce the activation of oncogenes ensuring the survival of colon cancer cells (64).

Conversely, the analysis performed with PICRuST on A children's saliva revealed an abundance of fatty acid biosynthesis and catechol degradation pathways, in line with previous reports (65). The catecholamines are catabolic intermediates of various aromatic compounds, which contribute to Acetyl-CoA production. Acetyl-CoA, is also crucial for the cross-talk between multiple biological processes including, energy storage, membrane biosynthesis, and generation of signaling molecules that are produced in response to physiological cell processes (66, 67). Consequently, dysregulation of fatty acid synthesis can induce or promote disease development (68, 69).

In conclusion, Our results showed that saliva from active children had an enrichment of species and genera mainly associated with a healthier profile. On the contrary, the genera and the species enriched in the saliva from sedentary group could be linked non-communicable diseases. Nevertheless, our indirect observations need to be clarified by further (and possibly larger) studies aimed at understanding how an active lifestyle can modulate the composition of both oral and gut microbiota. Moreover, the minimum volume of physical exercise required to determine changes in oral microbiota composition remains to be assessed.

Data availability statement

The datasets presented in this study can be found in online repositories. The names of the repository/repositories and accession number(s) can be found in the article/[Supplementary material](#).

Ethics statement

The studies involving human participants were reviewed and approved by Ethics committee on human research of the University of Turin (9 March 2020: Protocol #134691) and Naples (17 January 2020: Protocol #376/19) approved the study. Written informed consent to participate in this study was provided by the participants' legal guardian/next of kin.

References

1. Grice EA, Segre JA. The human microbiome: our second genome. *Annu Rev Genom Hum Genet.* (2012) 13:151–70. doi: 10.1146/annurev-genom-090711-163814
2. Hou K, Wu ZX, Chen XY, Wang JQ, Zhang D, Xiao C, et al. Microbiota in health and diseases. *Signal Transduct Target Ther.* (2022) 7:135. doi: 10.1038/s41392-022-00974-4
3. Yatsunenko T, Rey FE, Manary MJ, Trehan I, Dominguez-Bello MG, Contreras M, et al. Human gut microbiome viewed across age and geography. *Nature.* (2012) 486:222–7. doi: 10.1038/nature11053
4. Aya V, Flórez A, Perez L, Ramírez JD. Association between physical activity and changes in intestinal microbiota composition: a systematic review. *PLoS One.* (2021) 16:e0247039. doi: 10.1371/journal.pone.0247039
5. Mailing LJ, Allen JM, Buford TW, Fields CJ, Woods JA. Exercise and the gut microbiome: a review of the evidence, potential mechanisms, and implications for human health. *Exerc Sport Sci Rev.* (2019) 47:75–85. doi: 10.1249/JES.0000000000000183
6. Choi JJ, Eum SY, Rampersaud E, Daunert S, Abreu MT, Toborek M. Exercise attenuates PCB-induced changes in the mouse gut microbiome. *Environ Health Perspect.* (2013) 121:725–30. doi: 10.1289/ehp.1306534
7. Evans CC, LePard KJ, Kwak JW, Stancukas MC, Laskowski S, Dougherty J, et al. Exercise prevents weight gain and alters the gut microbiota in a mouse model of high fat diet-induced obesity. *PLoS One.* (2014) 9:e92193. doi: 10.1371/journal.pone.0092193
8. Lambert JE, Myslicki JP, Bomhof MR, Belke DD, Shearer J, Reimer RA. Exercise training modifies gut microbiota in normal and diabetic mice. *Appl Physiol Nutr Metab.* (2015) 40:749–52. doi: 10.1139/apnm-2014-0452
9. Petriz BA, Castro AP, Almeida JA, Gomes CP, Fernandes GR, Kruger RH, et al. Exercise induction of gut microbiota modifications in obese, non-obese and hypertensive rats. *BMC Genomics.* (2014) 15:511. doi: 10.1186/1471-2164-15-511
10. Queipo-Ortuño MI, Seoane LM, Murri M, Pardo M, Gomez-Zumaquero JM, Cardona F, et al. Gut microbiota composition in male rat models under different nutritional status and physical activity and its association with serum leptin and ghrelin levels. *PLoS One.* (2013) 8:e65465. doi: 10.1371/journal.pone.0065465
11. Estaki M, Pither J, Baumeister P, Little JP, Gill SK, Ghosh S, et al. Cardiorespiratory fitness as a predictor of intestinal microbial diversity and distinct metagenomic functions. *Microbiome.* (2016) 4:42. doi: 10.1186/s40168-016-0189-7

Author contributions

AM, SO, and PBu: conceptualization. DV, DP, CC, and AC: methodology. VL and MC: software, validation, and formal analysis. PBr and CL: investigation and resources. AM, DV, and VL: data curation and writing—original draft preparation. AM, SO, MC, and PBu: writing—review and editing. PBu: visualization and supervision. AM, SO, FS, and PBu: project administration and funding acquisition. All authors contributed to the article and approved the submitted version.

Funding

This study was funded by the grant PRIN 2017_Prot.2017RS5M44 to PBu, AM, SO, and FS.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The reviewer GC declared a past co-authorship with the author CC to the handling editor.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Supplementary material

The Supplementary material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fnut.2023.1226891/full#supplementary-material>

12. Clarke SF, Murphy EF, O'Sullivan O, Lucey AJ, Humphreys M, Hogan A, et al. Exercise and associated dietary extremes impact on gut microbial diversity. *Gut*. (2014) 63:1913–20. doi: 10.1136/gutjnl-2013-306541
13. Allen AP, Dinan TG, Clarke G, Cryan JF. A psychology of the human brain-gut-microbiome axis. *Soc Personal Psychol Compass*. (2017) 11:e12309. doi: 10.1111/spc3.12309
14. Petersen LM, Bautista EJ, Nguyen H, Hanson BM, Chen L, Lek SH, et al. Community characteristics of the gut microbiomes of competitive cyclists. *Microbiome*. (2017) 5:98. doi: 10.1186/s40168-017-0320-4
15. Hughes RL, Holscher HD. Fueling gut microbes: a review of the interaction between diet, exercise, and the gut microbiota in athletes. *Adv Nutr*. (2021) 12:2190–215. doi: 10.1093/advances/nmab077
16. Durk RP, Castillo E, Márquez-Magaña L, Grosicki GJ, Bolter ND, Lee CM, et al. Gut microbiota composition is related to cardiorespiratory fitness in healthy young adults. *Int J Sport Nutr Exerc Metab*. (2019) 29:249–53. doi: 10.1123/ijsem.2018-0024
17. Noce A, Tranchita E, Marrone G, Grazioli E, Di Lauro M, Murri A, et al. The possible role of physical activity in the modulation of gut microbiota in chronic kidney disease and its impact on cardiovascular risk: a narrative review. *Eur Rev Med Pharmacol Sci*. (2023) 27:3733–46. doi: 10.26355/eurrev_202304_32172
18. Barton W, Penney NC, Cronin O, Garcia-Perez I, Molloy MG, Holmes E, et al. The microbiome of professional athletes differs from that of more sedentary subjects in composition and particularly at the functional metabolic level. *Gut*. (2018) 67:gutjnl-2016-313627–33. doi: 10.1136/gutjnl-2016-313627
19. Quiroga R, Nistal E, Estébanez B, Porras D, Juárez-Fernández M, Martínez-Flórez S, et al. Exercise training modulates the gut microbiota profile and impairs inflammatory signaling pathways in obese children. *Exp Mol Med*. (2020) 52:1048–61. doi: 10.1038/s12276-020-0459-0
20. Bai J, Hu Y, Bruner D. Composition of gut microbiota and its association with body mass index and lifestyle factors in a cohort of 7–18 years old children from the American gut project. *Pediatric Obes*. (2019) 14:e12480. doi: 10.1111/ijpo.12480
21. Moran-Ramos S, Lopez-Contreras BE, Villarruel-Vazquez R, Ocampo-Medina E, Macias-Kauffer L, Martínez-Medina JN, et al. Environmental and intrinsic factors shaping gut microbiota composition and diversity and its relation to metabolic health in children and early adolescents: a population-based study. *Gut Microbes*. (2020) 11:900–17. doi: 10.1080/19490976.2020.1712985
22. Santarossa S, Sitarik AR, Johnson CC, Li J, Lynch SV, Ownby DR, et al. Associations of physical activity with gut microbiota in pre-adolescent children. *Phys Act Nutr*. (2021) 25:24–37. doi: 10.20463/pan.2021.0023
23. Belström D, Holmström P, Nielsen CH, Kirkby N, Twetman S, Heitmann BL, et al. Bacterial profiles of saliva in relation to diet, lifestyle factors, and socioeconomic status. *J Oral Microbiol*. (2014) 6:6. doi: 10.3402/jom.v6.23609
24. Keller MK, Kressler CA, Belström D, Twetman S, Tanner ACR. Oral microbial profiles of individuals with different levels of sugar intake. *J Oral Microbiol*. (2017) 9:1355207. doi: 10.1080/20002297.2017.1355207
25. Acharya A, Chan Y, Kheur S, Jin LJ, Watt RM, Mattheos N. Salivary microbiome in non-oral disease: a summary of evidence and commentary. *Arch Oral Biol*. (2017) 83:169–73. doi: 10.1016/j.archoralbio.2017.07.019
26. Schmidt TS, Hayward MR, Coelho LP, Li SS, Costea PI, Voigt AY, et al. Extensive transmission of microbes along the gastrointestinal tract. *elife*. (2019) 8:e42693. doi: 10.7554/eLife.42693
27. Lupo C, De Pasquale P, Boccia G, Ungureanu AN, Moisé P, Mulasso A, et al. The Most active child is not always the fittest: physical activity and fitness are weakly correlated. *Sports (Basel)*. (2022) 11:3. doi: 10.3390/sports11010003
28. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. DADA2: high-resolution sample inference from Illumina amplicon data. *Nat Methods*. (2016) 13:581–3. doi: 10.1038/nmeth.3869
29. Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, et al. SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res*. (2007) 35:7188–96. doi: 10.1093/nar/gkm864
30. McMurdie PJ, Holmes S. Phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One*. (2013) 8:e61217. doi: 10.1371/journal.pone.0061217
31. Paradis E, Schliep K. Ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics*. (2019) 35:526–8. doi: 10.1093/bioinformatics/bty633
32. Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. Simple statistical identification and removal of contaminant sequences in marker-gene and metagenomics data. *Microbiome*. (2018) 6:226. doi: 10.1186/s40168-018-0605-2
33. Dhariwal A, Chong J, Habib S, King IL, Agellon LB, Xia J. Microbiome analyst: a web-based tool for comprehensive statistical, visual and meta-analysis of microbiome data. *Nucleic Acids Res*. (2017) 45:W180–8. doi: 10.1093/nar/gkx295
34. Chong J, Liu P, Zhou G, Xia J. Using microbiome analyst for comprehensive statistical, functional, and meta-analysis of microbiome data. *Nat Protoc*. (2020) 15:799–821. doi: 10.1038/s41596-019-0264-1
35. Segata N, Izard J, Waldron L, Gevers D, Miropolsky L, Garrett WS, et al. Metagenomic biomarker discovery and explanation. *Genome Biol*. (2011) 12:R60. doi: 10.1186/gb-2011-12-6-r60
36. Cecarini V, Gogoi O, Bonfili L, Veneruso I, Pacinelli G, De Carlo S, et al. Modulation of gut microbiota and neuroprotective effect of a yeast-enriched beer. *Nutrients*. (2022) 14:2380. doi: 10.3390/nu14122380
37. Kulecka M, Fraczek B, Mikula M, Zeber-Lubecka N, Karczmarski J, Paziewska A, et al. The composition and richness of the gut microbiota differentiate the top polish endurance athletes from sedentary controls. *Gut Microbes*. (2020) 11:1374–84. doi: 10.1080/19490976.2020.1758009
38. Kverka M, Zakostelska Z, Klimesova K, Sokol D, Hudcovic T, Hrnčir T, et al. Oral administration of *Parabacteroides distasonis* antigens attenuates experimental murine colitis through modulation of immunity and microbiota composition. *Clin Exp Immunol*. (2011) 163:250–9. doi: 10.1111/j.1365-2249.2010.04286.x
39. Olson CA, Vuong HE, Yano JM, Liang QY, Nussbaum DJ, Hsiao EY. The gut microbiota mediates the anti-seizure effects of the ketogenic diet. *Cells*. (2018) 7:31728–1741.e13. doi: 10.1016/j.cell.2018.04.027
40. Roncase EJ, Moon C, Chatterjee S, González-Pérez GE, Craik CS, O'Donoghue AJ, et al. Substrate profiling and high resolution co-complex crystal structure of a secreted C11 protease conserved across commensal Bacteria. *ACS Chem Biol*. (2017) 12:1556–65. doi: 10.1021/acschembio.7b00143
41. Wang N, Li R, Lin H, Fu C, Wang X. Enriched taxa were found among the gut microbiota of centenarians in East China. *PLoS One*. (2019) 14:e0222763. doi: 10.1371/journal.pone.0222763
42. Lei Y, Tang L, Liu S, Hu S, Wu L, Liu Y, et al. *Parabacteroides* produces acetate to alleviate heparanase-exacerbated acute pancreatitis through reducing neutrophil infiltration. *Microbiome*. (2021) 9:115. doi: 10.1186/s40168-021-01065-2
43. Du X, Xiang Y, Lou F, Tu P, Zhang X, Hu X, et al. Microbial community and short-chain fatty acid mapping in the intestinal tract of quail. *Animals (Basel)*. (2020) 10:1006. doi: 10.3390/ani10061006
44. Tan J, McKenzie C, Potamitis M, Thorburn AN, Mackay CR, Macia L. The role of short-chain fatty acids in health and disease. *Adv Immunol*. (2014) 121:91–119. doi: 10.1016/B978-0-12-800100-4.00003-9
45. Blaak EE, Canfora EE, Theis S, Frost G, Groen AK, Mithieux G, et al. Short chain fatty acids in human gut and metabolic health. *Benef Microbes*. (2020) 11:411–55. doi: 10.3920/BM2020.0057
46. Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, et al. A human gut microbial gene catalogue established by metagenomic sequencing. *Nature*. (2010) 464:59–65. doi: 10.1038/nature08821
47. Vatanen T, Franzosa EA, Schwager R, Tripathi S, Arthur TD, Vehik K, et al. The human gut microbiome in early-onset type 1 diabetes from the TEDDY study. *Nature*. (2018) 562:589–94. doi: 10.1038/s41586-018-0620-2
48. Bongiovanni T, Yin MOL, Heaney LM. The Athlete and Gut Microbiome: Short-chain Fatty Acids as Potential Ergogenic Aids for Exercise and Training. *Int J Sports Med*. (2021) 42:1143–1158. doi: 10.1055/a-1524-2095
49. Frampton J, Murphy KG, Frost G, Chambers ES. Short-chain fatty acids as potential regulators of skeletal muscle metabolism and function. *Nat Metab*. (2020) 2:840–848. doi: 10.1038/s42255-020-0188-7
50. Louis P, Flint HJ. Diversity, metabolism and microbial ecology of butyrate-producing bacteria from the human large intestine. *FEMS Microbiol Lett*. (2009) 294:1–8. doi: 10.1111/j.1574-6968.2009.01514.x
51. De Filippis F, Pasoli E, Ercolini D. Newly explored *Faecalibacterium* diversity is connected to age, lifestyle, geography, and disease. *Curr Biol*. (2020) 30:4932–4943.e4. doi: 10.1016/j.cub.2020.09.063
52. Everard A, Belzer C, Geurts L, Ouwerkerk JP, Druart C, Bindels LB, et al. Cross-talk between *Akkermansia muciniphila* and intestinal epithelium controls diet-induced obesity. *Proc Natl Acad Sci U S A*. (2013) 110:9066–71. doi: 10.1073/pnas.1219451110
53. Scheepers LE, Penders J, Mbakwa CA, Thijs C, Mommers M, Arts IC. The intestinal microbiota composition and weight development in children: the KOALA birth cohort study. *Int J Obes*. (2015) 39:16–25. doi: 10.1038/ijo.2014.178
54. Cataldi S, Bonavolontà V, Poli L, Clemente FM, De Candia M, Carvutto R, et al. The relationship between physical activity, physical exercise, and human gut microbiota in healthy and unhealthy subjects: a systematic review. *Biology (Basel)*. (2022) 11:479. doi: 10.3390/biology11030479
55. Canani RB, Costanzo MD, Leone L, Pedata M, Meli R, Calignano A. Potential beneficial effects of butyrate in intestinal and extraintestinal diseases. *World J Gastroenterol*. (2011) 17:1519–28. doi: 10.3748/wjg.v17.i12.1519
56. Machiels K, Joossens M, Sabino J, De Preter V, Arijis I, Eeckhaut V, et al. A decrease of the butyrate-producing species *Roseburia hominis* and *Faecalibacterium prausnitzii* defines dysbiosis in patients with ulcerative colitis. *Gut*. (2014) 63:1275–83. doi: 10.1136/gutjnl-2013-304833
57. Zeigler CC, Persson GR, Wondimu B, Marcus C, Sobko T, Modéer T. Microbiota in the oral subgingival biofilm is associated with obesity in adolescence. *Obesity (Silver Spring)*. (2012) 20:157–64. doi: 10.1038/oby.2011.305

58. Könönen E, Gursoy UK. Oral Prevotella species and their connection to events of clinical relevance in gastrointestinal and respiratory tracts. *Front Microbiol.* (2022) 12:798763. doi: 10.3389/fmicb.2021.798763
59. Bedree JK, Bor B, Cen L, Edlund A, Lux R, McLean JS, et al. Quorum sensing modulates the Epibiotic-parasitic relationship between *Actinomyces odontolyticus* and its *Saccharibacteria epibiont*, a *Nanosynbacter lyticus* strain, TM7x. *Front Microbiol.* (2018) 9:2049. doi: 10.3389/fmicb.2018.02049
60. He X, McLean JS, Edlund A, Yooseph S, Hall AP, Liu SY, et al. Cultivation of a human-associated TM7 phylotype reveals a reduced genome and epibiotic parasitic lifestyle. *Proc Natl Acad Sci U S A.* (2015) 112:244–9. doi: 10.1073/pnas.1419038112
61. Brinig MM, Lepp PW, Ouverney CC, Armitage GC, Relman DA. Prevalence of bacteria of division TM7 in human subgingival plaque and their association with disease. *Appl Environ Microbiol.* (2003) 69:1687–94. doi: 10.1128/AEM.69.3.1687-1694.2003
62. Zhu S, Han M, Liu S, Fan L, Shi H, Li P. Composition and diverse differences of intestinal microbiota in ulcerative colitis patients. *Front Cell Infect Microbiol.* (2022) 12:953962. doi: 10.3389/fcimb.2022.953962
63. Baj A, Moro E, Bistoletti M, Orlandi V, Crema F, Giaroni C. Glutamatergic signaling along the microbiota-gut-brain Axis. *Int J Mol Sci.* (2019) 20:1482. doi: 10.3390/ijms20061482
64. Hayes JD, Dinkova-Kostova AT, Tew KD. Oxidative stress in Cancer. *Cancer Cell.* (2020) 38:167–97. doi: 10.1016/j.ccell.2020.06.001
65. Cronan JE. Bacterial membrane lipids: where do we stand? *Annu Rev Microbiol.* (2003) 57:203–24. doi: 10.1146/annurev.micro.57.030502.090851
66. Harayama S, Rekik M, Ngai KL, Ornston LN. Physically associated enzymes produce and metabolize 2-hydroxy-2, 4-dienoate, a chemically unstable intermediate formed in catechol metabolism via meta cleavage in *Pseudomonas putida*. *J Bacteriol.* (1989) 171:6251–8. doi: 10.1128/jb.171.11.6251-6258.1989
67. Wakil SJ, Abu-Elheiga LA. Fatty acid metabolism: target for metabolic syndrome. *J Lipid Res.* (2009) 50:S138–43. doi: 10.1194/jlr.R800079-JLR200
68. Röhrig F, Schulze A. The multifaceted roles of fatty acid synthesis in cancer. *Nat Rev Cancer.* (2016) 16:732–49. doi: 10.1038/nrc.2016.89
69. Wang Y, Yu W, Li S, Guo D, He J, Wang Y. Acetyl-CoA carboxylases and diseases. *Front Oncol.* (2022) 12:836058. doi: 10.3389/fonc.2022.836058