

Hyperbolic polygonal billiards with finitely many ergodic SRB measures

GIANLUIGI DEL MAGNO[†], JOÃO LOPES DIAS[‡], PEDRO DUARTE[§] and JOSÉ PEDRO GAIVÃO[¶]

[†] *Universidade Federal da Bahia, Instituto de Matemática, Avenida Adhemar de Barros, Ondina, 40170–110 Salvador, BA, Brasil*
(e-mail: gdelmagno@ufba.br)

[‡] *Departamento de Matemática and CEMAPRE, ISEG, Universidade de Lisboa, Rua do Quelhas 6, 1200–781 Lisboa, Portugal*
(e-mail: jldias@iseg.ulisboa.pt)

[§] *Departamento de Matemática and CMAF, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, Edifício C6, Piso 2, 1749–016 Lisboa, Portugal*
(e-mail: pmduarte@fc.ul.pt)

[¶] *Departamento de Matemática and CEMAPRE, ISEG, Universidade de Lisboa, Rua do Quelhas 6, 1200–781 Lisboa, Portugal*
(e-mail: jpgaivao@iseg.ulisboa.pt)

(Received 21 February 2016 and accepted in revised form 19 October 2016)

Abstract. We study polygonal billiards with reflection laws contracting the angle of reflection towards the normal. It is shown that if a polygon does not have parallel sides facing each other, then the corresponding billiard map has finitely many ergodic Sinai–Ruelle–Bowen measures whose basins cover a set of full Lebesgue measure.

1. Introduction

The study of the asymptotic behaviour of billiards is an important subject in the theory of dynamical systems. Billiards exhibit a rich variety of statistical properties depending on the table geometry and the reflection law considered.

In this work, we are interested in polygonal tables. The billiard map of a polygonal billiard with the standard reflection law (the angle of reflection equals the angle of incidence) is conservative and non-chaotic: it preserves a measure that is absolutely continuous with respect to the Lebesgue measure and all its Lyapunov exponents are equal to zero.

A completely different dynamic arises when the reflection law is contracting [14], that is, when the angle of reflection measured from the normal is a contraction of the angle of

incidence. In this case, the billiard is a dissipative system: its map no longer preserves an absolutely continuous measure and may have attractors [1, 2, 8]. Indeed, if there are no period two orbits, then the billiard map has a uniformly hyperbolic attractor [9]. Notice that period two orbits correspond to collisions perpendicular to a pair of parallel sides of the billiard table. These orbits are parabolic and their union forms an attractor.

Contracting reflection laws appear in the study of the propagation of seismic waves [18] and are of interest for the design of bouncing robots [12].

For billiards in generic convex polygons with a strong contracting reflection law, we have recently proved the existence of countably many ergodic Sinai–Ruelle–Bowen measures (SRB), each one supported on a uniformly hyperbolic attractor [9]. This result is significantly extended in the current paper by enlarging the class of allowed polygons to include non-convex polygons and, more importantly, by removing any restriction on the contraction factor of the reflection law (cf. [8–11]). In addition, we establish that the basins of the ergodic SRB measures cover a set of full Lebesgue measure. The full result is the following theorem.

THEOREM 1.1. *For every polygon without parallel sides facing each other and every contracting reflection law, the corresponding billiard map has a hyperbolic attractor supporting finitely many ergodic SRB measures whose basins cover a set of full Lebesgue measure. Every ergodic SRB measure admits a decomposition into finitely many Bernoulli components, each component having exponential decay of correlations for Hölder observables. Every SRB measure of the billiard map is a convex combination of the ergodic SRB measures. Finally, the set of periodic points is dense in the attractor.*

Besides its specific interest, the previous result may prove useful in studying polygonal billiards with the standard reflection law because they lie at the boundary of the class of billiards considered in this paper.

A long standing conjecture of Palis [15] states there exists a dense set of dynamical systems such that each of them have finitely many attractors with ergodic SRB measures whose basins of attraction cover a set with full Lebesgue measure. Since polygons without parallel sides facing each other are dense in the space of all polygons, Theorem 1.1 verifies this conjecture for polygonal billiards with contracting reflection laws.

Polygonal billiards are piecewise smooth systems: they have discontinuities corresponding to trajectories reaching a corner of the table. Discontinuities may obstruct hyperbolicity in the sense that they may prevent the system from having local invariant manifolds or, at least, local invariant manifolds of uniform size. The local fractioning of the unstable manifolds produced by the discontinuities is measured by the branching number of the singular sets (see §3).

In order to prove Theorem 1.1, we first control the growth of the branching number, which is key to guaranteeing that the expansion along the unstable direction prevails over the fractioning of local unstable manifolds caused by the discontinuities. This allows us to extend the results of [9]. Then we apply a general theory of Pesin [16] and Sataev [17] for hyperbolic piecewise smooth maps to prove the existence of finitely many ergodic SRB measures. To get the exponential decay of correlations, we use results of

Chernov *et al* [4–7]. Finally, we adapt the ideas contained in [3, Proposition 4.2] to prove that almost every point is in the basin of some SRB measure.

The rest of the paper is organized as follows. In §2, we introduce the billiard maps and other basic definitions. The growth of the branching number is studied in §3. In §4, we establish the existence of finitely many ergodic SRB measures. Finally, in §5, we show that the basins of these measures cover the entire phase space up to a set of zero Lebesgue measure.

2. Billiard map

Let P be a polygonal domain (open and connected set) of \mathbb{R}^2 with d -sides and perimeter equal to one. The billiard in P with the specular reflection law is the flow on the unit tangent bundle of P generated by the motion of a free point-particle in P with specular reflection at ∂P : that is, the angle of reflection equals the angle of incidence. The corresponding billiard map Φ_P is the first return map on M , the set of unit vectors attached to ∂P and pointing inside P .

Each element of $x \in M$ can be identified with a pair (s, θ) , where s is the arc-length parameter of ∂P of the base point of x and θ is the angle formed by x with the positively oriented tangent to ∂P at s . Accordingly, we can write

$$M = S^1 \times [-\pi/2, \pi/2].$$

The domain of Φ_P does not coincide with the entire M . To specify it, we first introduce the sets V and S . Let $0 = s_1 < \dots < s_d < 1$ be the values of the arc-length parameter corresponding to the vertices of P . The set V is given by

$$V = \{s_1, \dots, s_d\} \times (-\pi/2, \pi/2),$$

whereas the set S is the subset of M consisting of elements whose forward trajectory hits a vertex of P at the first collision with ∂P . Define

$$N = V \cup S \quad \text{and} \quad N^+ = N \cup \partial M.$$

Both sets N and N^+ consist of finitely many smooth curves [9, Proposition 2.1].

The map Φ_P is defined on $M \setminus N^+$ and is a piecewise smooth map with singular set N^+ in the sense of Definition 4.1. Observe that $\Phi_P(x)$ is the unit vector corresponding to the first collision of the trajectory of $x \in M \setminus N^+$ with ∂P . For a detailed definition of Φ_P , we refer the reader to [9] for a polygonal table and to [5] for more general tables.

A *reflection law* is a function $f : (-\pi/2, \pi/2) \rightarrow (-\pi/2, \pi/2)$. For example, the specular reflection law corresponds to the identity function $f(\theta) = \theta$. Let $R_f : M \rightarrow M$ be the map $R_f(s, \theta) = (s, f(\theta))$. The billiard map for the polygon P with reflection law f is the map $\Phi_{f,P} : M \setminus N^+ \rightarrow M$ given by

$$\Phi_{f,P} = R_f \circ \Phi_P.$$

This map is just the first return map on M of the billiard flow in the polygon P with reflection law f (see Figure 1). We call $\Phi_{f,P}$ the *billiard map of P with reflection law f* .




FIGURE 1. Billiard flow with a contracting reflection law in a polygon without parallel sides facing each other.

If f is differentiable, then so is $\Phi_{f,P}$. In this case, the explicit expression for the derivative of $\Phi_{f,P}$ can be easily computed (for the derivative of Φ_P , see [5]) to be

$$D\Phi_{f,P}(s, \theta) = - \begin{pmatrix} \frac{\cos \theta}{\cos \bar{\theta}_1} & \frac{t(s, \theta)}{\cos \bar{\theta}_1} \\ 0 & f'(\bar{\theta}_1) \end{pmatrix}, \tag{2.1}$$

where $(\bar{s}_1, \bar{\theta}_1) = \Phi_P(s, \theta)$ and $(s_1, \theta_1) = \Phi_{f,P}(s, \theta)$, and $t(s, \theta)$ denotes the Euclidean distance in \mathbb{R}^2 between the points of ∂P with coordinates s and s_1 .

Given a differentiable reflection law f , we define

$$\lambda(f) = \sup_{\theta \in (-\pi/2, \pi/2)} |f'(\theta)|.$$

A differentiable reflection law is called *contracting* if $\lambda(f) < 1$. The simplest example of a contracting reflection law is $f(\theta) = \sigma\theta$ with $0 < \sigma < 1$ [1, 2, 8, 14].

Standing assumptions on f : we assume throughout the paper that f satisfies the following conditions.

- (1) f is a C^2 embedding from $[-\pi/2, \pi/2]$ to $[-\pi/2, \pi/2]$.
- (2) f is contracting.
- (3) $f(0) = 0$.
- (4) $f'(\theta) > 0$ for $\theta \in (-\pi/2, \pi/2)$.

Since $f' > 0$, all the entries of $D\Phi_{f,P}$ have the same sign. This simple fact will play an important role in the arguments presented in the next section.

3. Growth of the branching number

To simplify our notation, from now on we shall write Φ instead of $\Phi_{f,P}$. For $n \geq 1$, define

$$N_n^+ = N^+ \cup \Phi^{-1}(N^+) \cup \dots \cup \Phi^{-n+1}(N^+)$$

and

$$S_n^+ = S \cup \Phi^{-1}(S) \cup \dots \cup \Phi^{-n+1}(S).$$

The set N_n^+ contains all the points of M where the map Φ^n is not defined. Since $N = V \cup S$ and $\Phi^{-1}(\partial M) = \emptyset$, it follows that

$$N_n^+ = V \cup \partial M \cup S_n^+.$$




FIGURE 2. Sectors Δ_1, Δ_2 and Δ_3 sharing the same vertex $x \in N_n^+$.

As a direct consequence of the definition of S_n^+ ,

$$S_{n+k}^+ = S_n^+ \cup \Phi^{-n}(S_k^+) \quad \text{for } k > 0.$$

Hence $N_{n+k}^+ \setminus N_n^+ = S_{n+k}^+ \setminus S_n^+ \subset \Phi^{-n}(S_k^+)$ and

$$\Phi^n(N_{n+k}^+ \setminus N_n^+) \subset S_k^+. \tag{3.1}$$

Definition 3.1. A sector of order n with vertex $x \in N_n^+$ is a connected component Δ of $U \setminus N_n^+$ with $U \subset M$ being an open ball centred at a point $x \in N_n^+$ such that the closure of U intersects only smooth components of N_n^+ meeting at x (see Figure 2).

The smooth curves forming the boundary of Δ and meeting at x are called the *boundary curves* of Δ . A sector $\Delta' \subset \Delta$ of order greater than n with vertex x is called a *sub-sector* of Δ .

From the definition above, it follows that $\Phi^n : \Delta \rightarrow \Phi^n(\Delta)$ is a C^2 diffeomorphism for every sector Δ of order n . In particular, the first n th iterates of points of Δ visit the same sequence of sides of P . Moreover, if Γ is a boundary curve of Δ such that $\Gamma \not\subset \partial M$, then either $\Gamma \subset V$ or $\Phi^{k-1}(\Gamma) \subset S_1^+$ for some $1 \leq k \leq n$. In other words, all the trajectories starting at points of a boundary curve not contained in ∂M either emerge from the same vertex or hit the same vertex at the k th collision with ∂P .

Definition 3.2. A C^1 curve $t \mapsto \gamma(t) = (s(t), \theta(t)) \in M$ is called *increasing* if $s'(t)\theta'(t) \geq 0$ for every t , and it is called *strictly increasing* if $s'(t)\theta'(t) > 0$ for every t . A *decreasing* curve and a *strictly decreasing* curve are defined similarly by requiring that $s'(t)\theta'(t) \leq 0$ for every t and $s'(t)\theta'(t) < 0$ for every t , respectively. A curve $t \mapsto \gamma(t) = (s(t), \theta(t)) \in M$ is called *horizontal* if there exists a constant c such that $\theta(t) = c$ for every t .

Remark 3.3. The set $V \cup S_n^+$ consists of finitely many strictly decreasing C^2 curves [11, Proposition 2.3].

LEMMA 3.4. $\overline{S_n^+} \cap \partial M = \emptyset$ for $n \in \mathbb{N}$.

Proof. It can be easily checked that the case $n = 1$ holds true. Hence, in the rest of the proof, we can limit ourselves to consider the case when $n \geq 2$ and S_n^+ is replaced by

$R_n := S_n^+ \setminus S_1^+$. We argue by contradiction and suppose that $\overline{R_n} \cap \partial M \neq \emptyset$ for some $n \in \mathbb{N}$. Let $x \in \overline{R_n} \cap \partial M$. There are precisely two sectors of order n with vertex x that have a boundary curve lying on ∂M . Denote the one on the right-hand side of x by Δ . One boundary curve of Δ is contained in ∂M , whereas the other is contained in R_n .

Now, choose a point y in the interior of the boundary curve γ of Δ that is not contained in ∂M in such a way that the vertical segment γ_1 with one endpoint in y and the other one on ∂M is contained in the closure of Δ . Denote by γ_2 be the sub-curve of γ with endpoints x and y , and denote by z the endpoint of the segment γ_1 belonging to ∂M .

From the definition of S_n^+ , it follows that $\gamma_2' := \Phi(\gamma_2)$ is contained in a smooth component of S_k^+ for some $1 \leq k < n$. Since the smooth components of S_n^+ are strictly decreasing curves by Remark 3.3, so is γ_2' . The map Φ is C^2 differentiable on Δ , and so $\gamma_1' := \Phi(\gamma_1)$ is a C^2 curve. Since γ_1 is vertical and the entries of $D\Phi$ have the same sign (see (2.1)), it follows that γ_1' is strictly increasing.

Since $y \in \gamma_1 \cap \gamma_2$, it follows that $\Phi(y) \in \gamma_1' \cap \gamma_2'$. Moreover, since γ_1' and γ_2' are strictly monotone, both limits $\lim_{\gamma_1 \ni w \rightarrow z} \Phi(w)$ and $\lim_{\gamma_2 \ni w \rightarrow x} \Phi(w)$ exist. It is easy to see that these limits coincide because $x, z \in \partial M = S^1 \times \{-\pi/2, \pi/2\}$. In conclusion, the curves γ_1' and γ_2' intersect at both their endpoints. However, this is impossible because one curve is strictly increasing and the other is strictly decreasing. □

Definition 3.5. A sector Δ of order n is called *regular* if $\Phi^n|_\Delta$ admits an extension $\Psi_\Delta : \overline{\Delta} \rightarrow \Psi_\Delta(\overline{\Delta})$ that is a C^2 diffeomorphism, where $\overline{\Delta}$ is the closure of Δ .

LEMMA 3.6. Suppose that Δ is a regular sector of order n . If Γ is a boundary curve of Δ that is not contained in ∂M , then $\Psi_\Delta(\Gamma)$ is a C^2 increasing curve.

Proof. Define $F(s, \theta) = (s, f(-\theta))$ for $(s, \theta) \in M$ and let $N^- = F(V \cup S_1^+)$. Remark 3.3 combined with $f' > 0$ implies that N^- consists of strictly increasing C^2 curves.

Let Γ be as in the statement of the lemma. By the remark after Definition 3.1, either $\Gamma \subset V$ or $\Phi^i(\Gamma) \subset S_1^+$ for some $0 \leq i \leq n - 1$. Then it is not difficult to see that there exists $0 \leq k \leq n - 1$ such that $\Psi_\Delta(\Gamma) \subset \Phi^k(N^-)$. Since N^- consists of strictly increasing C^2 curves, to obtain the wanted conclusion it is enough to observe that if γ is a strictly increasing curve such that $\gamma \cap N_i^+ = \emptyset$, then $\Phi^i(\gamma)$ is strictly increasing curve as well. This is so, because all the entries of the matrix of $D\Phi^i$ have the same sign (see (2.1)). □

Definition 3.7. Let Δ be a sector with vertex x . A curve $\gamma : [a, b] \rightarrow M$ is called Δ -curve if $\gamma(a) = x$ and $\gamma(t) \in \Delta$ for $t \in (a, b]$.

In the next lemma, we give a necessary condition for a regular sector Δ to contain singular Δ -curves. By definition, if Δ is a sector of order n , then $\Delta \cap S_n^+ = \emptyset$. Hence, if there exists a Δ -curve γ contained in S_{n+k}^+ for some $k > 0$, then we must have $\gamma \subset S_{n+k}^+ \setminus S_n^+$.

LEMMA 3.8. Suppose that Δ is a regular sector of order n . If there exists a Δ -curve contained in S_{n+k}^+ for some $k > 0$, then there exists also a horizontal Δ -curve.

Proof. First of all, we observe that if x is the vertex of Δ , then $x \notin \partial M$ by Lemma 3.4. Now, let γ be the Δ -curve contained in S_{n+k}^+ . Then $\gamma \subset S_{n+k}^+ \setminus S_n^+$, and so (3.1) implies that $\Psi_\Delta(\gamma) \subset S_k^+$. Hence $\Phi_\Delta^n(\gamma)$ is a strictly decreasing curve passing through $\Psi_\Delta(x)$. Next, the region $\Psi_\Delta(\Delta)$ is bounded by the curves Γ_1 and Γ_2 , which are the images under Ψ_Δ of the boundary curves of Δ , passing through $\Psi_\Delta(x)$, and are increasing by Lemma 3.6.

Summarizing, $\Psi_\Delta(\Delta)$ is bounded by the increasing curves Γ_1 and Γ_2 and contains a strictly decreasing curve $\Psi_\Delta(\gamma)$. Moreover, all these curves pass through $\Psi_\Delta(x)$. It is now easy to see that $\Psi_\Delta(\Delta)$ must contain a horizontal curve χ passing through $\Psi_\Delta(x)$. Since horizontal curves are mapped by Ψ_Δ^{-1} into horizontal curves, we conclude that $\Psi_\Delta^{-1}(\chi)$ is a horizontal Δ -curve. □

Since sectors are bounded by strictly decreasing curves, among the sectors of a given order with a given vertex, there are only two whose closure contains horizontal curves passing through the vertex.

Definition 3.9. Let b_n be the maximum number of distinct regular sectors of order n with the same vertex. We call b_n the *branching number of order n* .

THEOREM 3.10. $b_n \leq (2n - 1)b_1$ for $n \geq 1$.

Proof. To prove the corollary, we show that $b_{n+1} - b_n \leq 2b_1$ for $n \geq 1$, which implies the wanted conclusion. This is achieved by estimating the maximum number of regular sectors of order n containing components of S_{n+1}^+ passing through their vertices, and the maximum number of such components.

Consider a regular sector Δ of order n with vertex x . Let a be the maximum number of Δ -curves contained in S_{n+1}^+ . Since Δ is of order n , these curves are indeed contained in $S_{n+1}^+ \setminus S_n^+$. By (3.1), their image under Ψ_Δ consists of an equal number of curves contained in S_1^+ and meeting at $\Psi_\Delta(x)$. Hence $a \leq b_1$. If $a > 0$, then Lemma 3.8 implies that there exist horizontal Δ -curves. Since there are only two sectors of order n with vertex x having this property, we conclude that $b_{n+1} - b_n \leq 2b_1$. □

Definition 3.11. A non-regular sector is called *primary* if it is not a sub-sector of a non-regular sector.

Remark 3.12. It is not difficult to see that if Δ is a non-regular sector of order n with vertex x , then either $x \in \partial M$ or there exists $0 \leq k < n$ such that Φ^k is a C^2 diffeomorphism on the closure of Δ , and $\Phi^k(x)$ is a tangential singularity. Moreover, every sub-sector of Δ is non-regular as well. If Δ is primary, then $k = n - 1$.

4. Existence of SRB measures

In this section, we prove our main result. It relies on results of Pesin and Sataev on the existence and properties of SRB measures for general hyperbolic piecewise smooth maps. For the convenience of the reader, we first state the results with the necessary definitions.

4.1. Hyperbolic piecewise smooth maps. Let \mathcal{M} be a smooth manifold with Riemannian metric ρ . The Lebesgue measure of \mathcal{M} generated by ρ is denoted by ν .

Let \mathcal{U} be a connected open subset of \mathcal{M} with compact closure. Finally, let \mathcal{N} be a subset of \mathcal{U} closed in the relative topology of \mathcal{U} .

Definition 4.1. A map $\mathcal{F} : \mathcal{U} \setminus \mathcal{N} \rightarrow \mathcal{M}$ is called a *piecewise smooth map* if \mathcal{F} is a C^2 diffeomorphism from $\mathcal{U} \setminus \mathcal{N}$ onto its image $\mathcal{F}(\mathcal{U} \setminus \mathcal{N})$. The set $\mathcal{N}^+ := \mathcal{N} \cup \partial\mathcal{M}$ is called the *singular set* of \mathcal{F} .

$$\text{Let } \mathcal{N}^- = \partial(\mathcal{F}(\mathcal{M} \setminus \mathcal{N}^+)).$$

Definition 4.2. Let $\mathcal{U}^+ = \{x \in \mathcal{U} : \mathcal{F}^n(x) \notin \mathcal{N}^+ \ \forall n \geq 0\}$ be the set of all elements of \mathcal{M} with infinite positive semi-orbit. Define

$$\mathcal{D} = \bigcap_{n \geq 0} \mathcal{F}^n(\mathcal{U}^+).$$

The set $\mathcal{A} = \overline{\mathcal{D}}$ is called the *generalized attractor* of \mathcal{F} .

Condition A1. There exist positive constants \tilde{A} and a such that for every $x \in \mathcal{U} \setminus \mathcal{N}^+$,

$$\|D^2\mathcal{F}(x)\| \leq \tilde{A}\rho(x, \mathcal{N}^+)^{-a} \quad \text{and} \quad \|D^2\mathcal{F}^{-1}(\mathcal{F}(x))\| \leq \tilde{A}\rho(x, \mathcal{N}^-)^{-a}.$$

A cone in $T_x\mathcal{M}$, $x \in \mathcal{U}$ with an axial linear subspace $P \subset T_x\mathcal{M}$ and angle $\alpha > 0$ is the set given by

$$C_\alpha(x, P) = \{v \in T_x\mathcal{M} : \angle(v, P) \leq \alpha\}.$$

Condition A2. The map \mathcal{F} is uniformly hyperbolic. Namely, there exist two constants $c > 0, \lambda > 1$ and cones

$$C^s(x) = C_{\alpha^s(x)}(x, P^s(x)) \quad \text{and} \quad C^u(x) = C_{\alpha^u(x)}(x, P^u(x))$$

with axial subspaces $P^u(x), P^s(x)$ and positive angles $\alpha^u(x), \alpha^s(x)$ for $x \in \mathcal{U} \setminus \mathcal{N}^+$ such that:

- (1) $T_x\mathcal{U} = P^u(x) \oplus P^s(x)$;
- (2) $\dim P^u(x)$ and $\dim P^s(x)$ are constant;
- (3) the angle between $C^u(x)$ and $C^s(x)$ is uniformly bounded away from zero;
- (4) $D\mathcal{F}(x)(C^u(x)) \subset C^u(\mathcal{F}(x))$ and $D\mathcal{F}^{-1}(\mathcal{F}(x))(C^s(\mathcal{F}(x))) \subset C^s(x)$ for $x \in \mathcal{U} \setminus \mathcal{N}^+$; and
- (5) if $x \in \mathcal{U}^+$ and $n \in \mathbb{N}$, then $\|D\mathcal{F}^n(x)v\| \geq c\lambda^n\|v\|$ for $v \in C^u(x)$, and $\|D\mathcal{F}^{-n}(\mathcal{F}^n(x))v\| \geq c\lambda^n\|v\|$ for $v \in C^s(x)$.

Definition 4.3. We say that a piecewise smooth map \mathcal{F} is *hyperbolic* or that \mathcal{A} is a *hyperbolic attractor* if \mathcal{F} satisfies Conditions A1 and A2.

Following [16], for every $\epsilon > 0$ and every $l \in \mathbb{N}$, we define

$$\begin{aligned} \hat{\mathcal{D}}_{\epsilon,l}^+ &= \{x \in \mathcal{U}^+ : \rho(\mathcal{F}^n(x), \mathcal{N}^+) \geq l^{-1}e^{-n\epsilon} \ \forall n \geq 0\}, \\ \mathcal{D}_{\epsilon,l}^- &= \{x \in \mathcal{A} : \rho(\mathcal{F}^{-n}(x), \mathcal{N}^-) \geq l^{-1}e^{-n\epsilon} \ \forall n \geq 0\}, \\ \mathcal{D}_{\epsilon,l}^+ &= \hat{\mathcal{D}}_{\epsilon,l}^+ \cap \mathcal{A} \end{aligned}$$

and

$$\mathcal{D}_\epsilon^\pm = \bigcup_{l \geq 1} \mathcal{D}_{\epsilon,l}^\pm, \quad \mathcal{D}_\epsilon^0 = \mathcal{D}_\epsilon^- \cap \mathcal{D}_\epsilon^+.$$

Roughly speaking, the sets $\mathcal{D}_{\epsilon,l}^+$ (respectively, $\mathcal{D}_{\epsilon,l}^-$) consists of points in \mathcal{D} whose forward (respectively, backward) orbit does not get too close to the singular set. These sets are compact.

The local stable manifold $W_{\text{loc}}^s(x)$ through a point $x \in \mathcal{U}^+$ is the set of all $y \in \mathcal{U}^+$ such that $\mathcal{F}^n(x)$ and $\mathcal{F}^n(y)$ belong to the same connected component of $\mathcal{M} \setminus \mathcal{N}^+$ for every $n \geq 0$ and $\rho(\mathcal{F}^n(x), \mathcal{F}^n(y)) \rightarrow 0$ as $n \rightarrow +\infty$. The local unstable manifold $W_{\text{loc}}^u(x)$ at $x \in \mathcal{D}$ is defined similarly by replacing \mathcal{F} with \mathcal{F}^{-1} .

Local stable and unstable manifolds exist at each point of \mathcal{D}_ϵ^0 for sufficiently small ϵ .

PROPOSITION 4.4. *If \mathcal{F} is a hyperbolic piecewise smooth map, then there exists $\epsilon_0 > 0$ such that for all $0 < \epsilon < \epsilon_0$, $W_{\text{loc}}^s(x)$ (respectively, $W_{\text{loc}}^u(x)$) is a C^1 embedded submanifold of uniform size $\delta_l > 0$ for every $l \in \mathbb{N}$ and for every $x \in \hat{\mathcal{D}}_{\epsilon,l}^+$ (respectively, $x \in \mathcal{D}_{\epsilon,l}^-$).*

Proof. This is [16, Proposition 4] for points in $\mathcal{D}_{\epsilon,l}^+$ (respectively, $\mathcal{D}_{\epsilon,l}^-$). In order to obtain the same result for points in $\hat{\mathcal{D}}_{\epsilon,l}^+$ it is enough to observe that the local stable manifold $W_{\text{loc}}^s(x)$ depends only on the forward orbit of x . □

Definition 4.5. Let \mathcal{F} be a hyperbolic piecewise smooth map. An invariant Borel probability measure μ on the attractor \mathcal{A} is called† SRB if $\mu(\mathcal{D}_\epsilon^0) = 1$ with $\epsilon > 0$, as in Proposition 4.4, and if the conditional measures of μ on the local unstable manifolds are absolutely continuous.

Definition 4.6. Given an invariant Borel probability μ on \mathcal{M} , its basin $B(\mu)$ is the set of points $x \in \mathcal{U}^+$ such that

$$\lim_{n \rightarrow +\infty} \frac{1}{n} \sum_{j=0}^{n-1} \varphi(\mathcal{F}^j(x)) = \int \varphi d\mu$$

for any continuous function $\varphi : \mathcal{M} \rightarrow \mathbb{R}$. We say that μ is a *physical measure* if $B(\mu)$ has positive ν -measure.

PROPOSITION 4.7. [16, Theorem 3] *Every ergodic SRB measure is a physical measure.*

Let $\mathcal{N}_\epsilon^+ \subset \mathcal{M}$ be the ϵ -neighbourhood of \mathcal{N}^+ for $\epsilon > 0$. The Lebesgue measure of a submanifold $W \subset \mathcal{M}$ is denoted by ν_W . A smooth submanifold $W \subset \mathcal{M}$ is called a *u-manifold* if the dimension of W is equal to the dimension of the unstable subspaces of \mathcal{F} and the tangent space of W at x is contained in $C^u(x)$ for every $x \in W$.

Condition H. There exist constants $C > 0, a \in (0, 1), \beta > 0$ and $\epsilon_0 > 0$ such that for every *u*-manifold W , every $n \geq 1$ and every $\epsilon \in (0, \epsilon_0)$,

$$\nu_W(W \cap \mathcal{F}^{-n}(\mathcal{N}_\epsilon^+)) \leq C\epsilon^\beta (a^n + \nu_W(W)).$$

Roughly speaking, this condition states that the relative measure of points in a *u*-manifold ending up in a small neighbourhood of \mathcal{N}^+ is of the same order as the size of the neighbourhood. To the best of our knowledge, Condition H was first introduced in [6] (see also [5]).

† These measures are called *Gibbs u-measures* in [16].

Theorem 4.8 below contains the main results of Pesin and Sataev concerning SRB measures for hyperbolic piecewise smooth maps. It states that if such a map \mathcal{F} satisfies Condition H, then \mathcal{F} admits SRB measures, each of them being a convex combination of finitely many ergodic SRB measures [17, Theorems 5.12 and 5.15]. Moreover, every ergodic SRB measure decomposes into finitely many Bernoulli components cyclically permuted by \mathcal{F} [16, Theorem 4], and the periodic orbits of \mathcal{F} are dense in the attractor \mathcal{A} [16, Theorem 11].

THEOREM 4.8. *Let \mathcal{F} be a hyperbolic piecewise smooth map satisfying Condition H. There exist finitely many ergodic SRB measures μ_1, \dots, μ_m concentrated on pairwise disjoint subsets $\mathcal{E}_1, \dots, \mathcal{E}_m$ of the attractor \mathcal{A} such that:*

- (1) *for every SRB measure μ , there exist $\alpha_1, \dots, \alpha_m \geq 0$ with $\sum_{i=1}^m \alpha_i = 1$ such that $\mu = \sum_i \alpha_i \mu_i$;*
- (2) *for each $i = 1, \dots, m$, there exist disjoint subsets $\mathcal{M}_{i,1}, \dots, \mathcal{M}_{i,k_i}$ with $k_i \in \mathbb{N}$ such that $\mathcal{E}_i = \bigcup_{j=1}^{k_i} \mathcal{M}_{i,j} \pmod{0}$, $\mathcal{F}(\mathcal{M}_{i,j}) = \mathcal{M}_{i,j+1}$ for $1 \leq j < k_i$, $\mathcal{F}(\mathcal{M}_{i,k_i}) = \mathcal{M}_{i,1}$ and the system $(\mathcal{F}^{k_i}|_{\mathcal{M}_{i,j}}, \mu_{i,j})$ with $\mu_{i,j}$ being the normalized restriction of μ_i to $\mathcal{M}_{i,j}$ is Bernoulli; and*
- (3) *the set of periodic points of \mathcal{F} is dense in \mathcal{A} .*

The system $(\mathcal{F}^{k_i}|_{\mathcal{M}_{i,j}}, \mu_{i,j})$ is called a *Bernoulli component* of \mathcal{F} .

We observe that Pesin obtained a weaker result than Conclusion (1) of Theorem 4.8. He proved the existence of SRB measures and that each SRB measure is a convex combination of countably many ergodic SRB measures.

Condition H does not appear in the works of Pesin and Sataev. However, this condition is equivalent to Conditions H3 and H4 assumed by Sataev. Pesin assumed similar but weaker conditions. Conditions H3 and H4 of Sataev are the following.

Condition H3. There exist positive constants B, β', ϵ_1 such that

$$\nu(\mathcal{F}^{-n}(\mathcal{N}_\epsilon^+)) < B\epsilon^{\beta'} \quad \text{for } n \geq 1 \text{ and } \epsilon \in (0, \epsilon_1).$$

A smooth submanifold $W \subset M$ is a *u-manifold* if the dimension of W is equal to the one of the unstable subspaces of \mathcal{F} and the tangent space of W is contained in $C^u(x)$ for every $x \in W$.

Condition H4. There exist positive constants β' and ϵ_1 such that, for every *u-manifold* W , there exist an integer $m = m(W)$ and a constant $B = B(W) > 0$ such that, for every $0 < \epsilon < \epsilon_1$:

- (1) $\nu_W(W \cap \mathcal{F}^{-n}(\mathcal{N}_\epsilon^+)) < \epsilon^{\beta'} \nu_W(W)$ for $n > m$; and
- (2) $\nu_W(W \cap \mathcal{F}^{-n}(\mathcal{N}_\epsilon^+)) < B\epsilon^{\beta'} \nu_W(W)$ for $n \geq 1$.

For completeness, we provide the proof of the equivalence between Condition H and Conditions H3 and H4.

LEMMA 4.9. *Condition H is equivalent to Conditions H3 and H4.*

Proof. The fact that H3 and H4 imply H is trivial. We prove the other direction of the equivalence. Since $a \in (0, 1)$ and $\nu_W(W)$ has a uniform upper bound in W , there exists

a constant B such that $C(a^n + \nu_W(W)) < B$ for every $n \geq 1$ and every u -manifold W . From H, it follows that if $\beta' = \beta$ and $\epsilon_1 = \epsilon_0$, then

$$\nu_W(W \cap \mathcal{F}^{-n}(\mathcal{N}_\epsilon^+)) < B\epsilon^{\beta'}$$

for every $n \geq 1$ and every u -manifold W . Condition H3 follows from the previous inequality by covering \mathcal{U} with a smooth family of u -manifolds and using Fubini's Theorem. Condition H4 follows directly from H by taking

$$\beta' \in (0, \beta), \quad \epsilon_1 = \min\{\epsilon_0, (2C)^{1/(\beta' - \beta)}\}$$

and

$$B(W) = \frac{1 + a/\nu_W(W)}{2}, \quad m(W) = \left\lfloor \frac{\log \nu_W(W)}{\log a} \right\rfloor,$$

where $\lfloor x \rfloor$ is the integer part of x . Indeed, if $\epsilon < \epsilon_1$ and $n > m(W)$, then

$$\begin{aligned} \nu_W(W \cap \mathcal{F}^{-n}(\mathcal{N}_\epsilon^+)) &\leq C\epsilon^\beta(a^n + \nu_W(W)) \\ &\leq 2C\epsilon^\beta \nu_W(W) \leq \epsilon^{\beta'} \nu_W(W). \end{aligned}$$

Moreover, for every $n \geq 1$,

$$\begin{aligned} \nu_W(W \cap \mathcal{F}^{-n}(\mathcal{N}_\epsilon^+)) &\leq C\epsilon^\beta(a^n + \nu_W(W)) \leq C\epsilon^\beta(a + \nu_W(W)) \\ &\leq 2C\epsilon^\beta B(W)\nu_W(W) \leq \epsilon^{\beta'} B(W)\nu_W(W). \end{aligned} \quad \square$$

4.2. *SRB measures for polygonal billiards.* Recall that $\Phi_{f,P}$ denotes the billiard map for the polygon P with a contracting reflection law f satisfying the conditions introduced at the end of §2. As before, we will simply write Φ for $\Phi_{f,P}$ when no confusion can arise. Also, recall that $N^+ = V \cup S \cup \partial M$ and $N^- = \partial(\Phi(M \setminus N^+))$.

The sets $M, N, N^\pm, D, D_\epsilon^\pm, A, \dots$ are the analogue for the billiard map Φ of the sets $\mathcal{M}, \mathcal{N}, \mathcal{N}^\pm, \mathcal{D}, \mathcal{D}_\epsilon^\pm, \mathcal{A}, \dots$ for a general piecewise smooth map \mathcal{F} . We also observe that, for Φ , the analogue of \mathcal{U} is the set $M \setminus \partial M$.

We say that a polygon P has *no parallel sides facing each other* if the endpoints of every straight segment contained inside P and joining orthogonally two sides of P are vertices of P . Notice that P has no parallel sides facing each other if and only if Φ has no periodic orbits of period two. The reflection law f does not play any role in the previous claim because we assumed that $f(0) = 0$.

PROPOSITION 4.10. *The map $\Phi_{f,P}$ is piecewise smooth satisfying Condition A1. Moreover, $\Phi_{f,P}$ satisfies Condition A2 if and only if P does not have parallel sides facing each other.*

Proof. The first part is a direct consequence of the fact that the standard billiard map satisfies Condition A1 (see [13, Theorem 7.2]) and that a reflection law f together with its inverse has bounded second derivatives. The second claim follows from [9, Corollary 3.4]. □

Remark 4.11. It is easy to see that the horizontal direction ($\theta = \text{constant}$) is always preserved by $D\Phi_{f,P}$. If $\Phi_{f,P}$ is uniformly hyperbolic, then the horizontal direction is indeed the expanding direction of $\Phi_{f,P}$ [9, Corollary 3.4].

We can now state the first part of our main result.

THEOREM 4.12. *If P does not have parallel sides facing each other, then the conclusions of Theorem 4.8 hold for $\Phi_{f,P}$ for every contracting reflection law f . Moreover, each Bernoulli component of $\Phi_{f,P}$ has exponential decay of correlations for Hölder observables.*

The proof of this theorem is given in §4.4.

4.3. Growth lemma. We introduce a new condition called n -step expansion and prove that it implies Condition H. Results of this type are called *growth lemmas* (for instance, see [5, §5]). We adopt this terminology. The n -step expansion condition was introduced in [6]. Rather than giving the most general formulation of this condition, we formulate it only for the billiard map Φ .

Definition 4.13. A horizontal open segment contained in $M \setminus \partial M$ is called an h -curve.

Definition 4.14. We say that Φ satisfies the n -step expansion condition if there exists $n \in \mathbb{N}$ such that

$$\beta(\Phi) := \liminf_{\delta \rightarrow 0} \sup_{\Gamma \in \mathcal{H}(\delta)} \sum_{\gamma \in \pi_0(\Gamma \setminus N_n^+)} \frac{1}{a_n(\gamma)} < 1, \tag{4.1}$$

where $\mathcal{H}(\delta)$ is the set of h -curves of length less than or equal to δ , $\pi_0(\Gamma \setminus N_n^+)$ is the set of connected components of $\Gamma \setminus N_n^+$ and $a_n(\gamma)$ is the least expansion coefficient of $D\Phi^n|_{(1,0)}$ on γ : that is,

$$a_n(\gamma) = \inf_{x \in \gamma} \|D_x \Phi^n|_{(1,0)}\|.$$

Given an h -curve γ , we denote by ℓ_γ the Lebesgue of γ . We will drop the index γ in ℓ_γ when no confusion can arise about which curve γ the measure ℓ refers to.

Recall that N_ϵ^+ denotes the ϵ -neighbourhood of N^+ .

THEOREM 4.15. (Growth lemma) *If Φ satisfies the n -step expansion condition, then there exist $\beta(\Phi) \leq a < 1$, $\epsilon_0 > 0$ and $C > 0$ such that, for any h -curve Γ , $r \geq 0$ and $0 < \epsilon < \epsilon_0$,*

$$\ell(\Gamma \cap \Phi^{-r}(N_\epsilon^+)) \leq C\epsilon(a^r + \ell(\Gamma)). \tag{4.2}$$

Proof. Since $\Phi(M \setminus N) \subset (-\lambda(f)\pi/2, \lambda(f)\pi/2)$ and $\lambda(f) < 1$, there exists a small ϵ_0 such that ϵ_0 -neighbourhood of ∂M does not intersect $\Phi(M \setminus N^+)$. Therefore, it is enough to prove (4.2) with N_ϵ^+ replaced by N_ϵ .

Choose $\delta > 0$ in such a way that

$$\zeta := \sup_{\Gamma \in \mathcal{H}(\delta)} \sum_{\gamma \in \pi_0(\Gamma \setminus N_1^+)} \frac{1}{a_1(\gamma)} < 1.$$

Notice that $\zeta^{1/n} \geq \zeta \geq \beta(\Phi)$. We call an h -curve *long* if its length is larger than or equal to δ ; otherwise we call it *short*.

Let $N_0^+ = \emptyset$ and consider an h -curve Γ . By Remark 3.3, the set $\Gamma \cap N_p^+$ consists of q_p elements for every $p \geq 0$. Hence $\Phi^p(\Gamma \setminus N_p^+)$ is a union of pairwise disjoint h -curves: that is,

$$\Phi^p(\Gamma \setminus N_p^+) = \bigcup_{i=1}^{q_p} \Gamma_{p,i}.$$

Clearly, $q_p \leq q_{p+1}$, and $q_0 = 1$ because $\Gamma_{0,1} = \Gamma$.

Write $r = mn + u \geq 0$ for $m \geq 0$ and $0 \leq u \leq n - 1$. Let $\Psi = \Phi^n$. Let $I_{k,l}$ be the set of indices $i \in \{1, \dots, q_{mn}\}$ such that:

- (1) $\Psi^{-m+k}(\Gamma_{mn,i}) \subset \Gamma_{kn,l}$; and
- (2) $\Psi^{-s}(\Gamma_{mn,i})$ is contained in a short h -curve for $1 \leq s \leq m - k - 1$.

Denote by \mathcal{L} the set of pairs of indices (k, l) with $k \in \{1, \dots, m\}$ and $l \in \{1, \dots, q_{kn}\}$ such that $\Gamma_{kn,l}$ is a long h -curve. The sets $\{I_{k,l}\}_{(k,l) \in \mathcal{L}}$ are disjoint and together with $I_{0,1}$ form a partition of $\{1, \dots, q_{mn}\}$. Then we can estimate $\ell(\Gamma \cap \Phi^{-r}(N_\epsilon))$ as

$$\begin{aligned} \ell(\Gamma \cap \Phi^{-r}(N_\epsilon)) &= \ell(\Gamma \cap \Psi^{-m}(\Phi^{-u}(N_\epsilon))) \\ &= \sum_{i=1}^{q_m} \ell(\Psi^{-m}(\Gamma_{mn,i} \cap \Phi^{-u}(N_\epsilon))) \\ &\leq \ell(\Lambda_{0,1}) + \sum_{(k,l) \in \mathcal{L}} \ell(\Psi^{-k}(\Lambda_{k,l})), \end{aligned}$$

where

$$\Lambda_{k,l} := \bigcup_{i \in I_{k,l}} \Psi^{-m+k}(\Gamma_{mn,i} \cap \Phi^{-u}(N_\epsilon)).$$

Since the restriction of Ψ^{-m+k} to $\Gamma_{mn,i}$ is affine,

$$\frac{\ell(\Psi^{-k}(\Lambda_{k,l}))}{\ell(\Psi^{-k}(\Gamma_{kn,l}))} = \frac{\ell(\Lambda_{k,l})}{\ell(\Gamma_{kn,l})}.$$

Since $\Gamma_{kn,l}$ is long for $(k, l) \in \mathcal{L}$, it follows that, for any $r \geq 0$ and any $0 < \epsilon < \epsilon_0$,

$$\ell(\Gamma \cap \Phi^{-r}(N_\epsilon)) \leq \ell(\Lambda_{0,1}) + \frac{1}{\delta} \sum_{(k,l) \in \mathcal{L}} \ell(\Lambda_{k,l}) \ell(\Psi^{-k}(\Gamma_{kn,l})).$$

We now estimate $\ell(\Lambda_{k,l})$. Let $t = m - k$. Given $i \in I_{k,l}$, let $i_s \in \{1, \dots, q_{(m-t+s)n}\}$ be the index defined by $\Psi^{-t+s}(\Gamma_{mn,i}) \subset \Gamma_{(m-t+s)n,i_s}$ for $1 \leq s \leq t$. Also, let $a(s, i_s)$ be the least expansion of Ψ along the curve $\Psi^{-1}(\Gamma_{(m-t+s)n,i_s})$ for $1 \leq s \leq t$. Thus

$$\begin{aligned} \ell(\Lambda_{k,l}) &\leq \sum_{i \in I_{k,l}} \ell(\Psi^{-t}(\Gamma_{mn,i} \cap \Phi^{-u}(N_\epsilon))) \\ &\leq \sum_{i \in I_{k,l}} \frac{\ell(\Gamma_{mn,i} \cap \Phi^{-u}(N_\epsilon))}{a(1, i_1) \cdots a(t, i_t)}. \end{aligned}$$

The uniform transversality between N and the horizontal direction implies that there exists a constant $C' > 0$ independent of Γ, r and i' such that

$$\ell(\Gamma_{r,i'} \cap N_\epsilon) \leq C' \epsilon.$$

Let $d > 0$ be the maximum number of intersection points of N with h -curves. Also, let $b > 0$ be the least expansion of Φ along the horizontal direction. Note that b is not necessarily greater than one and that d and b depend only on Φ . Then each $\Gamma_{mn,i}$ contains at most $(d + 1)^u$ curves $\Phi^{-u}(\Gamma_{r,i'})$. This, together the previous estimate, implies that

$$\ell(\Gamma_{mn,i} \cap \Phi^{-u}(N_\epsilon)) \leq \frac{(d + 1)^u}{b^u} C' \epsilon.$$

Define $I_1 = \{i_1 : i \in I_{k,l}\}$ and $I_s = \{i_{s-1}, i_s : i \in I_{k,l}\}$ for $1 \leq s \leq t$. Then

$$\begin{aligned} \sum_{i \in I_{k,l}} \frac{1}{a(1, i_1) \cdots a(t, i_t)} &= \sum_{j_1 \in I_1} \sum_{(j_1, j_2) \in I_2} \cdots \sum_{(j_{t-1}, j_t) \in I_t} \frac{1}{a(1, j_1) \cdots a(t, j_t)} \\ &= \sum_{j_1 \in I_1} \frac{1}{a(1, j_1)} \sum_{(j_1, j_2) \in I_2} \frac{1}{a(2, j_2)} \cdots \sum_{(j_{t-1}, j_t) \in I_t} \frac{1}{a(t, j_t)}. \end{aligned}$$

Since the curve $\Gamma_{(m-t+s-1)n, j_{s-1}} \supseteq \Psi^{-1}(\Gamma_{(m-t+s)n, j_s})$ is short, (4.1) implies that

$$\sum_{(j_{t-1}, j_t) \in I_t} \frac{1}{a(s, j_s)} \leq \zeta, \quad 2 \leq s \leq t.$$

The same argument does not necessarily apply to $\sum_{j_1 \in I_1} 1/a(1, j_1)$ because the curve $\Gamma_{kn,l} \supseteq \Psi^{-1}(\Gamma_{(m-t+1)n, j_1})$ may be long. However,

$$\sum_{j_1 \in I_1} \frac{1}{a(1, j_1)} \leq \frac{d+1}{b}.$$

Hence

$$\sum_{i \in I_{k,l}} \frac{1}{a(1, i_1) \cdots a(t, i_t)} \leq \frac{d+1}{b} \zeta^{t-1},$$

and so

$$\ell(\Lambda_{k,l}) \leq C'' \epsilon \zeta^{m-k}, \quad C'' = \frac{C'}{\zeta} \left(\frac{d+1}{b}\right)^{u+1}.$$

The above estimate implies that

$$\ell(\Gamma \cap \Phi^{-r}(N_\epsilon)) \leq C'' \epsilon \zeta^m + \frac{C'' \epsilon}{\delta} \sum_{k=1}^m \zeta^{m-k} \sum_{l=1}^{q_{kn}} \ell(\Psi^{-k}(\Gamma_{kn,l})).$$

Since

$$\sum_l \ell(\Psi^{-k}(\Gamma_{kn,l})) = \ell\left(\Psi^{-k}\left(\bigcup_{l=1}^{q_{kn}} \Gamma_{kn,l}\right)\right) \leq \ell(\Gamma),$$

$$\begin{aligned} \ell(\Gamma \cap \Phi^{-r}(N_\epsilon)) &\leq C'' \epsilon \zeta^m + \frac{C'' \epsilon \ell(\Gamma)}{\delta(1-\zeta)} \\ &\leq C'' \epsilon \zeta^{(r/n)-1} + \frac{C'' \epsilon \ell(\Gamma)}{\delta(1-\zeta)}, \end{aligned}$$

which implies the wanted conclusion. □

4.4. *Proof of Theorem 4.12.* In this subsection, we prove the n -step expansion for the billiard map and Theorem 4.12.

Since we assume that the polygon P does not have parallel sides facing each other, the map Φ is uniformly hyperbolic, by Proposition 4.10.

Define the least expansion rate of $D\Phi^n$ along the unstable direction by

$$A_n := \inf_{x \in M^+} \|D_x \Phi^n|_{(1,0)}\|.$$

Also define

$$\alpha(x) := \frac{\cos \theta(x)}{\cos \bar{\theta}_1(x)} \geq \cos\left(\frac{\pi}{2}\lambda(f)\right) \tag{4.3}$$

and

$$\alpha_n(x) := \|D_x \Phi^n|_{(1,0)}\| = \alpha(x) \cdots \alpha(\Phi^{n-1}(x)). \tag{4.4}$$

LEMMA 4.16. *Suppose that Δ is a primary non-regular sector of order n with vertex $x \notin \partial M$. If Δ' is a sub-sector of Δ of order $m > n$, or if $\Delta' = \Delta$ and $m = n$, then*

$$\lim_{\Delta' \ni y \rightarrow x} \alpha_m(y) = +\infty.$$

Proof. By Remark 3.12, the map Φ^{n-1} is a C^2 diffeomorphism on the closure of Δ and $\Phi^{n-1}(x)$ is a tangential singularity. Thus $\bar{\theta}_1(\Phi^k(x)) = \pm\pi/2$, and so $\alpha(\Phi^{n-1}(y)) \rightarrow +\infty$ as $\Delta \ni y \rightarrow x$. The claim now follows from (4.3), (4.4) and $\Delta' \subseteq \Delta$. \square

PROPOSITION 4.17. *The map Φ has n -step expansion for every sufficiently large n .*

Proof. Given a sector Δ of order $\leq n$ with vertex x , let

$$\Delta_\epsilon = \{y \in \Delta(x) : \text{dist}(y, x) < \epsilon\}$$

for $\epsilon > 0$. Denote by C_ϵ the union of all Δ_ϵ with Δ being a primary non-regular sector of order $\leq n$ with vertex not belonging to ∂M .

For a fixed $\Gamma \in \mathcal{H}(\delta)$,

$$\sum_{\gamma \in \pi_0(\Gamma \setminus N_n^+)} \frac{1}{a_n(\gamma)} = \sum_{\gamma'} \frac{1}{a_n(\gamma')} + \sum_{\gamma''} \frac{1}{a_n(\gamma'')},$$

where $\sum_{\gamma'}$ and $\sum_{\gamma''}$ denote the sum over the components of $\Gamma \setminus N_n^+$ intersecting the complement of C_ϵ and contained in C_ϵ , respectively. From the definition of b_n , it follows that there are at most b_n connected components of $\Gamma \setminus N_n^+$ contained in the complement of C_ϵ provided that δ is sufficiently small.

Since Φ is hyperbolic, there exist $c > 0$ and $\Lambda > 1$ such that $A_n \geq c\Lambda^n$ for every $n \in \mathbb{N}$. As b_n grows linearly in n (by Theorem 3.10), we can find $n_0 \in \mathbb{N}$ such that $A_n > b_n$ for every $n \geq n_0$. Choose $n \geq n_0$, which will be kept fixed throughout the rest of the proof. Now, N_n^+ consists of finitely many curves that either are disjoint or intersect pairwise at finitely many points (see the proof of [11, Proposition 2.3]). Thus, there exists $d(\epsilon) \rightarrow 0$ as $\epsilon \rightarrow 0$ such that any two distinct components of $C_\epsilon^c \cap N_n^+$ either have a distance greater than $d(\epsilon)$ or meet at a point that is not a tangential singularity. In view of the choice of n and the fact that $a_n(\gamma) \geq A_n$, there exists $0 < \eta < 1$ such that

$$\sum_{\gamma'} \frac{1}{a_n(\gamma')} \leq \frac{b_n}{A_n} = 1 - \eta, \tag{4.5}$$

provided that $\delta < d(\epsilon)$.

From Lemma 4.16, $a_n(\gamma'') \rightarrow +\infty$ as $\epsilon \rightarrow 0$ for every γ'' . Thus, by choosing ϵ sufficiently small, we can make sure that

$$\sum_{\gamma''} \frac{1}{a_n(\gamma'')} < \frac{\eta}{2}. \tag{4.6}$$

Combining (4.5) and (4.6), which do not depend on Γ ,

$$\sum_{\gamma} \frac{1}{a_n(\gamma)} < 1 - \frac{\eta}{2}$$

for every sufficiently small δ and for every $\Gamma \in \mathcal{H}(\delta)$. This implies (4.1) and completes the proof. \square

Proof of Theorem 4.12. First of all, we observe that, for $\mathcal{F} = \Phi$, Theorem 4.8 remains valid if Condition H is satisfied only by h -curves, since the local unstable manifolds of Φ are h -curves (see [11, Proposition 2.7]). By Proposition 4.10, the map Φ satisfies Conditions A1 and A2. By Proposition 4.17 and Theorem 4.15, Condition H holds for Φ and h -curves. Hence Theorem 4.8 applies to Φ .

We now prove the second claim of the theorem. Consider a Bernoulli component $(\Phi^{k_i}|_{\mathcal{M}_{i,j}}, \mu_{i,j})$ of Φ and let $\Psi = \Phi^{k_i}|_{\mathcal{M}_{i,j}}$. By Proposition 4.17, the n -step expansion condition holds true for Ψ for some $n \geq k_i$. It follows from [4, Proposition 10.1] (see also [6, Theorem 10]) that it is enough to establish the exponential decay of correlations for $(\Psi^n, \mu_{i,j})$. To do that, we apply a theorem of Chernov and Zhang [7, Theorem 1] to Ψ^n . This theorem has five hypotheses H.1–H.5. It is not difficult to see that H.1 and H2 follow from our Conditions A2 and A1, respectively. The finiteness of the number of smooth components of N^+ and N^- follows from Remark 3.3 and the first part of the proof of Lemma 3.6. Hypothesis H.3 is satisfied if we take as the Ψ^n -invariant class of smooth u -curves the set of all h -curves. Indeed, this class satisfies the three conditions of H.3: (i) the curvature of the h -curves is clearly uniformly bounded; (ii) the restriction of $D\Psi^n$ along h -curves has uniform distortion bounds because the restriction of Ψ^n to an h -curve is a piecewise affine map; and (iii) by Lemma 5.7 (see also [16, Proposition 10]), the stable holonomy is absolutely continuous. Since $(\Psi, \mu_{i,j})$ is Bernoulli, hypothesis H.4 is trivially satisfied. Thus, using [7, Theorem 1], we conclude that $(\Psi^n, \mu_{i,j})$ has exponential decay of correlations for Hölder observables. The same is true for $(\Psi, \mu_{i,j})$ by [4, Proposition 10.1]. \square

5. Basins of the ergodic SRB measures

Recall that $\Phi = \Phi_{f,P}$ denotes the billiard map for the polygon P with contracting reflection law f . The aim of this section is to prove the following theorem.

THEOREM 5.1. *Let P be a polygon without parallel sides facing each other. Then the union of the basins of the ergodic SRB measures of Φ has full Lebesgue measure.*

Let us briefly sketch the strategy to prove this theorem, which is based on [3, Proposition 4.2].

5.1. Outline of the strategy. We take any horizontal line segment γ (i.e., parallel to the unstable direction) and prove in Lemma 5.9 that a positive fraction of its points are in the basin of attraction of some ergodic SRB measure μ_0 . For this purpose, we consider the accumulation points of the Birkhoff averages $(1/n) \sum_{j=0}^{n-1} \Phi^j \text{Leb}_{\gamma}$, which start from the Lebesgue measure Leb_{γ} on the horizontal curve γ . Ergodic components of these limit points are SRB measures (Lemmas 5.6 and 5.8).

Now let A be the complement of the union of the basins of attraction of SRB measures and assume, by contradiction, that $\text{Leb}(A) > 0$. By Fubini’s Theorem, there exists a horizontal curve γ such that $\text{Leb}_\gamma(A) > 0$. Take any full density point $x \in \gamma \cap A$. Then, for every SRB measure μ_0 , the density of $\gamma \cap B(\mu_0)$ at x is zero. On the other hand, picking an ergodic SRB measure μ_0 such that $\text{Leb}_\gamma(B(\mu_0)) > 0$, there are sub-arcs $\gamma_n \subset \gamma$ accumulating on x such that the iterates $\Phi^n(\gamma_n) \subset B(\mu_0)$ converge to some non-trivial segment γ_∞ of unstable manifold in the attractor associated to μ_0 . Since the iterates $\Phi^n(\gamma_n)$ accumulate on γ_∞ , the stable manifolds of points in γ_∞ intersect transversally $\Phi^n(\gamma_n)$. Thus, using the holonomy along stable manifolds and bounded distortion of the unstable dynamics, we obtain by pull-back a positive fraction $\eta > 0$ such that $\text{Leb}_{\gamma_n}(B(\mu_0)) \geq \eta > 0$ for all large n . This contradicts the fact that $\gamma \cap B(\mu_0)$ has zero density at x .

For smooth systems, as in [3], the previous sketch proof can be easily implemented. In our setting, because basins of attraction are in general not open, and also because the sizes of local stable and unstable manifolds are not uniform, the sketch proof above requires elaboration. Three key ingredients for this purpose are Lemma 5.4, showing that most of the connected components of the iterates $\Phi^n(\gamma)$ of a horizontal curve γ have lengths above any small specified threshold, Lemma 5.5, saying that on any horizontal curve almost every point is forward regular in the sense of Pesin, and Lemma 5.7, showing that the holonomy along stable manifolds is Lipschitz when we fix the order $l \in \mathbb{N}$ of Pesin’s sets (controlling the size of stable manifolds).

Finally, we note that, in our setting, the unstable distortion is always zero because the corresponding dynamic is affine. Compared with [3], this simplifies our analysis slightly.

5.2. *Preparatory lemmas.* Choose $\epsilon > 0$ so that Proposition 4.4 and Theorem 4.15 hold. From now on, to simplify our notation, we will drop the index ϵ from the symbols $\hat{D}_{\epsilon,l}^+, D_{\epsilon,l}^-, D_{\epsilon,l}^0$.

Definition 5.2. Given a C^1 simple open curve γ in M and a point $x \in \gamma$, we say that γ has size δ around x if the length of the connected components of $\gamma \setminus \{x\}$ is greater than or equal to δ .

Definition 5.3. Given an h -curve γ , a point $x \in \gamma$ and $n \in \mathbb{N}$, let $R_\gamma^n(x)$ be the connected component of $\Phi^n(\gamma)$ containing $\Phi^n(x)$. Define

$$\gamma(n, \delta) = \{y \in \gamma : R_\gamma^n(y) \text{ has size } \delta \text{ around } \Phi^n(y)\}.$$

Let $\epsilon_0 > 0$ and $0 < a < 1$ be as in Condition H and let $c > 0$ and $\lambda > 1$ be as in Condition A2.

LEMMA 5.4. *There exists $\tilde{C} = \tilde{C}(\Phi) > 0$ such that, for every h -curve γ , there exists $n_0 = n_0(\gamma)$ such that if $n \geq n_0$ and $0 < \delta < c\epsilon_0$, then*

$$\ell(\gamma \setminus \gamma(n, \delta)) \leq \tilde{C}\ell(\gamma)\delta.$$

Proof. Let γ be an h -curve. If $x \in \gamma \setminus \gamma(n, \delta)$, then either $x \in N_n^+$ or $d(\Phi^n(x), \partial\Phi^n(\gamma \setminus N_n^+)) < \delta$. Let $B \subset \partial\Phi^n(\gamma \setminus N_n^+)$ be the image under Φ^n of the endpoints of γ that do

not belong to N_n^+ . Notice that this set consists of at most two points and the elements of $\partial\Phi^n(\gamma \setminus N_n^+) \setminus B$ are discontinuities of Φ^{-n} . Let B_δ denote the δ -neighbourhood of B . Since the horizontal direction coincides with the unstable direction and $\|D\Phi^n|_{E^u}\| \geq c\lambda^n$,

$$\gamma \setminus \gamma(n, \delta) \subset \bigcup_{i=0}^{n-1} (\gamma \cap \Phi^{-i}(N_{\delta\lambda^{-n+i}/c}^+) \cup (\gamma \cap \Phi^{-n}(B_\delta)).$$

Using (4.2),

$$\begin{aligned} \ell(\gamma \setminus \gamma(n, \delta)) &\leq \sum_{i=0}^{n-1} \ell(\gamma \cap \Phi^{-i}(N_{\delta\lambda^{-n+i}/c}^+)) + \ell(\gamma \cap \Phi^{-n}(B_\delta)) \\ &\leq \sum_{i=0}^{n-1} \frac{C\delta}{c\lambda^{n-i}} (a^i + \ell(\gamma)) + \frac{2\delta}{c\lambda^n} \\ &\leq \frac{c\delta}{c\lambda^n} \sum_{i=0}^{n-1} ((a\lambda)^i + \lambda^i \ell(\gamma)) + \frac{2\delta}{c\lambda^n} \\ &\leq \frac{C\delta}{c\lambda^n} \left(\frac{1 - (a\lambda)^n}{1 - a\lambda} + \frac{1 - \lambda^n}{1 - \lambda} \ell(\gamma) \right) + \frac{2\delta}{c\lambda^n} \\ &\leq \frac{C\delta}{c} \left(\frac{\lambda^{-n} - a^n}{1 - a\lambda} + \frac{\lambda^{-n} - 1}{1 - \lambda} \ell(\gamma) + \frac{2}{C\lambda^n} \right) \\ &\leq \frac{2C}{c} \ell(\gamma) \delta \end{aligned}$$

for all n such that the expression in parentheses in the penultimate inequality is smaller than $2\ell(\gamma)$. □

Let $\hat{D}^+ = \bigcup_{l \in \mathbb{N}} \hat{D}_l^+$.

LEMMA 5.5. For any h -curve γ , $\ell(\gamma) = \ell(\gamma \cap \hat{D}^+)$.

Proof. Since \hat{D}_l^+ is increasing in l , $\ell(\gamma \setminus \hat{D}_l^+) \rightarrow \ell(\gamma \setminus \hat{D}^+)$ as $l \rightarrow +\infty$. Now

$$\gamma \setminus \hat{D}_l^+ \subset \bigcup_{n=0}^{\infty} (\gamma \cap \Phi^{-n}(N_{l^{-1}e^{-\epsilon n}}^+)).$$

If l is sufficiently large, then, by (4.2),

$$\begin{aligned} \ell(\gamma \setminus \hat{D}_l^+) &\leq \sum_{n=0}^{\infty} \ell(\gamma \cap \Phi^{-n}(N_{l^{-1}e^{-\epsilon n}}^+)) \\ &\leq \frac{C}{l} \sum_{n=0}^{\infty} e^{-\epsilon n} (a^n + \ell(\gamma)) \leq \frac{C}{l} \cdot \frac{1 + \ell(\gamma)}{1 - e^{-\epsilon}} \xrightarrow{l \rightarrow +\infty} 0. \end{aligned} \quad \square$$

Given an h -curve γ , we call a γ -limit measure any weak- $*$ accumulation point of the averages

$$\mu_{\gamma,n} := \frac{1}{n} \sum_{j=0}^{n-1} \Phi_*^j \text{Leb}_\gamma,$$

where Leb_γ denotes the normalized Lebesgue measure of γ : that is, given any Borel set $A \subset M$,

$$\text{Leb}_\gamma(A) = \frac{\ell(\gamma \cap A)}{\ell(\gamma)}.$$

The next is the key result on the existence of SRB measures proved in [16]. We recall part of the proof, for the convenience of the reader.

LEMMA 5.6. *If γ is an h -curve, then any γ -limit measure μ is invariant and satisfies $\mu(D^0) = 1$. In addition, if $\gamma \cap D^- \neq \emptyset$, then μ is an SRB measure.*

Proof. Consider the function $\varphi : M \setminus N^+ \rightarrow \mathbb{R}$,

$$\varphi(x) := \text{dist}(x, N^+),$$

where the distance between x and N^+ is measured along the horizontal line through x . Since φ is bounded from above, for any probability measure ν , the integral $\int \log \varphi d\nu$ is well defined in $[-\infty, +\infty)$. Notice also that, for all sub-intervals I of some fixed compact interval containing the origin, $\int_I \log |x| dx \geq -\ell(I)$. Hence, by a change of coordinates,

$$\int_{\gamma \cap \Phi^{-j}(N_\epsilon^+)} \log \text{dist}(\Phi^j(x), N^+) dx \geq -\ell(\gamma \cap \Phi^{-j}(N_\epsilon^+))$$

and, by (4.2), there exists a constant $C > 0$ such that

$$\begin{aligned} \int \log \varphi d\mu_{\gamma,n} &= \int_{N_\epsilon^+} \log \varphi d\mu_{\gamma,n} + \int_{M \setminus N_\epsilon^+} \log \varphi d\mu_{\gamma,n} \\ &\geq -\frac{1}{n} \sum_{j=0}^{n-1} \frac{C\epsilon(a^j + \ell(\gamma))}{\ell(\gamma)} - \log(1/\epsilon) \\ &\geq -C\epsilon - \frac{C\epsilon}{n\ell(\gamma)(1-\eta)} - \log(1/\epsilon) > -\infty. \end{aligned}$$

Because this lower bound is uniform in n , the function $\log \varphi$ is μ -integrable, with $\int \log \varphi d\mu \geq -C\epsilon - \log(1/\epsilon)$. Similarly, $\log \varphi \circ \Phi^k$ is μ -integrable for every $k \in \mathbb{Z}$. Hence $\mu(\Phi(N^+)) = 0$.

To prove that μ is invariant, it is enough to see that, for any continuous function $\psi : M \rightarrow \mathbb{R}$,

$$\int \psi d\mu = \int \psi \circ \Phi d\mu. \tag{5.1}$$

If $\psi = 0$ on $\Phi(N^+)$, then the composition $\psi \circ \Phi$ is also continuous on M and hence

$$\lim_{k \rightarrow +\infty} \int \psi \circ \Phi d\mu_{\gamma,n_k} = \int \psi \circ \Phi d\mu.$$

On the other hand, a standard calculation gives

$$\left| \int \psi d\mu_{\gamma,n} - \int \psi \circ \Phi d\mu_{\gamma,n} \right| \leq \frac{2}{n} \|\psi\|_\infty.$$

Passing to the limit, we get (5.1).

In general, given any continuous function $\psi : M \rightarrow \mathbb{R}$, because $\mu(\Phi(N^+)) = 0$, we can approximate ψ by continuous functions ψ' vanishing on $\Phi(N^+)$ and, taking the limit, we obtain the invariance relation (5.1) for ψ . This proves that μ is invariant.

By Lemma 5.5, $\text{Leb}_\gamma(\hat{D}^+) = 1$. Hence, because $\Phi(\hat{D}^+) \subseteq \hat{D}^+$, $\mu_{\gamma,n}(\hat{D}^+) = 1$ for all $n \geq 1$. Thus, taking the limit, $\mu(\hat{D}^+) = 1$.

We are now left to prove that $\mu(D^-) = 1$. From Birkhoff's ergodic theorem, for μ -almost every $x \in M$, $\lim_{n \rightarrow +\infty} (1/n) \log \varphi(\Phi^{-n}x) = 0$. This implies that there exists $l \in \mathbb{N}$ such that $\varphi(\Phi^{-n}x) \geq l^{-1}e^{-n\epsilon}$ for all $n \in \mathbb{N}$. Hence

$$\text{dist}(\Phi^{-(n-1)}x, N^-) \geq \text{dist}(\Phi^{-n}x, N^+) = \varphi(\Phi^{-n}x) \geq l^{-1}e^{-n\epsilon},$$

which proves that $x \in D_l^-$ for some $l \in \mathbb{N}$. Thus $\mu(D^-) = 1$. The last claim follows from the proof of existence of SRB measures in [16, Theorem 1]. □

In the next lemma, we prove that the stable holonomy is Lipschitz continuous (c.f. [16, Proposition 10]).

LEMMA 5.7. *Given $l \in \mathbb{N}$, there are constants $\tilde{\delta} > 0$ and $C > 0$ such that if Γ and Γ' are h -curves whose distance is less than $\tilde{\delta}$, $x_1, x_2 \in \Gamma \cap \hat{D}_l^+$ with $|x_1 - x_2| < \tilde{\delta}$ and $x'_i := \Gamma' \cap W_l^s(x_i)$, then*

$$\frac{|x'_1 - x'_2|}{|x_1 - x_2|} \leq C.$$

Proof. The proof consists of a few steps.

(a) First, the slope of local stable manifolds is uniformly bounded away from zero. This follows easily from the expression for tangent space to the stable manifold in the proof of [9, Proposition 3.1].

(b) Let $\Gamma_n(x_1)$ denote the Φ^n pre-image of the connected component of $\Phi^n(\Gamma)$ that contains $\Phi^n(x_1)$. Let $\Gamma'_n(x_1)$ be the corresponding component of Γ' with respect to x'_1 . Denoting $\alpha_n = (\Phi^n|_{\Gamma_n(x_1)})'$ and $\alpha'_n = (\Phi^n|_{\Gamma'_n(x'_1)})'$, respectively,

$$\left| \log \frac{\alpha_n}{\alpha'_n} \right| \leq C\delta.$$

Notice that $\alpha_n(s_0, \theta_0) = \prod_{j=0}^{n-1} \rho(\theta_j)$, where $\rho(\theta) := \cos f(\theta) / \cos \theta$ and, for each $i \geq 1$, (s_i, θ_i) is the image of (s_{i-1}, θ_{i-1}) by the specular billiard.

Given another point $(s'_0, \theta'_0) \in W_{\text{loc}}^s(s_0, \theta_0)$, denote by θ'_i the analogous angles for (s'_0, θ'_0) . Since $\log \rho(\theta)$ is Lipschitz and $|\theta'_j - \theta_j|$ decay geometrically with j ,

$$\begin{aligned} \left| \log \frac{\alpha_n(s_0, \theta_0)}{\alpha_n(s'_0, \theta'_0)} \right| &\leq \sum_{j=0}^{n-1} |\log \rho(\theta_j) - \log \rho(\theta'_j)| \\ &\leq \sum_{j=0}^{n-1} C|\theta_j - \theta'_j| \lesssim C|\theta_0 - \theta'_0| = C\delta. \end{aligned}$$

(c) Write $x_{1,n} = \Phi^n(x_1)$, $x_{2,n} = \Phi^n(x_2)$, and consider the first $n \geq 1$ such that $x_{1,n}$ and $x_{2,n}$ do not belong to the same branch domain of Φ . Because $x_i \in \hat{D}_l^+$, and these two points are separated by a singular curve, $|x_{1,n} - x_{2,n}| \leq 2l^{-1}e^{-n\epsilon}$. Writing $x'_{1,n} = \Phi^n(x'_1)$

and $x'_{2,n} = \Phi^n(x'_2)$, we have $|x_{1,n} - x'_{1,n}| = 1/\alpha_n(x_1) \ll l^{-1}e^{-n\epsilon}$. Hence, combining this information with (a),

$$\frac{|x'_{1,n} - x'_{2,n}|}{|x_{1,n} - x_{2,n}|} \leq C.$$

(d) By parts (b) and (c) and the mean value theorem,

$$\begin{aligned} \frac{|x'_1 - x'_2|}{|x_1 - x_2|} &\asymp \left| \frac{\alpha_n}{\alpha'_n} \right| \frac{|x'_{1,n} - x'_{2,n}|}{|x_{1,n} - x_{2,n}|} \\ &\leq e^{C\delta} \frac{|x'_{1,n} - x'_{2,n}|}{|x_{1,n} - x_{2,n}|} \leq Ce^{C\delta}. \end{aligned} \quad \square$$

By Proposition 4.4, for every $l \in \mathbb{N}$, there exists $\delta_l > 0$ such that any point $x \in \hat{D}_l^+$ (respectively, $x \in D_l^-$) has a local stable (respectively, unstable) curve of size δ_l around x , denoted by $W_l^s(x)$ (respectively, $W_l^u(x)$). For any set $A \subset D_l^0$, define $W_l^s(A) = \bigcup_{x \in A} W_l^s(x)$.

LEMMA 5.8. *Let γ be an h -curve such that $\gamma \cap D^- \neq \emptyset$. If μ is a γ -limit measure, then there exists an ergodic component μ_c of μ such that $\text{Leb}_\gamma(B(\mu_c)) > 0$.*

Proof. By Lemma 5.6, μ is an SRB measure of Φ . According to Theorem 4.12, we may decompose μ into a finite number of ergodic components $\mu_i, i = 1, \dots, r$ such that μ is a convex combination of the ergodic measures μ_i : that is, $\mu = \sum_i \alpha_i \mu_i$. If $B := \bigcup_{i=1}^r B(\mu_i)$, then

$$\mu(B) = 1. \tag{5.2}$$

Given $l \leq l'$ in \mathbb{N} , define

$$\Omega_{l,l'}^0 := \left\{ x \in D_l^0 : \text{Leb}_{W_l^u(x)}(D_{l'}^0 \cap B) > 1 - \frac{\delta}{10} \right\},$$

where $\delta = \delta(l)$ is smaller than $\tilde{\delta}$, as in Lemma 5.7, and to be determined later. Notice that $\mu(D^0 \cap B) = 1$ due to Lemma 5.6 and (5.2). By the properties of the SRB measures (see [16, Proposition 9]), we know that for μ -almost every $z \in D^0$ there exists $l \in \mathbb{N}$ such that $z \in D_l^0$ and $\text{Leb}_{W_l^u(x)}(D^0 \cap B) = 1$. This implies that, for some $l' \geq l$, $\text{Leb}_{W_{l'}^u(x)}(D_{l'}^0 \cap B) > 1 - \delta/10$. Therefore

$$\mu\left(\bigcup_{l \leq l'} \Omega_{l,l'}^0\right) = 1.$$

Let $B(x, \delta)$ be the open ball of size δ around $x \in M$. For every $x \in \Omega_{l,l'}^0$, denote by $\Pi_{l,l'}(x)$ the union of all h -curves of size $\delta/10$ centred at $y \in W_{l'}^s(x) \cap B(x, \delta)$.

The set

$$U_{l,l'} := \bigcup_{x \in \Omega_{l,l'}^0} \Pi_{l,l'}(x)$$

is open and contains $\Omega_{l,l'}^0$. Choose $l \leq l'$ sufficiently large such that

$$\mu(\Omega_{l,l'}^0) > \frac{9}{10}.$$

Since μ is a γ -limit and $U_{l,l'}$ is open,

$$\liminf_n \mu_{\gamma,n}(U_{l,l'}) \geq \mu(U_{l,l'}) \geq \mu(\Omega_{l,l'}^0) > \frac{9}{10}.$$

Hence, for infinitely many $n \in \mathbb{N}$,

$$\text{Leb}_\gamma(\Phi^{-n}(U_{l,l'})) > \frac{9}{10}. \tag{5.3}$$

Notice that $\delta(l) \rightarrow 0$ as $l \rightarrow \infty$. According to Lemma 5.4, for any sufficiently large $l \in \mathbb{N}$, there exists $n_0 \in \mathbb{N}$ such that, for every $n \geq n_0$,

$$\text{Leb}_\gamma(\gamma(n, \delta(l))) > \frac{9}{10}.$$

This lower bound together with (5.3) implies that there exist $l \leq l'$ and $n \in \mathbb{N}$ such that

$$\text{Leb}_\gamma(\gamma(n, \delta(l)) \cap \Phi^{-n}(U_{l,l'})) > \frac{4}{5}.$$

Therefore, $\Phi^n(y) \in \Pi_{l,l'}(x)$ for some $x \in \Omega_{l,l'}^0$ and $y \in \gamma(n, \delta)$.

Recall that $R_\gamma^n(y)$ is the h -curve of size δ around $\Phi^n(y)$. Hence, we can find $z \in \gamma$ such that $\Phi^n(z) \in W_l^s(x)$ and an h -curve contained in $R_\gamma^n(y)$ of size $\frac{9}{10}\delta$ around $\Phi^n(z)$. Moreover, since $x \in \Omega_{l,l'}^0$, we know that,

$$\text{Leb}_{W_l^u(x)}(D_l^0 \cap B) > 1 - \frac{\delta}{10}.$$

This means that $W_l^s(W_l^u(x) \cap D_l^0 \cap B)$ has no ‘vertical gaps’ having size larger than $\delta/10$. The local stable manifolds of points in D_l^0 vary continuously. Thus, by choosing a sufficiently small $\delta(l)$ and using the absolute continuity of the stable holonomy, we can make sure that $R_\gamma^n(y)$ intersects $W_l^s(W_l^u(x) \cap D_l^0 \cap B)$ on a set of positive ℓ -measure. Notice that if $z \in B \cap D_l^0$, then $W_l^s(z) \subset B$. Hence $\text{Leb}_{R_\gamma^n(y)}(B) > 0$. This implies that $\text{Leb}_\gamma(B(\mu_i)) > 0$ for some $i = 1, \dots, r$ because the restriction of Φ^{-n} to $R_\gamma^n(y)$ is affine. □

LEMMA 5.9. *If γ is an h -curve, then there exists an ergodic SRB measure μ_c such that $\text{Leb}_\gamma(B(\mu_c)) > 0$.*

Proof. In order to apply Lemma 5.8, we produce a sub-limit Γ of iterates of γ such that $\Gamma \cap D^- \neq \emptyset$.

Let μ be a γ -limit measure. By Lemma 5.6, we can take $l \in \mathbb{N}$ such that $\mu(D_l^0) > \frac{9}{10}$. By Lemma 5.4, there exists $\delta > 0$, which we assume to be much smaller than the size of the local stable curves of points in D_l^0 , such that, for all large n ,

$$\text{Leb}_\gamma(\gamma(n, \delta)) > \frac{9}{10}.$$

Denote by λ_0 (respectively, λ_1) the restriction of the measure Leb_γ to $\gamma(n, \delta)$ (respectively, $\gamma \setminus \gamma(n, \delta)$) so that $\text{Leb}_\gamma = \lambda_0 + \lambda_1$. Setting $\mu_{\gamma,n}^0 = (1/n) \sum_{j=0}^{n-1} \Phi_*^j \lambda_0$ and $\mu_{\gamma,n}^1 = (1/n) \sum_{j=0}^{n-1} \Phi_*^j \lambda_1$, we also have $\mu_{\gamma,n} = \mu_{\gamma,n}^0 + \mu_{\gamma,n}^1$. Hence there exists weak-*

sublimits μ_0 of $\mu_{\gamma,n}^0$ and μ_1 of $\mu_{\gamma,n}^1$, respectively, such that $\mu = \mu_0 + \mu_1$. By construction, μ_0 has total mass $\geq 9/10$ while μ_1 has total mass $\leq 1/10$.

Given a set $\gamma_0 \subset M$, let us say that $z \in M$ is a γ_0 -limit point if there exists a sequence of points $x_k \in \gamma_0$ and a sequence of times $n_k \rightarrow +\infty$ such that $z = \lim_{k \rightarrow +\infty} \Psi^{n_k}(x_k)$. With this terminology, any point $z \in \text{supp}(\mu)$ is a γ -limit point and any point $z \in \text{supp}(\mu_0)$ is a $\gamma(n, \delta)$ -limit point. Notice that $\mu(\text{supp}(\mu_0)) \geq \mu_0(\text{supp}(\mu_0)) \geq 9/10$. Therefore $\mu(\text{supp}(\mu_0) \cap D_\Gamma^0) > 4/5$. Hence, if $z \in \text{supp}(\mu_0) \cap D_\Gamma^0$ and Γ is an h -curve $\Gamma \subset W_\Gamma^u(z)$ of size δ around z , then Γ is accumulated by forward iterates γ_n of γ with $\ell(\gamma_n) \geq \delta$.

Applying Lemma 5.8 to Γ , $\text{Leb}_\Gamma(B(\mu_c)) > 0$ for some ergodic SRB measure μ_c , which is an ergodic component of a Γ -limit measure.

Since Γ is accumulated by forward iterates γ_n of γ with $\ell(\gamma_n) \geq \delta$, the Lipschitz continuity of the stable holonomy (see Lemma 5.7) implies that $\text{Leb}_{\gamma_n}(B(\mu_c)) > 0$, and so $\text{Leb}_\gamma(B(\mu_c)) > 0$ because the restriction of Φ^{-n} along γ_n is affine. □

5.3. *Proof of Theorem 5.1.* Let μ_1, \dots, μ_r be the ergodic SRB measures of Φ . Define

$$B = B(\mu_1) \cup \dots \cup B(\mu_r) \quad \text{and} \quad A = M \setminus B.$$

Assume that $\text{Leb}(A) > 0$. We will derive a contradiction from this assumption.

By Fubini's Theorem, there is an h -curve γ such that $\ell(\gamma \cap A) > 0$. By Lemma 5.4, for some small $\delta > 0$ and some $n_0 \in \mathbb{N}$, $\ell(\gamma(n, \delta) \cap A) > 0$ for all $n \geq n_0$.

Take a Lebesgue density point $x \in \gamma(n, \delta) \cap A$ and consider a strictly increasing sequence $\{n_i\}$ such that the sequence of h -curves $\{R_\gamma^{n_i}(x)\}$ converges to some h -curve Γ of length $\geq \delta$. Notice that the space of all h -curves with $\ell(\Gamma) \geq \delta$ endowed with the Hausdorff distance is compact.

By Lemma 5.9, $\text{Leb}_\Gamma(B) > 0$. Hence, Lemma 5.5 gives an $l \in \mathbb{N}$ such that

$$\eta := \ell(\Gamma \cap \hat{D}_l^+ \cap B) > 0.$$

Since the local stable curves of points of $\Gamma \cap \hat{D}_l^+ \cap B$ have uniform size δ_l , they must intersect $R_\gamma^{n_i}(x)$ for all sufficiently large i . The Lipschitz continuity of the stable holonomy (see Lemma 5.7) implies that

$$\ell(R_\gamma^{n_i}(x) \cap B) \geq \frac{\eta}{2} \ell(R_\gamma^{n_i}(x)).$$

Consider the curve $\gamma' := \Phi^{-n_i}(R_\gamma^{n_i}(x))$. Since the restriction of Φ^{n_i} to γ' is affine, we conclude that

$$\ell(\gamma' \cap B) \geq \frac{\eta}{2} \ell(\gamma').$$

However, x is a density point of $A \cap \gamma$, and so the proportion

$$\frac{\ell(\gamma' \cap B)}{\ell(\gamma')}$$

can be made arbitrarily small by choosing a sufficiently small h -curve γ' around x . This fact contradicts the previous conclusion.

Acknowledgement. PD was partially supported by National Funding from FCT—Fundação para a Ciência e a Tecnologia under the project: UID/MAT/04561/2013. JPG was supported through the FCT/MEC grant SFRH/BPD/78230/2011. JPG and JLD were partially supported by the Project CEMAPRE—UID/MULTI/00491/2013 financed by FCT/MEC through national funds.

REFERENCES

- [1] A. Arroyo, R. Markarian and D. P. Sanders. Bifurcations of periodic and chaotic attractors in pinball billiards with focusing boundaries. *Nonlinearity* **22**(7) (2009), 1499–1522.
- [2] A. Arroyo, R. Markarian and D. P. Sanders. Structure and evolution of strange attractors in non-elastic triangular billiards. *Chaos* **22** (2012), 026107.
- [3] C. Bonatti and M. Viana. SRB measures for partially hyperbolic systems whose central direction is mostly contracting. *Israel J. Math.* **115** (2000), 157–193.
- [4] N. Chernov. Decay of correlations and dispersing billiards. *J. Stat. Phys.* **94**(3–4) (1999), 513–556.
- [5] N. Chernov and R. Markarian. *Chaotic Billiards (Mathematical Surveys and Monographs, 127)*. American Mathematical Society, Providence, RI, 2006.
- [6] N. Chernov and H.-K. Zhang. Billiards with polynomial mixing rates. *Nonlinearity* **18**(4) (2005), 1527–1553.
- [7] N. Chernov and H.-K. Zhang. On statistical properties of hyperbolic systems with singularities. *J. Stat. Phys.* **136**(4) (2009), 615–642.
- [8] G. Del Magno, J. Lopes Dias, P. Duarte, J. P. Gaivão and D. Pinheiro. Chaos in the square billiard with a modified reflection law. *Chaos* **22** (2012), 026106.
- [9] G. Del Magno, J. Lopes Dias, P. Duarte, J. P. Gaivão and D. Pinheiro. SRB measures for polygonal billiards with contracting reflection laws. *Comm. Math. Phys.* **329** (2014), 687–723.
- [10] G. Del Magno, J. Lopes Dias, P. Duarte and J. P. Gaivão. Ergodicity of polygonal slap maps. *Nonlinearity* **27**(8) (2014), 1969–1983.
- [11] G. Del Magno, J. Lopes Dias, P. Duarte and J. P. Gaivão. Ergodic properties of polygonal billiards with strongly contracting reflection laws. *Preprint*, 2015, arXiv:1501.03697.
- [12] L. H. Erickson and S. M. La Valle. Toward the design and analysis of blind, bouncing robots. *IEEE Int. Conf. on Robotics and Automation (Karlsruhe, Germany, 6–10 May 2013)*. IEEE, Piscataway, NJ, 2013.
- [13] A. Katok et al. *Invariant Manifolds, Entropy and Billiards; Smooth Maps with Singularities (Lecture Notes in Mathematics, 1222)*. Springer, Berlin, 1986.
- [14] R. Markarian, E. J. Pujals and M. Sambarino. Pinball billiards with dominated splitting. *Ergod. Th. & Dynam. Sys.* **30**(6) (2010), 1757–1786.
- [15] J. Palis. A global view of dynamics and a conjecture on the denseness of finitude of attractors. *Astérisque* **261** (2000), 339–351.
- [16] Y. B. Pesin. Dynamical systems with generalized hyperbolic attractors: hyperbolic, ergodic and topological properties. *Ergod. Th. & Dynam. Sys.* **12**(1) (1992), 123–151.
- [17] E. A. Sataev. Invariant measures for hyperbolic mappings with singularities. *Uspekhi Mat. Nauk* **47**(1(283)) (1992), 147–202, 240; Engl. trans. *Russian Math. Surveys* **47**(1) (1992), 191–251.
- [18] R. E. Sheriff and L. P. Geldart. *Exploration Seismology*. Cambridge University Press, Cambridge, 1995.