

# Intrinsic Properties of a Non-Symmetric Number Triangle

Isabel Cação and Helmuth R. Malonek  
CIDMA and Department of Mathematics  
University of Aveiro  
3810-193 Aveiro  
Portugal  
[isabel.cacao@ua.pt](mailto:isabel.cacao@ua.pt)  
[hrmalon@ua.pt](mailto:hrmalon@ua.pt)

M. Irene Falcão  
CMAT and Department of Mathematics  
University of Minho  
4710-057 Braga  
Portugal  
[mif@math.uminho.pt](mailto:mif@math.uminho.pt)

Graça Tomaz  
CIDMA and Department of Mathematics  
Polytechnic of Guarda  
6300-659 Guarda  
Portugal  
[gtomaz@ipg.pt](mailto:gtomaz@ipg.pt)

## Abstract

Several authors are currently working on generalized Appell polynomials and their applications in the framework of hypercomplex function theory in  $\mathbb{R}^{n+1}$ . A few years ago, two of the authors of this paper introduced a prototype of these generalized Appell polynomials, which heavily draws on a one-parameter family of non-symmetric number triangles  $\mathcal{T}(n)$ ,  $n \geq 2$ . In this paper, we prove several new and interesting properties of finite and infinite sums constructed from entries of  $\mathcal{T}(n)$ , similar to the ordinary Pascal triangle, which is not a part of that family. In particular, we obtain a recurrence relation for a family of finite sums, analogous to the ordinary Fibonacci sequence, and derive its corresponding generating function.

## 1 Introduction

Almost ten years ago, Falcão and Malonek [7] proved a set of basic properties of a one-parameter family of non-symmetric Pascal triangles. Such family, obtained as result of

studies on generalized Appell polynomials in the framework of hypercomplex function theory in  $\mathbb{R}^{n+1}$  [6], is given by the infinite array of rational numbers

$$T_s^k(n) = \binom{k}{s} \frac{\binom{n+1}{2}_{k-s} \binom{n-1}{2}_s}{(n)_k}, \quad n, k = 1, 2, \dots, \quad s = 0, 1, \dots, k, \quad (1)$$

where  $(a)_r := a(a+1) \cdots (a+r-1)$ , for any integer  $r \geq 1$  is the Pochhammer symbol with  $(a)_0 := 1, a \geq 0$ . If  $n = 1$ , then  $T_0^k(1) = 1$  and  $T_s^k(1) = 0, s > 0$ . We let  $\mathcal{T}(n), n \geq 2$ , denote the family of triangles whose elements are given by (1).

The paper is organized as follows: first we recall several results related to interesting properties of  $\mathcal{T}(n)$ , specifically those relating to sums of row entries and series along its main diagonal. Next, we highlight the significant role played by sums over entries in the anti-diagonals of  $\mathcal{T}(n)$ , which are similar to those found in the Fibonacci sequence within the ordinary Pascal triangle. Naturally, the interconnections between all entries of  $\mathcal{T}(n)$  and those along the main diagonal rely on several well-known and lesser-known combinatorial identities, which we use extensively throughout the paper. The final result of the paper is some surprising revelation about a property of such an analog of Fibonacci sequence for  $\mathcal{T}(n)$ .

## 2 Preliminary results

We recall here the most relevant properties of the family  $\mathcal{T}(n)$ . Relations between adjacent elements of  $\mathcal{T}(n)$  were obtained by Falcão and Malonek [7], while Cação et al [4] derived results concerning the main diagonal elements of the triangle, i.e., the sequence of numbers

$$\mathcal{T}_k(n) := T_k^k(n) = \frac{\binom{n-1}{2}_k}{(n)_k}, \quad k = 0, 1, 2, \dots, \quad n = 2, 3, \dots \quad (2)$$

The first result [4, Proposition 8] shows the relation between any element  $T_s^k(n)$  of the triangle  $\mathcal{T}(n)$  and the main diagonal elements  $\mathcal{T}_m(n), m = s, s+1, \dots, k$

**Theorem 1.** *For  $k = 0, 1, 2, \dots$  and  $r = 0, \dots, k$ , we have*

$$T_{k-r}^k(n) = (-1)^r \binom{k}{r} \sum_{s=0}^r \binom{r}{s} (-1)^s \mathcal{T}_{k-s}(n). \quad (3)$$

The next result [4, Proposition 11] concerns a recurrence satisfied by the sequence  $(D_k(n))_{k \geq 0}$  consisting of alternating partial sums of the main diagonal elements, i.e.,

$$D_k(n) := \sum_{s=0}^k (-1)^s \mathcal{T}_s(n), \quad (4)$$

whose first elements are

$$1, \frac{n+1}{2n}, \frac{3n+1}{4n}, \frac{(n+1)(5n+7)}{8n(n+2)}, \frac{n(11n+28)+9}{16n(n+2)}, \frac{(n+1)(7n(3n+16)+107)}{32n(n+2)(n+4)}, \frac{n(n(43n+281)+485)+151}{64n(n+2)(n+4)}, \dots$$

**Theorem 2.** *The sequence  $(D_k(n))_{k \geq 0}$ , satisfies the recurrence relation*

$$(n+1)D_{k+1}(n) - 2(k+n+1)D_{k+2}(n) + (2k+n+1)D_k(n) = 0, \quad (5)$$

*with initial conditions*

$$D_0(n) = 1, \quad D_1(n) = \frac{n+1}{2n}.$$

We end this section by observing that the triangular array  $T_s^k(n)$  can be easily written as a scaled integer triangle, for some values of  $n$ . In fact, it is easy to show, using the well-known properties of the Pochhammer symbol, that

$$T_s^k(2n+1) = \frac{\binom{n+k-s}{n} \binom{n+s-1}{n-1}}{\binom{k+2n}{2n}}.$$

Therefore, for odd values of  $n$ , the entries of the triangle  $\binom{k+n-1}{n-1} T_s^k(n)$  are integers, as illustrated in Table 1.

Triangles	First rows	OEIS link
$\binom{k+2}{2} T_s^k(3)$	$\begin{array}{cccc} 1 & & & \\ 2 & 1 & & \\ 3 & 2 & 1 & \\ 4 & 3 & 2 & 1 \end{array}$	triangle <a href="#">A004736</a>
$\binom{k+4}{4} T_s^k(5)$	$\begin{array}{cccc} 1 & & & \\ 3 & 2 & & \\ 6 & 6 & 3 & \\ 10 & 12 & 9 & 4 \end{array}$	triangle <a href="#">A104633</a>
$\binom{k+6}{6} T_s^k(7)$	$\begin{array}{cccc} 1 & & & \\ 4 & 3 & & \\ 10 & 12 & 6 & \\ 20 & 30 & 24 & 10 \end{array}$	triangle <a href="#">A103252</a>

Table 1: Some particular scaled triangles obtained from  $\mathcal{T}(n)$ , for odd  $n$ .

When  $n$  is even, determining the scale factor becomes more challenging (Cação et al. [3] provide further details). Table 2 illustrates the particular cases  $n = 2$  and  $n = 4$ , as well as the limiting case  $n = \infty$ .

### 3 Filling in the gap: Fibonacci-like sequences

The study conducted by Falcão and Malonek [7] did not take into consideration the relationships among the entries located along the rising diagonals of  $\mathcal{T}(n)$  (also called northeast diagonals [10] or anti-diagonals [2]). The aim of this paper is to fill in this gap with a surprising result presented in Theorem 3. From the qualitative point of view, it shows once

Triangles	First rows	OEIS link
$2^{m_k}T_s^k(2)$	$\begin{array}{cccc} 1 & & & \\ 3 & 1 & & \\ 5 & 2 & 1 & \\ 35 & 15 & 9 & 5 \end{array}$	$m_k = $ <a href="#">A283208</a>
$2^{M_k}T_s^k(4)$	$\begin{array}{cccc} 1 & & & \\ 5 & 3 & & \\ 7 & 6 & 3 & \\ 21 & 21 & 15 & 7 \end{array}$	$M_{k-1} = m_{k-1} + $ <a href="#">A050605</a>
$2^kT_s^k(\infty)$	$\begin{array}{cccc} 1 & & & \\ 1 & 1 & & \\ 1 & 2 & 1 & \\ 1 & 3 & 3 & 1 \end{array}$	Pascal triangle <a href="#">A007318</a>

Table 2: Other integer triangles obtained by scaling  $\mathcal{T}(n)$ .

more the particular nature of  $\mathcal{T}(n)$  as a combinatorial object arising from generalized Appell polynomials in hypercomplex analysis.

Consider the triangle  $\mathcal{L}(n)$  obtained after re-indexing of the triangle  $\mathcal{T}(n)$  as illustrated in Figure 1. The elements  $L_s^k(n)$  of  $\mathcal{L}(n)$  are given by

$$L_s^k(n) := T_s^{k-s}(n), \quad k = 0, 1, 2, \dots, \quad s = 0, 1, \dots, \lfloor \frac{k}{2} \rfloor, \quad n = 2, 3, \dots \quad (6)$$

Table 3 shows the first few lines of the array  $\mathcal{L}(n)$ , where each line has pairwise repeating lengths. In the limiting case  $n = \infty$ , one can recognize well-known scaled triangles, as illustrated in Table 4.

One of the most interesting patterns of the ordinary Pascal triangle is that the sum of the elements of its anti-diagonals gives rise to the Fibonacci sequence. Following this idea, we construct the sequence  $(S_k(n))_{k \geq 0}$

$$S_k(n) := \sum_{s=0}^{\lfloor \frac{k}{2} \rfloor} L_s^k(n), \quad (7)$$

consisting of the sum of the elements  $L_s^k(n)$  in the  $k$ -th rising diagonal of the array  $\mathcal{T}(n)$ .

The next result shows that the sequence  $(S_k(n))_{k \geq 0}$ , built according to the construction of the Fibonacci sequence from the elements of the ordinary Pascal triangle, is identical to the sequence  $(D_k(n))_{k \geq 0}$ , built of the alternating partial sums of the main diagonal elements of the triangle  $\mathcal{T}(n)$ .

**Theorem 3.** *The sequence  $(S_k(n))_{k \geq 0}$  given by (7) is identical to the sequence  $(D_k(n))_{k \geq 0}$ , given by (4).*




Figure 1: A re-indexing of the triangle  $\mathcal{T}(n)$ .

*Proof.* We first assume that  $k$  is even, i.e.,  $k = 2m$  ( $m \in \mathbb{N}_0$ ). Then, from (6) and (7) we have

$$S_{2m}(n) = \sum_{s=0}^m T_s^{2m-s} = \sum_{\ell=0}^m T_{m-\ell}^{m+\ell}.$$

We can use (3) with  $k = m + \ell$  and  $r = 2\ell$  to obtain

$$\begin{aligned} S_{2m}(n) &= \sum_{\ell=0}^m \binom{m+\ell}{2\ell} \sum_{s=0}^{2\ell} (-1)^s \binom{2\ell}{s} \mathcal{T}_{m+\ell-s}(n) \\ &= \sum_{\ell=0}^m \binom{m+\ell}{2\ell} \sum_{s=-\ell}^{\ell} (-1)^{\ell-s} \binom{2\ell}{\ell-s} \mathcal{T}_{m+s}(n). \end{aligned}$$

Reversing the order of summation, we get

$$S_{2m}(n) = \sum_{s=-m}^m (-1)^s \mathcal{T}_{m+s}(n) \Sigma_{s,m}$$

where

$$\Sigma_{s,m} := \sum_{\ell=|s|}^m (-1)^\ell \binom{m+\ell}{2\ell} \binom{2\ell}{\ell-s}.$$

$L_s^k(n)$	$s = 0$	$s = 1$	$s = 2$	$s = 3$	$\dots$
$k = 0$	1				
$k = 1$	$\frac{n+1}{2n}$				
$k = 2$	$\frac{n+3}{2^2 n}$	$\frac{n-1}{2n}$			
$k = 3$	$\frac{(n+3)(n+5)}{2^3 n(n+2)}$	$\frac{2(n-1)}{2^2 n}$			
$k = 4$	$\frac{(n+5)(n+7)}{2^4 n(n+2)}$	$\frac{3(n-1)(n+3)}{2^3 n(n+2)}$	$\frac{n-1}{2^2 n}$		
$k = 5$	$\frac{(n+5)(n+7)(n+9)}{2^5 n(n+2)(n+4)}$	$\frac{4(n-1)(n+5)}{2^4 n(n+2)}$	$\frac{3(n^2-1)}{2^3 n(n+2)}$		
$k = 6$	$\frac{(n+7)(n+9)(n+11)}{2^6 n(n+2)(n+4)}$	$\frac{5(n-1)(n+5)(n+7)}{2^5 n(n+2)(n+4)}$	$\frac{6(n^2-1)}{2^4 n(n+2)}$	$\frac{(n-1)(n+3)}{2^3 n(n+2)}$	
$k = 7$	$\frac{(n+7)(n+9)(n+11)(n+13)}{2^7 n(n+2)(n+4)(n+6)}$	$\frac{6(n-1)(n+7)(n+9)}{2^6 n(n+2)(n+4)}$	$\frac{10(n-1)(n+1)(n+5)}{2^5 n(n+2)(n+4)}$	$\frac{4(n^2-1)}{2^4 n(n+2)}$	
$\dots$					

Table 3: Values of  $L_s^k(n)$  in the first rising diagonals of the array  $\mathcal{T}(n)$ .

Triangles	First rows	OEIS link
$2^k L_s^k(\infty)$	1	<a href="#">A128099</a> Row sums are the Jacobsthal numbers <a href="#">A001045</a>
	1	
	1 2	
	1 4	
	1 6 4	
	1 8 12	
	1 10 24 8	
	1 12 40 32	
	1 14 60 80 16	
$2^{k-s} L_s^k(\infty)$	1	<a href="#">A011973</a> triangle of coefficients of (one version of) Fibonacci polynomials
	1	
	1 1	
	1 2	
	1 3 1	
	1 4 3	
	1 5 6 1	
	1 6 10 4	
	1 7 15 10 1	

Table 4: Some particular scaled triangles obtained from  $\mathcal{L}(\infty)$ .

Using the formula (cf. [8, Formula 5.21]) and replacing  $(r, m, k)$  by  $(m + \ell, 2\ell, \ell - s)$  we can

express  $\Sigma_{s,m}$  as

$$\Sigma_{s,m} = \sum_{\ell=|s|}^m (-1)^\ell \binom{m+\ell}{\ell-s} \binom{m+s}{\ell+s} = \sum_{\ell=|s|}^m (-1)^\ell \binom{m+s}{s+\ell} \binom{m+\ell}{m+s}.$$

We now use the relation

$$\sum_k (-1)^k \binom{\ell}{m+k} \binom{s+k}{n} = (-1)^{\ell+m} \binom{s-m}{n-\ell}, \quad \ell \in \mathbb{N}_0, \quad m, n \in \mathbb{Z}$$

(cf. [8, Formula 5.24]) with  $(k, \ell, m, n, s)$  replaced by  $(\ell, m+s, s, m+s, m)$  to conclude that

$$\Sigma_{s,m} = (-1)^m,$$

which leads to

$$S_{2m}(n) = \sum_{s=-m}^m (-1)^{m+s} \mathcal{T}_{m+s}(n) = \sum_{s=0}^{2m} (-1)^s \mathcal{T}_s(n) = D_{2m}.$$

The case where  $k$  is odd can be handled using similar arguments.  $\square$

By combining the relation (5) with the coincidence of both sequences  $(S_k(n))_{k \geq 0}$  and  $(D_k(n))_{k \geq 0}$ , we get immediately that the sequence  $(S_k(n))_{k \geq 0}$ , considered as the analog(7) of the Fibonacci sequence, can be characterized by a second-order recurrence with variable coefficients as follows:

**Corollary 4.** *For any integer  $n \geq 2$  the elements  $S_k(n)$  are recursively defined by*

$$S_{k+1}(n) = \frac{n+1}{2(k+n)} S_k(n) + \frac{2k+n-1}{2(k+n)} S_{k-1}(n), \quad \text{for } k \geq 1, \quad (8)$$

with initial conditions

$$S_0(n) = 1, \quad S_1(n) = \frac{n+1}{2n}. \quad (9)$$

The recurrence (8), which connects three consecutive terms of the sequence  $(S_k(n))_{k \geq 0}$ , enables the derivation of a differential equation, whose solution is the ordinary power series generating function of that sequence.

**Theorem 5.** *The generating function of the sequence  $(S_k(n))_{k \geq 0}$ ,  $n = 2, 3, \dots$  can be written in terms of the Gauss hypergeometric function as*

$$F_n(x) = {}_2F_1(1, 1; 1; x) {}_2F_1(1, \frac{n-1}{2}; n; -x) = \frac{{}_2F_1(1, \frac{n-1}{2}; n; -x)}{1-x}.$$

*Proof.* Let  $F_n$  be the ordinary power series generating function of the sequence  $(S_k(n))_{k \geq 0}$ , i.e.,

$$F_n(x) = \sum_{k=0}^{\infty} S_k(n)x^k.$$

We rewrite the recurrence relation (8) in the following form:

$$(3 - n - 2k)S_{k-2}(n) - (1 + n)S_{k-1}(n) + 2(n - 1)S_k(n) + 2kS_k(n) = 0.$$

Then, we multiply by  $x^k$  and sum over  $k \geq 2$  to obtain

$$\begin{aligned} (3 - n) \sum_{k=2}^{\infty} S_{k-2}(n)x^k - 2 \sum_{k=2}^{\infty} kS_{k-2}(n)x^k \\ - (1 + n) \sum_{k=2}^{\infty} S_{k-1}(n)x^k + 2(n - 1) \sum_{k=2}^{\infty} S_k(n)x^k + 2 \sum_{k=2}^{\infty} kS_k(n)x^k = 0, \end{aligned}$$

or, equivalently,

$$\begin{aligned} (3 - n)x^2F_n(x) - 2x^3F_n'(x) - 4x^2F_n(x) - (1 + n)(xF_n(x) + S_0(n)) \\ + 2(n - 1)(F_n(x) - S_1(n)x - S_0(n)) + 2(xF_n'(x) - xS_1(n)) = 0. \end{aligned}$$

By using the initial conditions (9) in the above equation, we get the first order linear differential equation

$$F_n'(x) + \frac{1}{x-1} \left( \frac{n+1}{2} - \frac{n-1}{x^2+x} \right) F_n(x) = \frac{n-1}{x-x^3}. \quad (10)$$

The use of the standard integrating factor method for solving (10) leads to

$$F_n(x) = \frac{(n-1)(1+x)^{\frac{n-1}{2}}x^{1-n}}{1-x} \int x^{n-2}(1+x)^{-\frac{n+1}{2}} dx. \quad (11)$$

Since  $(1+x)^{-a} = {}_2F_1(1, a; 1; -x)$  (cf. [1, Formula 15.1.8]), we have

$$\begin{aligned} \int x^{n-2}(1+x)^{-\frac{n+1}{2}} dx &= \int x^{n-2} \sum_{k=0}^{\infty} \binom{n+1}{k} \frac{(-x)^k}{k!} \\ &= \frac{x^{n-1}}{n-1} \sum_{k=0}^{\infty} \frac{(n-1)_k \left(\frac{n+1}{2}\right)_k}{(n)_k} \frac{(-x)^k}{k!} \\ &= \frac{x^{n-1}}{n-1} {}_2F_1\left(n-1, \frac{n+1}{2}; n; -x\right). \end{aligned} \quad (12)$$

Combining (11) and (12) we get

$$F_n(x) = \frac{(1+x)^{\frac{n-1}{2}} {}_2F_1\left(n-1, \frac{n+1}{2}; n; -x\right)}{1-x}.$$



The result follows by applying Euler's transformation

$${}_2F_1(a, b; c; -x) = (1+x)^{c-a-b} {}_2F_1(c-a, c-b; c; -x).$$

□

We observe that the ODE (10) can be written as

$$\frac{2}{n-1}F'_n(x) + \frac{1}{x-1}\left(\frac{n+1}{n-1} - \frac{2}{x^2+x}\right)F_n(x) = \frac{2}{x-x^3},$$

which gives in the limit case  $n = \infty$ ,

$$F_\infty(x) = -\frac{2}{x^2+x-2}.$$

Notice that  $F_\infty(x)$  is also the generating function of the sequence  $(2^{-k}J_{k+1})_{k \geq 0}$ , where  $J_k$  denotes Jacobsthal numbers. In fact,

$$F_\infty(x) = \frac{2}{3}\left(\frac{1}{1-x} + \frac{1}{2+x}\right) = \sum_{k=0}^{\infty} \frac{1}{3}\left(2 + \left(-\frac{1}{2}\right)^k\right)x^k,$$

and taking into account the Binet form of the Jacobsthal numbers  $J_k = \frac{1}{3}(2^k - (-1)^k)$  (cf. [10]), we obtain

$$F_\infty(x) = \sum_{k=0}^{\infty} 2^{-k}J_{k+1}x^k,$$

i.e.,  $S_k(\infty) = 2^{-k}J_{k+1}$ . Furthermore, we immediately conclude that

$$\lim_{k \rightarrow \infty} S_k(\infty) = \frac{2}{3}.$$

We can obtain explicit expressions for  $F_n(x)$  for certain values of  $n$ , by using properties of the Gauss hypergeometric function.

- $F_2(x) = \frac{2(-1 + \sqrt{1+x})}{(1-x)x}$
- $F_3(x) = \frac{-2x + 2(1+x)\log(1+x)}{(1-x)x^2}$
- $F_4(x) = \frac{2(8 - 8\sqrt{1+x} + x(12 + 3x - 8\sqrt{1+x}))}{(1-x)x^3}$
- $F_5(x) = \frac{-2x(6 + x(9 + 2x)) + 12(1+x)^2\log(1+x)}{(-1+x)x^4}$

Finally, we refer to some limiting properties of  $S_k(n)$ .

**Theorem 6.** *Consider the sequence  $(S_k(n))_{k \geq 0}$ ,  $n = 2, 3, \dots$ . Then*

$$\lim_{k \rightarrow \infty} S_k(n) = {}_2F_1\left(1, \frac{n-1}{2}; n; -1\right).$$

*Proof.* The result is an immediate consequence of the Theorem 3, along with equations (4) and (2). In fact,

$$S_k(n) = D_k(n) = \sum_{s=0}^k (-1)^s \mathcal{T}_s(n) = \sum_{s=0}^k (-1)^s \frac{\left(\frac{n-1}{2}\right)_s}{(n)_s} = \sum_{s=0}^k \frac{(-1)^s (1)_s \left(\frac{n-1}{2}\right)_s}{s! (n)_s},$$

which proves the desired result.

It is worth noting that the hypergeometric series  ${}_2F_1(1, \frac{n-1}{2}; n; -1)$  converges absolutely [1], since  $n - 1 - \frac{n-1}{2} > 0$ .  $\square$

It is well known that the ratio of two consecutive ordinary Fibonacci numbers converges to the golden ratio. Here the corresponding property can be obtained as an immediate consequence of Theorem 6. Specifically, we have

$$\lim_{k \rightarrow \infty} \frac{S_{k+1}(n)}{S_k(n)} = 1.$$

## 4 Final remarks

It was initially unclear what kind of result could be expected from constructing an analog of the Fibonacci sequence using hypercomplex tools and following the usual rules of the ordinary Pascal triangle. Specifically, we were uncertain about the result of summing the rising diagonal elements of  $\mathcal{T}(n)$ . This gap was left unaddressed in our previous work [7]. However, as explained in Section 3, Theorem 3 provides the answer to this question and leads to the main recurrence relation given in (8)-(9).

Concluding our final remarks, we would like to highlight the potential for fruitful and interesting connections between real, complex, and hypercomplex analysis when considering hypercomplex polynomials [5, 9, 11] from the specific discrete viewpoint of combinatorial relations.

## 5 Acknowledgments

This work was supported by Portuguese funds through the CMAT - Research Centre of Mathematics of University of Minho - and through the CIDMA - Center of Research and Development in Mathematics and Applications (University of Aveiro) and the Portuguese Foundation for Science and Technology (“FCT - Fundação para a Ciência e Tecnologia”), within projects UIDB/00013/2020, UIDP/00013/2020, UIDB/04106/2020 and UIDP/04106/2020.

## References

- [1] M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, US Government Printing Office, 10th printing, 1972.
- [2] L. Abrams, D. E. Fishking, and S. Valdes-Leon, Reflecting the Pascal matrix about its main antidiagonal, *Linear Multilinear Algebra* **47** (2000), 129–136.
- [3] I. Cação, M. I. Falcão, and H. R. Malonek, Hypercomplex polynomials, Vietoris’ rational numbers and a related integer numbers sequence, *Complex Anal. Oper. Theory* **11** (2017), 1059–1076.
- [4] I. Cação, M. I. Falcão, H. R. Malonek, and G. Tomaz, Non-symmetric number triangles arising from hypercomplex function theory in  $\mathbb{R}^{n+1}$ , in O. Gervasi, B. Murgante, et al., eds., *Computational Science and Its Applications—ICCSA 2022 Workshops*, Lect. Notes in Comp. Sci., Vol. 13377, Springer Cham, 2022, pp. 420–434.
- [5] R. Delanghe, F. Sommen, and V. Souček, Clifford algebra and spinor-valued functions: a function theory for the Dirac operator, *Mathematics and its Applications*, Vol. 53, Kluwer Academic, 1992.
- [6] M. I. Falcão, J. Cruz, and H. R. Malonek, Remarks on the generation of monogenic functions, in K. Gürlebeck and C. Könke, eds., *17th Inter. Conf. on the Appl. of Computer Science and Mathematics on Architecture and Civil Engineering*, Germany, 2006. Available at [https://e-pub.uni-weimar.de/opus4/files/2939/f81\\_pdfa.pdf](https://e-pub.uni-weimar.de/opus4/files/2939/f81_pdfa.pdf).
- [7] M. I. Falcão and H. R. Malonek, A note on a one-parameter family of non-symmetric number triangles, *Opuscula Math.* **32** (2012), 661–673.
- [8] R. L. Graham, D. E. Knuth, and O. Patashnik, *Concrete Mathematics: A Foundation for Computer Science*, Addison-Wesley Longman Publishing Co., Inc., 1994.
- [9] K. Gürlebeck, K. Habetha, and W. Sprößig, *Application of Holomorphic Functions in Two or Higher Dimensions*, Birkhäuser Verlag, 2016.
- [10] T. Koshy, *Fibonacci and Lucas Numbers with Applications*, Pure and Applied Mathematics: A Wiley Series of Texts, Monographs and Tracts, John Wiley & Sons, 2019.
- [11] H. R. Malonek, I. Cação, M. I. Falcão, and G. Tomaz, Harmonic analysis and hypercomplex function theory in co-dimension one, in A. Karapetyants, V. Kravchenko, and E. Lifyand, eds., *Modern Methods in Operator Theory and Harmonic Analysis—OTHA 2018*, Springer Proc. Math. Stat., Vol. 291, Springer Cham, 2019, pp. 93–115.

2010 *Mathematics Subject Classification*: Primary 11B37; Secondary 11B83, 11B39, 05A19, 33C05.

*Keywords*: *Fibonacci sequence, recurrence relation, hypercomplex function theory, hypergeometric function.*

---

(Concerned with sequences [A001045](#), [A004736](#), [A007318](#), [A011973](#), [A050605](#), [A103252](#), [A104633](#), [A128099](#), and [A283208](#).)

---