



University of Dundee

Evolutionary, structural and functional insights in nuclear organisation and nucleocytoplasmic transport in trypanosomes

Padilla-Mejia, Norma E.; Field, Mark C.

Published in:
FEBS Letters

DOI:
[10.1002/1873-3468.14747](https://doi.org/10.1002/1873-3468.14747)

Publication date:
2023

Licence:
CC BY

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Padilla-Mejia, N. E., & Field, M. C. (2023). Evolutionary, structural and functional insights in nuclear organisation and nucleocytoplasmic transport in trypanosomes. *FEBS Letters*. Advance online publication. <https://doi.org/10.1002/1873-3468.14747>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

REVIEW

Evolutionary, structural and functional insights in nuclear organisation and nucleocytoplasmic transport in trypanosomes

 Norma E. Padilla-Mejia¹  and Mark C. Field^{1,2} 

1 School of Life Sciences, University of Dundee, UK

2 Institute of Parasitology, Biology Centre, Czech Academy of Sciences, České Budějovice, Czechia

Correspondence

 N. E. Padilla-Mejia, School of Life Sciences,
 University of Dundee, Dundee DD1 5EH,
 UK

Tel: +44 01382-386255

 E-mail: nepadillamejia@dundee.ac.uk

 (Received 27 June 2023, revised 18
 September 2023, accepted 19 September
 2023, available online 15 October 2023)

doi:10.1002/1873-3468.14747

Edited by Nathalie Arhel

One of the remarkable features of eukaryotes is the nucleus, delimited by the nuclear envelope (NE), a complex structure and home to the nuclear lamina and nuclear pore complex (NPC). For decades, these structures were believed to be mainly architectural elements and, in the case of the NPC, simply facilitating nucleocytoplasmic trafficking. More recently, the critical roles of the lamina, NPC and other NE constituents in genome organisation, maintaining chromosomal domains and regulating gene expression have been recognised. Importantly, mutations in genes encoding lamina and NPC components lead to pathogenesis in humans, while pathogenic protozoa disrupt the progression of normal development and expression of pathogenesis-related genes. Here, we review features of the lamina and NPC across eukaryotes and discuss how these elements are structured in trypanosomes, protozoa of high medical and veterinary importance, highlighting lineage-specific and conserved aspects of nuclear organisation.

Keywords: evolutionary diversity; nuclear lamina; nuclear pore complex; nucleus; trypanosoma

Protozoan parasites cause a plethora of human, animal and plant diseases, many of which are fatal without intervention and have profound impact on economic activity and quality of life. The class Kinetoplastida (clade Discoba) [1] are flagellated protozoa possessing

a kinetoplast, an unusual configuration of mitochondrial DNA [2,3]. The Kinetoplastida include *Trypanosoma* and *Leishmania* species, causative agents of multiple diseases, with particular importance to tropical and subtropical countries, although outbreaks in

Abbreviations

BES, bloodstream expression site; BSF, bloodstream form; CF, cytoplasmic filaments; Cr, *Chlamydomonas reinhardtii*; CRWN, crowded nuclei protein; Dbp5, RNA helicase DEAD box protein 5; Esc1, establishes silent chromatin 1 (protein); FG-Nup, FG-repeats nucleoporin; FG-repeat, phenylalanine-glycine repeat; Gle1/2, mRNA export factor Gle1 or Gle2; Gp210, nuclear pore membrane glycoprotein 210; Hs, *Homo sapiens*; HAT, human African trypanosomiasis; IR, inner ring; LAD, lamin-associated domains; LECA, last eukaryotic common ancestor; Mex67, mRNA export factor 67; Mlp1/2, myosin-like protein 1 or 2; MR, *trans*-membrane scaffolding ring; mRNA, messenger RNA; mRNP, messenger ribonucleoprotein; Mtr2, mRNA transport regulator; NB, nuclear basket; Ndc1, nuclear division cycle protein 1 (nucleoporin); NE, nuclear envelope; NE81, 81 kDa nuclear envelope protein; Nic96, 96 kDa nucleoporin-interacting component; NLS, nuclear localization signal; NMCP, nuclear matrix constituent protein; NPC, nuclear pore complex; Nsp1, nucleoskeletal-like protein 1; Nup, nucleoporin; NUP-1, nucleoporin 1; NUP-2, nucleoporin 2; Nxf1, nuclear RNA export factor 1; Nxt1, NTF-2 related export protein 1; OR, outer ring; PCF, procyclic culture form; Pf, *Plasmodium falciparum*; POM, pore membrane protein; RBP10, RNA binding protein 10; Sc, *Saccharomyces cerevisiae*; Sec13, transport protein Sec13; Seh1, nucleoporin Seh1 (Sec13 homologue 1); Tb, *Trypanosoma brucei*; Tg, *Toxoplasma gondii*; Tpr, translocated promoter region protein; TREX1/2, three-prime repair exonuclease 1 or 2; tRNA, transfer RNA; TY1, epitope tag TY1; VSG, variant surface glycoprotein..

North America, Europe (especially the Mediterranean belt) and Oceania are occurring with increased frequency [4,5]. Human African Trypanosomiasis (HAT), or sleeping sickness, is caused by *Trypanosoma brucei gambiense* and *Trypanosoma brucei rhodesiense* and is fatal if untreated [5]. Veterinary infections, with a spectrum of severity and clinical symptoms, are caused by *T. evansi*, *T. equiperdum*, *T. congolense* and *T. vivax* and are distributed across Africa, Asia and Latin America; death of cattle and domestic animals due to trypanosomiasis causes huge economic loss in these regions [4,6]. *T. b. gambiense* and *T. b. rhodesiense* are transmitted by over 20 species of tsetse flies infected *via* blood feeding. The life cycle of *T. brucei* includes up to seven different forms developing in the mammal host and the tsetse fly [7], with the bloodstream form (BSF, mammal host) and the procyclic culture form (PCF, insect host) being important experimental models as they can be cultured *in vitro* and are amenable to cell and molecular biology techniques.

Trypanosoma species are characterised by an elongated cell body of 10–20 µm propelled by a single flagellum [8]. Differentiation in these protozoa is a complex process involving a series of physiological and biochemical changes, one of the most distinctive being the replacement of the surface coat of variant surface glycoprotein (VSG) in the BSF to procyclin in the PCF [7]. VSG is expressed at a very high level and switches between paralogous genes with high frequency to facilitate immune evasion *via* antigenic variation [9,10]. VSG genes are transcribed from bloodstream expression sites (BESs) located at subtelomeric regions. Only a single BES is active and this monoallelic expression is central for immune evasion [11]. Moreover, one of the striking features of *T. brucei* is the unusual transcription of VSG and procyclin by RNA Polymerase I [12].

Drugs have been developed against all human trypanosome pathogens, *albeit* with differing levels of efficacy due to toxicity, drug resistance and other factors [13,14]. When drugs are combined with efficient local healthcare, the prognosis is very good and for HAT has achieved spectacular progress; disease occurrence reached a historic prevalence of under 1000 cases in 2018, remaining below that during 2022, and declared effectively eliminated from multiple countries [15]. However, in other cases, such as the American trypanosome, *Trypanosoma cruzi*, the causative agent of Chagas disease, the impact on human health remains severe, affecting 6–8 million people in the American continent, with up to 100 million people at risk of infection [16,17]. Another example of kinetoplastid

is *Leishmania* species, the causative agents of leishmaniasis, a disease with a range of clinical manifestations and severities and present in at least 88 countries with more than 350 million people at risk [18]. These two pathogens demand ongoing campaigns for disease control including characterisation of biochemical and genetic mechanisms of pathogenesis/virulence and the identification of new drug modalities [13,14].

The nucleus encloses genomic DNA within the nuclear envelope (NE), a double lipid bilayer associated with a filamentous meshwork or lamina residing at the inner face, providing mechanical support and contributing to multiple functions. Transport between the nucleus and cytoplasm occurs through the nuclear pore complex (NPC), a massive macromolecular gateway constructed from several hundred subunits and with involvement in regulating gene expression and other functions. In evolutionary biology, the last eukaryotic common ancestor, or LECA, is considered the first eukaryote and is minimally defined by having a mitochondrion, flagellum and a nucleus [19–21]. Additionally, the LECA also possessed a NE, NPCs and a nuclear lamina, with additional subcellular structures such as a cytoskeleton and membranous endosomal and exocytic compartments. Complex metabolism and gene regulation were also present in the LECA [19–21]. Widespread ultrastructural and genome features are likely to have been inherited from the LECA [20]. After the LECA, different lineages adapted their nuclei such that the nuclear lamina and NPC likely co-evolved with diverging mechanisms of gene expression, transcription and RNA processing. As the LECA bore NPCs and a nuclear lamina [19], it is interesting to consider how these crucial nuclear structures have diversified across eukaryotes.

Trypanosomes likely diverged from other eukaryotes over a billion years ago [22], and this deep distinction is reflected in multiple aspects of their biology, including their own nuclear structures that possess taxa-specific components [23]. Thus, apart from their medical relevance, they are a valuable model to gain insight into evolutionarily eukaryotic structures and processes. Here, we discuss the biology of the trypanosome nucleus, focusing on the nuclear lamina and NPC, emphasising the importance of these structures in the biology of these pathogenic protozoa. We will make comparisons *versus* the canonical models of mammalian cells and other recurrent biological models such as fungi and plants to highlight how trypanosomatid structures have evolved and diverged. Here, we take the nuclear lamina and NPC as fascinating examples of evolution and diversity across eukaryotes, but also linked to mechanisms of pathogenesis.

The nuclear lamina

At the inner face of the NE the nucleus is surrounded by the nuclear lamina, a filamentous meshwork that in metazoa is constituted of coiled-coil lamin proteins [24] (Fig. 1). The lamina is the major structural element of the nucleus, influencing nuclear shape, size and structural integrity [25]. Furthermore, lamina-mediated mechano-signalling allows the transduction of mechanical signals from the cytoskeleton to the nucleoplasm by direct connection of these structures through NE-located *trans*-membrane protein complexes. The lamina also interacts directly with the NPC and multiple NE proteins, influencing NE organisation [26].

Mammalian lamins are coiled-coil proteins of ~ 60–80 kDa, and belong to the intermediate filament family; they are evolutionary conserved across Metazoa [27]. There are four major lamin subtypes, encoded in humans by the *LMNA*, *LMNB1*, *LMNB2* genes to express lamin A and lamin C (generated by alternative splicing), lamin B1 and lamin B2, respectively. Lamin mutations cause heritable diseases termed laminopathies with a range of cardiac, muscular–skeletal and other issues and over 600 lamina mutations are linked to pathogenesis [28,29]. These proteins consist of a short globular N-terminal domain (head) followed by

a central α -coiled-coil domain (rod) and a globular C-terminal domain containing a lamin-tail domain incorporating an immunoglobulin fold, a nuclear localisation signal and a C-terminal CaaX prenylation signal [30] (Fig. 1). In B-type lamins the C-terminus is farnesylated at the cysteine residue, which increases the affinity for membranes and is necessary for lamin B localisation and network organisation [31]. Pre-lamin A is prenylated but the modification is cleaved and lost during maturation, while lamin C lacks the CaaX signal and is hence never prenylated [30]. Lamins form homodimers that associate with other dimers in a strict head-to-tail fashion to form filaments [32]. In mammalian nuclei, each lamin isoform forms a separate meshwork [33] that interconnects and interacts in a complex manner. These connections are codependent as depletions in one lamin isotype impact the equilibrium of the entire lamin meshwork and nuclear morphology [26,30,34–36]. Furthermore, B-type lamins are attached to the inner nuclear membrane, while A-type lamins face closer towards the nucleoplasm. Interestingly, in a further diversification of function, NPC components interact more closely with lamin C compared with lamin A [36].

Beyond structural roles, in mammalian cells, lamins participate in controlling genome architecture, DNA




Fig. 1. Comparison of major lamina protein architectures across eukaryotes. Main features are shown for experimentally characterised lamins; structural motifs/domains of relevance are shown on the right represented by different colours. Information taken from Uniprot [146], Trypsins database [65] and literature [49,52,53,147,148]. Number of aminoacids is indicated under each protein name. Only mature protein features are shown, pre-lamin A/C contains a CaaX box that is lost from the mature protein. Notice that all proteins are rich in coiled-coils domains (teal).




Fig. 2. Characterisation of lamins and NPCs in eukaryotic lineages. A phylogenetic tree showing the presence or absence of lamin or lamin-like proteins across different taxa. Major groups are indicated in the legend at the right (grey box) and representative species are included. Branches in the same colour correspond to the same super kingdom, e.g., Metazoa, Fungi and Apusozoa all correspond to Opisthokonta. LECA indicates the position of the LECA. The tree is based on Burki *et al.* [149]. Circles next to the taxa name indicates presence/absence of lamin proteins; black circle, mammalian (metazoan) lamins found; red circle, metazoan lamins detected *in silico* only without experimental characterisation; grey circle, metazoan lamins absent when using bioinformatic approaches; black empty circle, taxa-specific proteins found exhibiting lamina properties. More of one of these descriptions can apply for certain groups, e.g., TSAR, Haptophyta and Amoebozoa include species that, by bioinformatic analysis, show presence or absence of metazoan lamins, e.g., in Amoebozoa, some species have a mammalian lamin homologue (*Dictyostelium discoideum*), while others lack those (*Acanthamoeba castellanii*). Divergent taxa-specific lamin proteins with lamina properties have been characterised in Discoba and Streptophyta, which lack canonical lamins. Presence/absence of lamins is based on published analysis [27,46]. Names of organisms in which the NPC has been fully or partially characterised are in bold.

replication and transcription by interactions with chromatin (via lamin-associated domains, LADs), chromatin-modifying enzymes and transcription factors [37,38]. Lamins A/C can shuttle to the nucleoplasm where they bind and regulate euchromatic regions and interact with transcription and cell cycle regulators [39]. In mammalian cells, lamins are phosphorylated differentially during the cell cycle, i.e., some residues are phosphorylated at the onset of mitosis with others modified during interphase and mitosis [40]. Phosphorylated lamins regulate gene expression through enhancer binding [41], expanding the roles of phosphorylated lamins during interphase. Importantly, lamins support the mechanism of cell division, as they solubilise during open mitosis to allow full breakdown of the NE in a highly regulated process that includes serine phosphorylation that induces lamin solubility and disassembly [42]. However, this is not a universal feature as in many organisms that possess lamins or lamin analogues, mitosis is closed or semi-closed and

the NE does not break down, e.g., *T. brucei* [43] and *Dictyostelium discoideum* [44,45].

Lamins were initially considered Metazoa-restricted, but lamin orthologs have been identified in Rhizaria, Alveolata, Amoebozoa and other lineages (Fig. 2) [27,46]. Lamins are not universal and even closely related organisms in the same taxa may, or may not, have lamin genes, e.g., in Amoebozoa, the species *D. discoideum* and *Acanthamoeba castellanii* have and lack a lamin ortholog, respectively [27,46]. Some organisms lack proteins homologous to lamins but possess proteins that partially mimic lamin functions, e.g., Esc1 in *Saccharomyces cerevisiae*, a protein localising to the nuclear periphery that interacts with the NPC and has chromatin-silencing properties [46–48]. In spite of the *in silico* identification of divergent lamin orthologs across a wide range of eukaryotes (Fig. 2), few lamin and lamina-like systems are experimentally characterised, including NE81 in *D. discoideum* (Amoebozoa), an orthologue of lamin B [44,45];

NMCP1/2 in plants (CRWN in *Arabidopsis thaliana*) [49–51] and NUP-1/NUP-2 in *T. brucei* [52,53] (Figs 1 and 2). Therefore, defining a lamin analogue no longer resides with sequence alone but rather needs to consider structural and functional similarities, such as coiled-coil organisation, nuclear peripheral distribution, regulation of nuclear shape and size and interaction and regulation of chromatin.

As homologues of Metazoan lamins have been found in several lineages (Amoebozoa, Rhizaria, Stramenopila, Alveolata, Haptophyta [27,46,54]; Fig. 2), lamins must have been present very early in eukaryotic evolution, *circa* the LECA [27,46,54]. It has also been suggested that lamins are polyphyletic, i.e., they had multiple origins during eukaryote evolution [27]. However, as the gene and species trees from lamins are concordant, this seems unlikely [46]. It is unknown if different lamina systems (namely mammalian lamins, plant NMCPs or trypanosoma NUP-1/NUP-2) originated by evolutionary divergence or convergence as while they contain a conserved coiled-coil/ α -helical rod structure there is extreme divergence in size and sequence among them [27,54,55]. Importantly, the LECA can be considered a single cell or a population [56]. As a population with an available pool of genes [56], it is possible that the LECA possessed all three types of lamin proteins, lamin-like, NMCP-like and NUP-1/NUP-2-like, but that only one was retained in different organisms [54,55]. Alternatively, trypanosomes and plants may have replaced lamins with NUP-1/NUP-2 or NMCP, respectively [54,55]. The restriction of the latter two systems to single lineages would support the replacement model.

In African trypanosomes, NUP-1/NUP-2 are lamina analogs taxonomically restricted to Kinetoplasts [46,52,53]. As mentioned, NUP-1 (406 kDa) [52] and NUP-2 (170 kDa) [53] are coiled-coil proteins lacking lamin domains [46] such as the lamin tail domain or the CaaX motif (Fig. 1). Originally detected by subnuclear fractionation and immuno-electron microscopy at the nucleoplasmic face of the NE [57], NUP-1 has an N-terminal region followed by a central rod containing a region of 17 repeats of 144 amino acids and finally a C-terminal domain bearing an NLS [52] (Fig. 1). NUP-2 possesses a central monopartite NLS and coiled-coil regions in the N-terminal and C-terminal regions [53] (Fig. 1). In *T. cruzi*, two different NUP-1 isoforms of different sizes are apparently expressed from a single gene, however, the functional relevance of this is unexplored [58]. Remarkably, NUP-1 is seven times larger than mammalian lamins and consequently spans a more extensive volume of the smaller trypanosome nucleus (Fig. 3A). Confocal microscopy suggests

that NUP-1 has a predominantly extended conformation, observed by imaging an N-terminal tagged NUP-1 and the central repeats, with resolution between the two signals, suggesting an extended conformation [52].

NUP-1 localises to the nuclear periphery during interphase, while NUP-2 localises to the nuclear periphery and nucleoplasm (Fig. 3B) [52,53,59]. Both have roles in nuclear integrity as depletion results in nuclear blebbing, abnormal nuclear enlargement with irregular boundaries and aberrant extensions of the NE [52,53], a common phenotype of mutated lamins in other systems [60]. Furthermore, NUP-1/NUP-2 have contacts with the NPC, as evidenced by interactions with TbNup98 and other nucleoporins [59,61], suggesting that the NPC may facilitate anchoring at the NE, but if these interactions have additional functions besides simple structural support is unknown.

NUP-1 also influences telomere positioning [52], and both NUP-1/NUP-2 regulate gene expression as they participate in silencing VSG genes in the bloodstream form, contributing to monoallelic VSG expression [52,53]. Furthermore, NUP-1/NUP-2 participate in repression of procyclin, the major antigen expressed in the insect stage [52,53]. Hence, the trypanosome lamina influences the differentiation and expression of pathogenesis-related genes. This is also supported by overexpression of NUP-1 fragments, which leads to the upregulation of important RNA-binding proteins, including RBP10 [59], a master regulator of differentiation between procyclin and bloodstream forms [62].

Trypanosoma brucei presents a highly organised 3D genome architecture and chromosome territories, which influence DNA accessibility and thus, homologous recombination and gene expression [63]. This includes the subtelomeric regions containing VSG genes that are folded into compact domains [63], resulting in a genome-wide configuration that changes after VSG gene switching [64]. If NUP-1/NUP-2 can influence such genome architecture through interactions with chromatin is still unexplored.

The assembly and geometry of the trypanosome lamina is also distinct from mammalian lamins, which form homodimers assembling strictly head-to-tail (Fig. 3C) [32]. By contrast NUP-1 has the flexibility to assemble in a head-to-head, tail-to-tail or head-to-tail manner, forming homophilic or heterophilic interactions between NUP-1 domains [59] (Fig. 3D), essentially a hub-and-spoke architecture. As occurs with mammalian lamins, NUP-1 and NUP-2 are codependent in maintaining nuclear shape and integrity, as depletion of either impacts the other, and with the N-terminal region of NUP-1 acting as the main anchor point for NUP-2 [53,59]. During the cell cycle, distinct




Fig. 3. NUP-1 features in *Trypanosoma brucei*. (A) Schematic highlighting differences in size between mammalian (adipocyte) (~10 μm) and trypanosome nuclei (~2 μm). Corresponding lamina-constituting proteins are shown as a white line at approximate scale and molecular weights in kilodaltons. The cartoon only considers the elongated, primary structure of the proteins. Length only is significant, not the width of the line. The schematic is to emphasise that NUP-1/NUP-2 are huge proteins in a smaller nucleus when compared to equivalent human structures. As the sizes and sequences of the proteins are radically different, the inference is that the overall structure/assembly of the lamina in each case is likely to be distinct (see C and D). The potential volume that each lamina protein can access is clearly very different between mammals and trypanosomes. This representation is based on protein and nucleus size only and does not consider aspects such as higher-order assembly nor secondary/tertiary structure. (B) NUP-1 and NUP-2 detected at the nucleus by immunofluorescence assays. *T. brucei* BSF cells showing nucleus (N) and kinetoplast (k) (the mitochondrial DNA). Cells were probed with an α-NUP-1 antibody (red) (top panel) or α-NUP-2-TY1 (green) (bottom panel). Scale bar 5 μm. Microscopy images taken by N.E. Padilla-Mejia. (C) Assembly of mammalian lamins. Lamins exist as a homodimer that assemble first in a parallel manner and then in an antiparallel way to form filaments. (D) Assembly of NUP-1 molecules. NUP-1 can form interactions with other NUP-1 molecules in a head-to-head, tail-to-tail (homophilic interactions) or head-to-tail (heterophilic interactions).

NUP-1 domains locate to specific nuclear subregions during interphase, mitosis and cytokinesis [59], with the C-terminal region present in the bridge linking the daughter nuclei at late mitosis, while the N-terminal region is absent from this region [59]. The central repeat region of NUP-1 is present at the nuclear periphery and nucleoplasm, with this latter location predominant during early mitosis [59]. These observations reflect the flexibility and elastic properties of

NUP-1 and suggest that NUP-1 domains act to support specific functions and engage with the mitotic machinery. This may reflect the high-molecular weight of NUP-1, although in many kinetoplastids the ortholog is considerably smaller than the African trypanosome NUP-1 [52,65] (A. Makarov & M. C. Field, unpublished results). NUP-1 also has multiple potential phosphorylation sites [65,66] and preliminary data suggests that mutations in those sites lead to

aberrations in NUP-1 dynamics (A. Makarov, L. Koreny & M. C. Field, unpublished data). Moreover, mammalian lamins have additional post-translational modifications, such as sumoylation [67]; whether such modifications occur and are functionally relevant for NUP-1/NUP-2 remains to be explored.

In summary, the extended coiled-coil proteins NUP-1 and NUP-2 constitute major components of the trypanosoma lamina, despite no obvious sequence relationship to mammalian lamins. NUP-1/NUP-2 function similarly to Metazoan lamins, supporting roles in maintaining nuclear morphology, interactions with the NPC and chromatin, cell division, and regulation of gene expression, including genes related to development.

Conclusions and perspectives: nuclear lamina

One of the crucial scaffolds in the cell, the nuclear lamina is an exemplar for molecular evolution and diversity. The identity of the nuclear lamina in many eukaryotic clades is awaiting discovery and characterisation, including many protozoa of medical importance. Interestingly, although lamins, NMCPs and NUP-1/NUP-2 systems are different in sequence, the overall functions of the corresponding lamina are highly conserved, acting always as an architectural scaffold of the nuclear structure with a predominant role in chromatin and gene expression regulation. There are many unresolved questions, for example, how do NUP-1/NUP-2 influence genome architecture in trypanosomes and regulate expression of VSGs and procyclins? Moreover, as the primary structures of lamin analogues are divergent and taxa-specific, it remains to be uncovered what, if any, analogues are present in many pathogens, including the malaria parasite and its relatives, *Toxoplasma* and *Cryptosporidium*.

The nuclear pore complex

The NPC is a macromolecular channel allowing bidirectional trafficking between the nucleus and the cytoplasm. The NPC consists of ~30 different nucleoporins (Nups) with multiple copies of each, adding up a molecular mass of ~50 to 120 MDa in yeasts and mammals, respectively, due to stoichiometry and structural variation [68,69]. The NPC is built from multiple subcomplexes arranged in rings: (a) cytoplasmic filaments (CF); (b) outer rings (OR), one embedded in the cytoplasmic face of the NE and one nuclear ring at the inner face of the NE; (c) a *trans*-membrane scaffold ring (MR), an anchoring point for NPC

subcomplexes; (d) a central pore, that contains highly mobile phenylalanine-glycine (FG)-repeat proteins creating a permeability barrier regulating cargo transport and (e) the nuclear basket (NB) [68,70] (Fig. 4A). A substantial proportion of the NPC consists of FG-repeat proteins, intrinsically disordered proteins that interact with cargo and cargo receptors and are located at the central pore, the CF and the NB [68,70]. One of the hallmarks of NPC is an octagonal cylindrical structure. The NPC has an eight-fold rotational symmetry around the central transport channel (Fig. 4A). Due to its complex character, we consider the structure and function of the NPC separately. In the NPC structure section, we will refer constantly to architecture in humans and yeast as most structural characterisation comes from studies in these organisms. We will then compare with trypanosomes and other parasites.

NPC structure

High-resolution structures of NPCs from mammals, *S. cerevisiae*, *Xenopus laevis*, *D. discoideum* and *Chaetomium thermophilum* are available [71–82], while the protein components have been identified for *Schizosaccharomyces pombe* [83,84], *Chlamydomonas reinhardtii* [85], *A. thaliana* [86], *Malus domestica* [87], *Tetrahymena thermophila* [88] and *T. brucei* [61,89] (Fig. 2). These data have allowed evolutionary reconstruction [23,90], highlighting that overall morphology, organisation and architecture of NPC subcomplexes are conserved but also demonstrating that lineage-specific subunit stoichiometry is present within NPCs. Thus, NPC architectural diversity is produced through gains and losses of components together with varying stoichiometry.

The NPC contains two outer rings (ORs), one at the cytoplasmic face and one at the nucleoplasmic face (Fig. 4A). The major building block of the outer ring is the Y-complex (Fig. 4B) which owes its name to its ‘Y’ shape [72,74,79]. In metazoan NPCs (human and *Xenopus*), the cytoplasmic and nuclear outer rings are constituted by two staggered rings of eight Y-complexes (Fig. 4B). Eight of this Y-complex doublet forms a ring arrangement, i.e., 32 copies of the complex per NPC [80]. The Y-complex is a source of great variability in the NPC across the eukaryotes due to two features; (a) the number of Y-complex copies per NPC and (b) the number of nucleoporins constituting this complex (Fig. 4B,C) [78,80,82]. For example, in *S. cerevisiae*, these outer rings are composed by a single ring of eight Y-complexes, i.e., 16 copies [77] (Fig. 4B). In the green algae *C. reinhardtii*, the cytoplasmic ring contains a




Fig. 4. NPC features in eukaryotes. (A) Schematic of the NPC structure showing rings and components. Views from the NE plane, cytoplasm and nucleoplasm are presented. (B) Schematic showing how the outer rings (OR, burgundy) are configured in different organisms. The NPC in *Saccharomyces cerevisiae* has one OR at the cytoplasmic face of the NE and one OR at the nucleoplasmic face. The *Homo sapiens* NPC bears two ORs at each side while the algae *Chlamydomonas reinhardtii* has two ORs at the nucleoplasmic side and only one at the cytoplasmic side. Evidence points that *Trypanosoma brucei* has only two ORs but awaits confirmation. (C) Schematic showing localization of the Y-complex (burgundy) and comparison of its architecture in four different organisms; *S. cerevisiae*, *H. sapiens*, *C. reinhardtii* and *T. brucei*. Left, homologue subunits are depicted in the same burgundy colour, taxa-specific subunits in other colours. Right, Coulson plot showing the number and different homologues of the components of the Y-complex in the mentioned species. (D) Schematic highlighting the different composition of the nuclear basket in organisms in panel C. Figure shows the number and different homologues of the components of the NB in *S. cerevisiae*, humans, *C. reinhardtii* and *T. brucei*.

single ring of eight Y-complexes whereas the nucleoplasmic ring consists of 16 copies forming two rings [85] (Fig. 4B). A more divergent arrangement is present in *Schi. pombe*, in which the Y-complexes are of differing composition between the nuclear and cytoplasmic copies [23,73,83,84] (Fig. 4B). In *T. brucei*, evidence points to one ring of Y-complex per outer ring (Fig. 4B) [61] but this awaits confirmation. Variability in the Y-complex is also introduced by distinct component composition, with 6–10 nucleoporins present depending on species (Fig. 4C). In *S. cerevisiae*, there are seven subunits: ScNup84, ScNup85, ScNup120, ScNup133, ScNup145C, ScSec13 and ScSeh1 [73,77,91], but in humans nine subunits, including orthologs to yeast components together with two additional subunits, HsNup43 and HsNup37 (Fig. 4C) [79,80]. In *T. brucei*, the Y-complex contains eight subunits, seven orthologs

to the yeast cohort, *albeit* that trypanosomes lack a Seh1 ortholog, but replace this with an ortholog of HsNup43, TbNup41 (Fig. 4C) [61].

Moreover, the outer rings connect to the nuclear basket (NB), a structure that protrudes from the NPC into the nucleoplasm (Fig. 4A). The NB is an essential platform for RNA export by docking export factors and facilitating transport [68,70]. Across eukaryotes, the NB is constituted by a different number of subunits, for example, in humans it is constituted by three subunits, HsNup153, HsTpr and HsNup50 [92,93], while in yeast it is constituted by five subunits, ScNup1, ScNup60, ScMlp1/2 and ScNup2 [73,77], and in algae NPC basket contains only two subunits [85] (Fig. 4D). As some of the Nups present in these platforms are dynamic, exact functions remain unclear. For example, in mammalian cells it is not known if

HsNup153 is essential for Tpr attachment to the NPC as evidence is contradictory [94]. Moreover, apart from RNA export, the basket Nups have been implicated in chromatin remodelling and control of gene expression [70]. Regarding RNA export, in eukaryotes this process is helped by different complexes; Mex67-Mtr2, TREX and TREX-2 [23,95,96], which interact with NPC components to coordinate the export. The NB has also been suggested to act as a quality control point for RNA export, as it distinguishes between normal and aberrant mRNAs, although the mechanisms for this are unclear [23,97,98].

In trypanosomes, the NB consists of at least two proteins, TbNup92 and TbNup110 (Fig. 4D) which are Kinetoplastid restricted and divergent from NB components in humans and yeast [61,99]. Additional proteins that are likely NB components have also been identified by proteomics, and which also bear predominantly coiled-coil structures (E. R. Butterfield, S. O. Obado, S. R. Scutts, W. Zhang, B. T. Chait, M. P. Rout & M. C. Field, unpublished results). Export factors are also divergent across taxa and classical mammalian complexes such as TREX and TREX-2, regulating mRNA splicing, export and aiding in translocation of mRNPs to the NPC, are essentially absent from trypanosomes [61,100]. Again, this suggests that the complexes interacting with the NPC evolved distinctly in trypanosomes. A remarkable example of this is the recent characterisation of Mex67 paralogues in trypanosomes [101,102]. Mex67-Mtr2 is the main mature mRNP carrier in yeast (Nxf1-Nxt1 complex in humans), essential for translocation of the mRNP to the FG-Nups in the NPC [103]. *T. brucei* is the first protozoa identified to have multiple paralogues of Mex67; TbMex67, TbMex67b and TbMex67L, all encoded by different genes [102,104]. TbMex67 and TbMex67b localise to the nucleolus and the NPC and interact with multiple NPC components and are required for mRNA export. By contrast TbMex67L localises exclusively to the nucleolus and instead interacts with ribosomal biogenesis factors and ribosomal components [102]. Furthermore, paralogues Mex67 and Mex67b interact with the small GTPase Ran, suggesting a Ran-dependent mechanism of RNA export [61,102,104]. This represents another hallmark of trypanosomes, as in animals, fungi and plants, the bulk mRNA export occurs in a Ran-independent manner [61,102].

The NPC outer rings also connect to the cytoplasmic filaments (CFs), that establish transport directionality and provide docking sites for transport factors contributing to the mRNA export process (Fig. 4A) [72]. Residing at the OR at the cytoplasmic side, they are mainly constituted by the yeast Nup82 complex [73,91].

This complex recruits the RNA helicase Dbp5 and associated factors such as Gle1 and Gle2 [105] that release the mRNA to the cytoplasm. In yeast, the Nup82 complex is constituted by three proteins, ScNup82, ScNup159 and ScNup42, while in humans it is constituted by HsNup214, HsNup88 and HsNup62 [80]. In *T. brucei*, the equivalent Nup76 complex is constituted of TbNup76, TbNup140 and TbNup149 [61]. Of these, only TbNup76 shows homology to ScNup82, while the other two subunits are apparently taxon specific [23,61]. Regarding associated factors, only Gle2 has been found in trypanosomes [23,61], with the possibility that taxa-specific proteins interacting with the CF for the release of the mRNAs from the NPC are yet to be discovered.

The inner ring is the structural core and most conserved NPC element (Figs 4A and 5A). IRs extend from the NE membrane to the central channel, shaping and stabilising the NE and providing an anchor for remaining NPC core components and FG Nups. There are three subcomplexes constituting the IR extensively studied in yeast; the ScNup49-ScNup57-ScNsp1 heterotrimer and two heterodimers ScNup188-ScNup192 and ScNup157-ScNup170 [106]. ScNup49-ScNup57-ScNsp1 constitute part of the central transport channel and form a diffusion barrier with their disordered FG repeats [106]. These interact with ScNic96, an evolutionary highly conserved protein that anchors channel Nups with the ScNup192-ScNup188 subcomplex (both α -solenoid). A deeper layer of the IR, interacting with the membrane ring and ScNic96, is the β - α -structured subcomplex ScNup157-ScNup170. A difference between organisms is the presence of extra copies in humans of Nup155 (ortholog of ScNup157/ScNup170) forming additional pillar structures in the vertebrate NPC [77,79] and present only in the nuclear IR in the algae *C. reinhardtii* [85]. *T. brucei* presents orthologs for all of these proteins, namely, TbNup225, TbNup181 (α -solenoids), TbNup119, TbNup144 (β - α -structured), TbNup96 (ortholog of ScNic96) and FG-Nups, TbNup53a, TbNup53b and TbNup62 (orthologs of ScNup57, 47 and Nsp1, respectively) [23,61], showing again the conservation of these NPC elements across eukaryotes.

By contrast, the *trans*-membrane ring has a diverse arrangement, with little sequence conservation across eukaryotes [23,91]. The MR is composed of pore membrane proteins (POMs), four identified in humans and yeast, but only Ndc1 and POM33 have orthologs between species, with the remaining POMs apparently taxa-specific [23]. Moreover, the exact functions of individual POMs remain elusive, beyond contributions to stabilising connections between the NE and the IR [80].




Fig. 5. Heterogeneity in the NPC. (A) Levels of conservation along the NPC among eukaryotes. Green, high conservation, red low conservation. (B) Structural variants of the NPC present in the same cell detected in *Saccharomyces cerevisiae* [116]; Form I corresponds to the canonical NPC, Form II includes an extra OR in the nucleoplasmic side and Form III lacks the NB and is detected in proximity to the nucleolus.

So far, no orthologs for POMs have been detected in trypanosomes [23,61] or other pathogens.

Apart from Kinetoplastid parasites, characterisation of NPC components in several medically important protozoa is in progress, including the Alveolata *Toxoplasma gondii* [107], *Plasmodium berghei* and *Plasmodium falciparum* [108,109] (Fig. 2), and emphasise divergence as several Nups are significantly larger than homologues in animals and fungi. For example, in *P. falciparum*, one of the more virulent species causing malaria, nucleoporin PfSec13 is 90 kDa *versus* animal/fungal Sec13 at 35 kDa. This configuration has been generated by fusion at the C-terminal region with a region similar to yeast Nup145C [110]. In *To. gondii*, TgNup302 is also significantly larger than the yeast homologue Nup100 (302 kDa *versus* 100 kDa, respectively) and bears an autocatalytic domain that generates two polypeptides of 150 and 170 kDa respectively [107]. These observations suggest that Apicomplexan NPCs assemble by distinct pathways from animals and fungi. Interestingly, the localisation and number of NPCs varies considerably between developmental stages in *Plasmodium*, ranging from three to ~60 NPC per nucleus in erythrocytic stages (ring, trophozoite and schizonts) [111]. This may reflect the transcriptional state of the cell and/or metabolic activity as it clearly does in metazoan cells [112,113].

The evolution of the NPC is associated with the protocoatomer hypothesis, which proposes that a huge number of membrane-associated complexes, including the NPC, have a common evolutionary origin as they share deep architectural similarities [23,54]. Protocoatomers are membrane-deforming proteins constituted of β -propeller and α -solenoid domains, whose secondary structure (but not sequence) is highly retained between many subcellular compartments. There are two distinct subfamilies; Type I, featuring an N-terminal β -propeller

followed by a continuous α -solenoid and Type II, constituted by an α -solenoid but with a break in the α -solenoid [23,54]. NPCs contain proteins from both subfamilies, suggesting NPC architecture may have been established late during eukaryogenesis, but also implying that a basic nucleocytoplasmic transport system is ancient and was established pre-LECA [114]. Bioinformatic analysis of NPC subunits show that these are not conserved in sequence among eukaryotes but that the proteins retain recognisable conserved domains and secondary structures. A minor proportion of Nups are lineage specific, e.g., TbNup140 and TbNup149 in *T. brucei*, a potential indication of functional diversification [19,23,54].

All these studies and observations indicate a common structural design for the NPC across eukaryotes [91,100,114], represented in Fig. 4A, but that different taxa have adaptations to this generalised plan. As mentioned, the eukaryotic NPC architectural diversity is produced through gains and losses of components together with varying stoichiometry. In terms of conservation, the IR is the most conserved NPC substructure (Fig. 5A) while subunits in the basket and cytoplasmic filaments are less conserved. A rationale for divergent proteins in the trypanosome NB and CF is that they have adapted to interact codependently with an also divergent mRNA export pathway (Fig. 5A). An alternate view, expressed by one of us recently (Mark C. Field in [115]), is that there are so many differences within RNA processing pathways and NPC composition between trypanosomes and other lineages that this may reflect a more fundamental distinction, and that trypanosomes could be examples of an earlier eukaryotic state [115]; however, the validity of either argument remains unestablished. Apart from variations in the NPC architecture among eukaryotes, variations in composition have been detected among tissues and even in the same cell.

Nuclear pore heterogeneity in the same cell

Studies in yeast have highlighted that the NPC does not have a uniform structure, but rather it can vary even within a single cell, as NPCs of at least three distinct compositions coexist in the same cell at the same time in *S. cerevisiae* [116–118]. Form I is predominant and has the canonical arrangement of two outer rings (OR). Form II contains three ORs, two at the nucleoplasmic side and with enrichment of nucleoporin Nup188 in the nuclear basket (normally in the inner ring). Form III contains the canonical two ORs but lacks nuclear basket subunits and is abundant (but not exclusive) in NPCs proximal to the nucleolus (Fig. 5B). So far, Form III are the only of these forms functionally characterised in *S. cerevisiae* [117,119], highlighting that basket-less NPCs are the default assembly state and that baskets only assemble in an mRNA/mRNA-dependent manner [117,119]. Active RNA Polymerase II transcription and the presence of mature, processed mRNAs are requisites for the assembly of the basket Nups [119]. Importantly, the basket serves as a quality control platform to ensure that non-processed mRNAs stay in the nucleus [119]. The proximity of basket-less NPCs to the nucleolus was suggestive of selective transport of ribosomal subunits through these NPCs for ribosomal production, however, it was later shown that this transport can occur through nuclear pores located far from the nucleolus [120]. Thus, the purpose of basket-less nuclear pores in proximity to the nucleolus is still to be determined. Moreover, it is still unclear how Forms I and II compositions may impact function. These observations highlight that the NPC is heterogeneous even in a single cell; whether there are different NPC compositions in parasites engaging in different cell functions is unknown.

Nuclear pore heterogeneity among tissues

In mammals and plants, a tissue-specific expression of NPC components has been described [121,122], as protein and mRNA levels vary across cell types of healthy and pathological tissues [121]. Interestingly, this tissue-dependent expression is linked to cell differentiation, tissue development and function [121]. Protein and mRNA expression profiles in human tissue culture cells indicate cell type-specific changes to expression of certain NPC components, i.e., cell-specific changes have been observed for HsNup50, Tpr (nuclear basket), HsAladin, HsNup214 (cytoplasmic filaments), HsGp210, HsPOM121 (membrane ring) and HsNup37 (inner ring) [121]. Changes to Nup expression across

tissues may indicate variation in NPC assembly, and hence is suggestive of the existence of specialised nuclear pores with cell type-specific functions [121]. Moreover, these differences in expression, together with loss-of-function mutations in Nup genes, can cause tissue-specific phenotypes [121]. For example, in humans, HsNup155 is highly expressed in the heart and when mutated, leads to cardiac disease [123]. How these differences in stoichiometry NPC subunits impact normal and aberrant functions is an important area of current investigation.

NPC functions

The main function of the NPC is nucleocytoplasmic trafficking but the structure supports roles in additional activities, such as modulation of chromatin, regulation of gene expression (gene activation or silencing; epigenetic gene regulation; regulation of transcription and mRNA export) and transcriptional activity (assembly of transcriptional complexes). Moreover, the NPC subunits can participate in transcriptional memory, i.e., the cellular ability to enhance transcriptional response upon an external cue, which can be maintained epigenetically [124]. As discussed, the NPC represents a docking site for RNA export complexes through the NE and CF. Interestingly, the NPC influences development and differentiation in *Drosophila* as well as hormonal and immunological signalling in plants [124]. Moreover, consistent with many functions occurring in the nucleoplasm, some Nups shuttle between the NPC and the nuclear interior. In several organisms, including yeasts, humans, *Drosophila* and trypanosomes, soluble dynamic Nups bind transcriptionally active genes and recruit them to the nuclear periphery to regulate gene expression [68,70].

In recent years, different diseases have been linked to mutations in the genes of Nups, disorders that have been termed ‘nucleoporopathies’ [72]. These diseases can also arise through alterations in the NPC composition, in subunit levels (sequestration, aggregation or overexpression) or in nucleocytoplasmic transport [72]. Such alterations lead to a range of pathologies, including neurodegenerative diseases (e.g., dementia, Huntington’s, Alzheimer, Parkinson’s) [125], autoimmune disorders (Systemic lupus erythematosus, rheumatic disease), cancer [126], renal [127,128] and ovary dysfunction [129,130].

Trypanosome Nups perform conserved functions observed in higher eukaryotes, as they participate in lamina anchoring (TbNup98) [52,59], tRNA trafficking (TbNup62 and TbNup53a) [131] and mRNA

processing through interactions with the *trans*-splicing machinery [132]. Similar functions are being unravelled for other parasites, as in *To. gondii*, in which TgNup302 participates in trafficking of the ribosomal RNAs and also interacts with histone modification enzymes, suggest a role in gene regulation [107]. Trypanosome Nups also actively participate during mitosis, as TbNup92 (located in the NB) associates with the spindle pole body during cell division to ensure correct chromosome segregation [99] while TbNup144 plays a role in mitosis checkpoints [131]. These roles reflect conservation of Nup functions in cell division across eukaryotes. As happens in mammals, some trypanosome Nups, namely, TbNup53b and TbNup92, are dynamic and shuttle between the NPC and nucleoplasm to fulfil roles in the nuclear interior to interact with the *trans*-splicing machinery and to contribute to chromatid segregation, respectively [99,132].

The dynamics of NPC and NE components during nuclear division remain essentially uncharacterised in parasites. Although open and close mitosis seem distinct, the mechanisms underpinning disassembly of the NPC are highly similar and occur in a synchronised manner. In *Aspergillus nidulans* and *Schi. pombe*, during anaphase the intact daughter nuclei are only connected by a narrow bridge from where NPC components gradually disassemble; peripheral NB Nups detach first from the NPC, followed by central scaffold Nups and finally, POMs [133–136]. In *T. brucei*, evidence points towards a similar mechanism, where NB Nups (TbNup92) [99] are first lost from the mitotic bridge, followed by structural Nups (TbNup158, TbNup53b) [89,132], however, the exact mechanisms for NPC disassembly during mitosis in trypanosomes remain to be explored in detail.

Nups in parasites can openly contribute to regulating expression and interestingly, do so with genes related to developmental stage or pathogenesis. In *T. brucei*, TbNup53b contributes to regulation of expression of procyclins, normally expressed only in the procyclic stage, but depletion of TbNup53b results in the expression of procyclins in the bloodstream form [132]. How Nups in trypanosomes contribute to regulate expression is unknown, but in other eukaryotes is finely orchestrated by Nups in collaboration with the nuclear lamina, epigenetic (histone) marks and RNA processing factors, for example, plants [137], *S. cerevisiae* [138], *Schi. pombe* [139,140] and *Drosophila* [141,142]. Less explored are the roles of NPC components in virulence and infection, e.g., *Leishmania*, an intracellular protozoan parasite, uses the metalloprotease GP63 to degrade host Nups and nuclear receptors, altering the composition of the infected host cell [143,144], while

Theileria amulata, a bovine intracellular pathogen, recruits NPC components of the host cell to the surface to form porous structures of unknown function [145].

Conclusions and perspectives: the NPC

One of the largest macromolecule complexes in the cell, the NPC is a superb exemplar of molecular evolution, diversity and adaptation. Each lineage possesses an NPC which has evolved and adapted to meet specific biological constraints. Although the NPC core is mostly conserved, remaining NPC components have adapted to interact with transcription, mRNA processing and mRNA export machineries that are frequently highly specialised between lineages. Nevertheless, there are many unsolved questions about the NPC components in parasites, for example, about their precise role in gene expression and transcription regulation or whether and how those components may, or may not, influence virulence or development. Moreover, while the overall functions of the NPC are highly conserved, there are significant differences in the mechanisms by which these are supported, providing a complex tangle of convergent and divergent evolutionary pathways.

Acknowledgements

Our laboratory is supported by the Wellcome Trust (204697/Z/16/Z to MCF). We thank Erin Butterfield and Alex Makarov for fruitful discussions and comments.

References

- Schoch CL, Ciuffo S, Domrachev M, Hottel CL, Kannan S, Khovanskaya R, Leipe D, McVeigh R, O'Neill K, Robbertse B *et al.* (2020) NCBI taxonomy: a comprehensive update on curation, resources and tools. *Database (Oxford)* **2020**, baaa062.
- Englund PT, DiMaio DC and Price SS (1977) A nicked form of kinetoplast DNA in *Leishmania tarentolae*. *J Biol Chem* **252**, 6208–6216.
- Simpson L (1972) The kinetoplast of the hennoflagellates. *Int Rev Cytol* **32**, 139–207.
- Desquesnes M, Gonzatti M, Sazmand A, Thevenon S, Bossard G, Boulange A, Gimonneau G, Truc P, Herder S, Ravel S *et al.* (2022) A review on the diagnosis of animal trypanosomoses. *Parasit Vectors* **15**, 64.
- Pays E, Radwanska M and Magez S (2023) The pathogenesis of African trypanosomiasis. *Annu Rev Pathol* **18**, 19–45.
- Buscher P, Gonzatti MI, Hebert L, Inoue N, Pascucci I, Schnauffer A, Suganuma K, Touratier L and Van

- Reet N (2019) Equine trypanosomiasis: enigmas and diagnostic challenges. *Parasit Vectors* **12**, 234.
- 7 Luzak V, Lopez-Escobar L, Siegel TN and Figueiredo LM (2021) Cell-to-cell heterogeneity in trypanosomes. *Annu Rev Microbiol* **75**, 107–128.
 - 8 Hill KL (2003) Biology and mechanism of trypanosoma cell motility. *Eukaryot Cell* **2**, 200–208.
 - 9 Lamont GS, Tucker RS and Cross GA (1986) Analysis of antigen switching rates in *Trypanosoma brucei*. *Parasitology* **92** (Pt 2), 355–367.
 - 10 Borst P (1986) Discontinuous transcription and antigenic variation in trypanosomes. *Annu Rev Biochem* **55**, 701–732.
 - 11 Navarro M and Gull K (2001) A pol I transcriptional body associated with VSG mono-allelic expression in *Trypanosoma brucei*. *Nature* **414**, 759–763.
 - 12 Rudenko G, Bishop D, Gottesdiener K and Van der Ploeg LH (1989) Alpha-amanitin resistant transcription of protein coding genes in insect and bloodstream form *Trypanosoma brucei*. *EMBO J* **8**, 4259–4263.
 - 13 Rao SPS, Manjunatha UH, Mikolajczak S, Ashigbie PG and Diagana TT (2023) Drug discovery for parasitic diseases: powered by technology, enabled by pharmacology, informed by clinical science. *Trends Parasitol* **39**, 260–271.
 - 14 De Rycker M, Wyllie S, Horn D, Read KD and Gilbert IH (2023) Anti-trypanosomatid drug discovery: progress and challenges. *Nat Rev Microbiol* **21**, 35–50.
 - 15 WHO (2023) WHO Interim Guidelines for the Treatment of Gambiense Human African Trypanosomiasis. WHO, Geneva.
 - 16 WHO (2023) Chagas disease (American trypanosomiasis). [https://who.int/news-room/fact-sheets/detail/chagas-disease-\(american-trypanosomiasis\)](https://who.int/news-room/fact-sheets/detail/chagas-disease-(american-trypanosomiasis)). Accessed June 2023.
 - 17 PAHO (2023) Chagas disease. <https://www.paho.org/en/topics/chagas-disease>. Accessed June 2023.
 - 18 Torres-Guerrero E, Quintanilla-Cedillo MR, Ruiz-Esmenjaud J and Arenas R (2017) Leishmaniasis: a review. *F1000Res* **6**, 750.
 - 19 Koumandou VL, Wickstead B, Ginger ML, van der Giezen M, Dacks JB and Field MC (2013) Molecular paleontology and complexity in the last eukaryotic common ancestor. *Crit Rev Biochem Mol Biol* **48**, 373–396.
 - 20 Eme L, Tamarit D, Caceres EF, Stairs CW, De Anda V, Schon ME, Seitz KW, Dombrowski N, Lewis WH, Homa F *et al.* (2023) Inference and reconstruction of the heimdallarchaeal ancestry of eukaryotes. *Nature* **618**, 992–999.
 - 21 Lopez-Garcia P and Moreira D (2015) Open questions on the origin of eukaryotes. *Trends Ecol Evol* **30**, 697–708.
 - 22 Lukes J, Skalicky T, Tyc J, Votypka J and Yurchenko V (2014) Evolution of parasitism in kinetoplastid flagellates. *Mol Biochem Parasitol* **195**, 115–122.
 - 23 Makarov AA, Padilla-Mejia NE and Field MC (2021) Evolution and diversification of the nuclear pore complex. *Biochem Soc Trans* **49**, 1601–1619.
 - 24 Aebi U, Cohn J, Buhle L and Gerace L (1986) The nuclear lamina is a meshwork of intermediate-type filaments. *Nature* **323**, 560–564.
 - 25 Stiekema M, van Zandvoort M, Ramaekers FCS and Broers JLV (2020) Structural and mechanical aberrations of the nuclear lamina in disease. *Cell* **9**, 1884.
 - 26 Turgay Y, Eibauer M, Goldman AE, Shimi T, Khayat M, Ben-Harush K, Dubrovsky-Gaupp A, Sapra KT, Goldman RD and Medalia O (2017) The molecular architecture of lamins in somatic cells. *Nature* **543**, 261–264.
 - 27 Kollmar M (2015) Polyphyly of nuclear lamin genes indicates an early eukaryotic origin of the metazoan-type intermediate filament proteins. *Sci Rep* **5**, 10652.
 - 28 Evangelisti C, Rusciano I, Mongiorgi S, Ramazzotti G, Lattanzi G, Manzoli L, Cocco L and Ratti S (2022) The wide and growing range of lamin B-related diseases: from laminopathies to cancer. *Cell Mol Life Sci* **79**, 126.
 - 29 Shin JY and Worman HJ (2022) Molecular pathology of laminopathies. *Annu Rev Pathol* **17**, 159–180.
 - 30 Tenga R and Medalia O (2020) Structure and unique mechanical aspects of nuclear lamin filaments. *Curr Opin Struct Biol* **64**, 152–159.
 - 31 Nmezi B, Xu J, Fu R, Armiger TJ, Rodriguez-Bey G, Powell JS, Ma H, Sullivan M, Tu Y, Chen NY *et al.* (2019) Concentric organization of A- and B-type lamins predicts their distinct roles in the spatial organization and stability of the nuclear lamina. *Proc Natl Acad Sci USA* **116**, 4307–4315.
 - 32 Strelkov SV, Herrmann H and Aebi U (2003) Molecular architecture of intermediate filaments. *Bioessays* **25**, 243–251.
 - 33 Shimi T, Kittisopikul M, Tran J, Goldman AE, Adam SA, Zheng Y, Jaqaman K and Goldman RD (2015) Structural organization of nuclear lamins A, C, B1, and B2 revealed by superresolution microscopy. *Mol Biol Cell* **26**, 4075–4086.
 - 34 Shimi T, Pfliegerhaer K, Kojima S, Pack CG, Solovei I, Goldman AE, Adam SA, Shumaker DK, Kinjo M, Cremer T *et al.* (2008) The A- and B-type nuclear lamin networks: microdomains involved in chromatin organization and transcription. *Genes Dev* **22**, 3409–3421.
 - 35 Goldberg MW, Huttenlauch I, Hutchison CJ and Stick R (2008) Filaments made from A- and B-type lamins differ in structure and organization. *J Cell Sci* **121**, 215–225.
 - 36 Xie W, Chojnowski A, Boudier T, Lim JS, Ahmed S, Ser Z, Stewart C and Burke B (2016) A-type lamins form distinct filamentous networks with differential

- nuclear pore complex associations. *Curr Biol* **26**, 2651–2658.
- 37 Madsen-Osterbye J, Bellanger A, Galigniana NM and Collas P (2022) Biology and model predictions of the dynamics and heterogeneity of chromatin-nuclear lamina interactions. *Front Cell Dev Biol* **10**, 913458.
 - 38 Vester K, Preussner M, Holton N, Feng S, Schultz C, Heyd F and Wahl MC (2022) Recruitment of a splicing factor to the nuclear lamina for its inactivation. *Commun Biol* **5**, 736.
 - 39 Naetar N, Georgiou K, Knapp C, Bronshtein I, Zier E, Fichtinger P, Dechat T, Garini Y and Foisner R (2021) LAP2alpha maintains a mobile and low assembly state of A-type lamins in the nuclear interior. *Elife* **10**, e63476.
 - 40 Liu SY and Ikegami K (2020) Nuclear lamin phosphorylation: an emerging role in gene regulation and pathogenesis of laminopathies. *Nucleus* **11**, 299–314.
 - 41 Ikegami K, Secchia S, Almakki O, Lieb JD and Moskowitz IP (2020) Phosphorylated lamin A/C in the nuclear interior binds active enhancers associated with abnormal transcription in progeria. *Dev Cell* **52**, 699–713.e11.
 - 42 Ottaviano Y and Gerace L (1985) Phosphorylation of the nuclear lamins during interphase and mitosis. *J Biol Chem* **260**, 624–632.
 - 43 Ogbadoyi E, Ersfeld K, Robinson D, Sherwin T and Gull K (2000) Architecture of the *Trypanosoma brucei* nucleus during interphase and mitosis. *Chromosoma* **108**, 501–513.
 - 44 Kruger A, Batsios P, Baumann O, Luckert E, Schwarz H, Stick R, Meyer I and Graf R (2012) Characterization of NE81, the first lamin-like nucleoskeleton protein in a unicellular organism. *Mol Biol Cell* **23**, 360–370.
 - 45 Batsios P, Graf R, Koonce MP, Larochele DA and Meyer I (2019) Nuclear envelope organization in *Dictyostelium discoideum*. *Int J Dev Biol* **63**, 509–519.
 - 46 Koreny L and Field MC (2016) Ancient eukaryotic origin and evolutionary plasticity of nuclear lamina. *Genome Biol Evol* **8**, 2663–2671.
 - 47 Andrulis ED, Zappulla DC, Ansari A, Perrod S, Laiosa CV, Gartenberg MR and Sternglanz R (2002) Esc1, a nuclear periphery protein required for Sir4-based plasmid anchoring and partitioning. *Mol Cell Biol* **22**, 8292–8301.
 - 48 Faure G, Jezequel K, Roisne-Hamelin F, Bitard-Feildel T, Lamiabile A, Marcand S and Callebaut I (2019) Discovery and evolution of new domains in yeast heterochromatin factor Sir4 and its partner Esc1. *Genome Biol Evol* **11**, 572–585.
 - 49 Ciska M, Hikida R, Masuda K and Moreno Diaz de la Espina S (2019) Evolutionary history and structure of nuclear matrix constituent proteins, the plant analogues of lamins. *J Exp Bot* **70**, 2651–2664.
 - 50 Ciska M, Masuda K and Moreno Diaz de la Espina S (2013) Lamin-like analogues in plants: the characterization of NMCP1 in *Allium cepa*. *J Exp Bot* **64**, 1553–1564.
 - 51 Ciska M, Masuda K and Moreno Diaz de la Espina S (2018) Characterization of the lamin analogue NMCP2 in the monocot *Allium cepa*. *Chromosoma* **127**, 103–113.
 - 52 DuBois KN, Alsford S, Holden JM, Buisson J, Swiderski M, Bart JM, Ratushny AV, Wan Y, Bastin P, Barry JD *et al.* (2012) NUP-1 is a large coiled-coil nucleoskeletal protein in trypanosomes with lamin-like functions. *PLoS Biol* **10**, e1001287.
 - 53 Maishman L, Obado SO, Alsford S, Bart JM, Chen WM, Ratushny AV, Navarro M, Horn D, Aitchison JD, Chait BT *et al.* (2016) Co-dependence between trypanosome nuclear lamina components in nuclear stability and control of gene expression. *Nucleic Acids Res* **44**, 10554–10570.
 - 54 Rout MP, Obado SO, Schenkman S and Field MC (2017) Specialising the parasite nucleus: pores, lamins, chromatin, and diversity. *PLoS Pathog* **13**, e1006170.
 - 55 Graf R, Batsios P and Meyer I (2015) Evolution of centrosomes and the nuclear lamina: Amoebozoan assets. *Eur J Cell Biol* **94**, 249–256.
 - 56 O'Malley MA, Leger MM, Wideman JG and Ruiz-Trillo I (2019) Concepts of the last eukaryotic common ancestor. *Nat Ecol Evol* **3**, 338–344.
 - 57 Rout MP and Field MC (2001) Isolation and characterization of subnuclear compartments from *Trypanosoma brucei*. Identification of a major repetitive nuclear lamina component. *J Biol Chem* **276**, 38261–38271.
 - 58 Picchi GF, Ferreira AM, Souza FS, Lourenco EE, Arauco PR, Lorusso A, Bordignon J, Krieger MA, Goldenberg S and Fragoso SP (2011) *Trypanosoma cruzi*: identification of DNA targets of the nuclear periphery coiled-coil protein TcNUP-1. *Exp Parasitol* **127**, 147–152.
 - 59 Padilla-Mejia NE, Koreny L, Holden J, Vancova M, Lukes J, Zoltner M and Field MC (2021) A hub-and-spoke nuclear lamina architecture in trypanosomes. *J Cell Sci* **134**, jcs251264.
 - 60 Funkhouser CM, Sknepnek R, Shimi T, Goldman AE, Goldman RD and Olvera de la Cruz M (2013) Mechanical model of blebbing in nuclear lamin meshworks. *Proc Natl Acad Sci USA* **110**, 3248–3253.
 - 61 Obado SO, Brillantes M, Uryu K, Zhang W, Ketaren NE, Chait BT, Field MC and Rout MP (2016) Interactome mapping reveals the evolutionary history of the nuclear pore complex. *PLoS Biol* **14**, e1002365.
 - 62 Mugo E and Clayton C (2017) Expression of the RNA-binding protein RBP10 promotes the bloodstream-form differentiation state in *Trypanosoma brucei*. *PLoS Pathog* **13**, e1006560.

- 63 Muller LSM, Cosentino RO, Forstner KU, Guizetti J, Wedel C, Kaplan N, Janzen CJ, Arampatzi P, Vogel J, Steinbiss S *et al.* (2018) Genome organization and DNA accessibility control antigenic variation in trypanosomes. *Nature* **563**, 121–125.
- 64 Faria J, Luzak V, Muller LSM, Brink BG, Hutchinson S, Glover L, Horn D and Siegel TN (2021) Spatial integration of transcription and splicing in a dedicated compartment sustains monogenic antigen expression in African trypanosomes. *Nat Microbiol* **6**, 289–300.
- 65 Shanmugasundram A, Starns D, Bohme U, Amos B, Wilkinson PA, Harb OS, Warrenfeltz S, Kissinger JC, McDowell MA, Roos DS *et al.* (2023) TriTrypDB: an integrated functional genomics resource for kinetoplastida. *PLoS Negl Trop Dis* **17**, e0011058.
- 66 Urbaniak MD, Martin DM and Ferguson MA (2013) Global quantitative SILAC phosphoproteomics reveals differential phosphorylation is widespread between the procyclic and bloodstream form lifecycle stages of *Trypanosoma brucei*. *J Proteome Res* **12**, 2233–2244.
- 67 Li Y, Jiang X, Zhang Y, Gao Z, Liu Y, Hu J, Hu X, Li L, Shi J and Gao N (2019) Nuclear accumulation of UBC9 contributes to SUMOylation of lamin A/C and nucleophagy in response to DNA damage. *J Exp Clin Cancer Res* **38**, 67.
- 68 Beck M and Hurt E (2017) The nuclear pore complex: understanding its function through structural insight. *Nat Rev Mol Cell Biol* **18**, 73–89.
- 69 Callam HG and Tomlin SG (1950) Experimental studies on amphibian oocyte nuclei. *Proc R Soc Lond B Biol Sci* **137**, 367–378.
- 70 Lin DH and Hoelz A (2019) The structure of the nuclear pore complex (an update). *Annu Rev Biochem* **88**, 725–783.
- 71 Amlacher S, Sarges P, Flemming D, van Noort V, Kunze R, Devos DP, Arumugam M, Bork P and Hurt E (2011) Insight into structure and assembly of the nuclear pore complex by utilizing the genome of a eukaryotic thermophile. *Cell* **146**, 277–289.
- 72 Bley CJ, Nie S, Mobbs GW, Petrovic S, Gres AT, Liu X, Mukherjee S, Harvey S, Huber FM, Lin DH *et al.* (2022) Architecture of the cytoplasmic face of the nuclear pore. *Science* **376**, eabm9129.
- 73 Fernandez-Martinez J, Kim SJ, Shi Y, Upla P, Pellarin R, Gagnon M, Chemmama IE, Wang J, Nudelman I, Zhang W *et al.* (2016) Structure and function of the nuclear pore complex cytoplasmic mRNA export platform. *Cell* **167**, 1215–1228.e25.
- 74 Fontana P, Dong Y, Pi X, Tong AB, Hecksel CW, Wang L, Fu TM, Bustamante C and Wu H (2022) Structure of cytoplasmic ring of nuclear pore complex by integrative cryo-EM and AlphaFold. *Science* **376**, eabm9326.
- 75 Huang G, Zhan X, Zeng C, Liang K, Zhu X, Zhao Y, Wang P, Wang Q, Zhou Q, Tao Q *et al.* (2022) Cryo-EM structure of the inner ring from the *Xenopus laevis* nuclear pore complex. *Cell Res* **32**, 451–460.
- 76 Huang G, Zhan X, Zeng C, Zhu X, Liang K, Zhao Y, Wang P, Wang Q, Zhou Q, Tao Q *et al.* (2022) Cryo-EM structure of the nuclear ring from *Xenopus laevis* nuclear pore complex. *Cell Res* **32**, 349–358.
- 77 Kim SJ, Fernandez-Martinez J, Nudelman I, Shi Y, Zhang W, Raveh B, Herricks T, Slaughter BD, Hogan JA, Upla P *et al.* (2018) Integrative structure and functional anatomy of a nuclear pore complex. *Nature* **555**, 475–482.
- 78 Kim SJ, Fernandez-Martinez J, Sampathkumar P, Martel A, Matsui T, Tsuruta H, Weiss TM, Shi Y, Markina-Inarrairaegui A, Bonanno JB *et al.* (2014) Integrative structure-function mapping of the nucleoporin Nup133 suggests a conserved mechanism for membrane anchoring of the nuclear pore complex. *Mol Cell Proteomics* **13**, 2911–2926.
- 79 Kosinski J, Mosalaganti S, von Appen A, Teimer R, DiGuilio AL, Wan W, Bui KH, Hagen WJ, Briggs JA, Glavy JS *et al.* (2016) Molecular architecture of the inner ring scaffold of the human nuclear pore complex. *Science* **352**, 363–365.
- 80 von Appen A, Kosinski J, Sparks L, Ori A, DiGuilio AL, Vollmer B, Mackmull MT, Banterle N, Parca L, Kastiritis P *et al.* (2015) *In situ* structural analysis of the human nuclear pore complex. *Nature* **526**, 140–143.
- 81 Zhang Y, Li S, Zeng C, Huang G, Zhu X, Wang Q, Wang K, Zhou Q, Yan C, Zhang W *et al.* (2020) Molecular architecture of the luminal ring of the *Xenopus laevis* nuclear pore complex. *Cell Res* **30**, 532–540.
- 82 Zhu X, Huang G, Zeng C, Zhan X, Liang K, Xu Q, Zhao Y, Wang P, Wang Q, Zhou Q *et al.* (2022) Structure of the cytoplasmic ring of the *Xenopus laevis* nuclear pore complex. *Science* **376**, eabl8280.
- 83 Asakawa H, Kojidani T, Yang HJ, Ohtsuki C, Osakada H, Matsuda A, Iwamoto M, Chikashige Y, Nagao K, Obuse C *et al.* (2019) Asymmetrical localization of Nup107-160 subcomplex components within the nuclear pore complex in fission yeast. *PLoS Genet* **15**, e1008061.
- 84 Asakawa H, Yang HJ, Yamamoto TG, Ohtsuki C, Chikashige Y, Sakata-Sogawa K, Tokunaga M, Iwamoto M, Hiraoka Y and Haraguchi T (2014) Characterization of nuclear pore complex components in fission yeast *Schizosaccharomyces pombe*. *Nucleus* **5**, 149–162.
- 85 Mosalaganti S, Kosinski J, Albert S, Schaffer M, Strenkert D, Salome PA, Merchant SS, Plitzko JM, Baumeister W, Engel BD *et al.* (2018) *In situ* architecture of the algal nuclear pore complex. *Nat Commun* **9**, 2361.
- 86 Tamura K, Fukao Y, Iwamoto M, Haraguchi T and Hara-Nishimura I (2010) Identification

- and characterization of nuclear pore complex components in *Arabidopsis thaliana*. *Plant Cell* **22**, 4084–4097.
- 87 Zhang C, An N, Jia P, Zhang W, Liang J, Zhang X, Zhou H, Ma W, Han M, Xing L *et al.* (2020) Genomic identification and expression analysis of nuclear pore proteins in *Malus domestica*. *Sci Rep* **10**, 17426.
- 88 Iwamoto M, Osakada H, Mori C, Fukuda Y, Nagao K, Obuse C, Hiraoka Y and Haraguchi T (2017) Compositionally distinct nuclear pore complexes of functionally distinct dimorphic nuclei in the ciliate *Tetrahymena*. *J Cell Sci* **130**, 1822–1834.
- 89 DeGrasse JA, DuBois KN, Devos D, Siegel TN, Sali A, Field MC, Rout MP and Chait BT (2009) Evidence for a shared nuclear pore complex architecture that is conserved from the last common eukaryotic ancestor. *Mol Cell Proteomics* **8**, 2119–2130.
- 90 Chopra K, Bawaria S and Chauhan R (2019) Evolutionary divergence of the nuclear pore complex from fungi to metazoans. *Protein Sci* **28**, 571–586.
- 91 Fernandez-Martinez J and Rout MP (2021) One ring to rule them all? Structural and functional diversity in the nuclear pore complex. *Trends Biochem Sci* **46**, 595–607.
- 92 Frosst P, Guan T, Subauste C, Hahn K and Gerace L (2002) Tpr is localized within the nuclear basket of the pore complex and has a role in nuclear protein export. *J Cell Biol* **156**, 617–630.
- 93 Krull S, Thyberg J, Bjorkroth B, Rackwitz HR and Cordes VC (2004) Nucleoporins as components of the nuclear pore complex core structure and Tpr as the architectural element of the nuclear basket. *Mol Biol Cell* **15**, 4261–4277.
- 94 Duheron V, Chatel G, Sauder U, Oliveri V and Fahrenkrog B (2014) Structural characterization of altered nucleoporin Nup153 expression in human cells by thin-section electron microscopy. *Nucleus* **5**, 601–612.
- 95 Derrer CP, Mancini R, Vallotton P, Huet S, Weis K and Dultz E (2019) The RNA export factor Mex67 functions as a mobile nucleoporin. *J Cell Biol* **218**, 3967–3976.
- 96 Stewart M (2019) Structure and function of the TREX-2 complex. *Subcell Biochem* **93**, 461–470.
- 97 Xie Y and Ren Y (2019) Mechanisms of nuclear mRNA export: a structural perspective. *Traffic* **20**, 829–840.
- 98 Schneider M, Hellerschmied D, Schubert T, Amlacher S, Vinayachandran V, Reja R, Pugh BF, Clausen T and Kohler A (2015) The nuclear pore-associated TREX-2 complex employs mediator to regulate gene expression. *Cell* **162**, 1016–1028.
- 99 Holden JM, Koreny L, Obado S, Ratushny AV, Chen WM, Chiang JH, Kelly S, Chait BT, Aitchison JD, Rout MP *et al.* (2014) Nuclear pore complex evolution: a trypanosome Mlp analogue functions in chromosomal segregation but lacks transcriptional barrier activity. *Mol Biol Cell* **25**, 1421–1436.
- 100 Obado SO, Rout MP and Field MC (2022) Sending the message: specialized RNA export mechanisms in trypanosomes. *Trends Parasitol* **38**, 854–867.
- 101 Dostalova A, Kaser S, Cristodero M and Schimanski B (2013) The nuclear mRNA export receptor Mex67-Mtr2 of *Trypanosoma brucei* contains a unique and essential zinc finger motif. *Mol Microbiol* **88**, 728–739.
- 102 Obado SO, Stein M, Hegedúsová E, Zhang W, Hutchinson S, Brillantes M, Glover L, Paris Z, Chait BT, Field MC *et al.* (2022) Mex67 paralogs mediate division of labor in trypanosome RNA processing and export. *bioRxiv*. doi: [10.1101/2022.06.27.497849](https://doi.org/10.1101/2022.06.27.497849) [PREPRINT].
- 103 Aibara S, Valkov E, Lamers MH, Dimitrova L, Hurt E and Stewart M (2015) Structural characterization of the principal mRNA-export factor Mex67-Mtr2 from *Chaetomium thermophilum*. *Acta Crystallogr F Struct Biol Commun* **71**, 876–888.
- 104 Hegedusova E, Kulkarni S, Burgman B, Alfonso JD and Paris Z (2019) The general mRNA exporters Mex67 and Mtr2 play distinct roles in nuclear export of tRNAs in *Trypanosoma brucei*. *Nucleic Acids Res* **47**, 8620–8631.
- 105 Folkmann AW, Noble KN, Cole CN and Wentz SR (2011) Dbp5, Gle1-IP6 and Nup159: a working model for mRNP export. *Nucleus* **2**, 540–548.
- 106 Stuwe T, Bley CJ, Thierbach K, Petrovic S, Schilbach S, Mayo DJ, Perriches T, Rundlet EJ, Jeon YE, Collins LN *et al.* (2015) Architecture of the fungal nuclear pore inner ring complex. *Science* **350**, 56–64.
- 107 Courjol F, Mouveaux T, Lesage K, Saliou JM, Werkmeister E, Bonabaud M, Rohmer M, Slomianny C, Lafont F and Gissot M (2017) Characterization of a nuclear pore protein sheds light on the roles and composition of the *Toxoplasma gondii* nuclear pore complex. *Cell Mol Life Sci* **74**, 2107–2125.
- 108 Ambekar SV, Beck JR and Mair GR (2022) TurboID identification of evolutionarily divergent components of the nuclear pore complex in the malaria model *Plasmodium berghei*. *mBio* **13**, e0181522.
- 109 Kehrer J, Kuss C, Andres-Pons A, Reustle A, Dahan N, Devos D, Kudryashev M, Beck M, Mair GR and Frischknecht F (2018) Nuclear pore complex components in the malaria parasite *Plasmodium berghei*. *Sci Rep* **8**, 11249.
- 110 Dahan-Pasternak N, Nasereddin A, Kolevzon N, Pe'er M, Wong W, Shinder V, Turnbull L, Whitchurch CB, Elbaum M, Gilberger TW *et al.* (2013) PfSec13 is an unusual chromatin-associated nucleoporin of *Plasmodium falciparum* that is essential for parasite proliferation in human erythrocytes. *J Cell Sci* **126**, 3055–3069.

- 111 Weiner A, Dahan-Pasternak N, Shimoni E, Shinder V, von Huth P, Elbaum M and Dzikowski R (2011) 3D nuclear architecture reveals coupled cell cycle dynamics of chromatin and nuclear pores in the malaria parasite *Plasmodium falciparum*. *Cell Microbiol* **13**, 967–977.
- 112 Maul GG, Deaven LL, Freed JJ, Campbell GL and Becak W (1980) Investigation of the determinants of nuclear pore number. *Cytogenet Cell Genet* **26**, 175–190.
- 113 Rabut G, Lenart P and Ellenberg J (2004) Dynamics of nuclear pore complex organization through the cell cycle. *Curr Opin Cell Biol* **16**, 314–321.
- 114 Field MC and Rout MP (2019) Pore timing: the evolutionary origins of the nucleus and nuclear pore complex. *F1000Res* **8**, F1000 Faculty Rev-369.
- 115 Field MC (2023) Deviating from the norm: nuclear organisation in trypanosomes. *Curr Opin Cell Biol* **85**, 102234.
- 116 Akey CW, Singh D, Ouch C, Echeverria I, Nudelman I, Varberg JM, Yu Z, Fang F, Shi Y, Wang J *et al.* (2022) Comprehensive structure and functional adaptations of the yeast nuclear pore complex. *Cell* **185**, 361–378.e25.
- 117 Galy V, Gadal O, Fromont-Racine M, Romano A, Jacquier A and Nehrbass U (2004) Nuclear retention of unspliced mRNAs in yeast is mediated by perinuclear Mlp1. *Cell* **116**, 63–73.
- 118 Niepel M, Molloy KR, Williams R, Farr JC, Meinema AC, Vecchietti N, Cristea IM, Chait BT, Rout MP and Strambio-De-Castillia C (2013) The nuclear basket proteins Mlp1p and Mlp2p are part of a dynamic interactome including Esc1p and the proteasome. *Mol Biol Cell* **24**, 3920–3938.
- 119 Bensidoun P, Reiter T, Montpetit B, Zenklusen D and Oeffinger M (2022) Nuclear mRNA metabolism drives selective basket assembly on a subset of nuclear pore complexes in budding yeast. *Mol Cell* **82**, 3856–3871.e6.
- 120 Delavoie F, Soldan V, Rinaldi D, Dauxois JY and Gleizes PE (2019) The path of pre-ribosomes through the nuclear pore complex revealed by electron tomography. *Nat Commun* **10**, 497.
- 121 Ori A, Banterle N, Iskar M, Andres-Pons A, Escher C, Khanh Bui H, Sparks L, Solis-Mezarino V, Rinner O, Bork P *et al.* (2013) Cell type-specific nuclear pores: a case in point for context-dependent stoichiometry of molecular machines. *Mol Syst Biol* **9**, 648.
- 122 Raices M and D'Angelo MA (2012) Nuclear pore complex composition: a new regulator of tissue-specific and developmental functions. *Nat Rev Mol Cell Biol* **13**, 687–699.
- 123 Zhang X, Chen S, Yoo S, Chakrabarti S, Zhang T, Ke T, Oberti C, Yong SL, Fang F, Li L *et al.* (2008) Mutation in nuclear pore component NUP155 leads to atrial fibrillation and early sudden cardiac death. *Cell* **135**, 1017–1027.
- 124 Pascual-Garcia P and Capelson M (2021) The nuclear pore complex and the genome: organizing and regulatory principles. *Curr Opin Genet Dev* **67**, 142–150.
- 125 Liu J and Hetzer MW (2022) Nuclear pore complex maintenance and implications for age-related diseases. *Trends Cell Biol* **32**, 216–227.
- 126 Nofrini V, Di Giacomo D and Mecucci C (2016) Nucleoporin genes in human diseases. *Eur J Hum Genet* **24**, 1388–1395.
- 127 Braun DA, Sadowski CE, Kohl S, Lovric S, Astrinidis SA, Pabst WL, Gee HY, Ashraf S, Lawson JA, Shril S *et al.* (2016) Mutations in nuclear pore genes NUP93, NUP205 and XPO5 cause steroid-resistant nephrotic syndrome. *Nat Genet* **48**, 457–465.
- 128 Miyake N, Tsukaguchi H, Koshimizu E, Shono A, Matsunaga S, Shiina M, Mimura Y, Imamura S, Hirose T, Okudela K *et al.* (2015) Biallelic mutations in nuclear pore complex subunit NUP107 cause early-childhood-onset steroid-resistant nephrotic syndrome. *Am J Hum Genet* **97**, 555–566.
- 129 Shore T, Levi T, Kalifa R, Dreifuss A, Rekler D, Weinberg-Shukron A, Nevo Y, Bialistok T, Moyal V, Gold MY *et al.* (2022) Nucleoporin107 mediates female sexual differentiation via Dsx. *Elife* **11**, e72632.
- 130 Weinberg-Shukron A, Renbaum P, Kalifa R, Zeligson S, Ben-Neriah Z, Dreifuss A, Abu-Rayyan A, Maatuk N, Fardian N, Rekler D *et al.* (2015) A mutation in the nucleoporin-107 gene causes XX gonadal dysgenesis. *J Clin Invest* **125**, 4295–4304.
- 131 Hegedusova E, Marsalova V, Kulkarni S and Paris Z (2022) Trafficking and/or division: distinct roles of nucleoporins based on their location within the nuclear pore complex. *RNA Biol* **19**, 650–661.
- 132 Holden JM, Koreny L, Obado S, Ratushny AV, Chen WM, Bart JM, Navarro M, Chait BT, Aitchison JD, Rout MP *et al.* (2018) Involvement in surface antigen expression by a moonlighting FG-repeat nucleoporin in trypanosomes. *Mol Biol Cell* **29**, 1100–1110.
- 133 De Souza CP, Osmani AH, Hashmi SB and Osmani SA (2004) Partial nuclear pore complex disassembly during closed mitosis in *Aspergillus nidulans*. *Curr Biol* **14**, 1973–1984.
- 134 Dey G, Culley S, Curran S, Schmidt U, Henriques R, Kukulski W and Baum B (2020) Closed mitosis requires local disassembly of the nuclear envelope. *Nature* **585**, 119–123.
- 135 Exposito-Serrano M, Sanchez-Molina A, Gallardo P, Salas-Pino S and Daga RR (2020) Selective nuclear pore complex removal drives nuclear envelope division in fission yeast. *Curr Biol* **30**, 3212–3222.e2.
- 136 Osmani AH, Davies J, Liu HL, Nile A and Osmani SA (2006) Systematic deletion and mitotic localization

- of the nuclear pore complex proteins of *Aspergillus nidulans*. *Mol Biol Cell* **17**, 4946–4961.
- 137 Zhu Y, Wang B, Tang K, Hsu CC, Xie S, Du H, Yang Y, Tao WA and Zhu JK (2017) An *Arabidopsis* nucleoporin NUP85 modulates plant responses to ABA and salt stress. *PLoS Genet* **13**, e1007124.
- 138 Dilworth DJ, Tackett AJ, Rogers RS, Yi EC, Christmas RH, Smith JJ, Siegel AF, Chait BT, Wozniak RW and Aitchison JD (2005) The mobile nucleoporin Nup2p and chromatin-bound Prp20p function in endogenous NPC-mediated transcriptional control. *J Cell Biol* **171**, 955–965.
- 139 Holla S, Dhakshnamoorthy J, Folco HD, Balachandran V, Xiao H, Sun LL, Wheeler D, Zofall M and Grewal SIS (2020) Positioning heterochromatin at the nuclear periphery suppresses histone turnover to promote epigenetic inheritance. *Cell* **180**, 150–164.e15.
- 140 Iglesias N, Paulo JA, Tatarakis A, Wang X, Edwards AL, Bhanu NV, Garcia BA, Haas W, Gygi SP and Moazed D (2020) Native chromatin proteomics reveals a role for specific nucleoporins in heterochromatin organization and maintenance. *Mol Cell* **77**, 51–66.e8.
- 141 Gozalo A, Duke A, Lan Y, Pascual-Garcia P, Talamas JA, Nguyen SC, Shah PP, Jain R, Joyce EF and Capelson M (2020) Core components of the nuclear pore bind distinct states of chromatin and contribute to polycomb repression. *Mol Cell* **77**, 67–81.e7.
- 142 Munafo M, Lawless VR, Passera A, MacMillan S, Bornelov S, Haussmann IU, Soller M, Hannon GJ and Czech B (2021) Channel nuclear pore complex subunits are required for transposon silencing in *Drosophila*. *Elife* **10**, e66321.
- 143 Isnard A, Christian JG, Kodiha M, Stochaj U, McMaster WR and Olivier M (2015) Impact of *Leishmania* infection on host macrophage nuclear physiology and nucleopore complex integrity. *PLoS Pathog* **11**, e1004776.
- 144 Guay-Vincent MM, Matte C, Berthiaume AM, Olivier M, Jaramillo M and Descoteaux A (2022) Revisiting *Leishmania* GP63 host cell targets reveals a limited spectrum of substrates. *PLoS Pathog* **18**, e1010640.
- 145 Huber S, Bar A, Epp S, Schmuckli-Maurer J, Eberhard N, Humbel BM, Hemphill A and Woods K (2020) Recruitment of host nuclear pore components to the vicinity of *Theileria* Schizonts. *mSphere* **5**, e00709-19.
- 146 UniProt-Consortium (2023) UniProt: the universal protein knowledgebase in 2023. *Nucleic Acids Res* **51**, D523–D531.
- 147 Gabler F, Nam SZ, Till S, Mirdita M, Steinegger M, Soding J, Lupas AN and Alva V (2020) Protein sequence analysis using the MPI bioinformatics toolkit. *Curr Protoc Bioinformatics* **72**, e108.
- 148 Blunt EL, Shandler JA, Hughes EJ, Sussman H, Christopherson RC and Richards EJ (2020) Coordination of NMCP1- and NMCP2-class proteins within the plant nucleoskeleton. *Mol Biol Cell* **31**, 2948–2958.
- 149 Burki F, Roger AJ, Brown MW and Simpson AGB (2020) The new tree of eukaryotes. *Trends Ecol Evol* **35**, 43–55.