

University of Groningen

Genomic profiling of post-transplant lymphoproliferative disorders using cell-free DNA

Veltmaat, Nick; Zhong, Yujie; Montes de Jesus, Filipe; Tan, Geok Wee; Bult, Johanna A.A.; Terpstra, Martijn M.; Mutsaers, Pim G.N.J.; Stevens, Wendy B.C.; Mous, Rogier; Vermaat, Joost S.P.

Published in:
Journal of Hematology and Oncology

DOI:
[10.1186/s13045-023-01500-x](https://doi.org/10.1186/s13045-023-01500-x)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2023

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Veltmaat, N., Zhong, Y., Montes de Jesus, F., Tan, G. W., Bult, J. A. A., Terpstra, M. M., Mutsaers, P. G. N. J., Stevens, W. B. C., Mous, R., Vermaat, J. S. P., Chamuleau, M. E. D., Noordzij, W., Verschuuren, E. A. M., Kok, K., Kluiver, J. L., Diepstra, A., Plattel, W. J., van den Berg, A., & Nijland, M. (2023). Genomic profiling of post-transplant lymphoproliferative disorders using cell-free DNA. *Journal of Hematology and Oncology*, 16, Article 104. <https://doi.org/10.1186/s13045-023-01500-x>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

CORRESPONDENCE

Open Access



Genomic profiling of post-transplant lymphoproliferative disorders using cell-free DNA

Nick Veltmaat¹, Yujie Zhong², Filipe Montes de Jesus³, Geok Wee Tan², Johanna A. A. Bult¹, Martijn M. Terpstra⁴, Pim G. N. J. Mutsaers⁵, Wendy B. C. Stevens⁶, Rogier Mous⁷, Joost S. P. Vermaat⁸, Martine E. D. Chamuleau⁹, Walter Noordzij³, Erik A. M. Verschuuren¹⁰, Klaas Kok⁴, Joost L. Kluiver², Arjan Diepstra², Wouter J. Plattel¹, Anke van den Berg² and Marcel Nijland^{1*}

Abstract

Diagnosing post-transplant lymphoproliferative disorder (PTLD) is challenging and often requires invasive procedures. Analyses of cell-free DNA (cfDNA) isolated from plasma is minimally invasive and highly effective for genomic profiling of tumors. We studied the feasibility of using cfDNA to profile PTLD and explore its potential to serve as a screening tool. We included seventeen patients with monomorphic PTLD after solid organ transplantation in this multi-center observational cohort study. We used low-coverage whole genome sequencing (lcWGS) to detect copy number variations (CNVs) and targeted next-generation sequencing (NGS) to identify Epstein-Barr virus (EBV) DNA load and somatic single nucleotide variants (SNVs) in cfDNA from plasma. Seven out of seventeen (41%) patients had EBV-positive tumors, and 13/17 (76%) had stage IV disease. Nine out of seventeen (56%) patients showed CNVs in cfDNA, with more CNVs in EBV-negative cases. Recurrent gains were detected for 3q, 11q, and 18q. Recurrent losses were observed at 6q. The fraction of EBV reads in cfDNA from EBV-positive patients was 3-log higher compared to controls and EBV-negative patients. 289 SNVs were identified, with a median of 19 per sample. SNV burden correlated significantly with lactate dehydrogenase levels. Similar SNV burdens were observed in EBV-negative and EBV-positive PTLD. The most commonly mutated genes were *TP53* and *KMT2D* (41%), followed by *SPEN*, *TET2* (35%), and *ARID1A*, *IGLL5*, and *PIM1* (29%), indicating DNA damage response, epigenetic regulation, and B-cell signaling/NFκB pathways as drivers of PTLD. Overall, CNVs were more prevalent in EBV-negative lymphoma, while no difference was observed in the number of SNVs. Our data indicated the potential of analyzing cfDNA as a tool for PTLD screening and response monitoring.

Keywords Post-transplant lymphoproliferative disorder, Cell-free DNA, Genomic profiling, Liquid biopsy, Epstein-Barr virus, Copy number variation, Single nucleotide variants, Low-coverage whole genome sequencing, Next generation sequencing

*Correspondence:

Marcel Nijland
m.nijland@umcg.nl

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

To the editor

Post-transplant lymphoproliferative disorder (PTLD) is a major complication after solid organ transplantation (SOT) [1]. While immunosuppressive therapy has been associated with early Epstein-Barr virus (EBV) driven PTLD, late-onset PTLD often lack EBV and have more genomic aberrations [2–4]. Diagnosis of PTLD is challenging due to its variable presentation. Serial monitoring of plasma EBV DNA levels and [18F]FDG PET/CT have limited sensitivity and specificity [5–7]. Based on recent successes in other B-cell lymphomas, analysis of plasma derived cell-free DNA (cfDNA) offers a promising minimally invasive approach for PTLD detection and disease monitoring [8, 9].

We investigated the feasibility of genomic profiling of PTLD in 17 patients with monomorphic PTLD by cfDNA analysis. Copy number variations (CNVs) were detected using low-coverage whole-genome sequencing (lcWGS). Targeted next-generation sequencing (NGS) was used for identifying EBV DNA load and somatic single nucleotide variants (SNVs) in cfDNA using a targeted panel including the EBV BamHI-W repeat region and *LMPI*, as well as the coding regions of 72 genes commonly mutated in B-cell lymphoma. SNVs and small insertions and deletions (indels) were called using an in-house pipeline. A detailed description of materials and methods can be found in Additional file 1.

Findings

The median age of the patients was 55 years (range 13–74). Median time between SOT and PTLD was 95 months (range 2–338). Most patients had stage IV disease ($n=13$, 76%) with a median metabolic tumor volume (MTV) of 302 mL (range 5–2070 mL). Lactate dehydrogenase (LDH) levels ranged from 210 to 5068 (Additional file 1: Table S1). PTLD tissue was EBV-positive in 7 out of 17 patients (41%). EBV copies in plasma quantified by qPCR were elevated (>5000 copies/mL) in 6 out of 15 (40%) evaluable patients.

The mean cfDNA yield of patients with PTLD (666 ng/mL, range 3–6049) was significantly higher compared to controls (21 ng/mL, range 6–54; $p=0.01$) (Additional file 1: Figure S2A), with the observed range of cfDNA levels in PTLD patients being consistent with observations in other malignancies [10]. In PTLD patients a moderate correlation was observed between cfDNA levels and MTV ($\rho=0.53$, $p=0.036$) and LDH levels ($\rho=0.57$, $p=0.019$) (Additional file 1: Figures S2B–C).

CNV analysis failed in 1 patient due to insufficient sequencing reads. CNVs were detected in cfDNA in 9 out of the 16 (56%) patients. The most frequent gains

involved 3q, 11q,18q and chromosome 21, while the most frequently lost region was 6q (Fig. 1A). lcWGS analysis of matched tumor samples revealed CNVs in all 5 patients. In general, more CNVs with higher amplitudes of gains/losses were observed in tissue as compared to the matched cfDNA samples, corresponding with a lower estimated tumor fraction (ETF) in cfDNA samples compared to tissue (Additional file 1: Figure S3).

The ETF was significantly correlated to LDH levels, but not to MTV (Fig. 1B, C). The number of CNVs per patient was much higher in EBV-negative patients compared to EBV-positive patients. This resulted in a higher mean fraction of genome altered (FGA) in EBV-negative tumors compared to EBV-positive tumors (0.152 vs 0.067), although not statistically significant ($p=0.095$) (Fig. 1D).

The median percentage of EBV reads (0.53%) was significantly higher (approximately 3-logs) in EBV-positive tumors, compared to EBV-negative tumors and controls ($p<0.001$). No significant difference was observed between EBV-negative PTLD and controls. A percentage of EBV reads above 0.0012% was indicative of an EBV-positive PTLD at the time of diagnosis (Additional file 1: Figure S4A). Six of the seven patients with EBV-positive tumors were tested positive in the diagnostic qPCR test. We observed a concordance between qPCR results and the EBV load as determined by NGS (Additional file 1: Figure S4B).

A total of 289 SNVs/InDels were identified in the 17 plasma samples with a median of 19 SNVs per sample (range 1–37). The number of SNVs was significantly correlated to LDH, but not MTV (Fig. 2B, C). The most frequently mutated genes were *TP53* and *KMT2D* (7/17, 41%), *SPEN* and *TET2* (6/17 cases (35%)), followed by *ARID1A*, *IGLL5* and *PIMI* (5/17, 29%) (Fig. 2A). We observed no difference in SNV burden and affected genes between EBV-negative and EBV-positive cases (Fig. 2D). Mutated genes and pathways in our study overlapped with findings of previously reported genes & pathways in PTLD [11].

In conclusion, the data of this study highlights the use of genomic profiling of plasma cfDNA analysis in patients with PTLD as a minimally invasive tool for potential screening strategies, genomic profiling and response monitoring. CNVs were successfully detected using lcWGS, while EBV status and the tumor mutational landscape could be captured using targeted NGS. EBV-negative PTLD had more CNVs compared to EBV-positive cases, suggesting a higher degree of genomic instability. Consequently, sequential EBV detection by EBV PCR and/or SNV analysis is the most suitable screening strategy for EBV-positive PTLD, while CNV and/or SNV profiling would be a good screening strategy




Fig. 1 Copy number aberrations found in cell-free DNA from PTLD patients. **A** Overview of copy number variations (CNVs) for individual PTLD patients, grouped by Epstein–Barr virus (EBV) status and sorted on fraction of genome altered (FGA). Chromosomal regions with gains are indicated in red and losses in blue. **B** A significant correlation between estimated tumor fraction (ETF) and lactate dehydrogenase (LDH) was observed. Dashed line represents cut-off value at 248 U/L discriminating clinically elevated LDH from normal LDH value. **C** Correlation between ETF and metabolic tumor volume (MTV). In panels **B** and **C**, the grey areas around the regression lines represent 95% CI and the Spearman coefficient is indicated with ρ . **D** FGA in PTLD patients categorized by Epstein–Barr virus (EBV) status, as determined by EBER-ISH, shows that EBV-negative patients have a higher FGA, although insignificant according to Wilcoxon Signed Rank test. Only PTLD samples with CNVs (FGA > 0) are shown in **B–D**

for EBV-negative cases. The utility of SNV analyses could contribute to tumor typing at diagnosis and response assessment. This study presents the first cfDNA analysis for PTLD, with limitations of small sample size and lack

of tissue biopsies in some patients. The value of ctDNA dynamics in a larger patient PTLD cohort is part of the ongoing observational NTR 7402 study [12].




Fig. 2 Single nucleotide variants in genes found in cfDNA of PTLD patients. **A** Waterfall plot displaying types of mutations in each plasma sample for each gene. Top- and right-side bar plots show total number of mutations found in a sample (tumor mutation burden, TMB) or a gene, respectively. Genes are sorted based on this number. Samples are grouped by Epstein Barr virus (EBV) status. EBV status, Ann-Arbor staging, Estimated tumor fraction (ETF) and fraction of genome altered (FGA) information is shown below. Pathway information per gene is indicated by color, left of the gene names. **B** A significant correlation between SNV load and lactate dehydrogenase (LDH) was observed. Dashed line represents cut-off value at 248 U/L discriminating clinically elevated LDH from normal LDH value. **C** Correlation between SNV load and metabolic tumor volume (MTV). In panels B and C, the grey areas around the regression lines represent the 95% CI and the Spearman coefficient is indicated with ρ . **D** The total number of SNVs per sample is shown, grouped by EBV status. EBV-negative samples show a slightly higher mean, though this difference is insignificant as tested by Wilcoxon Signed Rank test

Abbreviations

cfDNA Cell-free DNA
 CNV Copy number variation
 EBER ISH Epstein-Barr encoding region specific RNA in situ hybridization
 EBV Epstein-Barr virus
 ETF Estimated tumor fraction
 FGA Fraction of the genome altered
 IcWGS Low coverage whole genome sequencing

LDH Lactate dehydrogenase
 MTV Metabolic tumor volume
 NGS Next generation sequencing
 PTLD Post-transplant lymphoproliferative disorder
 qPCR Quantitative polymerase chain reaction
 R-CHOP Rituximab, cyclophosphamide, doxorubicin, vincristine, prednisone
 SNV Single nucleotide variants

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13045-023-01500-x>.

Additional file 1. Supplementary methods.

Acknowledgements

We thank the Center for Information Technology of the University of Groningen for their support and for providing access to the Gearshift high performance computing cluster.

Author contributions

NV wrote the manuscript, performed bio-informatic analyses on NGS data and interpreted the results. FMJ, GT, YZ contributed to manuscript text. MN, AvdB, AD, WP and JLK. revised and commented on the manuscript. FMJ analyzed [¹⁸F]FDG PET/CT scans. GT performed wet-lab experiments. YZ and MMT helped with ICGWS data analysis. PGNJM, WBSC, JSPV, RM, MEDC, EAMV and MN selected patients for this study. AD performed histological examination. MN supervised the project. All authors read and approved the final manuscript.

Funding

This work was supported by the Anita Veldman Foundation.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The study was conducted in accordance with the ethical principles of the Declaration of Helsinki and with the approval of the Medical Ethics Review Board of the University Medical Center Groningen (2018/437).

Consent for publication

Informed consent was waived for all retrospective patients and obtained for all prospectively included patients.

Competing interests

The authors declare no competing interests.

Author details

¹Department of Hematology, University Medical Centre Groningen, University of Groningen, Groningen, The Netherlands. ²Department of Pathology and Medical Biology, University Medical Centre Groningen, University of Groningen, Groningen, The Netherlands. ³Medical Imaging Center, Department of Nuclear Medicine and Molecular Imaging, University Medical Centre Groningen, University of Groningen, Groningen, The Netherlands. ⁴Department of Genetics, University Medical Centre Groningen, University of Groningen, Groningen, The Netherlands. ⁵Department of Hematology, Erasmus Medical Centre Rotterdam, Rotterdam, The Netherlands. ⁶Department of Hematology, Radboud University Medical Center Nijmegen, Nijmegen, The Netherlands. ⁷Department of Hematology, University Medical Center Utrecht, Utrecht, The Netherlands. ⁸Department of Hematology, Leiden University Medical Center, Leiden, The Netherlands. ⁹Department of Hematology, Amsterdam University Medical Center, Amsterdam, The Netherlands. ¹⁰Department of Pulmonology, University Medical Centre Groningen, University of Groningen, Groningen, The Netherlands.

Received: 26 July 2023 Accepted: 11 September 2023

Published online: 14 September 2023

References

- Francis A, Johnson DW, Teixeira-Pinto A, Craig JC, Wong G. Incidence and predictors of post-transplant lymphoproliferative disease after kidney

- transplantation during adulthood and childhood: a registry study. *Nephrol Dial Transplant*. 2018;33(5):881–9.
- Markouli M, Ullah F, Omar N, Apostolopoulou A, Dhillon P, Diamantopoulos P, et al. Recent advances in adult post-transplant lymphoproliferative disorder. *Cancers (Basel)*. 2022;14(23):5949.
- Morscio J, Finalet Ferreiro J, Vander Borghet S, Bittoun E, Gheysens O, Dierickx D, et al. Identification of distinct subgroups of EBV-positive post-transplant diffuse large B-cell lymphoma. *Mod Pathol*. 2017;30(3):370–81.
- Morscio J, Dierickx D, Ferreiro JF, Herreman A, Van Loo P, Bittoun E, et al. Gene expression profiling reveals clear differences between EBV-positive and EBV-negative posttransplant lymphoproliferative disorders. *Am J Transplant*. 2013;13(5):1305–16.
- San-Juan R, Manuel O, Hirsch HH, Fernández-Ruiz M, López-Medrano F, Comoli P, et al. Current preventive strategies and management of Epstein-Barr virus-related post-transplant lymphoproliferative disease in solid organ transplantation in Europe. Results of the ESGICH Questionnaire-based Cross-sectional Survey. *Clin Microbiol Infect*. 2015;21(6):604.
- Dierickx D, Habermann TM. Post-Transplantation Lymphoproliferative Disorders in Adults. *N Engl J Med*. 2018;378(6):549–62.
- Dharnidharka VR, Green M, Webber SA. Post-transplant lymphoproliferative disorders. Berlin: Springer; 2010.
- Rossi D, Diop F, Spaccarotella E, Monti S, Zanni M, Rasi S, et al. Diffuse large B-cell lymphoma genotyping on the liquid biopsy. *Blood*. 2017;129(14):1947–57.
- Spina V, Brusca A, Cuccaro A, Martini M, Di Trani M, Forestieri G, et al. Circulating tumor DNA reveals genetics, clonal evolution, and residual disease in classical Hodgkin lymphoma. *Blood*. 2018;131(22):2413–25.
- Yan Y, Guo Q, Wang F, Adhikari R, Zhu Z, Zhang H, et al. Cell-Free DNA: hope and potential application in cancer. *Front Cell Dev Biol*. 2021;9:639233.
- Menter T, Juskevicius D, Alikian M, Steiger J, Dirnhofer S, Tzankov A, et al. Mutational landscape of B-cell post-transplant lymphoproliferative disorders. *Br J Haematol*. 2017;178(1):48–56.
- Montes de Jesus F, Noordzij W, Glaudemans AWJM, Nijland M. Respon-evaluatie bij posttransplantatielymfomen door middel van 18F-FDG-PET/CT en circulerend tumor-DNA (MRD-PTLD). *Nederlands Tijdschrift voor Hematologie*. 2019;16(6):301–3.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

