# Erratum to: Transverse dental arch relationship at 9 and 12 years in children with unilateral cleft lip and palate treated with infant orthopedics: a randomized clinical trial (DUTCHCLEFT)

Noverraz, R.L.M.; Disse, M.A.; Ongkosuwito, E.M.; Kuijpers-Jagtman, A.M.; Prahl, C.

*Published in:*
Clinical Oral Investigations

*DOI:*
10.1007/s00784-015-1529-x

Clin Oral Invest (2015) 19:2351
DOI 10.1007/s00784-015-1529-x

ERRATUM

# Erratum to: Transverse dental arch relationship at 9 and 12 years in children with unilateral cleft lip and palate treated with infant orthopedics: a randomized clinical trial (DUTCHCLEFT)

**R. L. M. Noverraz[1] · M. A. Disse[1] · E. M. Ongkosuwito[2] · A. M. Kuijpers-Jagtman[2] · C. Prahl[1]**

Published online: 16 August 2015


**Erratum to: Clinical Oral Investigations**
**DOI 10.1007/s00784-015-1451-2**

The original version of the above article contained a mistake due to incorrect alignment of entries in Tables 11 and 14 during production. The entries are presented correctly below.

**Table 11** Evaluation of the total arch constriction in pairs

| | Center | Center | Mean | 95 % CI | | *P* |
|---|---|---|---|---|---|---|
| | | | | Lower | Upper | |
| 9 years | 1 | 2 | 1.63 | −2.09 | 5.34 | 0.842 |
| | 1 | 3 | −1.71 | −5.61 | 2.18 | 0.836 |
| | 2 | 3 | −3.34 | −6.73 | .05 | 0.054 |
| 12 years | 1 | 2 | 4.85 | 0.53 | 9.17 | 0.023* |
| | 1 | 3 | −4.39 | −4.02 | 4.95 | 1.000 |
| | 2 | 3 | 0.47 | −8.33 | −0.44 | 0.025* |

*Mean* mean total arch constriction score (points), 95*% CI* 95 % confidence interval of the difference

*$p<0.05$

**Table 14** Evaluation of age of hard palate closure with alveolar bone grafting in pairs

| Center | Center | Mean diff | 95 % CI | | *P* |
|---|---|---|---|---|---|
| | | | Lower | Upper | |
| 1 | 2 | −1.55 | −2.43 | 0.68 | 0.000*** |
| 1 | 3 | −0.99 | −1.88 | −0.11 | 0.023 |
| 2 | 3 | 0.56 | −0.20 | 1.32 | 0.218 |

*Mean diff* mean difference in age of hard palate closure with alveolar bone grafting (years), 95*% CI* 95 % confidence interval of the difference

***$p<0.001$

The online version of the original article can be found at http://dx.doi.org/10.1007/s00784-015-1451-2.

✉ C. Prahl
c.prahl@acta.nl

[1] Department of Orthodontics, Academic Centre for Dentistry Amsterdam (ACTA), Gustav Mahlerlaan 3004, 1081 LA Amsterdam, The Netherlands

[2] Department of Orthodontics and Craniofacal Biology and Cleft Palate Craniofacial Centre, Radboud University Medical Centre, Nijmegen, The Netherlands