


1-1-1998

## Some root invariants and Steenrod operations in $\text{Ext}_A(F_2, F_2)$

Robert R. Bruner

Wayne State University, robert.bruner@wayne.edu

Follow this and additional works at: <https://digitalcommons.wayne.edu/mathfrp>

 Part of the [Algebraic Geometry Commons](#), and the [Geometry and Topology Commons](#)

---

### Recommended Citation

Bruner, R. R. Some root invariants and Steenrod operations in  $\text{Ext}_A(F_2, F_2)$ . In M. Mahowald, S. Priddy, Eds. Homotopy Theory via Algebraic Geometry and Group Representations. Contemp. Math. 220 (1998), 27-33. Providence, RI: American Mathematical Society.

This Article is brought to you for free and open access by the Mathematics at DigitalCommons@WayneState. It has been accepted for inclusion in Mathematics Faculty Research Publications by an authorized administrator of DigitalCommons@WayneState.

# Some root invariants and Steenrod operations in $\text{Ext}_A(F_2, F_2)$

ROBERT R. BRUNER

ABSTRACT. We give the results of computations of root invariants in  $\text{Ext}$  over the Steenrod algebra through the 25-stem, with partial information through the 45-stem. This allows the computation of some new Steenrod operations as well.

These calculations were started to provide evidence for or against the algebraic Bredon-Löffler conjecture. This is the conjecture that

$$\eta_k^* : \text{Ext}_A^{s,t}(F_2, F_2) \longrightarrow \text{Ext}_A^{s,t}(\Sigma L_{-k}, F_2)$$

is a monomorphism for  $0 < t - s < k/2$ . Here  $L$  is the Laurent series ring  $F_2[x, x^{-1}]$  with its usual Steenrod algebra action,  $L_{-k}$  is the submodule of  $L$  consisting of those elements whose degree is greater than or equal to  $-k$ , and  $\eta_k : \Sigma L_{-k} \longrightarrow F_2$  is the nonzero homomorphism. The calculations of  $\eta_k^*$  showed [5] that the conjecture holds for  $k \leq 55$ . With those calculations in hand, we needed only calculate the homomorphisms

$$i_k^* : \text{Ext}_A^{s,t}(\Sigma^{1-k} F_2, F_2) \longrightarrow \text{Ext}_A^{s,t}(\Sigma L_{-k}, F_2)$$

induced by the projections onto the bottom dimensions,  $i_k : \Sigma L_{-k} \longrightarrow \Sigma^{1-k} F_2$ , to calculate all the root invariants through the 25 stem, and about half of those in the 26 through 45 stems. This extends the calculations of Mahowald and Shick [7], who found all the root invariants through the 16 stem. Our calculations were done by first using the programs described in [1, 2] to compute resolutions of the modules  $L_{-k}$  and the chain maps induced by  $\eta_k$  and  $i_k$ . We then used MAGMA to compute the induced maps of  $\text{Ext}$  and the root invariants. A calculation of  $\text{Ext}_A(F_2, F_2)$  for  $s < 40$  and  $t \leq 140$  with complete information on the product structure was calculated in the same manner: using MAGMA to process the

---

1991 *Mathematics Subject Classification*. Primary 55Q45, 55Q35, 55Q91; Secondary 55P42, 55P91, 55T15 .

output of the minimal resolution programs [1, 2]. A preliminary version of this can be found in [3], where the notation for elements of  $\text{Ext}_A(F_2, F_2)$  used here is defined. Charts showing the results of the Ext calculations (but not the induced homomorphisms) can be found at <http://www.math.wayne.edu/~rrb/cohom>.

By [6, Proposition 2.5], we may conclude that  $Sq^0(x) \in R(x)$  when  $|Sq^0(x)| = |R(x)|$ , that is, if  $k = n + s + 1$ . This allows us to use the calculations here to determine a number of Steenrod operations.

**THEOREM 1.** *In the cohomology of the mod 2 Steenrod algebra*

- (i)  $Sq^0(Ph_1) = h_2g$
- (ii)  $Sq^0(Pc_0) = h_1y$
- (iii)  $Sq^0(Pd_0) = gd_1$
- (iv)  $Sq^0(i) = h_2C$
- (v)  $Sq^0(k) = h_2h_5n + \epsilon h_3D_2$ , where  $\epsilon \in \{0, 1\}$
- (vi)  $Sq^0(q) = h_2Q_3$
- (vii)  $Sq^0(l) = h_0x_{6,47}$
- (viii)  $Sq^0(y) = h_4Q_3 + h_2x_1$

These are the  $Sq^0$ 's which are not 'tautologous'. In addition, we have 12 more which allow us to identify, in the resolution produced by the machine, the elements named. That is, we know by definition that  $e_1 = Sq^0(e_0)$ , for example. However, we have no way *a priori* to determine which of two elements in the 38 stem is  $e_1$  since we have no general mechanical method of computing Steenrod operations. Thus, the root invariant is quite useful, in that it unambiguously identifies  $e_1$  for us. The elements identified in this manner are  $n_1$ ,  $p_1$ ,  $e_1$ ,  $e_2$ ,  $f_1$ , and  $f_2$ . We also determine  $m_1$  and  $t_1$  up to  $F_2$  indeterminacy.

Finally, note that the root invariant  $R(h_2^2g) = r_1$  shows that the Strong Algebraic Bredon-Löffler Conjecture in [4] must have its  $(t - s)$ -intercept decreased by at least 3. This is an inessential modification, as the key observation in these calculations supporting the conjecture is the slope  $-1/2$ .

**CONJECTURE 2.** (*Strong Algebraic Bredon-Löffler Conjecture*) *The map  $\eta_k^*$  is a monomorphism if*

$$s < (k - n - 3)/2.$$

**KEY TO THE TABLES:** Each row in the table lists the filtration  $s$  and stem  $n$  of an element  $x \in \text{Ext}_A(F_2, F_2)$ , its root invariant  $R(x)$ , the dimension  $k$  of the cell on which it occurs, whether  $Sq^0(x) \in R(x)$ , and a basis for the indeterminacy in  $R(x)$ . The root invariant  $R(x)$  is in the  $n + k - 1$  stem. By [6, Proposition 2.5], we may test whether or not  $Sq^0(x) \in R(x)$  by checking whether or not  $|Sq^0(x)| = |R(x)|$ , that is,  $k = n + s + 1$ .

$k$	$s$	$n$	$x$	$R(x)$	$Sq^0 \in R(x)?$	Indeterminacy
2	1	0	$h_0$	$h_1$	yes	
3	2	0	$h_0^2$	$h_1^2$	yes	
4	3	0	$h_0^3$	$h_0^2 h_2$	yes	
8	4	0	$h_0^4$	$h_0^3 h_3$	no	
10	5	0	$h_0^5$	$Ph_1$	no	
11	6	0	$h_0^6$	$h_1 Ph_1$	no	
12	7	0	$h_0^7$	$h_0^2 Ph_2$	no	
16	8	0	$h_0^8$	$h_0^7 h_4$	no	
18	9	0	$h_0^9$	$P^2 h_1$	no	
19	10	0	$h_0^{10}$	$h_1 P^2 h_1$	no	
20	11	0	$h_0^{11}$	$h_0^2 P^2 h_2$	no	
24	12	0	$h_0^{12}$	$h_0^5 i$	no	
26	13	0	$h_0^{13}$	$P^3 h_1$	no	
27	14	0	$h_0^{14}$	$h_1 P^3 h_1$	no	
28	15	0	$h_0^{15}$	$h_0^2 P^3 h_2$	no	
32	16	0	$h_0^{16}$	$h_0^{15} h_5$	no	
34	17	0	$h_0^{17}$	$P^4 h_1$	no	
35	18	0	$h_0^{18}$	$h_1 P^4 h_1$	no	
36	19	0	$h_0^{19}$	$h_0^2 P^4 h_2$	no	
3	1	1	$h_1$	$h_2$	yes	
5	2	2	$h_1^2$	$h_2^2$	yes	
5	1	3	$h_2$	$h_3$	yes	
6	2	3	$h_0 h_2$	$h_1 h_3$	yes	
7	3	3	$h_0^2 h_2$	$h_1^2 h_3$	yes	
9	2	6	$h_2^2$	$h_3^2$	yes	
9	1	7	$h_3$	$h_4$	yes	
10	2	7	$h_0 h_3$	$h_1 h_4$	yes	
11	3	7	$h_0^2 h_3$	$h_1^2 h_4$	yes	
12	4	7	$h_0^3 h_3$	$h_0^2 h_2 h_4$	yes	
11	2	8	$h_1 h_3$	$h_2 h_4$	yes	
12	3	8	$c_0$	$c_1$	yes	
13	3	9	$h_1^2 h_3$	$h_2^2 h_4$	yes	
14	4	9	$h_1 c_0$	$h_2 c_1$	yes	
15	5	9	$Ph_1$	$h_2 g$	yes	
21	6	10	$h_1 Ph_1$	$r$	no	
20	5	11	$Ph_2$	$h_0^3 h_4^2$	no	
22	6	11	$h_0 Ph_2$	$q$	no	
23	7	11	$h_0^2 Ph_2$	$h_1 q$	no	
17	2	14	$h_3^2$	$h_4^2$	yes	
18	3	14	$h_0 h_3^2$	$h_1 h_4^2$	yes	
19	4	14	$d_0$	$d_1$	yes	

$k$	$s$	$n$	$x$	$R(x)$	$Sq^0 \in R(x)?$	Indeterminacy
20	5	14	$h_0d_0$	$h_0p$	yes	
25	6	14	$h_0^2d_0$	$y$	no	$h_1x$
17	1	15	$h_4$	$h_5$	yes	
18	2	15	$h_0h_4$	$h_1h_5$	yes	
19	3	15	$h_0^2h_4$	$h_1^2h_5$	yes	
20	4	15	$h_0^3h_4$	$h_0^2h_2h_5$	yes	
21	5	15	$h_1d_0$	$h_2d_1$	yes	
24	5	15	$h_0^4h_4$	$h_0^3h_3h_5$	no	
26	6	15	$h_0^5h_4$	$h_5Ph_1$	no	$h_0^2f_1$
27	7	15	$h_0^6h_4$	$h_1h_5Ph_1$	no	
28	8	15	$h_0^7h_4$	$h_0^2h_5Ph_2$	no	
19	2	16	$h_1h_4$	$h_2h_5$	yes	
23	6	16	$h_1^2d_0$	$h_1x$	yes	
24	7	16	$Pc_0$	$h_1y$	yes	
21	3	17	$h_1^2h_4$	$h_2^2h_5$	yes	
22	4	17	$e_0$	$e_1$	yes	
23	5	17	$h_0e_0$	$h_1e_1$	yes	
24	6	17	$h_0^2e_0$	$h_0^2f_1$	yes	
31	7	17	$h_0^3e_0$	$h_1^2h_5d_0$	no	
30	8	17	$h_1Pc_0$	$N$	no	
32	9	17	$P^2h_1$	$h_1h_5Pc_0$	no	
21	2	18	$h_2h_4$	$h_3h_5$	yes	
22	3	18	$h_0h_2h_4$	$h_1h_3h_5$	yes	
23	4	18	$f_0$	$f_1$	yes	
23	4	18	$h_0^2h_2h_4$	$h_1^2h_3h_5$	yes	
24	5	18	$h_0f_0$	$h_0^2c_2$	yes	
37	10	18	$h_1P^2h_1$	$R_1$	no	$h_0^2h_5i$
23	3	19	$c_1$	$c_2$	yes	
35	9	19	$P^2h_2$	$h_5Pd_0$	no	
36	10	19	$h_0P^2h_2$	$h_0^2h_5i$	no	
39	11	19	$h_0^2P^2h_2$	$h_1Q_1$	no	
25	4	20	$g$	$g_2$	yes	
26	5	20	$h_0g$	$h_1g_2$	yes	
28	6	20	$h_0^2g$	$h_0h_2g_2$	no	
25	3	21	$h_2^2h_4$	$h_3^2h_5$	yes	
27	5	21	$h_1g$	$h_2g_2$	yes	
27	4	22	$h_2c_1$	$h_3c_2$	yes	
31	8	22	$Pd_0$	$gd_1$	yes	
36	9	22	$h_0Pd_0$	$h_0^2Q_2$	no	
41	10	22	$h_0^2Pd_0$	$x_{10,27}$	no	$x_{10,28}, h_1X_1$
28	4	23	$h_4c_0$	$h_5c_1$	yes	

$k$	$s$	$n$	$x$	$R(x)$	$Sq^0 \in R(x)?$	Indeterminacy
29	5	23	$h_2g$	$h_3g_2$	yes	
30	6	23	$h_0h_2g$	$h_1h_3g_2$	yes	
31	7	23	$i$	$h_2C$	yes	
36	8	23	$h_0i$	$h_0^2D_2$	no	
38	9	23	$h_0^2i$	$h_0^2B_3$	no	
40	9	23	$h_0^2i + h_1Pd_0$	$h_0^7h_5^2$	no	
42	10	23	$h_0^3i$	$x_{10,32}$	no	$h_0^2h_3Q_2$
43	11	23	$h_0^4i$	$h_1x_{10,32}$	no	$h_0B_{23}$
44	12	23	$h_0^5i$	$h_0^2PD_2$	no	
30	5	24	$h_1h_4c_0$	$h_2h_5c_1$	yes	
43	10	24	$h_1^2Pd_0$	$B_5 + PD_2$	no	
44	11	24	$P^2c_0$	$x_{11,35} + h_0^2x_{9,40}$	no	
38	8	25	$Pe_0$	$x_{8,32} + x_{8,33}$	no	
39	9	25	$h_0Pe_0$	$h_1x_{8,32}$	no	$h_0^2x_{7,33}, h_0^8h_6$
40	10	25	$h_0^2Pe_0$	$h_0^2h_3Q_2$	no	
45	11	25	$h_0^3Pe_0$	$h_2B_5$	no	$h_0PA$
46	12	25	$h_1P^2c_0$	$h_2x_{11,35}$	no	
47	13	25	$P^3h_1$	$x_{13,34}$	no	$x_{13,35}$
41	6	26	$h_2^2g$	$r_1$	no	
36	7	26	$j$	$h_0A'$	no	
39	8	26	$h_0j$	$h_3Q_2$	no	$h_0h_2A, h_0^2A''$
40	9	26	$h_0^2j$	$h_0^2h_3D_2$	no	
51	13	27	$P^3h_2$	$e_0B_4$	no	$h_0^6x_{7,57}$
41	8	28	$d_0^2$	$G_{21}$	no	$h_0h_3A'$
42	9	28	$h_0d_0^2$	$h_1G_{21}$	no	$h_0h_2x_{7,40}$
44	10	28	$h_0^2d_0^2$	$h_0h_2G_{21}$	no	
37	7	29	$k$	$h_2h_5n$	yes	$h_3D_2$
40	8	29	$h_0k$	$h_0h_3A'$	no	
43	9	29	$h_0^2k$	$h_2G_{21}$	no	
33	2	30	$h_4^2$	$h_5^2$	yes	
34	3	30	$h_0h_4^2$	$h_1h_5^2$	yes	
35	4	30	$h_0^2h_4^2$	$h_1^2h_5^2$	yes	
36	5	30	$h_0^3h_4^2$	$h_0^2h_2h_5^2$	yes	
37	6	30	$r$	$r_1$	yes	
39	7	30	$h_0r$	$h_2^2H_1 + h_3A'$	no	
41	8	30	$h_0^2r$	$h_3x_{7,33}$	no	
44	9	30	$h_0^3r$	$h_0^2h_4D_2$	no	
49	10	30	$h_0^4r$	$P^2h_5^2$	no	$e_0A'$
50	11	30	$h_0^5r$	$h_1P^2h_5^2$	no	
51	12	30	$P^2d_0$	$x_{12,44}$	no	
33	1	31	$h_5$	$h_6$	yes	

$k$	$s$	$n$	$x$	$R(x)$	$Sq^0 \in R(x)?$	Indeterminacy
34	2	31	$h_0h_5$	$h_1h_6$	yes	
35	3	31	$h_0^2h_5$	$h_1^2h_6$	yes	
35	3	31	$h_1h_4^2$	$h_2h_5^2$	yes	
36	4	31	$h_0^3h_5$	$h_0^2h_2h_6$	yes	
37	5	31	$n$	$n_1$	yes	
40	5	31	$h_0^4h_5$	$h_0^3h_3h_6$	no	
42	6	31	$h_0^5h_5$	$h_6Ph_1$	no	
43	7	31	$h_0^6h_5$	$h_1h_6Ph_1$	no	$h_2^2Q_3$
40	8	31	$d_0e_0$	$h_3x_{7,33}$	yes	
44	8	31	$h_0^7h_5$	$h_0^6h_6Ph_2$	no	
45	9	31	$h_0d_0e_0$	$h_3G_{21}$	no	$h_0^2x_{7,53}$
48	9	31	$h_0^8h_5$	$h_0^7h_4h_6$	no	
46	10	31	$h_0^2d_0e_0$	$h_0x_{9,51}$	no	
50	10	31	$h_0^9h_5$	$h_6P^2h_1$	no	
51	11	31	$h_0^{10}h_5$	$h_1h_6P^2h_1$	no	$h_0gA'$
35	2	32	$h_1h_5$	$h_2h_6$	yes	
37	4	32	$d_1$	$d_2$	yes	
39	6	32	$q$	$h_2Q_3$	yes	
40	7	32	$l$	$h_0x_{6,47}$	yes	
43	8	32	$h_0l$	$x_{8,51}$	no	$h_0^2h_6Ph_2$
44	9	32	$h_0^2l$	$h_0^2x_{7,53}$	no	
37	3	33	$h_1^2h_5$	$h_2^2h_6$	yes	
38	4	33	$p$	$p_1$	yes	
39	5	33	$h_0p$	$h_1p_1$	yes	
45	7	33	$h_1q$	$x_{7,57}$	no	$h_1x_{6,53}, h_0^2h_6d_0$
37	2	34	$h_2h_5$	$h_3h_6$	yes	
38	3	34	$h_0h_2h_5$	$h_1h_3h_6$	yes	
39	4	34	$h_0^2h_2h_5$	$h_1^2h_3h_6$	yes	
41	6	34	$h_2n$	$h_3n_1$	yes	$h_6Ph_2$
43	8	34	$d_0g$	$h_1x_{7,53}$	yes	$h_0^2x_{6,53}$
44	9	34	$h_0d_0g$	$h_0^2m_1$	yes	
48	11	34	$Pj$	$h_0gA'$	no	
41	5	35	$h_2d_1$	$h_3d_2$	yes	
43	7	35	$m$	$m_1$	yes	$h_0^2h_6d_0$
44	8	35	$h_0m$	$h_0^2t_1$	yes	
47	9	35	$h_0^2m$	$h_3x_{8,51}$	no	
43	6	36	$t$	$t_1$	yes	$h_0^4h_4h_6$
41	3	37	$h_2^2h_5$	$h_3^2h_6$	yes	
43	5	37	$x$	$x_1$	yes	
44	6	37	$h_0x$	$h_0^2e_2$	yes	
45	7	37	$h_1t$	$h_2t_1$	yes	

$k$	$s$	$n$	$x$	$R(x)$	$Sq^0 \in R(x)?$	Indeterminacy
46	7	37	$h_0^2x$	$h_0h_2x_1$	no	
46	8	37	$e_0g$	$e_1g_2$	yes	
48	8	37	$h_0^3x$	$h_0^4f_2$	no	
41	2	38	$h_3h_5$	$h_4h_6$	yes	
42	3	38	$h_0h_3h_5$	$h_1h_4h_6$	yes	
43	4	38	$h_0^2h_3h_5$	$h_1^2h_4h_6$	yes	
43	4	38	$e_1$	$e_2$	yes	
44	5	38	$h_0^3h_3h_5$	$h_0^2h_2h_4h_6$	yes	
45	6	38	$h_1x$	$h_2x_1$	yes	
45	6	38	$y$	$h_4Q_3 + h_2x_1$	yes	
46	7	38	$h_0y$	$h_1h_4Q_3$	yes	$h_0^2h_6g$
48	8	38	$h_0^2y$	$h_0^5c_3$	no	
43	3	39	$h_1h_3h_5$	$h_2h_4h_6$	yes	
44	4	39	$h_5c_0$	$h_6c_1$	yes	
45	5	39	$h_1e_1$	$h_2e_2$	yes	
49	7	39	$h_1y$	$x_{7,74}$	no	
45	4	40	$h_1^2h_3h_5$	$h_2^2h_4h_6$	yes	
45	4	40	$f_1$	$f_2$	yes	
46	5	40	$h_0f_1$	$h_1f_2$	yes	
46	5	40	$h_1h_5c_0$	$h_2h_6c_1$	yes	
47	6	40	$h_0^2f_1$	$h_1^2f_2$	yes	
47	6	40	$h_5Ph_1$	$h_2h_6g$	yes	
45	3	41	$c_2$	$c_3$	yes	
46	4	41	$h_0c_2$	$h_1c_3$	yes	
47	5	41	$h_0^2c_2$	$h_1^2c_3$	yes	
49	4	44	$g_2$	$g_3$	yes	
49	3	45	$h_3^2h_5$	$h_4^2h_6$	yes	

## REFERENCES

1. Robert R. Bruner, "Calculation of large Ext modules", *Computers in Geometry and Topology* (M. C. Tangora, ed.), Marcel Dekker, New York, 1989, 79-104.
2. R.R.Bruner, "Ext in the nineties", pp. 71-90 in *Algebraic Topology*, Oaxtepec 1991, Contemp. Math. **146**, Amer. Math. Soc., Providence, 1993.
3. R.R.Bruner, "The Cohomology of the Mod 2 Steenrod Algebra", preprint, April 1997.
4. Robert R. Bruner, "Some remarks on the root invariant", *Proceedings of the Fields Institute Workshops on Homotopy Theory, January 1996*, Contemp. Math., to appear.
5. Robert Bruner and John Greenlees, "The Bredon-Löffler conjecture", *Exper. Math.* **4** (1995), 101-109.
6. M.E.Mahowald and D.C.Ravenel, "The root invariant in homotopy theory", *Topology* **32** (1993), 865-898.
7. M.E.Mahowald and Paul Shick, "Some root invariants in  $\text{Ext}_A(Z/2, Z/2)$ ", *Proceedings of the Northwestern Homotopy Theory Conference*, Contemp. Math. **19** (1983), 227-231.



MATHEMATICS DEPARTMENT, WAYNE STATE UNIVERSITY, DETROIT, MICHIGAN, 48202  
*E-mail address:* rrb@math.wayne.edu