



Corrigendum

Corrigendum to “APP-dependent alteration of GSK3 β activity impairs neurogenesis in the Ts65Dn mouse model of Down syndrome”
[Neurobiology of Disease 67 (2014) 24–36]

Stefania Trazzi, Claudia Fuchs, Marianna De Franceschi, Valentina Maria Mitrugno, Renata Bartesaghi, Elisabetta Ciani*

Department of Biomedical and Neuromotor Sciences, University of Bologna, 40126 Bologna, Italy



The authors regret that in this article, there was an unconscious error during image saving and subsequent preparation of Fig. 7A. Please find enclosed the correct images showing GSK3 β phosphorylation levels in the hippocampus of a control (left panel) and a DS (right panel) fetus.

A




Fig. 7. Altered GSK3 β phosphorylation in human fetuses with DS. **A**: Examples of phospho-GSK3 β Ser9 fluorescence immunohistochemistry at the level of the hippocampal dentate gyrus of a control (Euploid, gestational week GW19; n = 3) and a DS (Down, GW19; n = 3) fetus. The dashed line square indicates the region where the immunofluorescence was quantified. Scale bar: 20 μ m.

DOI of original article: <https://doi.org/10.1016/j.nbd.2014.03.003>

* Corresponding author.

E-mail address: elisabetta.ciani@unibo.it (E. Ciani).

<https://doi.org/10.1016/j.nbd.2020.104769>

Available online 25 January 2020

0969-9961/ © 2020 Elsevier Inc. All rights reserved.