

Haemophilus influenzae periorbital cellulitis in a 95-year-old patient

Stefano Veraldi, Valentina Benzecry

Dipartimento di Fisiopatologia Medico-Chirurgica e dei Trapianti, Università degli Studi, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

Periorbital cellulitis (POC) is an acute bacterial infection of the eyelids. In the past, before the introduction of *Haemophilus influenzae* type B vaccination, POC was usually caused by this bacterium. Vaccination was introduced in 1985 and extended in 1990. Since then, most cases are caused by *Staphylococcus aureus* or Group A β -hemolytic *Streptococcus*. We present a case of POC caused by *H. influenzae* in a 95-year-old woman: to our knowledge, this is the oldest patient with POC reported in the literature.

Key words: elderly, *Haemophilus influenzae*, periorbital cellulitis, *Staphylococcus aureus*, vaccination

Received: July 12, 2020
Accepted: November 30, 2020

Correspondence

Stefano Veraldi

Dipartimento di Fisiopatologia Medico-Chirurgica e dei Trapianti, Università degli Studi, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, via Pace 9, 20122 Milan, Italy
E-mail: stefano.veraldi@unimi.it

Conflict of interest

The Authors declare no conflict of interest

Consent for publication

We certify that consent for publication of photographs was obtained

How to cite this article: Veraldi S, Benzecry V. *Haemophilus influenzae* periorbital cellulitis in a 95-year-old patient. Journal of Gerontology and Geriatrics Online First 2020;Dec 10. <https://doi.org/10.36150/2499-6564-N489>

© Copyright by Società Italiana di Gerontologia e Geriatria (SIGG)



OPEN ACCESS

This is an open access article distributed in accordance with the CC-BY-NC-ND (Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International) license. The article can be used by giving appropriate credit and mentioning the license, but only for non-commercial purposes and only in the original version. For further information: <https://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>

INTRODUCTION

Periorbital cellulitis (POC), or preseptal cellulitis, is an acute bacterial infection of the eyelids¹⁻¹⁵. In the past, before the introduction of *Haemophilus influenzae* type B vaccination, POC was usually caused by this bacterium. In the United States, vaccination was introduced in 1985 and extended in 1990. Since then, most cases are caused by *Staphylococcus aureus* or Group A β -hemolytic *Streptococcus*. We present a case of POC caused by *H. influenzae* in a 95-year-old woman: to our knowledge, this is the oldest patient with POC reported in the literature.

CASE REPORT

A 95-year-old Caucasian woman was admitted with a clinical diagnosis of herpes zoster. Her son stated that the dermatitis appeared suddenly three days before. It was diagnosed by the patient's general practitioner as herpes zoster. A therapy with valacyclovir (3 g/day) was started. Dermatological examination showed a severe erythematous edema, with tiny round vesicles and pustules, at the left eyelids (Fig. 1). The consistency of the lesion was soft-parenchymatous. The patient complained of burning sensation, pain, and lachrymation.

General physical examination showed mild arterial essential hypertension. Left laterocervical lymphadenopathy was observed. Fever was 38.6°C. Ophthalmological examination was impossible because the patient could not open her left eye at all.

Laboratory tests showed leucocytosis with neutrophilia, and increase in erythrocyte sedimentation rate and C-reactive protein. Blood culture was negative. Cytological examination of vesicles and pustules revealed the presence



Figure 1. Severe erythematous edema, with small vesicles and pustules, at the left eyelids.

of cocci and excluded a herpetic infection. Swab obtained from pustular lesions was positive for *Staphylococcus epidermidis*. Culture of needle aspirate of the swelling was positive for *H. influenzae*. Nasal swab was positive for *S. aureus*. It was impossible to perform conjunctival swab. X-rays and computerized axial tomography of the head were negative. A diagnosis of POC was made. According to the antibiogram results, the patient was treated with ceftriaxone (2 g i.v. for 10 days): both *H. influenzae* and *S. aureus* were sensitive to this antibiotic. The clinical picture improved within three days. Complete remission was observed seven days later. Follow up (six months) was negative.

DISCUSSION

POC occurs almost exclusively in children^{1-7,9,12-15} and is more frequent in males than in females^{13,14}. Cases in adults are extremely rare^{8,10,11}. Important predisposing factors of POC are local trauma^{3,4,12,13}. In two groups of patients, local trauma were responsible for penetration of bacteria in 20.9%¹² and 23.1%¹³ of cases, respectively). Other risk factors are impetigo or adjacent focus of infection, otitis, conjunctivitis (in 42.9% of patients in the previously cited study¹²), dacryostenosis¹²,

sinusitis^{2,3,5,7,9,12,13} (observed from 8%¹² to 14.5%⁹ to 19%⁵ to 24.8%¹³ to 81%⁷ of patients), and upper respiratory tract infection^{4,9,11}. It is possible that in our patient the sudden appearance of the swelling was caused by an arthropod bite or sting: in the study by Rimon et al.¹², insect bites and stings were responsible for 9.8% of cases of POC. However, in our patient it was impossible to discover the source of *H. influenzae*. In the past, the most frequently involved species in POC was *H. influenzae*^{1-7,9,11,12}. In United States, after the introduction in 1985 of vaccination for *H. influenzae* type B, the incidence of POC caused by this bacterium rapidly and deeply decreased^{5,7,9,12,13,15}. Other bacteria involved are *S. aureus*^{2,3,15}, Group A β -hemolytic *Streptococcus*^{2,3,15}, and *Streptococcus pneumoniae*^{2,3}. *S. aureus* and/or Group A β -hemolytic *Streptococcus* have been isolated also from conjunctival exudate as well as *H. influenzae* (in 13 out of 89 tear specimens in the study by Powell et al.⁴). Blood culture of *H. influenzae* is often negative⁵⁻⁷. After the introduction of vaccination, *H. influenzae* positive blood cultures became ever more rare: 1 positive out of 34 patients⁶ and 2 positive out of 133 cultures⁷. POC is characterized by the rapid appearance of usually monolateral erythematous edema, with smooth surface. The consistency is parenchymatous-hard. Sometimes, vesicles, blisters and pustules, as in our

patient, appear. Local pain, fever, and general malaise are common. A dreadful complication is meningitis^{1,5}. Recurrences are possible. Laboratory tests show leucocytosis with neutrophilia and increase erythro sedimentation rate, C-reactive protein and α 1-acid glycoprotein. Differential diagnosis includes orbital cellulitis, a severe infection involving the contents of the orbit, and herpes zoster. Therapy is based on systemic antibiotics, selected according the results of antibiogram, and incision and drainage¹⁰.

References

- 1 Simpson GT, McGill TI, Healy GB. Hemophilus influenzae type B soft tissue infections of the head and neck. *Laryngoscope* 1981;91:17-29. <https://doi.org/10.1288/00005537-198101000-00004>
- 2 Shapiro ED, Wald ER, Brozanski BA. Periorbital cellulitis and paranasal sinusitis: a reappraisal. *Pediatr Infect Dis* 1982;1:91-4. <https://doi.org/10.1097/00006454-198203000-00005>
- 3 Jackson K, Baker SR. Periorbital cellulitis. *Head Neck Surg* 1987;9:227-34. <https://doi.org/10.1002/hed.2890090406>
- 4 Powell KR, Kaplan SB, Hall CB, et al. Periorbital cellulitis. Clinical and laboratory findings in 146 episodes, including tear counter current immunoelectrophoresis in 89 episodes. *Am J Dis Child* 1988;142:853-7. <https://doi.org/10.1001/archpedi.1988.02150080059024>
- 5 Schwartz GR, Wright SW. Changing bacteriology of periorbital cellulitis. *Ann Emerg Med* 1996;28:617-20. [https://doi.org/10.1016/s0196-0644\(96\)70083-5](https://doi.org/10.1016/s0196-0644(96)70083-5)
- 6 Dudin A, Othman A. Acute periorbital swelling: evaluation of management protocol. *Pediatr Emerg Care* 1996;12:16-20. <https://doi.org/10.1097/00006565-199602000-00005>
- 7 Barone SR, Aiuto LT. Periorbital and orbital cellulitis in the Haemophilus influenzae vaccine era. *J Pediatr Ophthalmol Strabismus* 1997;34:293-6. PMID: 9310918
- 8 Anzai S, Sato T, Takayasu S, et al. Periorbital necrotizing cellulitis. *Int J Dermatol* 1998;37:799. <https://doi.org/10.1046/j.1365-4362.1998.00526.x>
- 9 Ambati BK, Ambati J, Azar N, et al. Periorbital and orbital cellulitis before and after the advent of Haemophilus influenzae type B vaccination. *Ophthalmology* 2000;107:1450-3. [https://doi.org/10.1016/s0161-6420\(00\)00178-0](https://doi.org/10.1016/s0161-6420(00)00178-0)
- 10 Duarte Reis M, Freitas IP, Sousa Coutinho V, et al. Facial and periorbital cellulitis with orbital involvement. *J Eur Acad Dermatol Venereol* 2002;16:156-8. <https://doi.org/10.1046/j.1468-3083.2002.00422.x>
- 11 Wang CC, Kuo HY, Chiang DH, et al. Invasive Haemophilus influenzae disease in adults in Taiwan. *J Microbiol Immunol Infect*. 2008;41:209-14. PMID: 18629415
- 12 Rimon A, Hoffer V, Prais D, et al. Periorbital cellulitis in the era of Haemophilus influenzae type B vaccine: predisposing factors and etiologic agents in hospitalized children. *J Pediatr Ophthalmol Strabismus* 2008;45:300-4. <https://doi.org/10.3928/01913913-20080901-14>.
- 13 Moubayed SP, Vu TT, Quach C, et al. Periorbital cellulitis in the pediatric population: clinical features and management of 117 cases. *J Otolaryngol Head Neck Surg* 2011;40:266-70. PMID: 21518652
- 14 Crosbie RA, Nairn J, Kubba H. Management of paediatric periorbital cellulitis: our experience of 243 children managed according to a standardised protocol 2012-2015. *Int J Pediatr Otorhinolaryngol* 2016;87:134-8. <https://doi.org/10.1016/j.ijporl.2016.06.025>
- 15 Williams KJ, Allen RC. Paediatric orbital and periorbital infections. *Curr Opin Ophthalmol* 2019;30:349-55. <https://doi.org/10.1097/ICU.0000000000000589>