

ELEMENTARY PARTICLES AND FIELDS Experiment

The HADES-at-FAIR Project^{*}

K. Lapidus^{1),2)***}, G. Agakishiev³⁾, A. Balandin⁴⁾, R. Bassini⁵⁾, C. Behnke⁶⁾,
A. Belyaev³⁾, A. Blanco⁷⁾, M. Böhmer⁸⁾, P. Cabanelas⁹⁾, N. Carolino⁷⁾, J. C. Chen¹⁾,
S. Chernenko³⁾, J. Díaz¹⁰⁾, A. Dybczak⁴⁾, E. Epple¹⁾, L. Fabbietti¹⁾, O. Fateev³⁾,
P. Finocchiaro¹¹⁾, P. Fonte^{7),12)}, J. Friese⁸⁾, I. Fröhlich⁶⁾, T. Galatyuk^{6),13)}, J. A. Garzón⁹⁾,
R. Gernhäuser⁸⁾, A. Gil¹⁰⁾, K. Göbel⁶⁾, M. Golubeva²⁾, D. González-Díaz^{14),15)},
F. Guber²⁾, M. Gumberidze¹⁶⁾, S. Harabasz⁴⁾, K. Heidel¹⁷⁾, T. Heinz¹⁴⁾, T. Hennino¹⁶⁾,
R. Holzmann¹⁴⁾, P. Huck⁸⁾, J. Hutsch¹⁷⁾, A. Ierusalimov³⁾, I. Iori^{5),18)}, A. Ivashkin²⁾,
M. Jurkovic⁸⁾, B. Kämpfer^{17),19)}, M. Kajetanowicz⁴⁾, T. Karavicheva²⁾, I. Koenig¹⁴⁾,
W. Koenig¹⁴⁾, B. W. Kolb¹⁴⁾, G. Korcyl⁴⁾, G. Kornakov⁹⁾, R. Kotte¹⁷⁾, A. Kozuch^{4),20)},
E. Krebs⁶⁾, R. Krücken⁸⁾, H. Kuc^{4),16)}, W. Kühn²¹⁾, A. Kugler²²⁾, A. Kurepin²⁾,
A. Kurilkin³⁾, P. Kurilkin³⁾, V. Ladygin³⁾, R. Lalik¹⁾, J. S. Lange²¹⁾, M. Liu²¹⁾, T. Liu¹⁶⁾,
L. Lopes⁷⁾, M. Lorenz⁶⁾, G. Lykasov³⁾, L. Maier⁸⁾, A. Malakhov³⁾, A. Mangiarotti⁷⁾,
J. Markert⁶⁾, V. Metag²¹⁾, B. Michalska⁴⁾, J. Michel⁶⁾, C. Müntz⁶⁾, R. Münzer¹⁾,
L. Naumann¹⁷⁾, M. Palka⁶⁾, Y. Parpottas²³⁾, V. Pechenov¹⁴⁾, O. Pechenova⁶⁾, A. Pereira⁷⁾,
J. Pietraszko⁶⁾, W. Przygoda⁴⁾, B. Ramstein¹⁶⁾, C. Rehnisch⁶⁾, A. Reshetin²⁾,
P. Rosier¹⁶⁾, A. Rustamov¹⁴⁾, A. Sadovsky²⁾, P. Salabura⁴⁾, T. Scheib⁶⁾, A. Schmah^{1),24)},
H. Schuldes⁶⁾, E. Schwab¹⁴⁾, J. Siebenon¹⁾, V. Smolyankin²⁵⁾, M. Sobiella¹⁷⁾,
Yu. G. Sobolev²²⁾, S. Spataro^{21),26)}, B. Spruck²¹⁾, H. Ströbele⁶⁾, J. Stroth^{6),14)},
C. Sturm¹⁴⁾, A. Tarantola⁶⁾, K. Teilab⁶⁾, V. Tiflov²⁾, P. Tlusty²²⁾, M. Traxler¹⁴⁾, R. Trebacz⁴⁾,
A. Troyan³⁾, H. Tsertos²³⁾, E. Usenko²⁾, T. Vasiliev³⁾, S. Visotski²⁵⁾, V. Wagner²²⁾,
M. Weber⁸⁾, C. Wendisch¹⁷⁾, J. Wüstenfeld¹⁷⁾, S. Yurevich¹⁴⁾, and Y. Zanevsky³⁾
(HADES Collaboration)

Received March 31, 2011

Abstract—After the completion of the experimental program at SIS18 the HADES setup will migrate to FAIR, where it will deliver high-quality data for heavy-ion collisions in an unexplored energy range of up to 8 A GeV. In this contribution, we briefly present the physics case, relevant detector characteristics and discuss the recently completed upgrade of HADES.

DOI: 10.1134/S1063778812050146

*The text was submitted by the authors in English.

¹⁾Excellence Cluster “Origin and Structure of the Universe”, Garching, Germany.

²⁾Institute for Nuclear Research, Russian Academy of Science, Moscow.

³⁾Joint Institute of Nuclear Research, Dubna, Russia.

⁴⁾Smoluchowski Institute of Physics, Jagiellonian University, Kraków, Poland.

⁵⁾Istituto Nazionale di Fisica Nucleare, Sezione di Milano, Italy.

⁶⁾Institut für Kernphysik, J. W. Goethe-Universität, Frankfurt, Germany.

⁷⁾LIP—Laboratório de Instrumentação e Física Experimental de Partículas, Coimbra, Portugal.

⁸⁾Physik Department E12, Technische Universität München, Garching, Germany.

⁹⁾Departamento de Física de Partículas, Universidad de Santiago de Compostela, Spain.

¹⁰⁾Instituto de Física Corpuscular, Universidad de Valencia—CSIC, Spain.

¹¹⁾Istituto Nazionale di Fisica Nucleare—Laboratori Nazionali del Sud, Catania, Italy.

¹²⁾also at ISEC Coimbra, Portugal.

¹³⁾also at ExtreMe Matter Institute EMMI, Darmstadt, Germany.

¹⁴⁾GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany.

¹⁵⁾also at Technische Universität Darmstadt, Germany.

1. INTRODUCTION

The High Acceptance Di-Electron Spectrometer (HADES) is a modern multi-purpose detector currently operating at the SIS18 heavy-ion synchrotron in the region of kinetic beam energies of up to $2 A$ GeV for nucleus–nucleus collisions [1]. The main components of the experimental setup are a superconducting magnet, four planes of Multiwire Drift Chambers used for the tracking of charged particles, a Time-of-Flight wall and a hadron blind RICH detector.

At SIS18, HADES has accomplished a number of successful experimental campaigns. Among them are systematic investigations of the di-electron production in light colliding systems at $1\text{--}2 A$ GeV [2, 3] and in nucleon–nucleon reactions [4].

Primarily constructed for the electron–positron pair spectroscopy, HADES is proven to be perfectly suited for hadron measurements, including strangeness production in nucleus–nucleus [5–7] and in proton–proton collisions [8].

The experimental program at SIS18 is not finished—in 2011–2012 HADES will investigate heavy colliding systems: AuAu and AgAg at bombarding energies of 1.25 and $1.65 A$ GeV, respectively. Measurements of di-electron and hadron production in pion-induced reactions are also foreseen.

2. HADES AT SIS100

In heavy-ion collisions at beam energies delivered by the heavy-ion synchrotron SIS100 within the Facility for Antiproton and Ion Research (FAIR), the nuclear compression is expected to be significantly higher as compared to SIS18. The increase of the beam energy will furthermore cause a transition from

the below-threshold to above-threshold production of the light vector mesons (ρ , ω , ϕ). On the basis of thermal models one expects two orders of magnitude higher vector meson yields at SIS100 as compared with SIS18 [9]. Since the key point of the HADES program is the investigation of the vector meson spectral functions at high baryonic densities, these conditions are perfectly suited for the desired goal. Thus, a migration to the SIS100 will be a natural continuation of the HADES experimental program.

The HADES detector ensemble possesses a set of unique features for the measurements at SIS100. It is a running experiment with a well understood performance, which can deliver high-quality data shortly after installation in the cave. Moreover, the detector and the analysis software will be further tested and tuned with upcoming heavy-ion runs at SIS18.

It is planned that HADES will be installed in a common cave with the CBM experiment, performing measurements at beam energies of up to $8 A$ GeV. Therefore, it will provide a bridge to higher energies (up to $40 A$ GeV) to be explored by the CBM experiment.

3. ACCEPTANCE AND OCCUPANCY CONSIDERATIONS

As a first assessment of the HADES performance at SIS100, the detector acceptance and occupancy were studied by Monte-Carlo simulations.

3.1. *Di-Electron Acceptance*

In order to understand the acceptance for di-electrons in heavy-ion collisions at a kinetic beam energy of $8 A$ GeV, the pairs originating from the direct e^+e^- decay of ω mesons using the Monte-Carlo generator PLUTO [10] were studied. Individual leptons were propagated through the HADES acceptance filter and the pairs with laboratory opening angles less than 9° were rejected. Figure 1 illustrates in transverse momentum versus rapidity coordinates the distributions of di-electrons in the full phase space (upper row) and in the HADES acceptance (lower row). 21% of the di-electrons, coming from ω mesons produced in CC collisions at $8 A$ GeV, are in the HADES acceptance. This number has to be compared with 33% for the same system at $2 A$ GeV. From the distributions shown in Fig. 1 it also follows that the acceptance of HADES with its present geometrical configuration still covers the mid-rapidity region at $8 A$ GeV. Taking into account the much larger yields of vector mesons anticipated at the higher energy one can expect significantly higher di-electron rates in the corresponding region of

¹⁶⁾Institut de Physique Nucléaire (UMR 8608), CNRS/IN2P3—Université Paris Sud, France.

¹⁷⁾Institut für Strahlenphysik, Forschungszentrum Dresden-Rossendorf, Dresden, Germany.

¹⁸⁾also at Dipartimento di Fisica, Università di Milano, Italy.

¹⁹⁾also at Technische Universität Dresden, Germany.

²⁰⁾also at Państwowa Wyższa Szkoła Zawodowa, Nowy Sacz, Poland.

²¹⁾II.Physikalischs Institut, Justus Liebig Universität Giessen, Germany.

²²⁾Nuclear Physics Institute, Academy of Sciences of Czech Republic, Rez.

²³⁾Department of Physics, University of Cyprus, Nicosia, Cyprus.

²⁴⁾also at Lawrence Berkeley National Laboratory, USA.

²⁵⁾Institute of Theoretical and Experimental Physics, Moscow, Russia.

²⁶⁾also at Dipartimento di Fisica Generale, Università di Torino, Italy.

**E-mail: kirill.lapidus@ph.tum.de




Fig. 1. Transverse momentum versus rapidity distributions (upper row) for electron–positron pairs coming from the direct ω decays at beam kinetic energies of 2 A GeV (left column) and 8 A GeV (right column) and the HADES acceptance (lower row).




Fig. 2. Transverse momentum versus rapidity distributions (upper row) of charged hadrons (left: π^+ , middle: π^- , right: protons) produced in CC collisions at a beam kinetic energy of 8 A GeV and the HADES acceptance (lower row).

invariant masses for 8 A GeV (SIS100) as compared to 2 A GeV (SIS18), compensating by far the slight decrease in acceptance.

3.2. Hadron Acceptance

The acceptance of the HADES detector for charged hadrons was studied in a similar way. As an input, hadron events generated with the UrQMD transport code were used. The detector acceptance for the charged hadrons produced in CC collisions at a kinetic beam energy of 8 A GeV is illustrated on Fig. 2. At this energy, 50% of π^+ , 52% of π^- , and 15% of protons are inside the HADES acceptance.

Thus, the detector that was originally designed to study symmetric systems at bombarding energies close to 2 A GeV, where the acceptance of light particles is symmetric around midrapidity, provides an adequate acceptance for di-electrons and hadrons also at 8 A GeV.

3.3. Occupancy of the Detector

The transition to higher energies will be accompanied by the increase of the charged particle multiplicity in the forward part of the spectrometer. The maximum system size, possible to be studied at

SIS100, was evaluated. The performed simulations have shown that the charged particle multiplicity grows by a factor of 14 when going from CC collisions at 2 A GeV to AuAu collisions at 1.5 A GeV. Already in the latter case one expects a 20% double hit probability in the RPC detector (described below) that limits the tracking performance. According to our estimates, the hit occupancy in AuAu collisions at 1.5 A GeV corresponds to the NiNi colliding system at 8 A GeV bombarding energy.

4. THE UPGRADE OF THE APPARATUS

For the measurements at SIS100, HADES will benefit from the upgrade, performed in 2009–2010 with the aim to make the detector capable of measurements with really heavy ions at SIS18. The requirements for the upgrade were that the detector is able to cope with high charged particle multiplicities, typical for central collisions of gold nuclei, and accept a minimum trigger rate of 20 kHz even for heavy-ion collisions.

A Forward Wall detector consisting of 280 individual scintillators was installed, extending the HADES acceptance to the region of low polar angles ($0.33^\circ < \theta < 7.17^\circ$). This detector was already used in 2007 for the spectator proton detection in a $d\mu$ experiment [11]. It is essential for the upcoming heavy-ion runs at SIS18 and SIS100, since it allows the reconstruction of the reaction plane, necessary, e.g., for the analysis of particle flow.

One of the most important steps in the detector upgrade was the replacement of the low-granularity scintillator Time-of-Flight detector TOFin to be the high-granularity timing Resistive Plate Chambers (tRPCs) [12]. This detector offers a time resolution better than 100 ps with an efficiency close to 100%. With 1200 individually shielded 4-gap tRPCs, the RPC TOF wall enables HADES to handle multiplicities of up to 200 charged particles per event and it further improves the lepton and charged hadron identification.

Investigations of colliding systems with maximal nucleon numbers aiming at high statistics data samples required also an upgrade of the HADES data acquisition (DAQ) system.

The key component of the newly constructed system is a trigger and readout board (TRB) being a multi-purpose electronic device with on-board DAQ functionality [13].

With the upgraded setup the HADES Collaboration aims to collect high statistics data samples that allow to perform multi-differential (as a function of collision centrality, transverse momentum, invariant mass, etc.) studies of di-electron and hadron production.

5. AN ELECTROMAGNETIC CALORIMETER FOR HADES

Currently, another upgrade project is intensively investigated—the extension of the HADES particle identification capabilities by an electromagnetic calorimeter. The proposed calorimeter will open the possibility for the neutral pseudoscalar meson (π^0, η) reconstruction via their decays into two photons. The π^0 multiplicity is a natural choice for the absolute normalization of the measured di-electron spectra. The measurement of the η -meson yield is of particular importance, since its Dalitz decay is a significant source of the di-electrons for the invariant masses above $140 \text{ MeV}/c^2$. Whereas at beam energies of 1–2 A GeV the measurements of the η yield in heavy-ion collisions were performed by the TAPS Collaboration [14], at higher energies, in the SIS100 range, no data on the η production exist.

Another important function of the calorimeter is to improve the lepton–pion separation at high values of the particle momentum ($p > 0.5 \text{ GeV}/c$). For this purpose, a cut on the ratio of the deposited energy in the calorimeter to the measured momentum in the HADES tracking system will be employed.

We consider to build an electromagnetic calorimeter using lead-glass blocks from the end-cap calorimeter of the former OPAL experiment [15]. In total, ~900 modules will be arranged in six sectors of the HADES setup. The calorimeter will substitute the existing SHOWER detector and will be installed behind the tRPCs.

Dedicated in-beam tests of the lead-glass modules were already performed at CERN (electron and pion beams) and MAMI (γ beam), confirming the expected energy resolution of $\simeq 5\%/\sqrt{E/\text{GeV}}$ and demonstrating a good potential for the lepton–pion separation.

Recently, a calorimeter simulation software was developed, making use of the GEANT3 program [16] and an external Cherenkov photon transport code [17, 18]. The simulations have shown the feasibility of the π^0 and η reconstruction and the lepton/pion separation in heavy-ion collisions.

6. SUMMARY

In this contribution, plans of the HADES Collaboration for the future investigations of relativistic heavy-ion collisions at SIS100 were outlined. The newly updated experimental setup is ready to perform investigations of the compressed nuclear matter at FAIR exploiting both lepton and hadron probes.

ACKNOWLEDGMENTS

The HADES Collaboration gratefully acknowledges the support by the BMBF through grants nos. 06MT9156, 06GI146I, 06FY171, 06FY91001, and 06DR9059D (Germany), by GSI (TKrue 1012, GI/ME3, OF/STR), by the Helmholtz Alliance HA216/EMMI, by the Excellence Cluster Universe (Germany), by grants GA AS CR IAA100480803 and MSMT LC 07050MSMT (Czech Republic), by grant KBN5P03B 140 20 (Poland), by INFN (Italy), by CNRS/IN2P3 (France), by grants MCYT FPA2000-2041-C02-02 and XUGA PGID FPA2009-12931 T02PXIC20605PN (Spain), by grant UCY-10.3.11.12(Cyprus), by INTAS grant no. 06-1000012-8861 and EU contract RII3-CT-506078.

REFERENCES

1. G. Agakishiev et al. (HADES Collab.), Eur. Phys. J. A **41**, 243 (2009).
2. G. Agakishiev et al. (HADES Collab.), Phys. Lett. B **663**, 43 (2008).
3. G. Agakishiev et al. (HADES Collab.), Phys. Rev. Lett. **98**, 052302 (2007).
4. G. Agakishiev et al. (HADES Collab.), Phys. Lett. B **690**, 118 (2010).
5. G. Agakishiev et al. (HADES Collab.), Phys. Rev. C **80**, 025209 (2009).
6. G. Agakishiev et al. (HADES Collab.), Phys. Rev. C **82**, 044907 (2010).
7. G. Agakishiev et al. (HADES Collab.), arXiv:1010.1675 [nucl-ex].
8. J. Siebenson et al., arXiv:1009.0946 [nucl-ex].
9. A. Andronic, P. Braun-Munzinger, and J. Stachel, Nucl. Phys. A **772**, 167 (2006).
10. I. Fröhlich et al., PoS **ACAT2007**, 076 (2007).
11. K. Lapidus et al. (HADES Collab.), arXiv:0904.1128 [nucl-ex].
12. H. Alvarez-Pol et al. (HADES Collab.), Nucl. Instrum. Methods Phys. Res. A **535**, 277 (2004).
13. I. Fröhlich et al., IEEE Trans. Nucl. Sci. **55**, 59 (2008).
14. R. Averbeck et al. (TAPS Collab.), Z. Phys. A **359**, 65 (1997).
15. M. Akrawy et al., Nucl. Instrum. Methods Phys. Res. A **290**, 76 (1990).
16. <http://consult.cern.ch/writeup/geant/>
17. A. V. Aref'ev et al., Instrum. Exp. Tech. **51**, 511 (2008).
18. I. G. Alekseev et al., Instrum. Exp. Tech. **51**, 491 (2008).