



Letter to the Editor

Antonio Mastroianni¹
Valeria Vangeli¹
Maria Vittoria Mauro²
Roberto Manfredi³
Sonia Greco¹

Turicella otitidis central venous-related bacteremia during pediatric acute lymphoblastic leukemia

¹Infectious & Tropical Diseases Unit, "Annunziata" Hospital, Cosenza, Italy

²Microbiology & Virology Unit, "Annunziata" Hospital, Cosenza, Italy

³Infectious Diseases Institute, Alma Mater Studiorum University of Bologna, Bologna, Italy

Article history

Received: 4 November 2022; Revision Requested: 9 May 2023; Revision Received: 12 May 2023; Accepted: 1 June 2023; Published: 18 July 2023

Sir,

We read with great interest the article by De Frutos M *et al.* [1] about 5 cases of otitis media due to *Turicella otitidis*, some of them complicated, as a case of mastoiditis that required surgery.

We agree with the authors [1] that some microorganisms isolated only anecdotally can become relevant pathogens, especially in immunosuppressed patients and therefore it is necessary to update the epidemiological surveillance with the description of new cases [1]. A suggestive report of central venous catheter (CVC)-related *T. otitidis* bacteremia is described and commented on the ground of an updated literature review. *T. otitidis* is a non-fermenting Gram-positive rod isolated almost exclusively from ear exudate fluids, although its clinical significance in both acute and chronic otitis is still controversial [2-5]. Isolated since two decades [2], its genome has been sequenced [3]. *T. otitidis* habitat seems limited to the ear and surrounding areas, but it has been retrieved as an infrequent cause of bacteremia [2-5], especially in children with hematologic malignancies [2,4]. We discuss a rare case of CVC-related *T. otitidis* bacteremia in a child with acute leukemia, based on the available literature data.

A 11-year old child suffering from acute lymphoblastic leukemia was hospitalized due to fever, in the absence of organ signs of localization, and in particular of acute or chronic otitis, mastoiditis, sinusitis. Laboratory workup at hospital admission showed low hemoglobin of 9 gm% and high leukocyte count of $22.86 \times 10^9/L$ with predominant neutrophil (82%), alanine aminotransferase level of 96 U/L (normal range, 21-72 U/L) and C-reactive protein raised at 158 mg/L.

During the last previous outpatient check-up, seven days

before hospital admission, results of laboratory examinations showed a white blood cell count of $11.3 \times 10^9/L$ with neutrophil ratio of 60%, hemoglobin of 11 gm% and a level of C reactive protein of 5 mg/L.

The patient had a temperature of 39.4°C, a heart rate of 123 beats/min, a respiratory rate of 24 breaths/min, a blood pressure of 90/60 mmHg, and a SaO₂ of 94%.

Microbial isolation has been obtained from the CVC blood culture after a 48-hour incubation in a Bactec Ped Plus /aerobes vial (Becton Dickinson Italia s.p.a.). Direct microscopy and Gram stain pointed out pleomorphic, asporogenous Gram-positive bacilli with a palisade appearance. Appropriate culture media (Agar Columbia with 5% mutton blood by BioMerieux Italy s.p.a.) and Agar chocolate PVX (by BioMerieux Italy s.p.a.) maintained in a CO₂ thermostat, allowed the growth of small rounded granular greyish-creamy transparent colonies of a 1.5 mm diameter. The identification was performed by the automated system Vitek2-VitekMS (by BioMerieux Italy s.p.a.) while the in vitro antimicrobial susceptibility assay by Kirby-Bauer was carried out on Mueller Hinton agar plus 5% defibrinated horse blood (by BioMerieux Italy s.p.a.). According to the European Committee on Antimicrobial Susceptibility Testing (EUCAST) guidelines, the following MICs were obtained: benzilpenicillin 1.5 mg/L, ciprofloxacin 2mg/L, gentamicin 1.5 mg/L, vancomycin MIC 0.38 mg/L, clindamycin 0.5 mg/L, tetracycline MIC 1.5 mg/L, linezolid MIC 1 mg/L and rifampicin MIC 0.004 mg/L. No other microbiological tests were requested, in addition to peripheral blood cultures, because the boy did not present a precise organic symptomatology which would have suggested further pertinent microbiological investigations.

Systemic intravenous rifampicin in association with vancomycin CVC lock therapy and the prompt CVC removal, allowed a prompt resolution of local and systemic signs and symptoms of infection, in the absence of complications and recurrences, as observed in the subsequent, prolonged follow-up. Very few cases of systemic *T. otitidis* infection have

Correspondence:
Prof. Roberto Manfredi, MD
Infectious Diseases Institute, Alma Mater Studiorum University of Bologna
S. Orsola Teaching Hospital, Via Massarenti, 11, I-40138 Bologna, Italy
Phone: +39 051 2143618 - Fax: +39 051 343500
E-mail: roberto.manfredi@unibo.it

been reported until now [2-5], usually in cancer patients [1], with only one case described in a child with a malignant hemopathy like ours [4]. Clinicians should be aware of the opportunistic potential of *T. otitidis*, especially when immunocompromised hosts are of concern. In our experience, a timely CVC removal seemed more effective when compared with systemic-local antibiotic treatment based on the *in vitro* sensitivity testing.

FUNDING

None to declare

CONFLICT OF INTEREST

Authors declare no conflict of interest

REFERENCES

1. De Frutos M, López-Urrutia L, Aragón R, Vegas AM, Vázquez M, Eiros Bouza JM. *Turicella otitidis*, aportaciones a su posible papel en la etiología de la patología infecciosa del oído [*Turicella otitidis*, contributions to its role in the etiology of ear infections]. *Rev Esp Quimioter*. 2018 Jun;31(3):278-281. Spanish. PMID: 29696957
2. Li D, Shan W, Dai C, Kan J, Li M, Yin M. Bacteremia Caused by *Turicella otitidis* in a patient with diffuse large B-cell lymphoma. *Clin Lab*. 2020;66. doi: 10.7754/Clin.Lab.2019.190641.
3. Greninger AL, Kozyreva V, Truong CL, Graves M, Chaturvedi V. Draft genome sequence of *Turicella otitidis* TD1, isolated from a patient with bacteremia. *Genome Announc*. 2015;17;3:e01060-15. doi: 10.1128/genomeA.01060-15.
4. Loiez C, Wallet F, Fruchart A, Husson MO, Courcol RJ. *Turicella otitidis* in a bacteremic child with acute lymphoblastic leukemia. *Clin Microbiol Infect*. 2002;8:758-9. doi: 10.1046/j.1469-0691.2002.00474.x.
5. von Graevenitz A, Funke G. *Turicella otitidis* and *Corynebacterium auris*: 20 years on *Infection*. 2014;42:1-4. doi: 10.1007/s15010-013-0488-x.