

## Assessment of the specificity of cardiopulmonary response during tethered swimming using a new snorkel device

Marco Pinna · Raffaele Milia · Silvana Roberto · Elisabetta Marongiu · Sergio Olla · Andrea Loi · Matteo Ortu · Gian Mario Migliaccio · Filippo Tocco · Alberto Concu · Antonio Crisafulli

Received: 10 April 2012 / Accepted: 27 July 2012  
© The Physiological Society of Japan and Springer 2012

**Abstract** This study aimed at comparing maximal oxygen uptake ( $VO_{2max}$ ), maximal heart rate ( $HR_{max}$ ), and anaerobic threshold (AT) obtained from tethered swimming (SW) and three other testing procedures: cycling (CY), running (RU), and arm cranking (AC). Variables were assessed in 12 trained male swimmers by a portable gas analyzer connected to a modified snorkel system to allow expired gases collection during swimming. Athletes exhibited a higher  $VO_{2max}$  during the SW test as compared to the CY and the AC tests. There was no significant difference in  $VO_{2max}$  between the SW and the RU test, but the Bland and Altman plot highlighted a poor agreement between results. Moreover, AT occurred at higher workloads during SW in comparison to the other tests. These results do not support the use of any unspecific testing procedures to estimate  $VO_{2max}$ ,  $HR_{max}$ , and AT for swimming.

**Keywords** Exercise · Oxygen uptake · Anaerobic threshold · Ergometry

### Introduction

The measurement of certain physiological parameters, such as maximal oxygen uptake ( $VO_{2max}$ ), heart rate (HR) at a

given workload, and anaerobic threshold (AT), is a useful tool to assess athletes' physical capacity and to follow the progression of conditioning during training periods. However, such measurements should be performed under conditions similar to those experienced by athletes during real training sessions or competitions and using specific ergometers, otherwise results can be misleading. This assumption is particularly critical in swimming, as standard laboratory testing for work capacity usually employs the cycle-ergometer, the treadmill, and the arm-crank, which impose movements that are unrelated to swimming. Moreover, the technical limitations of the routine measurement of expired gases in a pool explain why the research dealing with  $VO_{2max}$  during this sport activity is not abundant.

To measure oxygen uptake ( $VO_2$ ) during swimming, a number of methods have been developed: the use of the Douglas bag to gather expired gas, thereby extrapolating  $VO_2$  [14, 15, 24, 29, 32];  $VO_2$  calculation by means of back-extrapolation [22, 33, 34]; and finally the recent use of commercially available respiratory snorkels able to calculate  $VO_2$  together with other respiratory parameters such as pulmonary ventilation (VE) and carbon dioxide production ( $VCO_2$ ) [25–27, 30]. This latter method allows breath-by-breath analysis of gas exchange, thereby allowing the description of oxygen uptake kinetics during specific swimming exercise [25, 26]. At present, this latter method is probably more reliable and comfortable for swimmers than the Douglas bag and the back-extrapolation, since the first suffers from technical limitation due to the fact that the bag used to gather expired gases is somewhat cumbersome. Furthermore, it is usually connected to the athlete by two tubes with unidirectional valves, which renders the apparatus not very comfortable to wear. On the other hand, the back-extrapolation method is

M. Pinna · R. Milia · S. Roberto · E. Marongiu · S. Olla · M. Ortu · F. Tocco · A. Concu · A. Crisafulli (✉)  
The Department of Medical Sciences, Sports Physiology Lab,  
University of Cagliari, Via Porcell 4, 09124 Cagliari, Italy  
e-mail: crisafulli@tiscali.it

A. Loi · G. M. Migliaccio  
Regional School of Sport of Sardinia, Italian Olympic  
Committee, Sardinia, Italy

indirect and thus less accurate than methods employing direct gas exchange measures.

Limited previous research focusing on physical capacity during swimming led to conflicting results. Early reports in trained swimmers concluded that  $VO_{2\max}$  during swimming was similar or lower than during running [12, 14, 15]. Furthermore, higher  $VO_{2\max}$  values were found in triathletes during both cycling and running in comparison with swimming [17, 18], while these differences were not present in swimmers [30]. Taking into account these conflicting results, further research on gas exchange during swimming is warranted. Moreover, to the best of our knowledge, none has to date assessed the specificity of gas exchange measure in swimmers by using the respiratory snorkel method. In detail, none has compared the swimmers'  $VO_2$  response assessed with the respiratory snorkel with other standard laboratory test procedures for work capacity, such as cycle-ergometry, treadmill, and arm-crank. In fact, the past research conducted with the snorkel method focused mainly on accumulated oxygen deficit, oxygen uptake kinetics, and oxygen cost of swimming [3, 4, 25–27]. Thus, a study dealing with the specificity of competitive swimmers' cardiopulmonary responses assessed by the snorkel method during different exercise tests is needed.

Given the relative novelty of the snorkel device and the conflicting results arising from the past literature, the aim of this investigation was to study the difference in gas exchange responses provided by four different incremental tests in trained swimmers: three unspecific laboratory test procedures (i.e. arm-cranking, cycling, and running), and one specific swimming test in a pool. The goal was to determine whether any of the unspecific laboratory tests yielded results similar and interchangeable to those obtained from the specific swimming test and if so which.

## Methods

### Subjects

Twelve trained male swimmers were recruited to take part in this investigation. Their mean  $\pm$  standard error (SE) of age, mass, and height were  $19.2 \pm 0.8$  years,  $68.9 \pm 1.6$  kg, and  $177.3 \pm 1.7$  cm, respectively. All athletes were regularly involved in regional and national competitions and trained on average  $12.5 \pm 0.3$  h a week. None had any history of cardiac or respiratory disease or was taking any medication at the time of the study, and none showed any abnormalities on physical examination and on resting electrocardiogram. Written informed consent was obtained by all of the participants after they were informed about the methods and aims of the study, whose protocol was

approved by the local ethical committee and carried out according to the Declaration of Helsinki.

### Experimental design

Each participant underwent the following protocol randomly assigned to eliminate any order effect:

1. Unspecific tests: athletes underwent three laboratory incremental exercise tests up to exhaustion using three different ergometers:

(a) Cycling (CY) test: participants performed an incremental test on an electromagnetically braked cycle ergometer (Tunturi EL 400, Finland) to assess the maximum workload achievable ( $W_{\max}$ ). This test consisted of a linear increase of work load of  $30 \text{ W min}^{-1}$ , starting from  $40 \text{ W min}^{-1}$ , at a pedaling frequency of 60 rpm, up to exhaustion (i.e. the point when the participant was unable to maintain a pedaling rate of at least 50 rpm).

(b) Running (RU) test: participants ran on a motorized treadmill (Runrace; Technogym, Forlì, Italy) up to exhaustion. The test consisted of a linear increase in running velocity of  $1 \text{ km h}^{-1}$  every minute, starting from  $5 \text{ km h}^{-1}$ , up to exhaustion, which was considered as the exercise level at which the athlete was unable to maintain the running speed. The treadmill was set at a gradient of 1 % throughout exercise to compensate for the lack of air friction [19].

(c) Arm-cranking (AC) test: which consisted of a linear increase in work load ( $10 \text{ W min}^{-1}$ ), starting from  $20 \text{ W min}^{-1}$ , at a cranking frequency of 60 rpm, up to exhaustion, taken as the point at which the athlete was unable to maintain a cranking rate of at least 50 rpm.

All unspecific tests were preceded by at least 10 min of rest. The average variable value of the last 3 min of resting was considered as the baseline level. Tests were conducted between 1000 and 1400 hours in a temperature-controlled room (room temperature set at  $25 \text{ }^\circ\text{C}$ , relative humidity at 50 %).

2. Specific swimming (SW) test: this test consisted in tethered swimming, conducted between 1000 and 1400 hours in a 25-m indoor swimming pool, under


the same water condition (water temperature of 27 °C). The athlete was attached by a waist belt to an elastic rope connected to a digital dynamometer (PCE, FM1000, Germany) able to continuously provide a display of the values of the force being applied to the rope. The belt arrangement permitted the swimmers to kick freely. All swimmers performed front crawl with free stroke frequency. The other extremity of the dynamometer was fixed to the starting block of the pool (Fig. 1a). The tension applied to the elastic rope was constantly monitored on the display of the dynamometer, and continuous vocal feedback was provided to an assistant who moved a pole with a colored signal fixed at its extremity forward or backward immersed in the water. The tested swimmer was instructed to follow the colored signal so that the assistant could adjust the tension applied to the rope simply by moving forward or backward the pole. Using this method, an incremental test was developed, which started from a tension of 3 kg and was increased progressively by 1 kg  $\text{min}^{-1}$ . The test terminated when the athlete was no longer able to maintain his position and to follow the colored signal for more than 30 s. The dynamometer had a computer interface which allowed the registration of the tension applied to the rope throughout tests. Figure 1b shows an example of the tension applied during one recording. Workload was calculated in  $\text{kg m min}^{-1}$  and then converted to  $W$ , the same unit as those obtained from the CY and the AC tests. The instrument range of measure was

from 0 to 100 kg, with a resolution of 0.05 kg and an accuracy of 0.5 % ( $\pm 5$  N).

Throughout the CY, RU, AC, and SW tests, participants' expired gases were analyzed by a portable metabolic device (VO2000; MedicGraphics, USA), which provided a 3-breath average of  $\text{VO}_2$ ,  $\text{VCO}_2$ , and pulmonary ventilation (VE) through telemetric transmission. This system has been shown to be reliable and to have show good agreement compared to a standard metabolic cart for laboratory use [9, 23]. Moreover, the VO2000 could gather HR values from a chest belt connected to the VO2000 by telemetric transmission. During the CY, RU, and AC tests, athletes wore a face mask and a breathing valve connected to the VO2000 for expired gases analysis. This system was modified for tethered swimming. In this setting, athletes breathed through a low volume (about 200 mL) corrugated flexible plastic tube attached to the VO2000 breathing valve (Fig. 1c, d). This modified system allowed expired gases to be collected, and prevented water from entering the analyzing system.

Tests were held at least 1 week apart, and athletes were asked to have a meal at least 3 h before testing. Athletes were also asked to avoid caffeine and alcohol ingestion the day before tests were scheduled. The anaerobic threshold was determined using the  $V$ -slope method, which detects AT by using computerized regression analysis of the slopes of the  $\text{VCO}_2$  versus  $\text{VO}_2$  plot during exercise [6]. Achievement of  $\text{VO}_{2\text{max}}$  was considered as the attainment of at least 2 of the following criteria: (1) a plateau in  $\text{VO}_2$  despite increasing;

**Fig. 1** **a** Schematic representation of the tethered-swimming apparatus. **b** Example of the recording of the tension applied to the rope during the incremental swimming test. **c, d** The modified breathing valve to collect expired gases



(2) a respiratory exchange ratio (RER) above 1.10; and (3) a heart rate (HR)  $\pm 10$  beats  $\text{min}^{-1}$  of predicted maximum HR calculated as  $220 - \text{age}$  [16]. Prior to testing, the VO2000 was calibrated according to the manufacturer's instructions. For the calibration during the SW tests, the additional dead space of the plastic tube was taken into account by modifying the pre-set parameters of the VO2000.

Additional experiments to assess the modified expired gas analysis system

Since the additional dead space of the modified expired gas analysis system could affect the respiratory response during swimming, in a sub-group of the participants ( $n = 4$ , age  $20.1 \pm 4.4$  years, height  $179.2 \pm 2.3$  cm, and mass  $69.3 \pm 4.2$  kg), additional experiments were performed, consisting of two randomly assigned incremental tests to exhaustion on the same cycle ergometer used for the CY tests. In one test, athletes breathed through the modified gas analysis system, while in the other test, expired gasses were collected by the standard VO2000 system.

#### Calculation and data analysis

Data were averaged for 1 min and displayed as a function of % of maximal workload achieved ( $W_{\text{max}}$ ) during each test. Results are presented as mean  $\pm$  SE. Bland and Altman statistics [7] to assess agreement between two methods of measurement were carried out in order to evaluate agreement between variables responses during the SW test and the other unspecific tests. Since it has been previously described that the test–retest reproducibility of  $VO_{2\text{max}}$  in verification tests ranged between 3.5 and 7.1 %, depending on errors in measurements introduced by exercise protocols and by gas analyser devices, and that a threshold around 5 % is often used to compare different exercise protocols to assess the “true”  $VO_{2\text{max}}$  [2, 21, 31], we chose the 5 % difference as the limit to consider methods interchangeable in parameter assessment. The repeated measures analysis of variance (ANOVA), followed by Dunnet post hoc tests when appropriated to compare SW data to those from the other tests, were also performed in order to have further comparison between tests. Moreover, the two-way ANOVA for repeated measures (factors of time and condition) was conducted for the additional experiment session (i.e. when subjects breathed through the modified gas analysis system or through the standard VO2000 while cycling) in order to find out whether or not the modified expired gas analysis device affected the respiratory parameters.


Significance was set at a  $p$  value  $< 0.05$ . Descriptive statistics were carried out before the ANOVA to confirm the assumptions of normality by means of the Kolmogorov–Smirnov test. The alpha level was set at  $p < 0.05$ .

Statistics were calculated employing commercially available software (Graph-Pad Prism).


## Results

All participants fulfilled the selected criteria for  $VO_{2\text{max}}$  achievement in all tests performed. The  $W_{\text{max}}$  values achieved during the CY, RU, AC, and SW tests were  $312.5 \pm 9.3$  W  $\text{min}^{-1}$ ,  $15.2 \pm 0.4$  km  $\text{h}^{-1}$ ,  $90.1 \pm 10.1$  W  $\text{min}^{-1}$ , and  $8.9 \pm 0.3$  kg m  $\text{min}^{-1}$  (i.e.  $87.3 \pm 2.9$  W  $\text{min}^{-1}$ ), respectively.

Figure 2 illustrates the HR and  $VO_2$  time courses as a function of % of  $W_{\text{max}}$  throughout the tests and recovery. During all tests, variables increased almost linearly with respect to workload. Figure 3 shows the  $HR_{\text{max}}$  level reached during each test together with the Bland and Altman plots. ANOVA found that  $HR_{\text{max}}$  was lower during the SW test in comparison to the RU and the AC tests,



**Fig. 2** HR and  $VO_2$  time courses as a function of % of  $W_{\text{max}}$  throughout the various tests of the protocol and recovery



**Fig. 3** HR<sub>max</sub> level reached during each test together with the Bland and Altman plots. The dotted lines identify the limits of the 5 % difference between methods. *p* < 0.05 versus the SW test

whereas no difference was found between the SW and the CY tests. The Bland and Altman plot highlighted that limits of agreement of the difference in HR<sub>max</sub> between the SW and the other tests were widespread. The most evident difference was with respect to the RU test, with limits of agreement lying between +2.6 and -32.3 bpm. This difference was also evident with respect to the CY test (limits of agreement between +17.4 and -24.9 bpm), notwithstanding that ANOVA did not find any significant difference between these two tests. In terms of % difference, the mean difference between the SW and the CY tests was -2.2 %, with limits of agreement between +10.3 and -14.8 %. It should also be noticed that 7 of the 12 subjects studied had a difference between tests higher than 5 %.

Figure 4 exhibits the levels attained of VO<sub>2max</sub>. ANOVA showed that, during the SW test, athletes achieved higher VO<sub>2max</sub> levels as compared to the CY and the AC tests, while no difference was found between the SW and RU tests. Bland and Altman plots demonstrated that limits of agreement of the difference in VO<sub>2max</sub> between the SW and the other tests lay between +1,117 and -1,367, +1,012 and -241, +1,212 and -380 mL min<sup>-1</sup> for the RU, CY, and AC tests, respectively. Although ANOVA did not find any difference between the SW and RU tests, it should be considered that the limits of agreement between these two tests were wide. In terms of % difference, the mean difference between the SW and RU tests was -3.5 %, with limits of agreement between +30.8 and


-38 %. Moreover, 7 of the 12 studied subjects had a difference between tests higher than 5 %.

During the SW test, AT occurred at a higher % of W<sub>max</sub> in comparison with all the other tests (Fig. 5). The Bland and Altman plot showed that the limits of agreement for this parameter were between +45.5 and -12.4, +43.4 and -8.1, +41.9 and -16.3 % of W<sub>max</sub> for the RU, CY, and AC tests, respectively.


Figure 6 shows the results from the session of additional experiments to assess the modified gas exchange system. No significant difference was found in VO<sub>2</sub> and VCO<sub>2</sub> responses between the incremental test with the modified gas analysis system and the test performed with the standard VO2000 equipment. However, it should be highlighted that the use of the snorkel caused a slight VE and HR increment at sub-maximal workloads with respect to free breathing. Indeed, HR was higher at rest and during the first exercise steps (up to 40 % of maximum workload) while subjects breathed throughout the snorkel than while they breathed with the standard equipment. Similarly, VE was higher at rest up to 30 % of maximum workload when athletes used the modified snorkel.

**Discussion**


Since it is considerably simpler to obtain VO<sub>2</sub> measures in a laboratory setting, it is tempting to predict VO<sub>2</sub> for



**Fig. 4** VO<sub>2</sub>max level reached during each test together with the Bland and Altman plots. The *dotted lines* identify limits of the 5 % difference between methods. *p* < 0.05 versus the SW test



**Fig. 5** AT level reached during each test together with the Bland and Altman plots. The *dotted lines* identify limits of the 5 % difference between methods. *p* < 0.05 versus the SW test



**Fig. 6** Results from the session of additional experiments to assess the modified gas exchange system

swimming from laboratory procedures. One of the purposes of this study was to determine whether non-specific laboratory tests provided results similar to those obtained from specific swimming testing procedures. This purpose had the practical implication to determine whether any of the unspecific laboratory tests could be used interchangeably with specific testing in order to assess swimmers' physical capacity and if so which. Results from the present study do not support the use of any unspecific tests to estimate  $VO_2$  for swimming, as none of the three laboratory tests employed in the present study provided data similar to those obtained from the specific test. To the best of our knowledge, this study is the first to compare competitive swimmers' cardiopulmonary responses assessed with the respiratory snorkel with other standard laboratory test procedures for work capacity. Hence, these are the first results obtained with the snorkel method demonstrating the uniqueness of swimming testing with respect to laboratory procedures for exercise capacity.

In fact, tethered swimming induced a higher  $VO_{2max}$  response as compared to cycling and cranking, as suggested by the ANOVA results and further strengthened by Bland and Altman statistics. Moreover, even though ANOVA did not find any difference between the SW and RU tests, results from the Bland and Altman plot highlighted that the limits of agreement of the difference in  $VO_{2max}$  between these two testing procedures were wide. In detail, the Bland and Altman plot demonstrated that the limits of agreement of the difference in  $VO_{2max}$  between the SW and RU tests were between +1,117 and -1,367 mL min<sup>-1</sup>. In terms of % difference, this was between +30.8 and -38 %, with 7 of the 12 studied subjects showing a difference between tests higher than 5 %. It should be noted that the test-retest reproducibility of  $VO_{2max}$  in verification tests previously described in the scientific literature ranged between 3.5 and 7.1 %, depending on errors in measurements introduced by exercise protocols and by gas analyzer devices, and that a

threshold around 5 % is often used to compare different exercise protocols to assess the “true”  $VO_{2max}$  [2, 21, 31]. That is, the difference in  $VO_{2max}$  yielded by two methods must not exceed this threshold to consider the two methods equivalent in assessing this parameter. In the present investigation, only in 5 of the 12 subjects enrolled this threshold was exceeded, and this fact meant that, in most of the subjects, the RU and SW tests could not be used interchangeably. Taken together, these results indicated that the use of the RU test could lead to substantial over- or under-estimation of the  $VO_{2max}$  reached during real swimming.

Similarly, the laboratory tests also led to misleading results for  $HR_{max}$ , as this parameter was on average over-estimated by the RU and AC tests with respect to the SW test, as testified by ANOVA results. On average, this parameter was not significantly different between the CY and SW tests when looking at ANOVA alone. However, the limits of agreement of the  $HR_{maxSW} - HR_{maxCY}$  difference showed a great dispersion. Indeed, the limits of agreement were between +17.4 and -24.9 bpm (+10.3 and -14.8 % in terms of % difference), with 7 of the 12 studied subjects showing a difference between tests higher than 5 %. All these facts suggested that substantial errors could be made when extrapolating  $HR_{max}$  in swimming from the cycling test. An even more striking difference was found when comparing the workload levels at which AT occurred. In fact, during the SW test, AT was detected at a higher % of  $W_{max}$  (at about 82 %) compared to all the other tests.

Since swimming requires less muscle mass (predominantly upper body) than running and cycling, it is tempting to assume that running and cycling testing should lead to a higher  $VO_{2max}$  than swimming. However, in the present study, this was not the case as the results showed that swimmers had on average a similar  $VO_{2max}$  level during the SW and RU tests, whereas they had an even higher  $VO_{2max}$  during the SW test as compared to the CY test. This fact could be explained by taking into account that  $VO_{2max}$  is training-sensitive. Hence, swimming training could induce in swimmers some specific adaptations leading to  $VO_{2max}$  enhancement only in their specific activity. It has in fact been suggested that the specifics of training affects the metabolic response of the musculature employed, and thus the metabolic response is specific to the training mode [14, 15, 20]. This is in accordance with what has been recently found by Roels and co-workers [30], who reported that swimmers had a higher  $VO_{2max}$  when assessed during swimming than during cycling. However, it is in contrast to earlier studies reporting that, in elite swimmers,  $VO_{2max}$  was lower during swimming than during running [14, 15]. We cannot explain these conflicting results. However, it should be noted that the earlier studies quoted used the

Douglas bag technique. This apparatus is cumbersome and not very comfortable to wear. Hence, it is possible that the Douglas bag limited athletes' movements so that they did not produce their top performance. In contrast, in the study by Roels et al. [30], who reported results similar to those of the present investigation, a snorkel similar to that of our investigation was employed. Thus, the different techniques used to collect expiratory gasses may explain the different outcomes of the various studies.

Another fact deserving attention is that the most striking difference between the SW test and the unspecific tests was found for AT, which occurred at a mean of 82.2 % of  $W_{max}$  during swimming, a value about 15–20 % higher than those detected in the other tests. Likewise for  $VO_{2max}$ , this fact could be the consequence of the specific adaptations induced by swimming training that enhanced the metabolic response during the specific exercise activity. This finding suggests that AT is more sensitive than  $VO_{2max}$  in detecting the specificity of training and that AT should be preferentially used when evaluating swimmers' fitness status. This conclusion is in good agreement with the concept that  $VO_{2max}$  is a valuable tool to distinguish between fit and unfit subjects, but it is not sensitive enough to discriminate between subjects of homogenous performance levels, since it suffers from limitations due to the fact that adequate motivation of the exercising subject is necessary to appropriately determine  $VO_{2max}$ , while sub-maximal parameters such as AT are less sensitive to motivation [13]. Moreover, while  $VO_{2max}$  reflects the integration of several mechanisms (ventilation, cardiac output, peripheral muscle  $O_2$  extraction, etc.), AT mainly reflects type II (fast-twitch glycolytic) muscle fibers recruitment, which are sensitive to training. In fact, the  $VO_2$  slow component, which is supposed to be linked to progressive recruitment of type II muscle fibers, has been described as being reduced in swimmers with a higher ventilatory threshold [25], thereby suggesting that mechanisms related to muscle recruitment are more specific indicators of athletes' fitness status than  $VO_{2max}$ . Thus, AT appears to be more sensitive than  $VO_{2max}$  for detecting the specificity of training. It has in fact been reported that anaerobic and ventilatory thresholds are important variables distinguishing endurance performance in athletes homogeneous in terms of  $VO_{2max}$  [5], and that AT is a more useful indicator of aerobic endurance performance than  $VO_{2max}$  [1, 13].

As far as  $HR_{max}$  is concerned, this variable showed a clear tendency to be lower in response to the SW than during the other tests. This fact is not novel since most of the research dealing with comparison between swimming and other kinds of effort has led to similar results [15, 28, 30]. Several factors could be responsible for the phenomenon: the different body position that improves cardiac filling and stroke volume, thereby reducing HR at a given cardiac



output; the enhanced vagal tone due to water immersion; and the reduced muscle mass used when swimming.

### Limitations of the study

Some technical considerations about the present investigation need to be acknowledged. First, as concerns the validity of the method employed in the present study, it should be emphasized that, from a biomechanical point of view, tethered swimming is different from free swimming since, among other factors, it does not take into account some critical aspects such as active and passive drag [10, 11]. However, previous research has reported that tethered swimming yielded essentially identical  $VO_{2max}$  values as free and flume swimming, thus indicating that results from tethered swimming tests were highly correlated with those of free swimming [8]. Another fact was that the additional dead space of the snorkel apparatus is likely to cause hyperventilation. However, results from additional experiments conducted with and without the snorkel suggested that VE was slightly increased by the snorkel only at sub-maximal workloads as compared to free breathing. Similarly to what has been described for VE, the snorkel device also induced a slight HR increase only at submaximal workloads. In contrast, the  $VO_2$  and  $VCO_2$  responses were substantially unaffected by the apparatus. Taken together, these facts led us to assume that the dead space added by the snorkel did not influence the outcome of the present study, which mainly focused on variables at AT and at maximum workload, i.e. when the snorkel device did not appear to significantly affect physiological response. Furthermore, athletes' mean value of  $VO_{2max}$  achieved (54.6 mL/kg/min, range 46.5–66.7 mL/kg/min) and workload level at AT (82.2 % of  $W_{max}$ ) were similar to those reported by previous studies in competitive swimmers [25, 27, 30], thus suggesting that the method employed in the present study yielded results comparable to those obtained by using a different commercially available snorkel.

Taken together, our data seem to support the snorkel method as a valid tool to assess swimmers' physical capacity in the pool setting. This method, although expensive, has been previously employed to assess the specificity of oxygen uptake kinetics, oxygen deficit, and energy cost in swimming [3, 4, 25, 26]. The present investigation provides further evidence that it is also suitable for  $VO_{2max}$  and AT assessment.

In conclusion, the present investigation provides evidence that none of the laboratory tests utilized to measure oxygen uptake in swimmers yielded results comparable to those obtained by tethered swimming testing. This fact strengthens the concept that swimmers should be tested in their specific activity and that predicting  $VO_{2max}$  for

swimming from other exercise testing procedures should be avoided. Moreover, the most specific parameter appears to be the anaerobic threshold, which shows the most relevant difference between swimming and the other tests.

**Acknowledgments** This study was supported by the University of Cagliari, the Italian Ministry of Scientific Research, and the National Olympic Committee of Italy. The authors wish to thank Mr. Corrado Sorrentino, Mr. Cesare Floris, Mr. Marco Cara, and the Atlantide Elmas, Rari Nantes Cagliari, and Esperia Cagliari sport clubs for their collaboration.

**Conflict of interest** None.

### References

- Allen WK, Seals DR, Hurley BF, Ehsani AA, Hagberg JM (1985) Lactate threshold and distance running performance in young and older endurance athletes. *J Appl Physiol* 58:1281–1284
- Bagger M, Petersen PH, Pedersen PK (2003) Biological variation in variables associated with exercise training. *Int J Sports Med* 24:433–440
- Barbosa TM, Keskinen KL, Fernandes R, Colaco P, Carmo C, Vilas-Boas JP (2005) Relationship between energetic, stroke determinants, and velocity in butterfly. *Int J Sports Med* 26:841–846
- Barbosa TM, Fernandes R, Keskinen KL, Colaco P, Cardoso C, Silva J, Vilas-Boas JP (2006) Evaluation of the energy expenditure in competitive swimming strokes. *Int J Sports Med* 27:894–899
- Basset Dr Jr, Howley ET (2000) Limiting factors for maximum oxygen uptake and determinants of endurance performance. *Med Sci Sports Exerc* 32:70–84
- Beaver WL, Wasserman K, Whipp BJ (1986) A new method for detecting anaerobic threshold by gas exchange. *J Appl Physiol* 60:2020–2027
- Bland JM, Altman DG (1986) Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1:307–310
- Bonen A, Wilson A, Yarkony M, Belcastro AN (1980) Maximal oxygen uptake during free, tethered, and flume swimming. *J Appl Physiol* 48:232–235
- Byard AD, Dengel DR (2002) Validity of a portable metabolic measurement system. *Med Sci Sport Exerc* 34:S149
- Capelli C, Pendergast DR, Termin B (1998) Energetics of swimming at maximal speeds in humans. *Eur J Appl Physiol* 78:385–393
- Di Prampero PE (1986) The energy cost of human locomotion on land and in water. *Int J Sports Med* 7:55–72
- Dixon RW, Faulkner J (1971) Cardiac output during maximum effort running and swimming. *J Appl Physiol* 30:653–656
- Faude O, Kindermann W, Meyer T (2000) Lactate threshold concepts. *Sports Med* 39:469–490
- Holmér I, Astrand P (1972) Swimming training and maximal oxygen uptake. *J Appl Physiol* 33:510–513
- Holmér I, Lundin A, Eriksson B (1974) Maximum oxygen uptake during swimming and running by elite swimmers. *J Appl Physiol* 36:711–714
- Howley ET, Bassett DR, Welch HG (1995) Criteria for maximal oxygen uptake: review and commentary. *Med Sci Sport Exerc* 27:1292–1301
- Kohrt WM, Morgan DW, Bates B, Skinner JS (1987) Physiological responses of triathletes to maximal swimming, cycling, and running. *Med Sci Sport Exerc* 19:51–55

18. Kohrt WM, O'Connor JS, Skinner JS (1989) Longitudinal assessment of responses by triathletes to swimming, cycling, and running. *Med Sci Sport Exerc* 21:569–575
19. Jones AM, Doust JH (1996) A 1 % treadmill grade most accurately reflects the energetic cost of outdoor running. *J Sport Sci* 14:321–327
20. Magel JR, Foglia GF, McArdle WD, Gutin B, Pechar GS, Katch FI (1975) Specificity of swim training on maximum oxygen uptake. *J Appl Physiol* 38:151–155
21. Midgley AW, Carroll S (2009) Emergence of the verification phase procedure for confirming “true”  $VO_{2max}$ . *Scand J Med Sci Sports* 19:313–322
22. Montpetit R, Léger LA, Lavoie JM, Cazorla G (1981)  $VO_{2peak}$  during free swimming using the backward extrapolation of the  $O_2$  recovery curve. *Eur J Appl Physiol* 47:385–391
23. Olson TP, Tracy JE, Dengel DR (2003) Validity of a low-flow pneumotach and portable metabolic system for measurement of basal metabolic rate. *Med Sci Sport Exerc* 35:S143
24. Poujade B, Hautier CA, Rouard A (2002) Determination of the energy cost of front-crawl swimming in children. *Eur J Appl Physiol* 87:1–6
25. Reis JF, Alves FB, Bruno PM, Vleck V, Millet GP (2012) Effects of aerobic fitness on oxygen uptake kinetics in heavy intensity swimming. *Eur J Appl Physiol* 112:1689–1697
26. Reis JF, Alves FB, Bruno PM, Vleck V, Millet GP (2012) Oxygen uptake kinetics and middle distance swimming performance. *J Sci Med Sport* 15:58–63
27. Reis VM, Marinho DA, Barbosa FP, Reis AM, Guidetti L, Silva AJ (2010) Examining the accumulated oxygen deficit method in breakstroke swimming. *Eur J Appl Physiol* 109:1129–1135
28. Rennie DW, Di Prampero PE, Cerretelli P (1971) Effects of water immersion on cardiac output, heart rate and stroke volume of man at rest and during exercise. *Med Sport* 24:222–228
29. Ribeiro JP, Cadavid E, Baena J, Monsalvete E, Barna A, De Rose EH (1990) Metabolic predictors of middle-distance swimming performance. *Br J Sports Med* 24:196–200
30. Roels B, Schmitt L, Libicz S, Bentley D, Richalet J-P, Millet G (2005) Specificity of  $VO_{2max}$  and the ventilatory threshold in free swimming and cycle ergometry: comparison between triathletes and swimmers. *Br J Sports Med* 23:965–968
31. Scharag-Rosemberg F, Carlsohn A, Cassel M, Mayer F, Scharhag J (2011) How to test maximal oxygen uptake: a study on timing and testing procedure of a supramaximal verification test. *Appl Physiol Nutr Metab* 36:153–160
32. Smith DJ, Norris SR, Hogg JM (2002) Performance evaluation of swimmers. *Sports Med* 32:539–554
33. Zamparo P, Bonifazi M, Faina M, Milan A, Sardella F, Schena F, Capelli C (2005) Energy cost of swimming of elite long-distance swimmers. *Eur J Appl Physiol* 94:697–704
34. Zamparo P, Capelli C, Cautero M, Di Nino A (2000) Energy cost of front-crawl swimming at supra-maximal speeds and underwater torque in young swimmers. *Eur J Appl Physiol* 83:487–491