



26th International Conference on Nuclear Tracks in Solids, 26ICNTS

Radon Concentration by SSNTD in South-East Sicily buildings

G. Immè^{a,b}, R. Catalano^{a,b}, C. Gianino^{b,c}, R. Filincieri^a, G. Mangano^a, D. Morelli^{a,b,*}

^a*Dipartimento di Fisica e Astronomia Università degli Studi di Catania, Via S. Sofia, 64, Catania 95123, Italy*

^b*INFN Sezione di Catania, Via S. Sofia, 64, Catania 95123, Italy*

^c*Liceo Scientifico Statale "Enrico Fermi" - Ragusa*

Abstract

Radon levels in buildings vary widely from area to area also depending on local geology. Thus, it is important to assess the radon prone area of a geographic region on the basis of geological data and to search for any possible correlation between the local geology and the indoor radon concentrations. We report about indoor radon measurements in Ragusa, a municipality of the SE Sicily, placed in the Hyblean Plateau (northern region of the African Plate), carried out in collaboration with schools. The survey was performed using Solid State Nuclear Track Detectors (SSNTD), CR-39 type, and a well-established methodology for chemical etching and reading, developed at the Radioactivity Laboratory of the Department of Physics - University of Catania.

© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Peer-review under responsibility of the Scientific Committee of 26ICNTS

Keywords: Radon, SSNTD

1. Introduction

Inhalation of Radon and its progeny is one of the principal sources of radiation exposure of the population (UNSCEAR, 2000). High radon exposure can have effects on the public health, Radon, in fact, is considered as the second cause of lung cancer after smocking (NAS, 1999). So it is very important to plan Radon monitoring survey. In many countries all around the world mapping of radon risk area or radon-prone area are in progress in accordance to the International Committee on Radiological Protection Recommendations (ICRP, 1993). Since 2005 in the framework of a scientific dissemination INFN (National Institute for Nuclear Physics) project, while high school students are informed on natural radioactivity and its health effects, indoor radon activity mapping in dwellings and schools is performed in the eastern Sicily. During the project the involved students firstly receive explanation about

* Corresponding author. Tel.: +39 095 3785222; fax: +0-000-000-0000 .

E-mail address: daniela.morelli@ct.infn.it

radon and on nuclear particles detection techniques; they also assemble themselves radon detectors and place them in the classrooms at school and at home (Catalano et al., 2012). In this way it is possible to monitor vast areas.

We report about the results of a Radon survey carried out in Ragusa municipality from October 2012 till October 2013.

Ragusa is located in the central Hyblean Plateau in the south-east of Sicily (Fig. 1.). The Hyblean Plateau is part of the northern margin of the African plate and has remained a relatively undeformed foreland during the Neogene collisional process affecting the African-European convergent belt; it consists of circa 6000 m of carbonates and marls with intercalations of volcanic horizons, which have occurred at several episodes (Patacca et al., 1979; Bianchi et al., 1989; Ragg et al., 1999). The rock successions in which Ragusa lies are Oligo-Miocene carbonates.




Fig. 1. Radon survey area.

2. Indoor Radon measurements

Indoor radon measurements were performed using SSNTD Cr-39 Type. Measurements were carried out from October 2012 till October 2013 in fifty different buildings. Two sets of detector were exposed, the first one from October 2012 till May 2013, the second one from May 2013 till October 2013. In particular, in thirty-five buildings detectors were placed in the same sites in the two measurement sets to evaluate seasonal influences and to obtain radon concentration values as mean over one year as indicated by the European and national radon regulation.

After the exposure, the CR-39 dosimeters were chemically etched using a 6.25 M NaOH solution at 98°C for 1 hour. The detector reading was carried out by means a semiautomatic system composed of an optical microscope equipped with a CCD camera connected to a PC. A video acquisition software allowed to capture and store the images (Field Of View or FOV) from the microscope. The stored images were analyzed by means of ImageJ 1.29x (Image Processing and Analysis in Java) freeware software, in which an opportune routine was edited (Immè et al., 2013).

The obtained results are shown in Fig. 2.




Fig. 2. (a) Indoor Radon concentration in 50 dwellings (October 2012- May 2013); (b) Indoor Radon concentration in 35 dwellings (May 2013- October 2013)

All the data are lower than the National law limits (200 Bq/m³). In table 1 the detail of the obtained values is reported.

Table 1. Indoor Radon concentrations – statistical analysis on monitored sites

Exposition period	Min [Bq/m ³]	Max [Bq/m ³]	Avg [Bq/m ³]	S.D. [Bq/m ³]	Median [Bq/m ³]
October 2012-May 2013	12,71	168,63	43,51	30,59	35,6
May 2013- October 2013	14,11	156,88	41,06	26,39	36,3

No relevant differences were found between summer and winter radon concentration values. In Fig. 3. the data are reported for the two periods recorded in the same buildings.




Fig. 3. Comparison between Radon concentration recorded in the two survey periods in the same thirty-five buildings

Only few buildings shown a high seasonal variability possibly due to the inhabitant's lifestyle. The previous data were, also, analyzed with respect to the building materials. The results are shown in Table 2, where the mean radon value is reported according to the different materials and the highest and lowest values.

Table 2. Radon concentration according to building materials

Building Material	Mean [Bq/m ³]	Max [Bq/m ³]	Min [Bq/m ³]
Concrete	18	18	18
Cement	38	79	20
Block of limestone	79	122	38
Bricks	18	20	16
Bricks and cement	42	120	18
Stone	50	150	18

The obtained results have shown that there is an high variability in indoor radon concentration for each material.

3. Conclusion

A survey of radon concentration in Ragusa, a city placed in the south-east of Sicily was carried out in the period October 2012 – October 2013. The radon concentration was measured using solid state nuclear tracks detectors. The results of the present work have shown medium indoor radon concentrations with values around the previously estimated mean value of 35 Bq/m³ for the Sicily (Bochicchio et al. 1999) but lower respect to the Italian average of 70 Bq/m³ and much lower respect to our previous survey in the same area (Catalano et al., 2012). The differences are imputable to the larger detector distribution density inside a limited area. These results suggest that for a better indoor radon mapping it is necessary to optimize the detector distribution according to the geomorphology of the area.

Acknowledgements

We thank the students and the teachers of the Liceo scientifico “Enrico Fermi” of Ragusa for their participation to the project and help in the dosimeter placing.

References

- Abu-Jarad, F., 1998. Application of nuclear track detectors for radon related measurements, Nucl.Tracks Radiat. Meas. 15, 525-534
- Bianchi, F., Carbone, S., Grasso, M., Invernizzi, G., Lentini, F., Longaretti, G., Merlini, S., and Moscardini, F., 1989. Sicilia orientale: Profilo geologico Nebrodi-Iblei. Mem. Soc. Geol. It. 38, 429-458.
- Bochicchio, F., Campos Venuti, G., Piermattei, S., Torri, G., Nucitelli, C., Risica, S., Tommasino, L. 1999. Results of the National Survey on radon indoors in all the 21 Italian Regions. In: Proceedings of the workshop “Radon in the living environment” Athens 19-23 April 1999.
- Catalano, R., G. Immè, G. Mangano, D. Morelli, A. Rosselli Tazzer, 2012 , Indoor radon survey in Eastern Sicily, Rad. Meas. 47, 105-110.
- ICRP, Protection against ²²²Rn at home and at work, ICRP Publication 65, Annals 23 (1993), pp. 1-45.
- Immè, G., Morelli, D., Aranzulla, M., Catalano, R., Mangano, G. 2013, Nuclear track detector characterization for alpha-particle spectroscopy, Rad. Meas. 50, 253-257.
- National Academy of Science (NAS) Reports, 1999. Health Effects of Exposure to Radon, Committee on Health Risks of Exposure to Radon (BEIR VI). National Academy Press.
- Patacca, E., Scandone, P., Giunta, G., and Liguori, V., 1979. Mesozoic paleotectonic evolution of the Ragusa zone (southeastern Sicily). Geol. Rom. 18, 331-369.
- Ragg, R., Grasso, M., Muller, B., 1999. Patterns of tectonic stress in Sicily from borehole and breakout observations and finite element modeling. Tectonics 18, 669-685
- United Nations Scientific Committee on the Effects of Atomic Radiation (UNSCEAR), 2000. Sources and Effects of Ionizing Radiation. 2000 Report to the General Assembly, with Annex. United Nations. New York