



Huge myxoid chondrosarcoma expanded into the thoracic cavity with spinal involvement

Ming Lu^{1,2} · Zhongxin Zhou³ · Zixiong Lei^{1,2} · Haomiao Li^{1,2} · Stefano Boriani⁴

Received: 22 February 2018 / Revised: 20 May 2018 / Accepted: 26 June 2018 / Published online: 4 July 2018
© The Author(s) 2018

Abstract

Purpose En bloc resection is the treatment of choice of myxoid chondrosarcoma. These tumors can produce huge masses. Anatomical constraints limit the possibility to perform en bloc resection in the spine.

Methods A very huge myxoid chondrosarcoma (14.2 × 10.8 × 11.4 cm) arising from T2 to T5 and invading the whole higher left pleural cavity was observed. Surgical planning according to WBB staging system was performed.

Results The tumor was successfully submitted to en bloc resection achieving a tumor-free margin as demonstrated by the pathologist's report.

Conclusions A careful planning and a multidisciplinary collaboration make possible to perform en bloc resection even in apparently impossible cases.

Keywords En bloc spondylectomy · Spinal myxoid chondrosarcoma · WBB staging system · Multidisciplinary collaboration

Introduction

Myxoid variety of chondrosarcoma is known to produce huge masses [1–3]. This low-malignant tumor is radiore-sistant [4] and has a very high tendency to recurrence when submitted to intralesional excision [5–9, 13]. Chemotherapy has been proved ineffective and hence not recommended [10–12]. Its tendency to progress to higher level of malignancy at recurrence is also known. En bloc resection is therefore the treatment of choice [2, 3, 13–15]. Regional

constraints and the dimension of some spine tumor can make impossible to perform en bloc resection [16–20].

This case is reported to demonstrate that careful planning, including multidisciplinary competences, makes possible to perform en bloc resection even in huge tumors.

Case report

A 34-year-old female was admitted to our department presenting with complains of increasing pain at the back, with a mass on the left back and complete paralysis of the lower extremities for 1 month. The patient initially presented to another hospital 3 months prior to this with a mass on the left back. An open incisional biopsy was performed at the left back in her primary hospital prior to our workup, and the diagnosis from histopathological analysis was of a low-grade myxoid chondrosarcoma.

A physical examination showed that there was complete paralysis of the lower extremities and decreased sensation below the trunk, with dysfunction of the bladder and bowel movement (ASIA impairment scale: B). Plain radiography, computed tomography and magnetic resonance imaging demonstrated an expansive mediastinal mass lesion with thoracic vertebral bodies and ribs involved (Fig. 1). Further

Ming Lu, Zhongxin Zhou and Zixiong Lei have contributed equally to this work.

✉ Haomiao Li
lihaomiao1977@hotmail.com

¹ Department of Orthopedic, The Third Affiliated Hospital of Southern Medical University, Guangzhou 510630, China

² Academy of Orthopaedics, Guangdong Province, Guangzhou 510630, China

³ Department of Interventional Vascular Surgery, The Third Affiliated Hospital of Southern Medical University, Guangzhou 510630, China

⁴ GSpine4 Spine Surgery Unit, IRCCS Galeazzi Orthopaedic Institute, Milan, Italy




Fig. 1 **a** Coronal, **b** sagittal and **c** axial MRI images showing a huge expansive mediastinal mass lesion (14.2×10.8×11.4 cm) invading the thoracic vertebral bodies and ribs and spinal cord compression are demonstrated




Fig. 2 Chest CT scan, showing the adjacent important structures of the mediastinum. There is a large soft tissue mass within the mottled calcification

investigation revealed no other tumor lesions existed. In the present case, the mass had become involved with the spinal canal and adhered to the aorta, trachea and esophagus (Figs. 2, 3) (Enneking stage IB and WBB stage 1-7/A-D). Selective arterial embolization was performed 1 day prior to the surgery. En bloc resection with a tumor-free margin by one-stage posterior approach was carefully designed by a multidisciplinary team.

Surgical planning by posterior approach (WBB-based en bloc resection type 2a, Fig. 4) [21] and detachment of the spinal muscles from the posterior elements from T2 to T5 were performed. The whole procedure was carried out in the prone position. First, the right pleural cavity was opened by section of the T2–T5 ribs, ligation of the azygos, release of the aortic arch and the esophagus. A dissection of the mass was performed by section of the T2–T6 left ribs and the left scapula. Lateral mass screws were implanted

at C6, C7. Pedicle screws were implanted at the levels of T1, T6, T7 and T8. A wide right semi-laminectomy was performed by ultrasonic osteotome from T2 to T5, and the T2–T5 right nerve roots were sacrificed. Then, sectors 8-9-10 were piecemeal excised by high-speed burr to approach the spinal canal at levels T2–T5. A rod was connected on the right side, and osteotomy was performed by Tomita saw at T1/T2 and T5/T6 disk levels. At this time, the huge mass can be rotated on the longitudinal axis, allowing to separate the thecal sac from the mass in the spinal canal under direct vision. The T2–T5 left nerve roots are sectioned after ligation. Then, the whole mass is finally removed without any apparent surgical violation of the margins (Fig. 5a, c, d). The reconstruction was performed by four rods connection at bilateral C6, C7, T6 and T1, T7, T8, respectively. A titanium cage was inserted between the T1–T6 vertebral body, and two vertebral body screws were implanted at T1 and T6 to reinforce the fixation of the titanium cage. Reconstruction of the ribs defect was performed by using two rods connecting the screws and the defect ends in the left side (Figs. 5b, 6d). A chest tube was placed to re-expand the lung.

The operation took 15 h in total, with a total of 5500 mL of bleeding. Ventilator-assisted ventilation was used for 17 days after operation in intensive care unit. After respiratory function exercise for 1 month, she can breathe normally without hypoxic symptoms. Pulmonary infection occurred postoperatively, and etiological examination suggested *Acinetobacter Bauman* infection. Chest drainage sustained for 64 days. Anti-infective treatment regimen was based on the results of bacterial susceptibility. Imipenem combined with cefoperazone was used for the first month and then cefoperazone only until chest drainage was removed. A slowly skin sensation recovery was observed on the trunk postoperatively. No improvement in muscle




Fig. 3 **a** The sagittal T2-weighted MRI image demonstrates the epidural mass and spinal cord compression. **b, c** The sagittal T1-weighted MRI images show low signal intensity in the T2–T5 vertebral bodies and tumor mass




Fig. 4 WBB-based planning of the en bloc resection. Posterior approach only, patient in prone position (type 2b). Step I: completely release of the tumor mass from left and from right. Step II: definition of the approach to the canal: sectors 8–12. Step III piecemeal excision of sectors 8–12, release of the dural sac after section of the nerve roots. Step IV: En bloc tumor removal by careful clockwise rotation along the longitudinal axis

strength of the lower extremities and defecation functions was observed at 3 months postoperatively. Reconstructed postoperative CT scan at 3 months showed no signs of tumor recurrence and well-united bone at the reconstruction site (Fig. 6).


Discussion

From the earliest report by Lievre [22], Hamdi [23] and Stener [24, 25], en bloc resection was later on popularized by Roy Camille [26] and Tomita [27, 28] till more recent attempt to standardize the surgical planning according to the WBB staging system [21, 29]. En bloc resection with tumor-free margins remains the best oncological management of the spine, which aims to a better local control and longer survival rate.

Recurrence of chondrosarcoma usually occurs in 3–5 years postoperatively, and the relapse tends to be faster if an intralesional excision was performed [5, 30]. Thus, a worse prognosis could be achieved. These results were proved to be related to the inadvertent intraoperative contamination [5, 31]. Regional anatomical constraints and huge volumes remain, however, a severe limit to perform an oncological appropriate en bloc resection (it means resulting in tumor-free margins without any tumor contamination), and some reports in the literature of procedures including the separation of the tumor in two or more pieces with intralesional margins [16–20]. A long-term survival rate has only been observed in low-grade spinal chondrosarcoma patients after repeated intralesional excision surgeries, combined with radiotherapy [32].

Here, a case of en bloc resection of a huge myxoid chondrosarcoma is reported, resulting in a fully tumor-free margin resection. To our knowledge, this is one of the largest tumors ever resected by this technique. The peculiar steps were: the ligation of the azygos, the release of the aortic arch and of the esophagus, the opening of a window in the posterolateral vertebrae (T2–T5 in sectors

Fig. 5 **a** Photograph of the en bloc resected specimen. **b** Photograph demonstrates the reconstruction of the chest wall defect and spinal stability by screws and rods. **c, d** The X-ray images of the en bloc resected specimen



8-9-10) in order to release the dural sac. Once the osteotomies of the spine at the T1/T2 and T5/T6 disk levels were performed, the specimen could rotate, allowing to finalize the resection.


The goal of this surgical procedure was to remove the tumor en bloc with macroscopically healthy margins. Owing to its huge dimension and proximity to vital neurovascular structures, the surgical morbidity and functional impairment must be seriously considered. In the present case, multiple ribs and partial chest wall tissues were excised combined with the mass. The chest wall has both a protective structural around the vital organs of the body and functional role of respiratory movement. Large chest wall defects can cause devastating respiratory and circulatory functional consequences [33, 34]. Reconstruction of the chest wall defect was performed to recreate a

stable chest wall with adequate functional capacity. Fadel et al. [35] reported an incidence of pneumonia of 35.3%: 6 patients suffered from pneumonia postoperatively in 17 patients following tumor resection involving the pleura and ribs. The use of sensitive antibiotics in the treatment of pulmonary infection and appropriate ventilator-assisted respiratory is critical for postoperative recovery.

Conclusions

The purpose of this article is to remark the role of preoperative planning in the surgical technique of en bloc resection. However, a careful follow-up will be mandatory for detecting local recurrence, whose risk—due to the histologically proven appropriate margin—is low but not negligible. The procedure

Fig. 6 Reconstructed **a** sagittal, **b** coronal, **c** axial and **d** three-dimensional computed tomography scan 3 months postoperatively



was successfully performed thanks to a very careful preoperative planning and to a multidisciplinary collaboration.

Acknowledgements The authors are indebted to Mr. Carlo Piovani for his excellent drawing interpretation of our surgical planning.

Compliance with ethical standards

Conflict of interest statement None of the authors has any potential conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Liu G, Wu G, Ghimire P, Pang H, Zhang Z (2013) Primary spinal chondrosarcoma: radiological manifestations with histopathological correlation in eight patients and literature review. *Clin Imaging* 37(1):124–133. <https://doi.org/10.1016/j.clinimag.2012.02.010> **Epub 2012 Jun 8**
2. Quiriny M, Gebhart M (2008) Chondrosarcoma of the spine: a report of three cases and literature review. *Acta Orthop Belg* 74(6):885–890
3. Casadei R, Ricci M, Ruggieri P, Biagini R, Benassi S, Picci P, Campanacci M (1991) Chondrosarcoma of the soft tissues. Two different sub-groups. *J Bone Joint Surg Br* 73(1):162–168
4. Gelderblom H, Hogendoorn PCW, Dijkstra SD et al (2008) The clinical approach towards Chondrosarcoma. *Oncologist* 13:320–329

5. Boriani S, De Iure F, Bandiera S, Campanacci L et al (2000) Chondrosarcoma of the mobile spine: report on 22 cases. *Spine (Phila Pa 1976)* 25:804–812
6. Schoenfeld AJ, Hornicek FJ, Pedlow FX et al (2012) Chondrosarcoma of the mobile spine: a review of 21 cases treated at a single center. *Spine (Phila Pa 1976)* 37:119–126
7. Yang X, Wu Z, Xiao J et al (2012) Chondrosarcomas of the cervical and cervicothoracic spine: surgical management and long-term clinical outcome. *J Spinal Disord Tech* 25:1–9
8. Bergh P, Gunterberg B, Meis-Kindblom JM, Kindblom LG (2001) Prognostic factors and outcome of pelvic, sacral, and spinal chondrosarcomas: a center-based study of 69 cases. *Cancer* 91:1201–1212
9. Boriani S, Saravanja D, Yamada Y et al (2009) Challenges of local recurrence and cure in low grade malignant tumors of the spine. *Spine (Phila Pa 1976)* 34(22 Suppl):S48–S57
10. D'Ambrosio FG, Shiu MH, Brennan MF (1986) Intrapulmonary presentation of extraskelatal myxoid chondrosarcoma of the extremity: report of two cases. *Cancer* 58:1144
11. Saleh G, Evans HL, Ro JY, Ayala AG (1992) Extraskelatal myxoid chondrosarcoma: a clinicopathologic study of ten patients with long-term followup. *Cancer* 70:2827
12. Patel SR, Burgess MA, Papadopoulos NE, Linke KA, Benjamin RS (1995) Extraskelatal myxoid chondrosarcoma: long-term experience with chemotherapy. *Am J Clin Oncol* 18:161
13. Boriani S, Bandiera C, Biagini DF et al (2000) Chondrosarcoma of the Mobile Spine: report on 22 Cases. *Spine* 25(7):804–812
14. Prevedello DM, Cordiero JG, Koerbel A, Ditzel LF, Araujo JC (2004) Management of primary spinal chondrosarcoma: report of two cases causing cord compression. *Arq Neuropsiquiatr* 63(3):875–878
15. Xinghai Y, Zhipeng W, Jianru DF, Quan H, Wei Z et al (2012) Chondrosarcomas of the cervical and cervicothoracic spine: surgical management and long-term clinical outcome. *J Spinal Disord Tech* 25(1):1–9
16. Demura S, Kawahara N, Murakami H, Akamaru T, Kato S, Oda M et al (2012) Giant cell tumor expanded into the thoracic cavity with spinal involvement. *Orthopedics* 35(3):e453–e456
17. Rena O, Davoli F, Allegra G, Casadio C, Turello D (2014) Giant chordoma of the upper thoracic spine with mediastinal involvement: a surgical challenge. *Asian Spine J.* 8(3):353–356. <https://doi.org/10.4184/asj.2014.8.3.353> **Epub 2014 Jun 9**
18. Selvaraj A, Wood AJ (2003) Superior mediastinal chordoma presenting as a bilobed paravertebral mass. *Eur J Cardiothorac Surg* 23:248–250
19. Topsakal C, Bulut S, Erol FS, Ozercan I, Yildirim H (2002) Chordoma of the thoracic spine: case report. *Neurol Med Chir (Tokyo)* 42:175–180
20. Chen B, Yang Y, Chen L, Zhou F, Yang H (2014) Unilateral lateral mass fixation of cervical spinal low-grade chondrosarcoma with intralésional resection: a case report. *Oncol Lett* 7(5):1515–1518 **Epub 2014 Mar 7**
21. Boriani S, Bandiera S, Colangeli S, Ghermandi R, Gasbarrini A (2014) En bloc resection of primary tumors of the thoracic spine: indications, planning, morbidity. *Neurol Res* 36:566–576
22. Lièvre JA, Darcy M, Pradat P, Camus JP, Bénichou C, Attali P, Joubin M (1968) Giant cell tumor of the lumbar spine; total spondylectomy in 2 states. *Rev Rhum Mal Osteoartic* 35(3):125–130
23. Hamdi FA (1969) Prosthesis for an excised lumbar vertebra: a preliminary report. *Can Med Assoc J* 100(12):576–580
24. Stener B, Johnsen OE (1971) Complete removal of three vertebrae for giant-cell tumour. *J Bone Joint Surg Br* 53(2):278–287
25. Stener B (1971) Total spondylectomy in chondrosarcoma arising from the seventh thoracic vertebra. *J Bone Joint Surg Br* 53(2):288–295
26. Roy-Camille R, Saillant G, Bisserie M, Judet TH, Hautefort E, Mamoudy P (1981) Resection vertebrale totale dans la chirurgie tumorale au niveau du rachis dorsal par voie posterieure pure. *Rev Chir Orthop* 67:421–430
27. Tomita K, Kawahara N, Baba H, Tsuchiya H, Nagata S, Toribatake Y (1994) Total en bloc spondylectomy for solitary spinal metastasis. *Int Orthop* 18:291–298
28. Tomita K, Kawahara N, Baba H, Tsuchiya H, Fujita T, Toribatake Y (1997) Total en bloc spondylectomy: a new surgical technique for primary malignant vertebral tumors. *Spine* 22:324–333
29. Boriani S, Weinstein JN, Biagini R (1997) Primary bone tumors of the spine. Terminology and surgical staging. *Spine (Phila Pa 1976)* 22(9):1036–1044
30. York JE, Berk RH, Fuller GN, Rao JS, Abi-Said D, Wildrick DM, Gokaslan ZL (1999) Chondrosarcoma of the spine: 1954 to 1997. *J Neurosurg* 90(1 Suppl):73–78
31. Matsumoto Y, Takahashi Y, Harimaya K et al (2013) Dedifferentiated chondrosarcoma of the cervical spine: a case report. *World J Surg Oncol* 11:32
32. Ozaki T, Lindner N, Hillmann A et al (1996) Influence of intralésional surgery on treatment outcome of chondrosarcoma. *Cancer* 77:1292–1297
33. Bennett DT, Weyant MJ (2014) Extended chest wall resection and reconstruction in the setting of lung cancer. *Thorac Surg Clin* 24(4):383–390
34. Yang H, Tantai J, Zhao H (2015) Clinical experience with titanium mesh in reconstruction of massive chest wall defects following oncological resection. *J Thorac Dis.* 7(7):1227–1234
35. Fadel E, Missenard G, Chapelier A, Mussot S, Leroy-Ladurie F, Cerrina J, Darteville P (2002) En bloc resection of non-small cell lung cancer invading the thoracic inlet and intervertebral foramina. *J Thorac Cardiovasc Surg* 123(4):676–685