

## REVIEW ARTICLE

## IMMUNOTOXICITY AND SENSITIZING CAPACITY OF METAL COMPOUNDS DEPEND ON SPECIATION

M. DI GIOACCHINO, N. VERNA, L. DI GIAMPAOLO, F. DI CLAUDIO, M.C. TURI, A. PERRONE, C. PETRARCA, R. MARIANI-COSTANTINI<sup>1</sup>, E. SABBIONI<sup>2</sup> and P. BOSCOLO<sup>3</sup>  
*Allergy Related Diseases Unit and <sup>1</sup>Molecular Pathology Unit, Ageing Research Center (CeSI) "G. d'Annunzio" University Foundation, Chieti; <sup>2</sup>Joint Research Center of the European Commission, ECVAM, Ispra; <sup>3</sup>Dept of Neurosciences, "G. d'Annunzio" University, Chieti, Italy*

Immunotoxicity of metal compounds is an issue of great importance due to the recent industrial application of metals with unknown toxicity on the immune system and the discovery of metal intermediary compounds not sufficiently studied yet. In this report we show results of our study on the immunotoxicity of the following metals: the Platinum group elements (Platinum, Palladium, Rhodium), Titanium and Arsenic. We applied functional and non functional assays and investigated both innate and adaptive immune systems, in particular, cell proliferation, cytokine production by PBMCs and O<sub>2</sub> production by neutrophils. We obtained the following results: only some Ti compounds (Titanocene, Ti ascorbate and Ti oxalate) show immunotoxicity. Trivalent As compounds (Sodium arsenite and tetraphenyl arsonium chloride) are more immunotoxic than the other investigated As compounds. Genotoxicity of Pt group compounds is in the following order: Pt > Rh > Pd. Immunotoxicity of Pt group compounds is in the following order: Pd > Pt > Rh. Lymphocytes and macrophages show a different reaction of neutrophils to metal toxicity. We can conclude that these studies show that metal immunotoxicity depends on speciation. In general speciation provides additional and often essential information in evaluating metal toxicity. However, there are many difficulties in applying speciation in investigating toxicokinetic aspects to many metals, mainly due to the lack of information about the existence and significance of species and to the lack of analytical methods for measuring species in biological samples.

Immunotoxicity of metals compounds is an issue of a great importance due to the recent industrial application of metals with unknown toxicity on the immune system and the discovery of metal intermediary compounds not sufficiently studied yet. Metals have various interactions with the immune system: essential elements, toxic or sensitizing (Table I).

Some metals, like chromium, have multiple interactions: trivalent chromium is an essential element (1), hexavalent has a strong toxicity (2) and chromium salt (potassium dichromate) may sensitize humans. Vanadium (V) is an element with wide industrial applications and environmental release, and 10<sup>-4</sup> M NaVO<sub>3</sub> exerts inhibitory effects on PBMCs, while at 10<sup>-7</sup> M it exerts a stimulatory action with a slight shift of the immune response towards a Th2-type response (3). Inhibitory effects of cadmium on peripheral blood mononuclear cell proliferation and cytokine release are reversed by zinc and selenium salts (4). Arsenic (As) also shows multiple activity on the immune system depending on speciation and dose (5). In fact, 10<sup>-7</sup> M trivalent As and DMAs increase *in vitro* proliferation of Peripheral Blood Mononuclear Cells (PBMC), showing a stimulatory metabolic effect, so

considered to be an essential element (6-7). Results of experimental studies indicate the beneficial effect of As in drinking water with a content ranging from 12 to 50 µg/l (7), while higher levels, exceeding 200 µg/l are reported to exert toxic effects in humans (8).

Platinum (Pt) salts may induce allergic asthma in exposed workers, whereas halogenated Pt compounds [(NH<sub>4</sub>)<sub>2</sub>PtCl<sub>4</sub>; (NH<sub>4</sub>)<sub>2</sub>PtCl<sub>6</sub>] strongly inhibit lymphocyte proliferation and cytokine release *in vitro* (9-10). In previous studies, Pt allergy seemed to be confined to a small group of charged compounds containing reactive ligand systems (the most effective of which being chloride ligand systems). While metallic Pt in the metallic, oxide or other stable forms showed lower sensitizing effects (11). In the mouse popliteal lymph-node assay, finely dispersed Pt, and PtO<sub>2</sub>, PtCl<sub>2</sub> and [Pt(NH<sub>2</sub>)<sub>4</sub>]Cl<sub>2</sub> did not induce an immune response, while Na<sub>2</sub>PtCl<sub>6</sub> and Na<sub>2</sub>PtCl<sub>4</sub> induced an increase of cells (mainly CD4<sup>+</sup>) which express proliferating cell nuclear antigens (12-13). These different *in vivo* immune effects of Pt compounds were related to their capacity to modulate *in vitro* mechanisms of receptor-mediated endocytosis in murine Langerhans cells (12). Na<sub>2</sub>PtCl<sub>6</sub> and Na<sub>2</sub>PtCl<sub>4</sub> exerted a more

*Key words: immunotoxicity, PBMC culture, cytokines, metal speciation*

Mailing address: Prof. Mario Di Gioacchino,  
Allergy Related Diseases Unit,  
Ageing Research Center (CeSI),  
Via Colle dell'Ara,  
66100 Chieti, Italy  
Tel: ++39 871 358578 Fax: ++39 871 551615  
e-mail: m.digioacchino@unich.it

0394-6320 (2007)

Copyright © by BIOLIFE, s.a.s.

This publication and/or article is for individual use only and may not be further reproduced without written permission from the copyright holder. Unauthorized reproduction may result in financial and other penalties

potent contact sensitizing effect than  $[\text{Pt}(\text{NH}_3)_4]\text{Cl}_2$  in endocytosis assay. The results of this experimental study are in agreement with those of an investigation on workers in a plant producing auto-catalysts: workers exposed to chloroplatinate (IV) compounds showed a cumulative probability of 51% of being sensitized after 5 years of exposure, while those exposed to the tetra-ammine Pt(II) dichloride alone were not sensitized (14).

Among the sensitizing metals, nickel may cause different immune reactions and symptoms depending on the type of exposure (15). Inhaled Ni can cause allergic asthma (IgE mediated 1<sup>st</sup> type of immune reactions). In a large number of subjects skin contact with Ni results in an allergic contact dermatitis mediated by a 4<sup>th</sup> type cell-mediated immune reaction. Food ingested Ni may have clinically relevant immune effects: gastrointestinal symptoms and a flare up of skin lesions. Studies have shown a decrease of NK cell activity (16-17) and apoptosis of CD8<sup>+</sup> cells in the intestinal epithelium, where CD4<sup>+</sup>45RO<sup>+</sup> cells proliferate (18).

The *in vitro* effects of tellurium compounds were also determined: Te(IV) inhibited PBMC proliferation and IFN-gamma, IL-5 and TNF-alpha release from PBMC more than Te (VI) (19). In 1998, we started, in collaboration with the "European Center for the Validation of Alternative Methods", a study in order to determine the potential immunotoxicity of metal compounds. It was a mechanistic study that compared the effects of a large number of metals and their compounds. In this report we will discuss the main results obtained by the experiments on titanium, platinum, palladium, rhodium and arsenic.

For each metal, we chose the compounds present in the environment, forms employed in the industry of prevalent occupational importance, present in the food chain or utilized in the treatment of many pathologies like cancer. The oxalate of titanium has wide industrial applications, like the aero-spatial industry, the ascorbate is mainly used in agriculture, the titanocene is employed as chemotherapeutic agent and the dioxide is a biocompatible component of sunscreens. The Platinum group elements (platinum, palladium, rhodium), used for jewellery and for making catalytic converters, are released into the environment both by vehicular traffic and from industries. Many As compounds are part of the food chain and are ingested with food or metabolized in our body. Some metabolites of arsenic, recently detected, are synthesized after the absorption of food containing As. They seem to be responsible for the genotoxicity and cancerogenicity of As. Table II shows the complete list of utilized metal compounds.

#### MATERIALS AND METHODS

Various methods are employed in the study of immunotoxicity, and can be divided in non-functional,

investigating the amount of specific cells or substances of immune system, and functional, studying the activity of cells in response to various stimuli and the experimental models of immune diseases. In this study the following methods were utilized: Lymphocyte proliferation, NK cell function, O<sub>2</sub> production by neutrophils, lymphocyte subpopulations and cytokine production in cultured peripheral mononuclear cells (IL5, INF  $\gamma$  and TNF $\alpha$ ). Cell proliferation and cytokine production will be discussed in this report.

*Cell proliferation*. 100 ml aliquots of PBMC suspension were placed in each well on 96-microtiter plate. Cells were then incubated for 78 hours at 37°C in humidified atmosphere with 5% CO<sub>2</sub> and in the following conditions:

- without PHA (control samples);
- with 20 mg/ml PHA (control samples);
- containing 10<sup>-4</sup> or 10<sup>-7</sup> M metal compounds without PHA;
- containing 10<sup>-4</sup> or 10<sup>-7</sup> M metal compounds with PHA.

Sodium arsenite (III), Sodium arsenate (V) and DMAs concentrations were 10<sup>-4</sup>, 10<sup>-5</sup>, 5X10<sup>-5</sup>, 10<sup>-6</sup>, 5X10<sup>-6</sup>, 10<sup>-7</sup>, 5X10<sup>-7</sup>, 10<sup>-8</sup>, 5X10<sup>-8</sup>, 10<sup>-9</sup> M;

Pt, Pd, Rh, Ti compounds were added from the 10<sup>-3</sup> and 10<sup>-7</sup> M stock solutions and As compounds were added from the 10<sup>-3</sup>, 10<sup>-4</sup>, 5X10<sup>-4</sup>, 10<sup>-5</sup>, 5X10<sup>-5</sup>, 10<sup>-6</sup>, 5X10<sup>-6</sup>, 10<sup>-7</sup>, 5X10<sup>-7</sup>, 10<sup>-8</sup> M stock solutions previously prepared.

BrdU cell proliferation assay was used (Oncogene, Darmstadt, Germany) for the evaluation of cell proliferation. During the last 24 hours of culture BrdU was added to wells of the microtiter plate. Cells were fixed and permeabilized, while the DNA was denaturated by a 30-min-treatment with fixative/denaturing solution at room temperature. Anti-BrdU monoclonal antibody was pipetted into the wells and incubated for one hour. Unbound antibodies were washed away and horseradish peroxidase-coniugate goat anti-mouse was added for 30' at room temperature. Chromogenic substrate solution tetramethylbenzidine was then added to each well and incubated in the dark at room temperature for 15 min. Stop solution was added to each well in the same order as the previously added substrate solution. Absorbance in each well was measured using a spectrophotometric plate reader at dual wavelengths of 450-540 nm. Color intensity was proportional to the amount of incorporated BrdU and to the degree of cell proliferation.

*Cultures for measurement of cytokines*. Human PBMC were purified from EDTA-treated whole blood by Ficoll-Hypaque density gradient centrifugation. Cultures were set up in 1 ml/well 24-wells costar plastic plates, using 0.8 ml of PBMC containing 1 million cells/well in complete medium, at 37°C, in humidified atmosphere with 5% CO<sub>2</sub> for 48 hours and the previously described conditions:

- without PHA (control samples)
- with 20 mg/ml PHA (control samples)
- containing 10<sup>-4</sup> or 10<sup>-7</sup> M metal compounds without PHA
- containing 10<sup>-4</sup> or 10<sup>-7</sup> M metal compounds with PHA

Supernatants of cultures were stored at -70 °C in aliquots until analysis. IFN- $\gamma$  and TNF- $\alpha$  levels in supernatants were determined by Quantikine ELISA kits (R&D Systems, Minneapolis, MN, USA) following the manufacturer's instructions.

#### RESULTS

*Titanium Compounds*. As shown in a previous study (20), Ti dioxide (biocompatible material) did not show

**Table I.** Possible interactions between metals and Immune system.

	Examples
<b>Essential Metals</b>	Copper, Iron, Manganese, Iodine, Zinc, Selenium, Trivalent Chromium, Arsenic
<b>Toxic Metals</b> <i>At high dose</i> <i>At low dose</i>	Zinc, Copper, Cobalt, Cadmium, Mercury Lead, Hexavalent Chromium
<b>Sensitizing Metals</b> <i>Allergic Contact dermatitis</i> <i>Asthma and rhinitis</i>	Nickel, Chromium, Cobalt, Platinum, Mercury, Tungsten, Vanadium Copper, Zinc, Palladium, Tin, Iodine, Lithium, Gold Nickel, Chromium, Cobalt, Platinum, Mercury, Vanadium

**Table II.** Metal compounds utilized in the study.

Titanium	TiO <sub>2</sub> (Ti dioxide), (C <sub>2</sub> H <sub>5</sub> ) <sub>2</sub> TiCl <sub>2</sub> (Ti ascorbate), TiO(C <sub>2</sub> O <sub>4</sub> ) <sub>2</sub> (Ti oxalate), Titanocene
Arsenic	Sodium Arsenite (III), Sodium Arsenate (V), potassium-esa-fluorum-arsenate (V), Sodium-esa-fluorum-arsenate (V), MMAs (V), DMAs (V), Tetra-phenil-arsonium chloride (III)
Platinum	halogenated compounds as (NH <sub>4</sub> ) <sub>2</sub> PtCl <sub>6</sub> and (NH <sub>4</sub> ) <sub>2</sub> PtCl <sub>4</sub> ; chlorinated compounds as PtCl <sub>2</sub> and PtCl <sub>4</sub> ; NH <sub>2</sub> PtI <sub>6</sub> ; CisPt
Palladium	halogenated compounds as (NH <sub>4</sub> ) <sub>2</sub> PdCl <sub>6</sub> and (NH <sub>4</sub> ) <sub>2</sub> PdCl <sub>4</sub> ; chlorinated compound as PdCl <sub>2</sub>
Rhodium	halogenated (NH <sub>4</sub> ) <sub>2</sub> RhCl <sub>6</sub> ; chlorinated RhCl <sub>3</sub>

**Table III.** Proliferation of PHA stimulated PBMC in presence of different concentration (10<sup>-4</sup> M and 10<sup>-7</sup> M) of As compounds (Mann Whitney U test: \*p=0.05; \*\*p<0,01; \*\*\*p<0.001).


Without As	1433 ± 158	
Sodium Arsenate	904 ± 181 *	1389 ± 240
Sodium Arsenite	379 ± 202 ***	1592 ± 146 *
MMAs	1477 ± 312	1351 ± 339
DMAs	1309 ± 377	1847 ± 343 **
Tetra-phenyl-arsonium-chloride	176 ± 10 ***	1454 ± 393
Sodium-esa-fluorum-arsenate	1312 ± 315	1412 ± 310
Potassium-esa-fluorum-arsenate	1398 ± 141	1371 ± 168
	10 <sup>-4</sup> M	10 <sup>-7</sup> M

any immunotoxicity, while titanocene (anticancer therapy) inhibited cytokine release and Ti oxalate and ascorbate have a marked immunotoxicity (21). In particular, the Titanium compound employed in industry (oxalate) and the one used in agriculture (ascorbate), significantly inhibited the cell proliferation in PHA stimulated cultures, when added at maximal concentration. On the contrary, they do not seem to have any effect in PHA unstimulated cultures. In fact, Ti ascorbate and oxalate stimulation index was significantly reduced from the control without metals (Fig. 1).


Ti oxalate and ascorbate and Titanocene (used as chemotherapeutic agent) reduced the release of TNF, when used at their maximal concentration. Ascorbate showed

a highly significant effect on TNF release at lower concentration also, whereas the activity of oxalate and titanocene was near the limits of statistical significance. The sensibility of INF was quite different: ascorbate did not show any activity, whereas titanocene and Ti oxalate at maximal concentration strongly inhibited INF release. The inhibitory activity of titanocene on cytokine release was so important, that in some experiments both INF and TNF were undetectable.


**Arsenic Compounds.** The studied immune functions, cell proliferation and cytokine release, showed that the toxicity of 10<sup>-4</sup> MAs compounds was in the following rank order: Sodium Arsenite (III) > Tetra-phenyl Arsonium Chloride (III) > DMAs (III) and Sodium Arsenate (V) >




**Fig. 1.** PBMC proliferation in cultures with Ti compounds: Stimulation Index (Mann Whitney U test: \*  $p < 0.001$ ).



**Fig. 2.** Percentage, related to the control, of IFN-g and TNF-a release from PHA stimulated PBMC in presence of  $10^{-4}$  M As compounds (Mann Whitney U test: \*  $p < 0.05$ ; \*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$ .)



**Fig. 3.** PHA stimulated PBMC proliferation without and with  $10^{-4}$  M chemical compounds (Mann Whitney U test: \*  $p < 0.05$ ; \*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$ ).



**Fig. 4.** Percentage (related to the control cultures) of PHA-stimulated TNF- $\alpha$  release from PBMC in presence of  $10^{-4}$  and  $10^{-7}$  M Pd, Pt and Rh compounds. Values are mean  $\pm$  S.E.M. ( $n = 9$ ). TNF- $\alpha$  release in the control cultures is  $1319 \pm 133$  pg/ml (Man Whitney U test. Significant difference in relation to the control cultures without metal compounds : \* $p < 0.05$ ; \*\*\* $p < 0.001$ ).

other compounds.

Sodium arsenite and tetraphenyl arsonium chloride are more immunotoxic than the other As compounds investigated (Fig. 2). Sodium arsenite was more efficient in inhibiting lymphocyte proliferation than tetraphenyl arsonium chloride, which was more active in inhibiting cytokine production by PBMC. Sodium arsenate inhibited PBMC proliferation also at  $10^{-5}$  M concentration but did not inhibit TNF- $\alpha$  production at  $10^{-4}$  M. Sodium-esa-fluorum-arsenate at  $10^{-4}$  M did not inhibit PBMC proliferation but slightly inhibited TNF- $\alpha$  release. On the whole, each inorganic or organic As compound exerts a specific immunotoxic effect.

Trivalent compounds showed the greatest toxicity on cell proliferation at maximal concentration of  $10^{-4}$  M. On the contrary,  $10^{-7}$  M of arsenite and DMAs induced a stimulation (Table III). Studying a dose-response curve it was observed that the stimulation was not obtained by concentrations below  $10^{-8}$  and  $10^{-9}$  M. Many authors have described this phenomenon of "hormesis" and all of them sustain that  $10^{-6}$  - $10^{-7}$  are the stimulatory concentrations (22-23). The increased proliferation of PBMC in presence of  $10^{-7}$  M trivalent As and DMAs could be explained by a stimulatory metabolic effect of As (6-7). Arsenic compounds at  $10^{-4}$  M (in particular sodium arsenite > tetraphenyl arsonium chloride) inhibited cytokine release by PBMC, whereas sodium arsenate and potassium- and sodium-esa-fluorum arsenate showed lower activity.

Recent investigations, showing that genotoxic and cytotoxic effects of As may be produced by trivalent intermediary organic compounds (produced during metabolism of MMAs and DMAs), can explain the immunotoxicity of pentavalent As compounds (24-25). A careful investigation on the uptake and the metabolism of

metallic compounds is necessary in order to fully explain the activity of substances produced during the metabolism of metal compounds.

*Platinum group elements – PGE- (Platinum, Palladium and Rhodium).* Palladium compounds at  $10^{-4}$  M and  $(\text{NH}_4)_2\text{PtCl}_6$  inhibited spontaneous and PHA-stimulated PBMC proliferation, as well as TNF- $\alpha$ , INF- $\gamma$  and IL-5 release by PHA-stimulated PBMC. Other Pt or Rh salts did not have the same effects (Fig. 3). The effects of the ammonium hexa- and tetra-chloro-platinate and palladate compounds (IV and II oxidation state, respectively) were similar when PBMC proliferation and INF- $\gamma$  or IL-5 release were considered; however, TNF- $\alpha$  release was reduced by all the chlorinated Pd compounds but only by  $10^{-4}$  M  $(\text{NH}_4)_2[\text{PtCl}_6]$  among the six tested Pt derivatives.

Pt(IV)ACC and Pd(IV)ACC were generally more active than Pt(II)ACC and Pd(II)ACC. Moreover, these Pt and Pd salts were more active in inhibiting PBMC proliferation and TNF- $\alpha$  release than ammonium hexachlororhodate (RhIV). Ammonium hexachlororhodate was in turn less effective than Rh(III) trichloride in reducing both PBMC proliferation and cytokine release.

PtCl<sub>2</sub> did not exert any effect on PBMC, while the only immune effect of PtCl<sub>4</sub> was to enhance the release of IL-5, known marker of the Th-2 immune response (26).

Na<sub>2</sub>PtI<sub>6</sub> (a tetravalent iodide Pt compound) inhibited IFN- $\gamma$  and IL-5 release. This demonstrates that iodide Pt compounds, which may be synthesized in the environment (11), can be immunotoxic.

Our study demonstrates that Pd (II, IV) and Rh (III, IV) inhibit TNF- $\alpha$  release and that Pt exerts a similar effect only as ammonium hexachloroplatinate (IV) salt (Fig. 4). TNF- $\alpha$  is involved in the pathogenesis of septic

shock and rheumatoid arthritis, and represents a cytotoxic factor for many tumour cells also by inducing apoptosis (24). In this regard, the substantial lack of effects of Pt salts on TNF- $\alpha$  may account for the more elevated cancer-related genotoxicity of Pt compounds among those of the PGE (28).

The results obtained may contribute to explain the immune effects of PGE compounds by providing evidences on the importance of speciation in metal toxicity (29). The same results are in agreement with the present ones showing no immune effects of Pt dichloride. It has been reported that atopy is not an important risk factor for the development of sensitization and allergy to Pt salts in exposed workers (30). However, prick test reactions to Pt, Pd and Rh were found in about 15 % of workers in an Italian Pt refinery, who also showed positive reactions to common allergens (a marker of atopy) (31). Moreover, a HLA-DR3 phenotype was found more common among Pt-sensitized symptomatic workers exposed to Pt salts, with a significant association varying with the intensity of the exposure to the sensitizing agent (32).

A study performed by three centers of Dermatology in Germany reported a surprisingly high number of positive patch test to Pd compounds (33). It is not clear whether sensitization to Pd occurs because of exposure to the metal or because of cross-reactions with other elements. Ni-reactive T lymphocyte clones from donors reacted with Pd and Cu salts (34). This cross-reactivity could have been favored by the location of Ni, Pd and Cu in the periodic table and by their similarity in binding to histidine residues of surface peptides of major histocompatibility complex class II molecules. Positive patch test reactions to both Pd and Ni were found in patients with dental restorations (35-36) and cross-reactions among Pd, Pt, Rh and iridium were shown on prick testing in workers exposed to these metals (31). Cross-reactivity between Pd and Ni sulphate was also found using patch test techniques in laboratory animals, guinea pigs (37-38).

#### DISCUSSION

We can conclude that these studies show that metal immunotoxicity depends on speciation. These studies indicate that chemical speciation of Titanium, Arsenic and PGE salts is important for determining their immunotoxic effects. Moreover, Pd compounds have to be taken in greater consideration also in order to prevent a possible risk of cross-reactions in subjects sensitized to other metals. In general speciation provides additional and often essential information in evaluating metal toxicity. However, there are many difficulties to apply speciation in investigating toxicokinetic aspects to many metals mainly due to the lack of information about the existence and significance of species and to the lack of analytical methods for measuring species in biological samples.

#### REFERENCES

1. **Boscolo P., M. Di Gioacchino, P. Bavazzano, M. White and E. Sabbioni.** 1997. Effects of chromium on lymphocyte subsets and immunoglobulins from normal population and exposed workers. *Life Sci.* 60:1319.
2. **Tanigawa T., S. Araki, T. Araki and N. Mininato.** 1991. A decrease in Leu-11a negative lymphocytes in relation to natural killer cell activity in chromate workers *Br. J. Ind. Med.* 48:211.
3. **Di Gioacchino M., E. Sabbioni, L. Di Giampaolo, C. Schiavone, M.B. Di Sciascio, M. Reale, V. Nicola, N. Qiao, R. Paganelli, P. Conti and P. Boscolo.** 2002. *In vitro* effects of vanadate on human immune functions. *Ann. Clin. Lab. Sci.* 32:148.
4. **Boscolo P., L. Di Giampaolo, N. Qiao, M. Reale, M.L. Castellani, I. Lucci, P. Travaglini, M. Kouri, N. Verna, A.R. Volpe, M. Carmignani, R. Paganelli, and M. Di Gioacchino.** 2005. Inhibitory effects of cadmium on peripheral blood mononuclear cell proliferation and cytokine release are reversed by zinc and selenium salts. *Ann. Clin. Lab. Sci.* 35:115.
5. **Di Giampaolo L., M. Di Gioacchino, N. Qiao, P. Travaglini, A. D'Intino, M. Kouri, J. Ponti, M.L. Castellani, M. Reale, E. Gabriele and P. Boscolo.** 2004. »*In vitro*» effects of different arsenic compounds on PBMC (preliminary study). *G. Ital. Med. Lav. Ergon* 26:183.
6. **Anke M., U. Krause and B. Groppel.** 1987. The effect of arsenic deficiency on growth, reproduction, life expectancy and disease symptoms in animals. In *Trace Substances in Environmental Health, vol 21*, B.B. Hemphill, ed. Columbia University of Missouri, p. 533.
7. **Uthus E.O. and C.D. Seaborn.** 1996. Deliberations and evaluations of the approaches, endpoints and paradigms for dietary recommendations of the other trace elements. *J. Nutr.* 126:245.
8. **Cantor K.P.** 2001. Invited commentary: arsenic and cancer and the urinary tract. *Am. J. Epidemiol.* 153:419.
9. **Boscolo P., M. Di Gioacchino, R. Paganelli and E. Sabbioni.** 2002. Studies on the immune effects of metals. *Int. J. Immunopathol. Pharmacol.* 15:14.
10. **Di Gioacchino M., L. Di Giampaolo, N. Verna, M. Reale, M.B. Di Sciascio, A.R. Volpe, M. Carmignani, J. Ponti, R. Paganelli, E. Sabbioni and P. Boscolo.** 2004. *In vitro* effects of platinum compounds on lymphocyte proliferation and cytokine release. *Ann. Clin. Lab. Sci.* 34:195.
11. **Merget R. and G. Rosner.** 2001. Evaluation of the health risk of platinum group metals emitted from automotive catalytic converters. *Sci. Total Environ.* 270:165.
12. **Schuppe H.C., J. Kulig, U. Kuhn, U. Lempertz, P. Kind,**

- J. Knop and D. Becker.** 1997. Immunostimulatory effects of platinum compounds: correlation between sensitizing properties in vivo and modulation of receptor-mediated endocytosis in vitro. *Int. Arch. Allergy Immunol.* 112:125.
13. **Schuppe H.C., A.C. Ronnan, S. von Schmiedeberg, T. Ruzicka, E. Gleichmann and P. Grien.** 1998. Immunomodulation by heavy metal compounds. *Clin. Dermatol.* 16:149.
  14. **Linnet J.P. and E.G. Hughes.** 1999. 20 years of medical surveillance on exposure to allergenic and non-allergenic platinum compounds: the importance of chemical speciation. *Occup. Environ. Med.* 56:191.
  15. **Salsano F., C. Francia, I. Roumpedaki, M. Proietti, S. Pisarri, N. Verna, E. Gabriele, G. Di Gioacchino and M. Di Gioacchino.** 2004. Immune effects of nickel. *Int. J. Immunopathol. Pharmacol.* 17:63.
  16. **M. Di Gioacchino, L. Di Giampaolo, L. Balatsinou, F. Di Claudio, N. Verna.** 2005. Nickel allergy: the guilty cells. *Int. J. Immunopathol. Pharmacol.* (in press).
  17. **Paganelli R., B. Buttari, E. Camera, M.L. Dell'Anna, A. Mastrofrancesco, L. Di Giampaolo, M. Reale, C. Schiavone, N. Verna, M. Di Gioacchino, E. Sabbioni, P. Boscolo and M. Picardo.** 2003. *In vitro* effects of nickel sulphate on immune functions of normal and allergic subjects: a regulatory role of zinc. *J. Trace Elem. Med. Biol.* 17:11.
  18. **Di Gioacchino M., P. Boscolo, E. Cavallucci, N. Verna, F. Di Stefano, M. Di Sciascio, S. Masci, M. Andreassi, E. Sabbioni, D. Angelucci and P. Conti.** 2000. Lymphocyte subset changes in blood and gastrointestinal mucosa after oral nickel challenge, in nickel sensitized women. *Contact Dermatitis* 43:206.
  19. **Meliddo G., L. Di Giampaolo, E. Toto, M. Russi, P. Travaglini, M. Appignani, M. Di Gioacchino and P. Boscolo.** 2003. *In vitro* effects of hexavalent and tetravalent tellurium on immune functions. *G. Ital. Med. Lav. Ergon* 25:71.
  20. **Russi M., G. Meliddo, E. Toto, L. Di Giampaolo, P. Travaglini, M. Appignani, M. Di Gioacchino and P. Boscolo.** 2003. Comparative *in vitro* study of various titanium compounds on the immune system. *G. Ital. Med. Lav. Ergon* 25:69.
  21. **Di Giampaolo L., M. Di Gioacchino, R. Sabbioni, M. Reale, E. Toto, P. Conti, R. Paganelli and P. Boscolo.** 2004. *In vitro* comparative immune effects of different titanium compounds. *Int. J. Immunopathol. Pharmacol.* 17:115.
  22. **Vega L., P. Ostrosky-Wegman, T.I. Fortoul, C. Diaz, V. Madrid and R. Saavedra.** 2001. Sodium arsenite reduces proliferation of human activated T-cells by inhibition of the secretion in normal human epidermal keratinocytes. *Toxicol. Appl. Pharmacol.* 172:225.
  23. **Calabrese E.J. and L.A. Baldwin.** 1999. U-shaped dose responses in biology, toxicology and public health. *Annual Review of Public Health* 22:15.
  24. **Thomas D.J., M. Styblo and S. Lin.** 2001. The cellular metabolism and systemic toxicity of arsenic. *Toxicol. Appl. Pharmacol.* 176:127.
  25. **Mass M.J., A. Tennant, B.C. Roop, W.R. Cullen, M. Styblo, D.J. Thomas and A.D. Kligerman.** 2001. Methylated trivalent arsenic species are genotoxic. *Chem. Res. Toxicol.* 14:355.
  26. **Romagnani S.** 2000. T-cell subsets (Th1 versus Th2). *Ann. Allergy Asthma Immunol.* 85:9.
  27. **Fujiki H., M. Suganuma, S. Okabe, M. Kurusu, K. Imai and K. Nakachi.** 2002. Involvement of TNF- $\alpha$  in human cancer development, prevention and palliative care. *Mech. Ageing Dev.* 123:1655.
  28. **Migliore L., G. Frenzilli, C. Nesti, S. Fortaner and E. Sabbioni.** 2002. Cytogenic and oxidative damage induced in human lymphocytes by platinum, rhodium and palladium compounds. *Mutagenesis* 17:411.
  29. **Boscolo P., L. Di Giampaolo, M. Reale, M.L. Castellani, A. Ritavolpe, M. Carmignani, J. Ponti, R. Paganelli, E. Sabbioni, P. Conti and M. Di Gioacchino.** 2004. Different effects of platinum, palladium, and rhodium salts on lymphocyte proliferation and cytokine release. *Ann. Clin. Lab. Sci.* 34:299.
  30. **Niezborala M. and R. Garnier.** 1996. Allergy to complex platinum salts: a historical prospective cohort study. *Occup. Environ. Med.* 53:252.
  31. **Santucci B., C. Valenzano, M. de Rocco and A. Cristaudo.** 2000. Platinum in the environment: frequency of reactions to platinum-group elements in patients with dermatitis and urticaria. *Contact Dermatitis* 43:333.
  32. **Newman Taylor A.J., P. Cullinan, P.A. Lymphany, J.M. Harris, R.J. Dowdeswell and R.M. du Bois.** 1999. Interaction of HLA phenotype and exposure intensity in sensitisation to complex platinum salts. *Am. J. Respir. Crit. Care Med.* 160:435.
  33. **Brasch J. and J. Geier.** 1997. Patch test results in schoolchildren. *Contact Dermatitis* 37:286.
  34. **Pistor F.H., M.L. Kapsenberg, M.M. Meinardi, M.E. von Blomberg and R.J. Sheper.** 1995. Cross-reactivity of human nickel-reactive T-lymphocyte clones with copper and palladium. *J. Invest. Dermatol.* 105:92.
  35. **Shaffran R.M., F. J. Storrs and P. Schalock.** 1999. Prevalence of gold sensitivity in asymptomatic individuals with gold dental restorations. *Am. J. Dermatol.* 10:201.
  36. **Kielhorn J., C. Melber, D. Keller and I. Mangelsdorf.**

2002. Palladium – a review of exposure and effects to human health. *Ind. J. Hyg. Environ. Health* 205:417.
37. **Wahlberg J.E. and C. Liden.** 2000. Cross-reactivity patterns of cobalt and nickel with repeated open applications (ROATs) to the skin of guinea pigs. *Am. J. Contact Dermatitis* 11:42.
38. **Wahlberg J.E. and C. Liden.** 1999. Cross-reactivity patterns of palladium and nickel studied by repeated open applications (ROATs) to the skin of guinea pigs. *Contact Dermatitis* 41:145.