

Oral presentation

Open Access

## Controlling the Virus Output Via Urokinase Receptor and Integrin Signaling

Guido Poli\*‡

Address: AIDS Immunopathogenesis Unit & Vita-Salute San Raffaele University and Scientific Institute, Via Olgettina n. 58, 20132 Milan, Italy

Email: Guido Poli\* - poli.guido@hsr.it

\* Corresponding author ‡Presenting author

from 2005 International Meeting of The Institute of Human Virology  
Baltimore, USA, 29 August – 2 September 2005

Published: 8 December 2005

*Retrovirology* 2005, **2**(Suppl 1):S76 doi:10.1186/1742-4690-2-S1-S76

We have described that either urokinase-type plasminogen activator (uPA) inhibits HIV expression in monocytic U937 and U1 cell lines (*M. Alfano et al., PNAS, 2002, 99:8862-67*). We have observed that uPA inhibited HIV expression exclusively when both uPAR and CD18/CD11b (Mac-1) were co-expressed at the cell surface. A second interactor of uPAR, FPRL1, was abundantly expressed on the surface of both unstimulated and stimulated U1 cells; however, peptide antagonists of FPRL1 did not interfere with HIV expression from U1 cells. Incubation of U1 cells with Trojan peptides expressing RhoA domains reversed the anti-HIV activity of uPA. In addition to cell line infection, uPA inhibited *in vitro* infection of primary monocyte-derived macrophages and virus replication from monocytes of infected individuals cultivated *ex-vivo*. Thus, RhoA-dependent cytoskeleton rearrangement and intracellular vesicles formation may be related to virion budding and entrapment in intracytoplasmic vacuoles. This is the first report linking integrin activation to a negative control of HIV replication, at least in monocyte/macrophages.