

**850** Prevalence of programmed death ligand-1 (PD-L1) by demographic, disease and sample characteristics in unresectable, stage III NSCLC (PACIFIC)

D. Planchard<sup>1</sup>, M.C. Garassino<sup>2</sup>, L. Paz-Ares<sup>3</sup>, C. Faivre-Finn<sup>4</sup>, A. Spira<sup>5</sup>, Y. Gu<sup>6</sup>, C. Wadsworth<sup>7</sup>, J. Whiteley<sup>8</sup>, M. Scott<sup>8</sup>, A-M. Boothman<sup>8</sup>, M. Ratcliffe<sup>8</sup>, J. Walker<sup>8</sup>, P.A. Dennis<sup>9</sup>, S.J. Antonia<sup>9</sup>

<sup>1</sup>Department of Medical Oncology, Thoracic Group, Institut Gustave Roussy, Villejuif, France, <sup>2</sup>Thoracic Unit, Fondazione IRCCS - Istituto Nazionale dei Tumori, Milan, Italy, <sup>3</sup>CiberOnc, Universidad Complutense and CNIO, University Hospital 12 de Octubre, Madrid, Spain, <sup>4</sup>Clinical Oncology, The University of Manchester and The Christie NHS Foundation Trust, Manchester, UK, <sup>5</sup>Virginia Cancer Specialists Research Institute, Fairfax, VA, USA, <sup>6</sup>AstraZeneca, Gaithersburg, MD, USA, <sup>7</sup>AstraZeneca, Alderley Park, UK, <sup>8</sup>Precision Medicine and Genomics, IMED Biotech Unit, AstraZeneca, Cambridge, UK, <sup>9</sup>Thoracic Oncology, H. Lee Moffitt Cancer Center and Research Institute, Tampa, FL, USA

**Background:** PACIFIC (NCT02125461) was a randomised, placebo-controlled, phase 3 trial evaluating the immune checkpoint inhibitor durvalumab in patients (pts) with unresectable, Stage III non-small cell lung cancer (NSCLC) who did not have disease progression after concurrent chemoradiotherapy (cCRT). Both primary endpoints of progression-free survival and overall survival were met and significantly improved with durvalumab, with similar safety, versus placebo (Antonia et al, NEJM 2017; 2018). We report exploratory analyses of the prevalence of tumour PD-L1 expression by baseline pt, disease and sample characteristics and by response to prior treatment for pts in PACIFIC.

**Methods:** If available (provision of formalin-fixed paraffin-embedded tumour resection or biopsy samples was optional), archived pre-cCRT tumour tissue was tested retrospectively for PD-L1 tumour cell (TC) expression using the VENTANA PD-L1 (SP263) immunohistochemistry assay and scored at validated pre-specified ( $\geq 25\%$ ) and post-hoc ( $\geq 1\%$ ) cutoffs. Overall PD-L1 prevalence (regardless of treatment arm) was summarised by pt subgroups defined by various characteristics, and assessed using a Pearson's chi-squared test for between-group differences.

**Results:** Of 713 randomized pts, 451 (63.2%) were evaluable for PD-L1 status. Among PD-L1-evaluable pts, 67.2% (303/451) had TC  $\geq 1\%$  and 35.3% (159/451) had TC  $\geq 25\%$  (similar to previous reports in metastatic NSCLC). PD-L1 prevalence by various characteristics at the TC  $\geq 1\%$  cut-off are reported in the table.

**Conclusions:** There were no important differences noted in PD-L1 prevalence between relevant subgroups at the TC  $\geq 1\%$  or TC  $\geq 25\%$  cut-offs (latter data to be presented). PD-L1 status was unaffected by sample type or age or biopsy location, suggesting expression is stable from pre-cCRT diagnostic biopsies, and supports the use of either primary tumour or lymph node biopsies for PD-L1 testing.

**Clinical trial identification:** NCT02125461.

**Editorial acknowledgement:** Medical writing support, which was in accordance with Good Publication Practice (GPP3) guidelines, was provided by Andrew Gannon of Cirrus Communications, an Ashfield company, and was funded by AstraZeneca.

**Legal entity responsible for the study:** AstraZeneca.

**Funding:** AstraZeneca.

**Disclosure:** D. Planchard: Personal fees: AstraZeneca, Boehringer Ingelheim, BMS, MSD, Pfizer, Novartis, Roche, Celgene, outside the submitted work. M.C. Garassino: Personal fees: AstraZeneca, Roche, BMS, MSD outside the conduct of this study. L. Paz-Ares: Advisory board fees: BMS, Lilly, MSD, AstraZeneca, Roche, Pfizer, Novartis, Incyte, Merck, Boehringer Ingelheim. C. Faivre-Finn: Research funding: AstraZeneca, MSD. A. Spira: Advisory fees, institutional research support: AstraZeneca. Y. Gu, J. Whiteley, M. Scott, J. Walker: Employment, stock: AstraZeneca. C. Wadsworth, P.A. Dennis: Employment, stock: AstraZeneca, outside the conduct of the study. A-M. Boothman: Employment, stock options: AstraZeneca, outside the conduct of the study. M. Ratcliffe: Consultant fees: AstraZeneca, outside the conduct of the study. All other authors have declared no conflicts of interest.

**Table: 850**

Patient/disease/sample characteristic	Subgroup	Prevalence of PD-L1 TC $\geq$ 1%, % (n/N)	P value
Race	White	67.5 (206/305)	0.5882
	Black or African American	81.8 (9/11)	
	Asian	66.4 (85/128)	
	Other	50.0 (3/6)	
Region	Asia	64.3 (74/115)	0.2642
	Europe	64.9 (124/191)	
	North America and South America	72.4 (105/145)	
Patient age (years)	<65	66.5 (167/251)	0.7418
	$\geq$ 65	68.0 (136/200)	
Smoking status	Smoker	66.8 (276/413)	0.5956
	Non-smoker	71.1 (27/38)	
Sex	Male	65.3 (209/320)	0.1858
	Female	71.8 (94/131)	
Histology	Squamous	64.9 (150/231)	0.2973
	Non-squamous	69.5 (153/220)	
Disease stage	IIIA	69.2 (166/240)	0.5029
	IIIB	66.2 (131/198)	
ECOG/WHO performance status	Normal activity (0)	65.8 (150/228)	0.5236
	Restricted activity (1)	68.6 (153/223)	
EGFR status	Positive	67.7 (21/31)	0.1176
	Negative	68.6 (264/385)	
	Unknown	51.4 (18/35)	
Best response to prior therapy	Complete response	41.7 (5/12)	0.1611
	Partial response	67.7 (151/223)	
	Stable disease	68.1 (143/210)	
Biopsy sample location	Lung	67.0 (209/312)	0.8045
	Lymph node(s)	65.6 (63/96)	
Specimen type	Formalin-fixed paraffin-embedded block	69.2 (126/182)	0.4461
	Unstained slides	65.8 (175/266)	
Collection method	Biopsy	68.2 (279/409)	0.5879
	Resection	63.6 (21/33)	
Age of biopsy sample	$\leq$ 90 days	73.3 (88/120)	0.0902