



Sep 15th, 11:20 AM

## Methane Dynamics of Kentucky Bottomland Hardwood Forested Systems

Marissa Miles  
*Murray State University*

Jessica B. Moon  
*Murray State University, 1jmoon8@murraystate.edu*

Skylar Ross  
*Murray State University*

Jarred Asselta  
*Murray State University*

Niklas Klauss  
*Murray State University*

*See next page for additional authors*

**Right click to open a feedback form in a new tab to let us know how this document benefits you.**  
Follow this and additional works at: [https://uknowledge.uky.edu/kwrri\\_proceedings](https://uknowledge.uky.edu/kwrri_proceedings)



Part of the [Engineering Commons](#), [Life Sciences Commons](#), and the [Physical Sciences and Mathematics Commons](#)

---

Miles, Marissa; Moon, Jessica B.; Ross, Skylar; Asselta, Jarred; Klauss, Niklas; Carey, Rosie; Khatiwada, Kabi Raj; Masri, Bassil El; Stinchcomb, Gary; and Runkle, Benjamin R. K., "Methane Dynamics of Kentucky Bottomland Hardwood Forested Systems" (2023). *Kentucky Water Resources Annual Symposium*. 4.  
[https://uknowledge.uky.edu/kwrri\\_proceedings/2023/event/4](https://uknowledge.uky.edu/kwrri_proceedings/2023/event/4)

This Presentation is brought to you for free and open access by the Kentucky Water Research Institute at UKnowledge. It has been accepted for inclusion in Kentucky Water Resources Annual Symposium by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

---

## Presenter Information

Marissa Miles, Jessica B. Moon, Skylar Ross, Jarred Asselta, Niklas Klauss, Rosie Carey, Kabi Raj Khatiwada, Bassil El Masri, Gary Stinchcomb, and Benjamin R. K. Runkle

## Methane Dynamics of Kentucky Bottomland Hardwood Forested Systems\*

Marissa Miles<sup>1,2</sup>, Jessica B. Moon<sup>1,2</sup>, Skylar Ross<sup>2,3</sup>, Jarred Asselta<sup>3</sup>, Niklas Klaus<sup>1</sup>, Rosie Carey<sup>1</sup>, Kabi Raj Khatiwada<sup>4</sup>, Bassil El Masri<sup>2,3</sup>, Gary Stinchcomb<sup>5</sup>, Benjamin R. K. Runkle<sup>4</sup>

<sup>1</sup>Department of Biological Sciences, Murray State University

<sup>2</sup>The Watershed Studies Institute, Murray State University

<sup>3</sup>Department of Earth and Environmental Sciences, Murray State University

<sup>4</sup>Department of Biological and Agricultural Engineering, University of Arkansas

<sup>5</sup>Department of Earth Sciences, University of Memphis

[1,2mmiles6@murraystate.edu](mailto:1,2mmiles6@murraystate.edu)

Bottomland hardwood forests can hold large amounts of carbon in their tree biomass and their deep soil carbon pools. However, they can also produce a significant amount of methane (CH<sub>4</sub>) in their anaerobic soils. While some of this CH<sub>4</sub> is oxidized in microsites and overlying waters, tree stems and exposed root systems can be significant conduits of CH<sub>4</sub> to the atmosphere, bypassing oxidation processes. We aim to identify differences in CH<sub>4</sub> source-sink dynamics among tree species, across heights on stems, and across stem hydrogeomorphic settings within bottomland hardwood wetlands. We predict that the greatest stem methane emissions will be found in hydrogeomorphic settings with the highest and most consistent of the following: inundation, organic matter content, and temperature. Additionally, we expect hydrogeomorphic setting to be a greater factor in explaining stem CH<sub>4</sub> emissions than species-level differences. CH<sub>4</sub> fluxes are being collected from stems of five tree species monthly in three protected areas in western Kentucky (i.e., Clarks River National Wildlife Refuge, Murphy's Pond Nature Preserve, Kentucky Lake's Hancock Biological Station). Monthly samplings include floodplain measurements at five heights (i.e., 20, 40, 60, 120, and 180 cm) on *Acer rubrum* and *Liquidambar styraciflua*, terrace measurements at five heights on *Quercus stellata* and *Quercus pagoda*, and measurements at two heights (i.e., 40 cm and 120 cm) at three hydrogeomorphic positions (i.e., slough, lake edge, and pond edge) on *Taxodium distichum*. Initial sampling was only completed for the floodplain and slough, during a moderate-to-severe drought (October 2022). During these dry conditions, floodplain soils acted as CH<sub>4</sub> sinks, while stems acted as CH<sub>4</sub> sources. We found no significant differences in CH<sub>4</sub> emissions among species, and contrary to existing literature, we did not observe significant differences in CH<sub>4</sub> emissions across stem heights. As we continue sampling, we expect stems to continue being CH<sub>4</sub> emitters, with greater emissions from stems during the hottest, wettest seasons. Under these conditions we also expect to see significant differences in emissions across stem heights as methanogenesis in soils increases. Stem measurements will be combined with estimates of the tree surface area to compare the contribution of stems to soils in sink-source dynamics of these systems. Our study will be essential to develop and validate tree stem CH<sub>4</sub> schemes to improve land surface models for CH<sub>4</sub> flux predictions.

\*This presentation is based upon work supported by the U.S. Geological Survey under Grant/Cooperative Agreement No. G21AP10631-01, WRRR 104B Annual Grant Program.