



Durham E-Theses

A Study on Adaptive Strategies of Wild and Cultivated Linum Populations Across Western Europe and Their Implications for Linum Trait Development and Ecology.

HABEAHAN, HORASMAN, FEBRICO

How to cite:

HABEAHAN, HORASMAN, FEBRICO (2023) *A Study on Adaptive Strategies of Wild and Cultivated Linum Populations Across Western Europe and Their Implications for Linum Trait Development and Ecology.*, Durham theses, Durham University. Available at Durham E-Theses Online:
<http://etheses.dur.ac.uk/15197/>

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in Durham E-Theses
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full Durham E-Theses policy](#) for further details.

Academic Support Office, Durham University, University Office, Old Elvet, Durham DH1 3HP
e-mail: e-theses.admin@dur.ac.uk Tel: +44 0191 334 6107
<http://etheses.dur.ac.uk>

1 A Study on Adaptive Strategies of Wild and Cultivated *Linum*
2 Populations Across Western Europe and Their Implications for *Linum*
3 Trait Development and Ecology.

4
5 *Collections of studies on local adaptation, and effects of local climates on genotype and phenotype of*
6 *wild and cultivated Linum populations found across Western Europe with implications for ecology*
7 *and development agricultural traits.*

8
9 by


10
11 Horasman Febrico Habeahan (Rico)

12
13
14
15
16 Department of Biosciences

17 Durham University

18
19
20
21
22
23
24
25
26 Supervised by: Dr. A.C. Brennan, Prof. A.R. Hoelzel, Dr. C. Kidner

27 This thesis is submitted in candidature of the degree 'Doctor of Philosophy' (Ph.D) at the
28 Department of Biosciences, Durham University (2018-2022).



Royal
Botanic Garden
Edinburgh



UK Research
and Innovation

| | | |
|----|---|----|
| 30 | Table of Contents: | |
| 31 | DECLARATION | 1 |
| 32 | DEDICATION | 2 |
| 33 | ACKNOWLEDGEMENTS | 3 |
| 34 | LIST OF ABBREVIATIONS | 4 |
| 35 | ABSTRACT..... | 5 |
| 36 | CHAPTER 1: INTRODUCTION | 7 |
| 37 | 1.2 INTRODUCTION TO STUDY SPECIES | 10 |
| 38 | 1.2.1 <i>Linum usitatissimum</i> : Cultivated Flax | 11 |
| 39 | 1.2.2 <i>Linum bienne</i> : Wild Flax | 13 |
| 40 | 1.2.3 Production Values of <i>Linum</i> Biproducts:..... | 14 |
| 41 | 1.3 LITERATURE REVIEW OF <i>LINUM</i> GENETIC DIVERSITY | 16 |
| 42 | 1.3.1 Methods for studying population genomics in <i>Linum</i> | 17 |
| 43 | 1.3.2 Self-Incompatibility in <i>Linum</i> | 19 |
| 44 | 1.3.3 Population Genetics Analysis..... | 20 |
| 45 | 1.3.4 Genetic Expression in Flowering Time:..... | 22 |
| 46 | 1.3.5 Observation of Pollen Viabilities..... | 23 |
| 47 | CHAPTER 2: FLOWERING TIME IN <i>LINUM</i> DUE TO VERNALIZATION AND EXPRESSION OF FIVE | |
| 48 | FLOWERING TIME GENES..... | 25 |
| 49 | 2.2 STUDY AND METHODS..... | 29 |
| 50 | 2.2.1 Samples and Experiment | 29 |
| 51 | 2.2.2 Relative Gene Expression: RNA Extractions..... | 34 |
| 52 | 2.2.3 Relative Gene Expression: cDNA synthesis:..... | 34 |
| 53 | 2.2.4 Expression of Genes of interest and House Keeping genes:..... | 35 |
| 54 | 2.2.5 Primer Testing..... | 36 |
| 55 | 2.2.6 Quantitative Real-Time Procedure | 37 |
| 56 | 2.2.7 Primer Efficiency | 38 |
| 57 | 2.2.8 Real time qPCR of Vernalised and Non-vernalised Samples..... | 40 |

| | | |
|----|--|----|
| 58 | 2.2.9 Clean-ups and quality controls: | 40 |
| 59 | 2.2.10 Relative Gene Expression: Quantification..... | 40 |
| 60 | 2.3 STATISTICAL ANALYSES FOR RELATIVE GENE EXPRESSION | 42 |
| 61 | 2.4 RESULTS: FLOWERING TIME BETWEEN EXPERIMENTS FROM DIFFERENT YEARS | 44 |
| 62 | 2.4.1 Results: Vernalization and flowering time | 45 |
| 63 | 2.4.2 Results: Primer Efficiency Values | 46 |
| 64 | 2.4.3 Results: Comparison of RGE using one and multiple HKGs. | 46 |
| 65 | 2.4.4 Results: Relative Gene Expression in Response to Vernalization | 47 |
| 66 | 2.4.5 <i>LuGI1.1 (GIGANTEA)</i> Expression Response | 50 |
| 67 | 2.4.6 <i>LuCO1</i> and <i>LuCO2 (CONSTANS)</i> Expression Response | 52 |
| 68 | 2.4.7 <i>LuFT1</i> and <i>LuFT2 (FLOWERING LOCUS T)</i> Expression Response | 53 |
| 69 | 2.4.8 Gene Expression and Flowering Time | 55 |
| 70 | 2.4.9 Relative Gene Expression and Latitude | 57 |
| 71 | 2.4.10 Results: Relative Gene Expression and Climate | 58 |
| 72 | 2.5 DISCUSSION AND CONCLUSION | 59 |
| 73 | CHAPTER 3: THE QUANTIFICATION OF POLLEN AND POLLEN TUBES TO REVEAL POLLEN | |
| 74 | GERMINATION VIABILITY IN <i>LINUM</i> UNDER DIFFERENT TEMPERATURE TREATMENTS. | 65 |
| 75 | 3.1. Pollen Structure | 66 |
| 76 | 3.2 STUDY MODELS AND METHODS | 69 |
| 77 | 3.2.1 Pollen Viability <i>In Vitro</i> | 70 |
| 78 | 3.2.2 Pollen Treatments..... | 71 |
| 79 | 3.2.3 Pollen and Pollen Tube Observation..... | 72 |
| 80 | 3.2.4 Pollen Counts | 73 |
| 81 | 3.2.5 Pollen Tube Counts | 75 |
| 82 | 3.3 ANALYSIS | 77 |
| 83 | 3.3.1 Analysis: Pollen and Pollen Tube Counts | 77 |
| 84 | 3.3.2 Analysis: Proportion of Pollen Tubes | 77 |
| 85 | 3.3.3 Analysis: Population of Origin Variables | 78 |

| | | |
|-----|--|-----|
| 86 | 3.4 RESULTS..... | 80 |
| 87 | 3.4.1 Results: <i>In vitro</i> pollen germination..... | 80 |
| 88 | 3.4.2 Results: Test for Normality | 80 |
| 89 | 3.4.3 Results: Pollen and Pollen Tubes Count..... | 81 |
| 90 | 3.4.4 Results: Pollen Tube Reaching the Ovary | 84 |
| 91 | 3.4.5 Results: Pollen Tube Proportion Over Pollen Count for Different <i>Linum</i> Species | 85 |
| 92 | 3.4.6 Results: Propotion of Pollen Tubes Reaching Ovary for Different <i>Linum</i> Species | 87 |
| 93 | 3.4.7 Results: Pollen and Latitude of Origin..... | 89 |
| 94 | 3.4.8 Results: Pollen and Local Climate | 93 |
| 95 | 3.5 DISCUSSION AND CONCLUSSION | 96 |
| 96 | CHAPTER 4: POPULATION STRUCTURE AND DIVERGENCE WITHIN LINUM SAMPLES ORIGINATING | |
| 97 | ACROSS WESTERN EUROPE LATITUDE RANGE..... | 102 |
| 98 | 4.2 STUDY AND METHODS..... | 106 |
| 99 | 4.2.1 Modified RAD Seq Protocol | 106 |
| 100 | 4.2.2 DNA Extraction Procedures..... | 106 |
| 101 | 4.2.3 The Modified CTAB protocol..... | 107 |
| 102 | 4.2.4 DNA Quality and Quantity..... | 108 |
| 103 | 4.2.5 Double-Digest RAD Sequencing | 108 |
| 104 | 4.2.5i RAD – Digestion:..... | 108 |
| 105 | 4.2.5ii RAD – Anneal adapter and Ligation | 109 |
| 106 | 4.2.5iii RAD – Pooling and Cleaning Ligated products | 109 |
| 107 | 4.2.5iv RAD – PCR Amplification to Generate Illumina Sequencing Libraries..... | 110 |
| 108 | 4.2.5v RAD – Pippin size selection Preparation | 111 |
| 109 | 4.2.5vi RAD – qPCR Quantification..... | 111 |
| 110 | 4.2.5vii RAD – Sequencing..... | 112 |
| 111 | 4.3 ANALYSIS: PROCESSING RADSEQ..... | 113 |
| 112 | 4.3.1 Analysis: Post RAD-Sequencing..... | 113 |
| 113 | 4.3.2 Analysis: Demultiplexing Raw Reads..... | 113 |

| | | |
|-----|--|-----|
| 114 | 4.3.3 Analysis: Mapping to a Reference Genome..... | 114 |
| 115 | 4.3.4 Analysis: Processing Mapped Reads | 115 |
| 116 | 4.3.5 Analysis: Processing Data in SambaR..... | 115 |
| 117 | 4.3.6 Analysis: Data Management in SambaR | 116 |
| 118 | 4.3.7 Analysis: Genetic Analysis in SambaR | 120 |
| 119 | 4.3.8 Analysis: Ancestry Coefficients | 121 |
| 120 | 4.3.9 Analysis: Using different Genome reference..... | 121 |
| 121 | 4.3.10 Analysis: Removing Heterozygosity | 124 |
| 122 | 4.3.11 Analysis: Filtering for Heterozygous Alleles..... | 125 |
| 123 | 4.3.12 De_novo mapping..... | 125 |
| 124 | 4.4 RESULTS: IMPLICATIONS OF POPULATION ANALYSIS..... | 127 |
| 125 | 4.4.1 Results: Genetic Diversity | 128 |
| 126 | 4.4.1 i Genome-wide Diversity..... | 128 |
| 127 | 4.4.1ii Multi-Locus Heterozygosity..... | 130 |
| 128 | 4.4.1iii Numberof private alleles..... | 131 |
| 129 | 4.4.1iv Variation in Segregating Sites (Tajima’s D)..... | 131 |
| 130 | 4.4.2 Results: Genetic Structure | 133 |
| 131 | 4.4.2i Principle Coordinate Analysis (PcoA) | 135 |
| 132 | 4.4.2ii Sequence Dissimilarity | 137 |
| 133 | 4.4.2iii DAPC Analysis..... | 139 |
| 134 | 4.4.2iv Landscape and Ecological Association | 142 |
| 135 | 4.4.3 Results: Population Divergence | 145 |
| 136 | 4.4.3i Population Dissimilarity | 146 |
| 137 | 4.4.3ii Minor Allele Frequency | 148 |
| 138 | 4.5 DISCUSSION AND CONCLUSSION..... | 151 |
| 139 | 4.5.1 Genetic structure | 151 |
| 140 | 4.5.2 Population Diversity and Divergence..... | 153 |
| 141 | 4.5.3 Breeding Strategies in Western European <i>L. bienne</i> | 154 |

| | | |
|-----|---|-----|
| 142 | 4.5.4 Genetic variation in Western European <i>L. bienne</i> | 155 |
| 143 | CHAPTER 5: VARIATION IN TRAITS BETWEEN VERNALIZED AND NON-VERNALIZED <i>LINUM</i> | 157 |
| 144 | 5.2 METHODS..... | 160 |
| 145 | 5.2.1 Data Analysis | 161 |
| 146 | 5.3 RESULTS..... | 163 |
| 147 | 5.3.1 Plant Traits vs. Days to Flowering | 163 |
| 148 | 5.3.2 Plant traits vs. Latitude | 168 |
| 149 | 5.3.3 Plant traits vs Climate Variables (pc1) | 171 |
| 150 | 5.3.4 Seed Area vs Latitude..... | 173 |
| 151 | 5.3.5 Seed Area vs Climate Variable (pc1) | 175 |
| 152 | 5.4 DISCUSSIONS AND CONCLUSIONS | 176 |
| 153 | 5.4.1 Number of Days to Flower and Plant Heights..... | 176 |
| 154 | 5.4.2 Number of Days to Flower and Stem Numbers | 177 |
| 155 | 5.4.3 Number of Days to Flower and Bud Numbers | 177 |
| 156 | 5.4.4 Effects of Vernalization in Plant Traits | 178 |
| 157 | 5.4.5 Effects of Climate in Plant traits for wild <i>L. bienne</i> | 179 |
| 158 | 5.4.6 Effects of Climates in Seed Areas for wild <i>L. bienne</i> | 180 |
| 159 | GENERAL DISCUSSION..... | 182 |
| 160 | APPENDICES | 189 |
| 161 | BIBLIOGRAPHY | 270 |
| 162 | | |
| 163 | | |

164

DECLARATION

165

166 Materials contained within this thesis have not previously been submitted, published or reviewed
167 for a degree at Durham University and/or other institutions. The research, methods, and analyses in
168 this thesis were conducted by the author, unless otherwise stated in the thesis. References are made
169 where the author have used, in part or in full, materials which were already published in the
170 literatures.

171

172 *“Copyright of this thesis rests solely with the author. Quotations and or materials contained within*
173 *this thesis should not be published without the author’s consent and information derived from this*
174 *thesis should be appropriately acknowledged.”*

175

DEDICATION

176

177 I dedicate this work to my mother, father and brother, my mentors/supervisors both past and
178 present and my friends who have helped me and are the foundations to the person I am today.

179

180

181

182 “In the sciences, the authority of thousands of opinions is not worth as much as one tiny spark of
183 reason in an individual man” – Galileo Galilei

184

185 ACKNOWLEDGEMENTS

186

187 Works and studies conducted during this thesis would not have been possible without the
188 support of many people and institutions within my time as a PhD student at the Department of
189 Biosciences in Durham University. I would like to first thank all the institutions which have
190 contributed to the completion of this thesis both technically and financially. I would like to thank the
191 Department of Biosciences at Durham University for providing the correct space and technical
192 equipment needed for the completion of this thesis. I would like to thank the UK Research and
193 Innovation (UKRI) and especially the Biotechnology and Biological Sciences Research Council (BBSRC)
194 for financial provision during my time at Durham University both in forms of laboratory research
195 grants and stipend support towards my living costs while being a student at Durham University. I
196 would like to also extend my thanks to the Royal Botanic Garden Edinburgh (RBGE) as our BBSRC
197 CASE Partner and the financial provision which I have received as part of my research grant from the
198 RBGE.

199 I would like to thank and acknowledge my current supervisor, Dr. Adrian Brennan for his
200 excellent patience and his professional guidance throughout the work of this thesis. I hope to one
201 day be as much of a guide and an excellent supervisor as Dr. Adrian Brennan to me during my time
202 at Durham University, you have made my PhD a richer experience with your extensive knowledge
203 and guidance, despite restrictions caused by the Covid-19 pandemic. I would like to extend this for
204 my other supervisors, Prof. Rus Hoelzel of Durham University and Dr. Catherine Kidner of the Royal
205 Botanic Garden Edinburgh. I would like to further acknowledge Dr. Pilar Suarez-Montes (CONACYT
206 fellow) for her kind help and insights during her time as a Postdoctoral research assistant at Durham
207 University and for her contributions during the chapter two and chapter five of this thesis. I would
208 like to thank all the good friends I have in Laboratory 19 (past and present) of the Department of
209 Biosciences for all the support you have shown me.

210 Finally, I want to acknowledge Dr. Rocio Perez-Barrales and Beatrice Landoni (Portsmouth
211 University) as our excellent collaborators during my time at Durham University. I want to thank you
212 for sharing wild flax materials along with your knowledge for the enrichment of this thesis. I hope
213 the outcome of this thesis will have further implications within your own future research and
214 collaborations.

215

216

217 LIST OF ABBREVIATIONS

218

219 AP1 = APETALA 1 (Gene)

220 CO = CONSTANS (Gene)

221 FLC = FLOWERING LOCUS C (Gene)

222 FLM = FLOWERING LOCUS M (Gene)

223 FRI = FRIGIDA (Gene)

224 FT = FLOWERING LOCUS T (Gene)

225 FUL = FRUITFUL (Gene)

226 GAPDH = Glyceraldehyde-3-phosphate dehydrogenase (Gene)

227 GI = GIGANTEA (Gene)

228 GOI = Gene of interest

229 HKGs = House Keeping Genes

230 LFY = LEAFY (Gene)

231 NGS = Next Generation Sequencing

232 NRT = No Reverse Transcriptase

233 NTC = No Template Controls

234 PCR = Polymerase Chain Reaction

235 RADSeq = Restriction-site Associated DNA Sequencing

236 RGE = Relative Gene Expression

237 RT-qPCR = Real-time quantitative polymerase chain reactions

238 S0 = Selfing 0 (Wild selfing generation)

239 S1 = Selfing 1 (First selfing Generation)

240 SD = Standard Deviation

241 SEP3 = Sepallata 3

242 SNPs = Single Nucleotide Polymorphisms

243 SOC = Suppressor of Overexpression of Constans

244 SVP = SHORT VEGETATIVE PHASE (Gene)

245 UBI = Ubiquitin (Gene)

246 VRN = Vernalization

247 **ABSTRACT**

248

249 Local adaptation plays a major part in plant survival and reproduction. *Linum* represents a
250 genus of potential study models which provide insights for both applied and evolutionary biology. As
251 a flowering plant, they may have evolved adaptations to achieve optimal flowering time. Most
252 flowering plants have developed their own strategies to flower in specific habitats. There are
253 interests from both evolutionary, and agricultural points of view regarding flowering development.
254 In evolutionary terms, flowering time may affect offspring and population fitness. In agriculture,
255 faster flowering time is a desirable trait for production. For this reason, *Linum* is a versatile model to
256 study. *Linum usitatissimum* (cultivated flax) is useful in several industries. Their wild predecessors,
257 the wild flax (*Linum bienne*) is relatively less studied in comparison to their cultivated relative. As
258 their predecessor, we suggest that implications of local adaptation in the wild flax, may aid the
259 development of their cultivar relatives. We examined both wild and cultivar type to determine local
260 adaptation strategies, particularly in terms of flowering.

261 In chapter 2 of this thesis, we examined three flowering time genes and two duplicate genes.
262 *Linum* is a temperate plant and as such requires vernalization. Vernalization is the process in which
263 plants require colder temperatures to induce the flowering process. Plants that require vernalization
264 often flower earlier when in colder temperatures than those that have not experienced cold
265 induction. We studied expression of five flowering time genes for implication of local adaptation
266 after treatment to vernalization. *L. bienne* appeared to express genes differently in comparison to
267 the cultivars. The expression of *Linum* FLOWERING LOCUS T (*LuFT*) revealed a positive correlation
268 with number of days to flower. This potentially identifies FLOWERING LOCUS T as one of the
269 important genes regulating vernalization in *Linum*. Our result revealed variation in relative flowering
270 time gene expressions. Wild and cultivated *Linum* demonstrate different relationships between
271 flowering time and environmental variables.

272 In chapter 3, we quantified the viability of pollen, an important part in the transfer of the
273 male gamete in flowering plants, under different temperature treatments. *Linum* is an established
274 temperate plant. Sensitivity to temperature changes maybe more predominant in temperate plants
275 as seasonal changes would reflect a challenge to flower in temperate environments. In this chapter,
276 *Linum* pollen was treated under different temperatures to observe their ability to germinate. This is
277 important to determine whether temperature plays a major part in affecting the viability of pollen,
278 which in turn plays a major role in the formation of seeds. This chapter revealed a reduction in the
279 number of pollen tubes formed under different treatments and across the two *Linum* species. In

280 addition to this, correlations to local climates were also observed, with variation in trends across the
281 temperature treatments.

282 In chapter 4 of this thesis, we examined population genetics of wild *L. bienne* samples
283 originating from different latitudes across western Europe. We examined Wild flax (*L. bienne*)
284 populations across western Europe to provide insights into their genetic structure and diversity. This
285 population analysis will develop our understanding of adaptation in wild *Linum* in response to their
286 environment. A double-digest RAD sequencing (DDRadSeq) protocol was utilized to look at variation
287 in SNPs across different populations. Both *L. bienne* and *L. usitatissimum* samples were sequenced
288 and aligned to an *L. usitatissimum* whole genome. Genetic structuring of our *Linum* samples were
289 revealed across Western Europe. Cultivars in our collection revealed to be more genetically related
290 to the Northern accessions of our wild samples.

291 In chapter 5 we summarized plant architectures. In this chapter traits and their relationship
292 to the latitude were summarized in relation to the requirement for vernalization. Four traits were
293 measured: overall height of the plants after first flowering, the number of stems of the plants after
294 first flowering, and the number of flower buds on the plants at first flowering. In addition to this,
295 seed size was examined in terms of its area. We also examined correlation between traits and
296 environmental variables. We found that there was a relationship between traits when no-
297 vernalization occurred. With vernalization, these relationships became less significant and, in some
298 cases, not significant. This illustrates that vernalization influences the relationship of traits beyond
299 flowering initiation. There were also suggestions that the traits measured correlate with latitude
300 under no-vernalization treatments. For vernalized individuals, the correlation for the traits measured
301 and latitude was not significant. Seed sizes were strongly correlated with both latitude and climatic
302 variables in all cases of the treatments. The findings suggest there are effect of environmental
303 variables in these measured traits, which suggests differentiation within wild *Linum* species.

304

305

306

307

308 CHAPTER 1: INTRODUCTION

309

310 For living organisms, adaptation to the local environment is important for species survival
311 and the success of their offspring. This adaptation to the environment is often referred to as “local
312 adaptation” (Brandon R N, 2014). Local adaptation often results in a higher mean fitness of a
313 population in their native environment. Large numbers of reciprocal studies and garden experiments
314 have illustrated this in plants (Lascoux M. et al., 2016). Local adaptation usually arises from biotic
315 and abiotic factors which represent a selection pressure that organisms must adapt to (Rúa M. et al.,
316 2016). Studies also suggests that local adaptation could be climate-driven, with suggestion that
317 locally adapted tree population managing to track climate change due to local adaptation (Moran E.,
318 2020). It is well established that local adaptation arises from selection by the local environment
319 which favours specific phenotypes. This can translate into a genotype-by-environment interaction in
320 response to biotic and abiotic factors (Sork V., 2018).

321 More generally, local adaptation is understood to be pervasive and is a key evolutionary
322 process that contributes towards the success of a population (Sork V., 2018; Meek M. et al., 2023). It
323 is generally a result of divergent selection on traits which can lead to reproductive isolation and even
324 speciation (White N. and Butlin R., 2021). One outcome of local adaptation is the maintenance of
325 ecologically important genes (Whitlock M., 2015). Adaptation to local environment may provide
326 insight into improving survival rate for populations not on the brink of extinction, to maximize their
327 likelihood of long-term persistence (Bay R. et al., 2018).

328 Meta-analysis of plant population suggested that locally adapted populations of plants tend
329 to perform better than foreign plants when at the site of their origin (Leimu M. and Fischer M.,
330 2008). In addition to this, they also suggested that population size have a positive role on the ability
331 of a population to locally adapt (Leimu M. and Fischer M., 2008). In plants, responses to selection
332 pressure in their environment may lead to differences in phenotypic traits, such as the size and
333 shapes of their leaves (Dudley S., 1996). Local adaptation of species to the environment have also
334 been previously observed in germination behaviour of plants (Donohue K. et al., 2010). Variation in
335 local adaptation can affect biodiversity of a given environment (Atkins K.E. et al., 2010) (Adams, J.,
336 2009).

337 There are various ways in which plants can adapt to their local environment, this is often
338 illustrated not only in phenotypic, but also genotypic changes of a plant’s biological make-up. A
339 study in various alpine landscapes found that diversity in the grass species *Poa alpina*, was affected
340 by their localities. This along with variation in phenotypes was found as a result of local adaptations

341 (Stöcklin J. et al., 2009). Local adaptation along with variation in breeding strategies may suggest
342 significance in an ever-changing environment. This is of interest for different species under different
343 environments. Evolutionary processes such as genetic selection and drift may provide opportunities
344 for further study to the importance of ecological genetics to plant adaptations, whereby genetic
345 mechanisms can be outlined for a specific species to understand their evolutionary processes and
346 ecological implications (Anderson J. et al., 2011). Differences in each gene pool can be used to detect
347 processes like genetic drift and speciation. In turn, this can influence performance of different
348 populations. Local adaptation studies in aquatic plants, reveal variation in performance between
349 Northern and Southern populations which alludes to different genetic performance due to genetic
350 drift and inbreeding (L. Santamaría *et al.*, 2003). Population genetic studies of different *Linum*
351 population could potentially reveal genetic factors which could contribute to performance of
352 different populations. This can then be utilised to underpin genetic loci to manipulate *Linum* under
353 different environments. This is particularly of interests in agriculture.

354 In the flowering plants, flowering strategies often form an important set of adaptations to
355 the local environment. For example, in *Mimulus guttatus* populations, selection to flower at the
356 correct time causes them to exhibit local adaptation to the environment (Hall M. et al., 2007). One
357 environmental factor influencing flowering is the sensitivity of an individual's flowering time to the
358 availability of sunlight in their local area (photoperiod sensitivity). Environmental selection on
359 flowering times was observed in *Campanulastrum americanum* to influence reproduction under
360 different natural light conditions (Galloway F., 2012). Responses to selection in flowering time is of
361 interest as flowering affects reproductive mechanisms of plants in the wild and affects offspring
362 health and survival. In the perennial plants, a single genomic region mediates local adaptation. This
363 region contains the floral integrators *FT* (Wang J. et al., 2018). Genetically, early/late flowering
364 populations is suggested to show differences in their expression of flowering time genes (Reeves
365 P.H. et al., 2000). This in turn reveals genetic differences in flowering time strategies for different
366 populations. In the model *Arabidopsis thaliana*, flowering time gene network and expression under
367 different treatments reveals several pathways, with photoperiod and vernalization pathways being
368 the most well established (Putterill J. et al., 2004). Studies reveal *CONSTANS (CO)* and *FLOWERING*
369 *LOCUS C (FLC)* genes are key to photoperiod and vernalization perception and regulates *FLOWERING*
370 *LOCUS T (FT)* (Kinmonth-Schultz *et al.*, 2021). With the suggestion that the floral integrator *FT* is in
371 the genomic region that could potentially mediate local adaptation, it is of interest to observe
372 variation in flowering-time genes and phenotypes.

373 For agriculture, the importance of local adaptation may reveal conditions for optimal yield of
374 a cultivar type. This is of interest to stakeholders in which diverse agroecosystems can be optimized

375 to prioritize climate adaptation responses in an ever-changing environment (Lee et al., 2014). This
376 could be important in sustainable agriculture, which is of interest to stakeholders in the agricultural
377 sector (Öhlund et al., 2015). In cultivated Maize (*Zea mays*) for example, genetic variants were found
378 to be associated with flowering time, an important adaptation mechanism in temperate plants (Li et
379 al., 2016). In cultivated apple trees (*Malus domestica*), genetic mapping of different cultivar types
380 suggested a cold perception mechanism which was linked to the flowering time network of genes
381 (Allard et al., 2016). It is of interest then to observe whether local adaptations to environmental
382 variables in cultivated lines is affecting the performance of such lines in terms of their sustainable
383 production in certain environments and to find optimal treatments for important cultivars to yield
384 products optimally.

385 As a crop, factors such as plant height, number of stems and as well as seed sizes are of
386 interest. It is of interest to look at whether seed size contributes to differences in reproduction as
387 well as differences in physical formation of seeds based on environmental conditions. It has been
388 suggested that seed size is regulated, and that viability of the embryo is enhanced when the seed
389 contains a substantial amount of starch and protein for the seedlings (Sundaresan V., 2005). This
390 suggests an advantage to having bigger seeds. In the model *Arabidopsis*, it has been shown that
391 environment affected seed size and that an observation was made whereby variance in progenies'
392 morphology were affected by parental environment (Brown et al., 2019). This potentially extends to
393 development across generations. When looking at plant heredity and performing genetic studies,
394 seed sizes are one factor of interest in relation to parental growth conditions. Looking at wild *Linum*
395 *bienne* grown in different environment may reveal the parental effect regarding morphology
396 structure such as seed sizes. Seed sizes may also be affected by the requirement of seed to “sense
397 cold”, otherwise known as vernalization. In addition, population genetics study may reveal
398 structuring and local adaptation mechanisms, previously unknown in *Linum*.

399


400

401 1.2 INTRODUCTION TO STUDY SPECIES

402 In this thesis, two species of the flowering plant in the genus *Linum* were studied. *Linum* is a
403 genus of flowering plants which contains more than 200 identified species, distributed in the
404 subtropics and temperate zones (Öhlund et al., 2015; Muravenko et al., 2010). *Linum* is found in
405 temperate into the subtropical regions with species such as the blue flaxes *L. lewisii* and the *L.*
406 *narbonense* often cultivated as garden ornamentals (Ionkova et al., 2013; Addicott, 1977). In
407 horticulture, prolific flowering makes them of interest in ornamental uses (Tork D. et al., 2022.
408 Geographically, *Linum* is distributed across Northern Americas, and the European continents.
409 Another cultivated species which is important industrially includes the *L. usitatissimum* (Jhala A. et
410 al., 2010). The wild ancestor of this cultivar is thought to be the pale flax *L. bienne* (Uysal et al.,
411 2010). Allaby *et al* (2005), reveals that there is a single domestication event for *Linum*, and they
412 suggest that the *sad2* locus of *Linum* was subject to artificial selection in the cultivars. The wild
413 relative is significant because it represents a primary gene pool for development of the cultivar.
414 Previous study reveals that there is a genetic differentiation between the two species and suggests
415 that 48% of *L. bienne* alleles were unique (Soto-Cerda et al., 2014). With this suggestion, it is also
416 possible to map *L. bienne* sequences to the cultivars to identify alleles favourable to the cultivars
417 (Soto-Cerda *et al.*, 2014). In this thesis, both wild (*L. bienne*) and cultivar (*L. usitatissimum*)
418 individuals will be examined.

419 Morphologically, *L. usitatissimum* tends to have taller stems when compared to *L. bienne*.
420 They also tend to have larger seeds, capsules, and flowers. The pale flax (*L. bienne* Mill) was
421 previously shown to vary in flowering phenology (Uysal et al., 2011) and the cultivated flax, *L.*
422 *usitatissimum* has been extensively studied in terms of its phenology with studies suggesting
423 development of certain phenotypes to optimise yield (Rehman et al., 2014). Wild and cultivated flax
424 differ in their stem structure seeds (Figure 1) .

425



426 Figure 1. Collection of wild and cultivated S1 (selfing) *L. usitatissimum* and *L. bienne* grown for >3 months
 427 period. Cultivars (in red boxes) are growing taller, sturdier stems than wild relatives. There is variation in plant
 428 heights and number of stems in our *Linum* samples (1A). Wild (*L. bienne*) and cultivar (*L. usitatissimum*)
 429 relatives also have varying seed sizes. Wild relatives (1B) tend to be smaller in length and width than the
 430 cultivars (1C). Each square is 1mm in area for Figures 1B and 1C.

431

432 The availability of *L. usitatissimum* whole genome sequence (Wang et al., 2012) means that
 433 studies looking at implications in the cultivars due to domestication processes were possible and
 434 thus the ability to infer some genetic consequences from domestication were revealed (Fu, 2012; Fu,
 435 2011). Although the availability of a whole genome is true for *L. usitatissimum*, there is currently no
 436 published whole genome for wild species of *Linum*.

437

438 1.2.1 *Linum usitatissimum*: Cultivated Flax

439 *Linum usitatissimum*, otherwise known as the “common cultivated flax” is an agriculturally
 440 important plant that serves multiple purposes (oil and fibre morphotypes). Therefore, *Linum* is of
 441 economic significance in agriculture. In terms of their morphology, You F. et al., 2017 found that

442 fibre morphotypes have greater straw weight, plant height and protein content than the oil
443 morphotypes while the oil morphotypes have greater seed weight, seed oil content, and branching
444 capability (You F. et al., 2017). Measures of oil and stem fibre content of several oil/fibre
445 morphotypes also shows variability, suggesting variation within this species for industrial use
446 (Diederichsen and Ulrich., 2009; Rozhmina T. et al., 2021).

447 Historically, *Linum usitatissimum* has been cultivated for its fibres and seeds, with evidence
448 of domestication in the Middle East, revealed by the diversity of the *sad2* gene locus (Allaby et al.,
449 2005). Plant domestication is an important process which enables human civilizations to utilize plant
450 materials for advancements (Diamond, 2002). In terms of agriculture, traits for larger seed (size and
451 weight) and fibre properties have been selected for, when considering flax as cultivars with
452 development for said traits still of interest to this day (Rahimi M. et al., 2011; Yan L. et al., 2014; Guo
453 D. et al., 2020). As a result of domestication, genes for seed sizes, flowering time, and capsule
454 dehiscence are revealed to be artificially selected (Zhang J. et al., 2020). More recently, qualities of
455 cell wall, stem strength and fibre properties are studied for using flax fibres as an environmentally
456 friendly composite alternative (Goudenhoft C. et al., 2019). Selective breeding in a trait may imply
457 better selection for that trait, however in long term, the consequences to those selected lines in
458 terms of survival and loss of functions in other genes have not been tested yet. It has been shown
459 that the temperate region of the world started to cultivate flax thousands of years ago.
460 Archaeological records illustrate that flax were cultivated during the Viking and Medieval age
461 (Ejstrud, et al., 2011) (Nag, et al., 2015). This suggests a process of long-term artificial selection in *L.*
462 *usitatissimum* with implications in several field of interests for agriculture. We are interested in
463 flowering time for this study. We suggest that selection for faster flowering in the cultivars may
464 result in loss of signalling in flowering time genes.

465 *Linum* seeds produces oleic compounds composed of triglycerides, and these are particularly
466 rich in α -linolenic acid, which gives linseed oil its ability to polymerize into solids. This is useful in
467 productions of industrial resins and solvents (Vereshchagin, & Novitskaya, 1965; McCullough, et al.,
468 2011), suggesting significance in the industry sector. In addition to this, recent research has found
469 that linseed oil has a positive effect on reducing the level of cholesterol in rabbits, illustrating
470 potential for linseed in the health food sector (Króliczewska, et al., 2018). Molecularly, it is suggested
471 that *Linum usitatissimum* seeds contains several biologically active components that are useful both
472 medically and industrially. This includes the presence of lignin in linseed, useful in resin production
473 (Del Rio et al., 2011; Touré & Xueming, 2010). These lignans have shown to be medically beneficial in
474 relation to their antioxidant properties (Hosseinian, et al., 2006). Fatty acid profiling has also
475 revealed that linseeds provide significant nutritional values (El-Beltagi, et al., 2007), suggesting

476 values in food security. Other potential uses of cultivated *Linum* species is their biogeographic trait
477 variation. This could be useful for ornamental purposes. In recent study of the ornamental *L. lewisii*,
478 traits such as flowering indeterminacy, seed mass and stem numbers suggest its potential use in field
479 restoration and agriculture (Innes, et al., 2022). These findings reveal that not only the common
480 cultivated flax was cultivated early on for their fibres, but also research into flax biproducts and
481 genetics is still ongoing with a strong potential in multiple fields such as industry, agriculture, the
482 health food sectors and horticulture. With these in mind, it is then of interest to optimise *Linum* crop
483 production. Examples of research in this area includes seed priming which influences crop growth
484 and development (Rehman H. *et al.*, 2014), and experiments to employ plant growth regulators
485 (giberellic acid) for optimal yield and growth in *L. usitatissimum* (Rastogi A. *et al.*, 2013). We propose
486 that, information into local adaptations and genetic implications of *Linum* wild crop relatives could
487 aid these efforts to further improve *Linum* as a crop plant. This may be done by exploring genetic
488 and phenotypic responses to environment changes to benefit agriculture. To this day, research for
489 implementation of wild relatives towards genetic engineering cultivated flax is relatively scarce. For
490 studies in this thesis, 18 varieties of *Linum usitatissimum* seeds were received from IPK World
491 Collection (<https://www.ipk-gatersleben.de/en/>). A list of cultivars and their morphotypes can be
492 found in appendix 1.

493

494 1.2.2 *Linum bienne*: Wild Flax

495 *Linum bienne* is a biennial plant and a wild relative of *Linum usitatissimum* (Gill KS, 1966). Its
496 common name is the “pale flax”, considered the wild forebear of the cultivated *Linum usitatissimum*.
497 A study by Allaby *et al.*, reveals this by exploring *sad2* loci from 30 accessions of wild and cultivated
498 flax. They found phylogenetic evidence that the wild type *L. bienne* was first domesticated for oil and
499 that there is an artificial selection of the loci, indicative of cultivation of the wild types *L. bienne*
500 (Allaby, et al., 2005). Further studies also revealed that *sad2* locus is a candidate domestication locus
501 associated with increased unsaturated fatty acid production in cultivated flax (Fu Y.B. et al., 2012).
502 This suggests implication for seed selection when *Linum* was first domesticated as opposed to fibre.
503 In the literature, there are suggestions that *L. bienne* are widely distributed around Western Europe,
504 the Mediteranean basins, North Africa and into Iran and the Caucasus. This represents its wide
505 distribution in sub-temperate into temperate zones (Zohary and Hopf, 1993). Just like the cultivar *L.*
506 *usitatissimum*, this wild progenitor has the same number of chromosome ($n=15$) and they’re both
507 homostylous, in contrast to some distylous relatives such as the *Linum Tenue* (Gutiérrez-Valencia *et*
508 *al.*, 2022).

509 Previously the wild *L. bienne* were considered as sub-species of *L. usitatissimum* (Uysal et al.,
510 2011), but now they are considered as the closest relative to *L. usitatissimum*. Phylogenetic studies
511 have illustrated that *L. bienne* is the closest relative of *L. usitatissimum* (Uysal, et al., 2010). Further
512 phylogenies from studies of *Linum* orthologs also suggests that *L. bienne* is the closest relatives of *L.*
513 *usitatissimum* (McDill J. et al., 2009; Sveinsson S. et al., 2011). This species represents a wider gene
514 pool, which can be of aid for *Linum usitatissimum* development in agriculture.

515 *L. bienne* is native to the Mediterranean and parts of Western Europe and are spread as far
516 north as the Scandinavian countries. Naturally, growing in different environments compared to
517 *Linum usitatissimum*, the *biene* represent a wider gene pool, which could be of interest in the
518 enhancement of *Linum usitatissimum*. Differences between wild populations could shed light on the
519 evolutionary trends of this species as well as their cultivar relatives. In the literature, population
520 genetic study of wild relative of flax is relatively scarce. Studies conducted in this thesis will suggest
521 some implications about the importance of wild relatives for future studies looking at the genetics of
522 *Linum*. Across Western Europe wild *L. bienne* are distributed across a range of latitude. For studies in
523 this thesis, collection of seeds was essential. Seeds from the wild were first collected in 2016 by Dr.
524 Adrian Brennan of Durham university and Dr. Rocío Pérez-Barrales of University of Granada. The
525 plants collected throughout western Europe. Different populations with different number of
526 individuals were used in each aspect of studies in this thesis. We used this collection in the
527 corresponding chapters.

528

529 1.2.3 Production Values of *Linum* Biproducts:

530 With the above put into consideration, the current observation is that flax output is
531 economically significant. One of the biproducts of *Linum* is its fibre. Flax fibre production in 2020
532 stands at 976,113 tonnes (FAOSTAT data, 2022). In 2016 alone flax fibre output reached a
533 production value of 535 million USD. This value increased more in 2020, with flax world trade value
534 estimated at 726 million USD (Flax Fibers | OEC, 2022). The world's largest flax fibre exporter in 2020
535 is France, with an estimate of 51.4% in total world export value. The export share in the UK only
536 stands at 0.12% with an export value of 870 thousand USD. The world's top importer of flax fibres is
537 China with 47.8% total import value which stands at 347 million USD. The UK imports 0.14% of total
538 flax fibre import value, with an import value of 1.03 million USD (Flax Fibers | OEC, 2022). The export
539 value growth from the year 2000 – 2020 in the UK is -76.7%, with export values, massively reduced
540 from 3.74 million USD at the start of this period. This reveals, that at least in the UK, current export
541 value in flax fibre is decreasing. This being the case, in developing countries demands for flax fibre

542 are increasing. In India alone, the import value increased from 5.93 million USD in the year 2000 to
543 47.7 million USD in 2020, an increase of 703%. Other countries such as China reveals an import value
544 growth of 254% between the year 2000 and 2020 while Vietnam shows a >1000% increase (Flax
545 Fibers | OEC, 2022). This new trend in demand of flax fibre in developing countries could provide a
546 reason for temperate countries such as the UK to increase export of flax fibres.

547 Other major biproduct of *Linum* cultivars is linseed. In 2020, the total world production of
548 linseed is 3,367,331 tonnes, with Kazakhstan producing the most at 105,8247 tonnes (FAOSTAT data,
549 2022). In the same year the trade value of linseed worldwide is 980 million USD. Major exporters of
550 linseed include Canada (25.8%), Russia (23.5%) and Kazakhstan (21.7%). Major importers of linseed
551 include China (26.8%), Belgium (22.7%) and Germany (8.75%). In the UK, export of linseed grew
552 57.7% between the years 2000 – 2020. The total export value of linseed in the UK in 2020 is 7.28
553 million USD. The biggest increase is seen in Kazakhstan between the years 2000 – 2020 (Flax Fibers |
554 OEC, 2022). The increase in export of linseed illustrates the economical values linseed represents
555 worldwide in terms of industry and agriculture. Potential to improve cultivar types will be of huge
556 interest to the economic values represented in *Linum* biproducts derived from flax fibres and
557 linseeds.

558

559

560 1.3 LITERATURE REVIEW OF *LINUM* GENETIC DIVERSITY

561 *Linum* underwent events of genome duplication and domestication. With *Linum*
562 domestication, it is worth considering the consequences that may have come with this process.
563 While vital to human development, domestication of wild plants often leads to a genetic
564 disadvantage because of divergence from the wild that could be caused by selective breeding,
565 associated with the process of domestication. This is often due to selection for “preferred traits”
566 (Purugganan & Fuller, 2009). Selected traits, either directly or indirectly selected, may in turn have
567 genetic implications within the domesticated populations. Fu Y. B., (2012) revealed that cultivated
568 flax have 27% reduction of nucleotide diversity when compared to the wild pale flax (*L. Bienne*),
569 perhaps because of domestication. Genetic effects such as the pleiotropic effects (Conner, 2002) and
570 linkage disequilibrium (Falconer, D.S., 1996) may also have occurred because of selective breeding in
571 domestication. This could lead to further genetic consequences such as reduced genetic diversity,
572 and increased chances of genetic drift (Rauf, et al., 2010).

573 Genetic diversity analyses and studies with different *Linum usitatissimum* cultivars are
574 widely available (Nag, et al., 2015; Diederichsen, 2001). These resources can be used for breeding
575 programmes in *L. usitatissimum*. Past studies have identified morphological traits-based diversity in
576 *L. usitatissimum* seed variation (seed colour, seed weight, and seed oil contents). They suggest
577 indirect selection for seed weight and seed colours (Diederichsen and Raney, 2006). Seven linseed
578 genotypes have previously found to have quantitative traits differences between seed traits
579 (Nôžková J, et al., 2014), suggesting variation in *L. usitatissimum* genotypes. Zhang et al, (2020)
580 conducted a genome wide association study for *L. usitatissimum* and suggested that during flax
581 domestication, genes relevant to flowering, dehiscence, oil production, and plant architecture were
582 artificially selected and that selection in these genes may shape their morphology . There is a
583 suggestion here of artificial selection in *L. usitatissimum*. In addition, molecular markers are also
584 widely available for *L. usitatissimum* genetic studies. Cloutier et al, (2009) have defined 83 Single-
585 Sequence Repeat (SSR) motifs for 23 *L. usitatissimum* accessions. In addition to this, another study
586 defined 28 SSR markers for the study of genetic fingerprinting in *L. usitatissimum*, which is useful in
587 assessing genetic purity (Pali V et al., 2014). A consensus of genetic and physical maps of *L.*
588 *usitatissimum* is also available under high resolution (74% of the estimated flax genome) (Cloutier S.
589 et al., 2012). These genetic resources don't consider wild relatives, which may be of further interest
590 in this area, and when looking at breedable traits. Initial studies looking at the wild relatives *Linum*
591 *bienne*, illustrates that they represent a wider genetic diversity from which the cultivar is derived
592 (Uysal et al., 2010). Soto-Cerda et al also suggested genetic association mapping is possible using
593 wild relatives of *Linum* to identify favourable alleles within the wild (Soto-Cerda et al., 2014).

594

595 1.3.1 Methods for studying population genomics in *Linum*

596 Within this study, we compare genetic diversity of wild *Linum bienne* samples, relative to the
597 cultivars. With regards to genetic analysis, it is worth noting Next-Gen sequencing (NGS) techniques.
598 NGS makes it possible to study local adaptation at a population level without a whole genome
599 sequence. This is of interest when looking at population genetics of wild relatives of agricultural
600 plants (Park & Kim, 2016). In relation to our study model, NGS is valuable in terms of gaining insights
601 into the genetic diversity of *L. bienne* and *L. usitatissimum*. Population analysis of wild relatives may
602 also be informative in the development of crop relatives. Several NGS techniques are available, and
603 the sequence data can be mapped de-novo or to a reference genome. This is relatively faster and
604 cheaper than producing a whole genome for the wild relatives.

605 With next generation sequencing technologies available, several *Linum* molecular markers
606 (microsatellites) have been developed (Fu & Peterson, 2010). It was shown that molecular markers
607 are transferable in many species of *Linum* (Soto-Cerda, et al., 2011). These transferable molecular
608 markers, often referred to as simple sequence repeats (SSRs), have been useful for data
609 enhancement for genetic and evolutionary studies. Because of this transferability, genetic and
610 evolutionary studies can compare between wild and cultivars of flax. Numerous species in the *Linum*
611 genus represent an interest to study their evolution, while the cultivar (*L. usitatissimum*) represents
612 an interest in the development of *Linum* as a crop through study of their genetic makeups and local
613 adaptation mechanisms such as self-incompatibility mechanisms (SI) and inbreeding depression
614 which their wild relatives may be able to reveal.

615 For whole genome sequencing, Illumina's Hi-C sequencing has become a mainstream
616 sequencing technique. It is preceded by the 3C method (Downes et al., 2021). The method is widely
617 used to examine organisation and conformation of chromosomes and secondary genetic structures
618 into chromosomal level organisation based on whole genome sequencing (Belton et al., 2012;
619 Lieberman-Aiden et al., 2009). The method involves the use of isoschizomer restriction enzymes
620 such as DpnII to digest cell nuclei isolates, recognising and cutting amino acid sites to generate an
621 overhang. These overhangs can then be used to to enable paired-end sequencing using adaptor
622 ligation (Belaghzal et al., 2017). However, to gain whole genome sequences often requires laborious
623 procedures that are relatively more expensive than reduced representation sequencing techniques
624 such as RAD Sequencing. This sequencing technique can help with identification of markers such as
625 single-nucleotide polymorphisms (SNP) that can be used to identify genetic signals such as
626 structuring between population and even identify novel loci for quantitative trait locus analyses.

627 The genome of *L. Usitatissimum* is estimated at 370Mb in size (Wang et al., 2012). Whole
628 genome sequences are publicly available down to chromosomal levels for the cultivated *L.*
629 *usitatissimum* having first been assembled using shotgun sequencing technologies in 2012 (Wang Z.
630 et al., 2012). The contig assembly contained 302 Mb of non-redundant sequence, representing an
631 estimated 81% genome coverage (Wang Z. et al., 2012). The initial genome is now referred to as CDC
632 Bethune v1 (Sa et al., 2021). The most widely used *L. usitatissimum* whole genome is the CDC
633 Bethune v2 (Sa et al., 2021; You et al., 2018). For this whole genome contigs contains a total N50 of
634 6.64Mb summing up 316 Mb of reads and a 97% coverage of annotated genes, considerably higher
635 than the previous CDC Bethune v1 (You et al., 2018). Another genome was assembled using long
636 reads for the Atlant variety of *L. usitatissimum*. Using a combined Oxford Nanopore and Illumina
637 sequencing technique, whole genome with a total length of 361.7 Mb, N50 of 350 kb, and 97.40%
638 completeness was recently achieved for the Atlant variant of the *L. usitatissimum* (Dmitriev et al.,
639 2021). The availability of this whole genome also suggests that continuity of these assemblies is
640 relatively poor as revealed by the relatively small contig N50. The largest contig N50 was found to be
641 365Kb (Sa et al., 2021; Dmitriev et al., 2021). They suggest that events such as recent whole-genome
642 duplication may result to the collapse of homologs and repeat sequences during the assembly
643 process, both under short and long reads sequencing (Sa et al., 2021; Dmitriev et al., 2021).

644 For the wild *L. bienne*, whole genome sequences are under development and not publicly
645 available. However, past studies using inter simple sequence repeats (ISSR) and RADSeq markers
646 have shown that the wild *L. bienne* illustrates ancestral relationship with the cultivated *L.*
647 *usitatissimum* and there were indications that suggests the contribution of *L. bienne* to *L.*
648 *usitatissimum* genome through gene flow (Fu, 2012; Gutaker RM., 2014). Molecular and cytogenic
649 studies of both cultivated and wild *Linum* were also undergone using high-throughput ribosomal
650 RNA. The studies suggests that 5S rDNA and ITS phylogeny is closely related between *L.*
651 *usitatissimum* and *L. bienne*. Both species have identical karyotypes and distribution of rDNA sites. In
652 both species, the coding sequences of 5S rDNA genes were similar (Bolsheva N. et al., 2017).
653 Comparison of ITS1 to 5.8S rDNA-ITS2 sequences showed that rDNA sequences were conservative in
654 all studied flax specimens (Bolsheva N. et al., 2017). In addition to this, Assessment of the number of
655 polymorphic sites in a genetic marker (*LuTFL* homologs) suggests mixed ancestry of the locus and
656 different copy of the locus suggests ancestry to the wild pale flax *L. bienne* (Gutaker RM., 2014).
657 More Northern populations are suggested to have high similarity to the *L. usitatissimum* (Gutaker
658 RM., 2014). This high similarity illustrates that whole genome for the cultivar is acceptable for use in
659 the wild *L. bienne* before the availability for its whole genome.

660 Other genetic events have also occurred as *Linum* species have undergone polyploidy
661 events. In plants, polyploidy is an event whereby whole genome duplication occurred increasing the
662 number of chromosomes. An example of this has also occurred in sugarcanes (Vilela et al., 2017).
663 Genetically, the genus *Linum* has experienced multiple polyploidy events. The last polyploidy event
664 occurring around 5-9 million years ago (mya) and a previous polyploidy event around 20-40 mya.
665 This reveals that every gene in the flax genome is potentially duplicated and multiplied through
666 polyploidy (Sveinsson et al., 2013). This needs to be accounted for when looking at their genetic
667 make-ups and for studying gene expression (Sveinsson et al., 2013). Polyploidy is important, as it is a
668 widely accepted to be a pervasive mechanism of plants and is often consequently followed by the
669 selective silencing of genes (Wendel & Adams, 2005). Studies have previously inferred that DNA
670 alterations in flax could induce changes that are heritable. These changes were also linked to
671 environmental factors (Schneeberger & Cullis, 1991). Polyploidy events can consequently rid of
672 genes, which were previously present in predecessors of cultivated flax. With the aid of cultivation,
673 breeding strategies of the genus, conditions may have changed and thus the genetic make-up of
674 cultivars is heritably different to those of the wild progenitor, representing a wider gene pool to be
675 studied.

676

677 1.3.2 Self-Incompatibility in *Linum*

678 In plants, control of outcrossing is often due to self-incompatibility (SI) genes at the S locus
679 (Newbigin E. et al., 1993). This locus is responsible for the prevention of self-fertilization and
680 therefore selfing in plants. Inactivation of the S-locus receptor kinase (SRK) and the S-locus cysteine-
681 rich protein (SCR) within this locus has led to the loss of self-incompatibility in some plants in the
682 *Brassicaceae* family (Sherman-Broyles, et al., 2007) and *Arabidopsis* (Suzuki, et al., 1999). Self-
683 Incompatibility genes is not yet fully understood in *Linum*, research focuses mainly on the formation
684 of hetero/homo styly. Differences in the S-genes expressed in the male and female reproductive
685 structure trigger self-rejection, controlling outcrossing, and leading to high heterozygosity
686 (Thompson & Kirch, 1992). Heterostyly in *Linum* means that there are morphological differences
687 between styles and anther filaments of *Linum* and it has been found as an outcrossing mechanism in
688 *Linum* (McDill, et al., 2009). Therefore, in *Linum*, polymorphisms of breeding organs contribute to
689 heterozygosity (Ebert, et al., 1989). In *Linum grandiflorum*, the S-locus controls the flower
690 morphology through regulations of transcriptional S-locus products (Ushijima, et al., 2011). This
691 reveals the connection of the S-locus regulation with the ability for *Linum* to promote heterozygosity
692 and therefore maintain genetic variation. The mating system of wild flax (*L. bienne*) is not yet fully
693 understood, although related species show distyly to ensure outcrossing, research regarding

694 polymorphisms in the wild relatives are still scarce to date. This thesis will explore aspects of *Linum*
695 reproduction by looking at pollen morphology, especially pollen viabilities measures such as the
696 availability of pollen to germinate under environmental stress, This will provide trait data to study
697 genetic mechanisms which may affect such morphology.

698

699 1.3.3 Population Genetics Analysis

700 In modern genetics, there is an interest in determining genotypes which represent
701 underlying phenotypes. These are often crucial to a study model with potentially many populations,
702 representing a wide gene pool. To do this, it is crucial to gain insight into genetic sequences of many
703 individuals relatively quickly and cheaply. For the purposes of sequencing, high capacity and
704 relatively low costing technology for sequencing were first achieved using microarrays-based
705 sequencing technologies (Schna et al., 1995). Later, widely used genetic markers were introduced
706 as “microsatellites”, also known as Single Sequence Repeats (SSRs) across the whole genome (Vieira
707 et al., 2016). This genotyping technology was a pre-cursor of Next-Generation Sequencing (NGS)
708 technologies such as RAD Sequencing. NGS provides sequence data for nucleic acids, DNA and RNA,
709 that can be analysed as biomarkers specific to regions of a genome. NGS also consists of several
710 techniques used today (Lemuth & Rupp, 2015). NGS genotyping technologies can often be an
711 alternative to the vastly more expensive and time consuming, whole genome strategies to look at a
712 study species. This is especially useful in studies involving species without publicly available genome
713 such as the wild relatives of *Linum* (Hu et al., 2021).

714 Peterson *et al* (2012) described one type of NGS genotyping technique as double-digest RAD
715 Sequencing or ddRADSeq. The protocol is a variation of the RAD sequencing protocol (Davey *et al*,
716 2010). DdRADSeq in comparison to RAD sequencing, uses a second restrictive digestion step to
717 improve the size selection step of the protocol (Peterson B. et al., 2012). This involves the use of a
718 second restriction enzyme and a second indexing step which allows for combinatorial indexing, more
719 specific to every individual (Peterson B. et al., 2012). When compared to a traditional RAD-
720 Sequencing technique, a double-digest RAD Sequencing can exclude regions which are flanked by a
721 very close or very distant recognition site. This then recovers libraries which consists of fragments
722 close to the specific target sequencing size (Peterson B. et al., 2012). The use of multiple enzymes
723 also contributes to a more diverse size selection availability in comparison to the traditional RAD
724 Sequencing technique (Peterson B. et al., 2012). In the study by Iguchi *et al* (2020), they were able to
725 identify diversity and selection mechanisms which leads to genetic variation in deep-sea amphipods

726 using a RAD sequencing protocol. This illustrates the significance of RAD sequencing for the use of
727 population diversity analysis, which is of interest when looking at our set of wild *L. bienne* samples.

728 ddRADSeq identifies Single Nucleotide Polymorphisms (SNPs) as genetic markers. SNPs
729 markers are simpler to genotype than microsatellites and are widely used in fine-scale population
730 genetic studies (Liu et al., 2019). The use of ddRADSeq have also been implied in herbarium samples
731 which have a relatively lower quality of preserved DNA, this protocol proves useful for obtaining
732 genetic datasets from a study species which is relatively fast to degrade such as plant leaf materials
733 (Jordon-Thaden et al., 2020). ddRADSeq have also been effective for genotyping organisms with a
734 larger genome size. A study using a non-model orchid species (genome size of 31.6Gbp) were viable
735 for ddRADSeq protocol, with inference of genetic diversity and differentiation (Gargiulo and Fay.,
736 2020). They also showed the requirement for a higher standard of quality (DNA extraction to library
737 preparation) when working with non-model organism of which their whole genome is not available
738 for reference (Gargiulo and Fay., 2020). This will prove as a challenge under the study model of this
739 thesis, as there is currently no whole genome publically available for the species *L. bienne*. This will
740 mean that any *de-novo* mapping of the extracted short sequences, will have to be of exceptionally
741 high quality. However, the use of *L. usitatissimum* whole genome is viable for reference to the later
742 mapping steps.

743 DdRADSeq uses indexing adaptors to “barcode” certain regions of a genome that have been
744 digested using two specific restriction enzymes. Subsequent bioinformatics analysis is then able to
745 highlight SNPs. The technique can be more cost effective than the previously developed SNP Chip
746 methods (Vieira et al., 2016). For any sequencing purposes, the first challenge is to extract a
747 sufficient amount of good quality DNA materials. There are multiple DNA extraction protocols
748 available to use. Nowadays, there are DNA extraction kits available to purchase such as Qiagen’s
749 Dneasy kits. For ddRADSeq purposes, the most used protocol is a CTAB DNA extraction protocol
750 (Jordon-Thaden et al., 2020). CTAB DNA extractions can also be modified to target for high
751 polysaccharides, which may be the case in plant material extractions (Porebski et al., 1997). For
752 ddRADSeq purposes, other studies have used between 0.1 – 0.8 µg of total DNA per sample (Jordon-
753 Thaden, et al., 2020). This relatively high amount of DNA may represent a challenge in the extraction
754 of *Linum* DNA materials. During this study, optimisation of ddRADSeq protocols for *Linum* will be
755 performed and DNA extraction protocols will be explored in the interest of extracting relatively high-
756 quality DNA required to build ddRADSeq libraries. To build ddRADSeq libraries, one of the most
757 important step is to select restriction enzymes which are specified for digestion at associated length
758 (250-500bp for this study). This would enable short reads over many individuals which are
759 subsequently barcoded. For the barcoding process, a ligation process can be applied to the digested

760 DNA materials. Adapters (single indexing) can be ligated to the DNA materials at both ends of the
761 restrictedly cut DNA materials. Libraries can then be pooled and sequenced.

762 When successful, ddRADSeq outputs forward and reverse reads of the library. Reads can
763 then be mapped either *de-novo* or to a reference genome before further analysis could take place
764 (Peterson B. et al., 2012). In the interest of illustrating population structure and diversification within
765 our wild and cultivated samples, population genetic analysis is essential. Population genetic analysis
766 is often useful for genotyping studies involving samples from a wide range of localities. Population
767 genetic studies often utilise mathematical models to evaluate how a population varies in terms of
768 their genetics (Servedio et al., 2014). As part of an evolutionary biology study, protocols into
769 population genetics may infer adaptation, speciation, and structuring across populations which may
770 infer biological conclusion such as local adaptation, inbreeding depression, and potential transfer of
771 genetic material from one population to another in the sample set (Hoban et al., 2016).

772 Often local adaptation can lead to population genetic structures, with population forming
773 specific niche to the local environments (Leimu & Fischer, 2008). In plants this can be an important
774 mechanism in which biotic interactions can often lead to responses to environmental changes such
775 as elevational changes (Grassein et al., 2014). For *Linum*, observations of population genetics for the
776 wild relatives are scarce compared to wild relatives of other species such as the wild crop relatives of
777 cabbage, *Brassica oleracea*. Population genetics studies in *Brassica oleracea* have revealed
778 population structuring along Western Europe for their wild relatives (Mittell et al., 2020). The
779 structuring can suggest diversity in wild relatives of cultivars, which represent a wider genepool for
780 the improvement of cultivar relatives. While this is explored in the *Brassicaceae*, it hasn't been
781 explored in the wild relatives of flax, *L. bienne*. In another example, population genetic analyses
782 suggests gene flow and hybridization in 12 plant species suggesting extinction due to introgression of
783 cultivar types to the wild (Ellstrand N. et al., 2003). This adds to the relevance of population genetic
784 analysis, especially in illustrating genetic relation in wild and cultivar relatives of plants. In this study,
785 initial population structure analysis will look to answer the question of "Can genetic distance infer
786 population structuring within the Western wild samples of *L. bienne* and its cultivar relatives?" Other
787 relationships in population genetics will also be investigated for evidence of genetic diversity within
788 our samples.

789

790 1.3.4 Genetic Expression in Flowering Time:

791 Within the model *Arabidopsis*, various environmental cues are known to affect the floral
792 regulatory network genes with light and temperature being the most significant (Li et al., 2015).

793 Timing of flowering in plants is often determined by environmental and endogeneous factors.
794 Environmental factors that have been shown to influence flowering time includes availability of
795 nutrients, ambient temperatures, drought, salinity, and the presence of pathogenic microbes (Cho L.
796 et al., 2017). Further, genetic expression of certain genes in the network affects other genes. As an
797 example, expression in genes such as *SOC1* have been shown to affect further genes in the pathway
798 such as *AP1* and *LFY*. There is an importance in quantitative input to each gene in the network for
799 expression in flowering times (Leal Valentim et al., 2015).

800 To look at specific gene expression within plants, specific genes can be targeted using
801 primers and real-time qPCR studies (Higuchi et al., 1992). Primers are relatively short stretches of
802 nucleic acid sequences which can uniquely identify a specific region of the whole genome (Cox and
803 Doudna, 2015). They complement and amplify specific nucleic acids using apolymerase enzyme.
804 Therefore, these primers are useful for studies looking at specific gene expressions.

805 With additional information, and access to sequencing technologies; this thesis will aim to
806 better understand flowering responses in *Linum* by generating genetic expression data. In the
807 studies performed in this thesis, both *L. usitatissimum* and *L. bienne* genetic materials were
808 collected, tested, and analysed regarding differences in species and population localities. It is of
809 interest to find out whether there are differences in genetic expression of flowering time related
810 genes in response to environmental factors such as temperature. How would these changes in
811 genetic expression measure up against population localities, and climate within those localities?
812 Within this thesis, we hypothesize that differences in relative gene expression are present in our
813 *Linum* collection and that this is correlated to population localities and climatic variables, reflective
814 of local adaptation.

815

816 1.3.5 Observation of Pollen Viabilities

817 In addition to observing variation in flowering time due to response to the environment, we
818 wanted to measure other adaptation in flowering for *Linum*. Since most studies already looked at
819 hetero/homo styly, we approached this using the viability or *Linum* pollen. In addition to flowering
820 time, we investigated if temperature affects the number of pollen, pollen tube formation and its
821 ability to germinate the ovule. Pollens are an important part of a plant's breeding strategy and act as
822 a male counterpart to the ovule. Plants often show differences in pollen response related to
823 differences in breeding systems of said plants (Hanley et al., 2008). In addition, climate has been
824 observed to affect pollen performance in plants (Jan Ejsmond et al., 2011; lovane M. et al., 2022).

825 Temperature acts as a major climatic factor which may limit the germination of pollen in plants. It is
826 deemed that temperature changes affects distribution of angiosperm species as the reproductive
827 stage is proven to be vulnerable to temperature. In the cultivated peach, it is observed that
828 increases of temperature to 20°C from their original 5.7°C in the field results in reduction in
829 receptivity of stigmas to pollens (Hedhly A. et al., 2005). They suggested that effects of temperature
830 on male and female organs of a flowering plants may be species-specific and could provide plants
831 with a level of plasticity to withstand environmental effects, such as ambient temperature, to make
832 sure they achieve fertilization (Hedhly A. et al., 2005). An older study in *Trifolium repense*, suggests
833 that growth of pollen tubes is affected by temperature and that pollen tube penetration into the
834 ovary is also observed to be negatively affected by colder and warmer temperatures (Chen and
835 Gibson, 1973). This effect in pollen viability due to temperatures could even be observed in plants
836 originating in warmer climates. In a species of groundnut (*Arachis hypogea*) it is observed that
837 warmer air temperatures (>35°C) results in failure of the setting of pods and pegs due to its lower
838 pollen viability. They found that for pollen germination and tube growth rate were less predictive in
839 discriminating genotypes for higher temperature than other parameters such as pollen tube length
840 and pollen germination rates (Kakani V. et al., 2002). This suggests plasticity in pollen viability
841 parameters for discriminating genotypes against temperatures in plants. In tomatoes (*Lycopersicon*
842 *esculentum*), it is suggested that temperatures affect *in-vitro* pollen germination and hybridization of
843 populations originating from different altitudes (Zamir D. et al., 1981; Maisonneuve and Den Nijs,
844 1984). Studies regarding pollen germination sensitivity to temperature stimulus can be achieved *in-*
845 *vivo* or *in-vitro*, with *in-vitro* procedures proving to be more elusive in plants (Boavida and
846 McCormick, 2007).

847 In several *Linum* species, it has been recently suggested that *in-vitro* germination is possible to
848 achieve up to 50% pollen germination rates (Lyakh and Soroka, 2021). However, they didn't include
849 the species *L. usitatissimum* or *L. bienne* within their study. Using the various sample populations
850 across western Europe that were in our collection, it is with interest to look at whether pollen
851 viability as pollen performance measure were affected under different treatments and whether
852 populations correlate with geographic and climatic variables, such as latitude, temperature, and
853 precipitation. We hypothesize that geographic and climatic variables affect pollen performance and
854 viability in our *Linum* collection and therefore its potentially affecting plasticity within population
855 fertilization strategy.

856 CHAPTER 2: FLOWERING TIME IN LINUM DUE TO VERNALIZATION AND
857 EXPRESSION OF FIVE FLOWERING TIME GENES.


858

859 In plants, it is widely accepted that environmental and climatic variables affect gene
860 expression changes which may in turn affect their performance as a response to changes in the
861 environment (Bigot et al., 2018; Elfving et al., 2011). Flowering initiation is of interest in population
862 survival as flowering is vital for reproduction and fitness in flowering plants. Flowering initiation also
863 have a major influence in seed dispersal and maternal effects (Galloway and Burgess, 2009;
864 Giménez-Benavides, Escudero and Iriondo, 2007). In species with agricultural significance, early
865 flowering types are actively selected for advantageous benefits such as late season drought
866 avoidance and better fruit development (Lotz L., 1990; Shavrukov et al., 2017; Ibrahim et al., 2018).
867 In the model *Arabidopsis thaliana*, environmental changes in photoperiod and temperature affects
868 induction of the FLOWERING LOCUS T (*FT*) gene in their leaves, affecting activation of downstream
869 floral meristem genes (Song et al., 2013). Expression of *FT* has been linked to internal and
870 environmental factors such as plant age, secretion of phytohormones (gibberellic acid) and ambient
871 temperature (Song et al., 2013). Furthermore, Lee et al suggests that the alteration of *FT* activities,
872 under different temperatures, regulate expression of downstream the floral meristem gene
873 SEPALLATA 3 (*SEP3*) (Lee et al., 2012).

874 Several studies have identified key flowering time genes that act as primary determinants of
875 the flowering-time network in the model *Arabidopsis thaliana* (Ballerini and Kramer., 2011; Welch S.
876 et al., 2004; Figure 1). Amongst these genes, the most rigorously studied are, the floral pathway
877 integrators such as FLOWERING LOCUS T (*FT*) and the SUPPRESSOR OF OVEREXPRESSION OF
878 CONSTANS (*SOC*), with FLOWERING LOCUS C (*FLC*) also studied (Sasaki et al., 2017; Li et al., 2015;
879 Valentim et al., 2015). In the model *Arabidopsis*, it has been illustrated that, *FT*, and *SOC* loci
880 expression activates downstream flowering meristem identity genes such as LEAFY(*LFY*),
881 APETALA1(*AP1*), SEPALLATA3 (*SEP3*) and FRUITFULL(*FUL*). This is because both *FT* and *SOC* are
882 considered transcription factor loci, and in the *Arabidopsis* flowering network, they are considered
883 as floral integrators (Welch S. et al., 2004; Lee and Lee, 2010). Furthermore, Ballerini et al (2011)
884 suggested that there is an element of conservation of *FT* homologs within flowering plants. Although
885 this is the case, they suggest that there is complexity in their regulation and evolution that is still of
886 interest in different angiosperm species to this day (Ballerini and Kramer., 2011). This suggests the
887 importance of *FT* as a floral regulator and the potential expression variation it may have in different
888 species, potentially due to environmental stimulus. In addition, there are two pathways that are

889 worth looking at in response to light and temperature regulation of *FT*. These are the Photoperiod
 890 and Vernalization pathway. The expression of photoperiod genes *CONSTANS* (*CO*) and *GIGANTEA*
 891 (*GI*) can activate *FT* due to photoperiodicity (Kurokura et al., 2017; Song Y. et al., 2014) and the
 892 vernalization gene *FRIGIDA* (*FRI*) is responsible in determining variation of flowering time due to
 893 vernalization requirements, defined as the requirement for seeds to sense cold to flower faster
 894 (Shindo et al., 2005). Finally, the downstream meristem genes function as an initiator for floral organ
 895 developments. Upregulation in these flowering meristem genes signals groups of meristem cells to
 896 develop into flowering cells, instead of shoot cells (Teper-Bamnolker and Samach, 2005), thus
 897 resulting in floral intitation after complex regulations of several genes in response to the
 898 environment.

899



900 Figure 1. An overview of the flowering time network in *Arabidopsis thaliana* adapted from the literature (Sharma N. et al.,
 901 2020; Leitjen W. et al., 2018; Chen et al., 2018; Ballerini and Kramer., 2011; Welch S. et al., 2004) Underlined are the genes
 902 tested. We also highlighted pathways that affects *FT* (pathways in brown).

903

904 Comparative genomics have also been used to model flowering time pathways in other
 905 species such as the temperate grasses (*Brachypodium distachyon*) (Higgins et al., 2010). They
 906 suggest that flowering time pathways in *Brachypodium distachyon* are highly like the model
 907 *Arabidopsis*. Reviews from Leitjen et al (2018) suggests that despite the overall conservation of these
 908 flowering time network genes, there is evidence of divergence of flowering time regulation in both

909 the model *Arabidopsis thaliana* and several crop species due to environmental factors such as
910 temperature. With this, it can be assumed that different plant populations express varied responses
911 to flowering due to the environment. To what extent and which direction each gene is affecting each
912 plant population is still of research interest to this day, especially in non-model, cultivated plants. As
913 an addition, this variation due to flowering responses doesn't just occur in the model *Arabidopsis*.
914 Flowering initiation in non-model organisms such as the common sunflower (*Helianthus annuus*),
915 have been suggested to be "exploited" by natural selection through genetic expression and tissue-
916 specific expressions of the *SOC1* genes affected by environmental factors. One of these factors is
917 photoperiodicity. Sunflower (*Helianthus annuus*) has shown diversity in photoperiodic responses,
918 due to changes in genetic expression modulated through natural selection (Blackman, et al., 2011).
919 Blackman et al, 2011, also suggests the occurrence of convergent evolution of photoperiod
920 responses in cultivated and wild types of sunflowers, which suggests phenotypic plasticity is weakly
921 constrained by genetic regulation (Blackman et al., 2011). In Japanese wild radish (*Raphanus*
922 *sativus*), recent research suggest that northern accessions were more sensitive to exposure to
923 prolonged cold conditions, in the sense that they require colder conditions to flower. This was
924 suggested by FLC expression, which is part of the vernalization pathway (see below). The Southern
925 wild radish population seem to be more sensitive to photoperiodicity, suggested by the expression
926 of photoperiodic genes (Han Q. et al., 2021). This is observed to correlate with multiple flowering
927 time gene expressions (Han Q, et al., 2021).

928 Vernalization is an adaptation, often found in temperate plants, which sense cold to allow
929 optimal timing of flowering initiation in spring (Kim *et al.*, 2009). Vernalization is the requirement for
930 seeds to be cooled from germination for individuals to flower earlier. In the model *Arabidopsis*
931 vernalization is widely studied. The vernalization network has been shown to progress around the
932 MADS-domain protein complex comprising of several flowering time genes forming complexes with
933 the MADS-protein SVP. One of these genes is the *FLC* (Flowering Locus C) which acts as a repressor
934 of *SOC1* (Chen et al., 2018). The closely related *FLM* (Flowering Locus M) also have been shown to
935 form complexes with the MADS-protein SVP. These proteins have shown to mediate response to
936 changes in ambient temperature by forming a complex with more specifically the β form of *FLM* at
937 lower temperatures. This in turn represses flowering at colder temperatures (Lee J. et al., 2013;
938 Chen et al., 2018; Posé et al., 2013). At relatively higher temperature, the MADS-protein SVP
939 degrades and the complex with *FLM- β* is not formed, thus producing reduced repressive complex,
940 allowing flowering initiation to proceed. Several vernalization related loci play an important part in
941 vernalization sensitivity and interact with other pathways leading to flowering initiation (Blázquez,
942 Koornneef and Putterill, 2001) (Chandler, Wilson and Dean, 1996). In this chapter, we will investigate

943 natural variation in flowering time responses for the temperate plant genus *Linum*. Both wild (*L.*
944 *bienne*) and cultivar types (*L. usitatissimum*) will be observed in terms of their phenotypic and
945 genetic responses to flowering initiation due to temperature changes in relation to vernalization.
946 This will imply potential adaptation in vernalization responses for both wild and cultivar types. The
947 main environmental variable tested will be temperature, as this corresponds more towards
948 vernalization requirements. However, expression of flowering time genes that may regulate
949 photoperiodicity were also observed.

950 Studies of *Arabidopsis thaliana* have also established that the latitude differences in
951 populations is associated with co-variation in growth and flowering time (Debieu et al., 2013a). In
952 respect of the wide distribution of *L. bienne* (as discussed in chapter 1), it is of interest to look at
953 differences in genetic expression from a flowering time perspective, based on phenotypic and
954 genotypic results. In *Linum*, vernalization requirement has been little studied, with studies mainly
955 focused on the cultivar, *L. usitatissimum* (Darapuneni et al., 2014a). The wild relative (*L. bienne*) is of
956 interest, especially for questions regarding climates, as the wild population are likely to be locally
957 adapted to their environments (Landoni et al., 2022). This adaptation can be observed as clines of
958 flowering initiation times within the wild species. Several genes are of interest when looking at the
959 development of *Linum* against latitude which are part of the flowering genes network. Gutaker et al.
960 have previously identified the alteration and expression of the flowering gene *LuTFL1*, which could
961 reflect latitudinal adaptation and crop selection for fibre production in northern populations
962 (Gutaker et al., 2019). However, other flowering time genes were not explored in this case. Other
963 flowering time genes could involve genes that are in the photoperiodicity network. Several flowering
964 time genes in cultivated *Linum* such as *LuCO* (CONSTANS) and *LuGI* (GIGANTEA) were previously
965 explored by Sun et al (2019) where they found variation in the expression of these genes under
966 several experiments with long days (22 °C/16 °C (12 h/12 h) for a 16 h/8 h photoperiod at a light
967 intensity of 300 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) and short days (Similar to long days treatment but with
968 10 h/14 h photoperiod instead) which revolves around photoperiodicity (Sun et al., 2019).

969 Under temperate wild *Arabidopsis* accessions, it has been recently observed that different
970 expression of genes of interest in this thesis correlates with the flowering time of the temperate
971 accessions (Kinmonth-Schultz et al., 2021). The main aim for this chapter is to find out whether
972 expression of important flowering time genes, such as Flowering Locus C (FLC) and Constans (CO)
973 influence the flowering initiation of different, vernalized, and non-vernalized *Linum* populations
974 reflecting varying environmental conditions across the range. The hypothesis was that flowering
975 initiation requirements based on vernalization differ within the *Linum* collection dependent on
976 source location and type (wild or cultivated); and that different flowering behaviour is supported by

977 differences in genetic expression of flowering time genes. During this experiment, vernalized and
978 non-vernalized individuals were examined to explore phenotypic and genotypic differences both
979 between species and within species. We asked: “How do flowering time responses to vernalization
980 differ across wild populations in western Europe and are there differences between wild and
981 cultivars?”.

982 Differences in vernalization requirement within different plant species have been observed
983 to subsequently affect their initial flowering time, this can be observed in species that requires both
984 long or short days to flower (Adhikari, Buirchell and Sweetingham, 2012) (Ream et al., 2014). These
985 flowering time interactions have been related to source population latitude, climate, and species. To
986 look for climatic variables, The R package BioClim available at,
987 <https://rdr.io/github/jjvanderwal/climates/man/bioclim2.html>. Various climatic parameters can be
988 used to summarize climate using PCA (principal component analysis) to extract major climatic factors
989 that may contribute to any differences in vernalization requirement observed in this experiment
990 involving relative gene expression. In the wild, it is important that flowering initiation responds
991 appropriately to local climate. Regarding vernalization, the expectation is that the more northern a
992 plant population is localized, the more dependent it is on vernalization cues such as colder
993 temperatures and wind speed. Thus, a colder climate of origin will be associated with greater
994 relative gene expression differences when grown at different temperatures.

995 There is an expectation that selection on flowering time genes is stronger in the wild, leading
996 to different vernalization responses in different latitudinal range. This implies interests in genetically
997 quantifying variation in vernalization requirements both between and within the two *Linum* species.
998 The expectation is for there to be genetic differences between the flowering time genes tested here
999 when comparing cultivars with the wild samples. These differences are then expected to be
1000 attributed to environmental variables such as local climate.


1001

1002

1003 2.2 STUDY AND METHODS

1004 2.2.1 Samples and Experiment

1005 We set out to investigate relationship between vernalization and genetic expression of
1006 flowering time genes using the following workflow:




1007 Figure 2. A workflow summary of the study methods in this chapter from the glasshouse to analyses in
 1008 computing facilities.

1009


1010 To measure phenotypic and genetic differences between representative *Linum* populations;
 1011 controlled growth of plant materials was conducted. Samples for this experiment were collected as
 1012 described in sections “1.2.1 *Linum usitatissimum*: Cultivated flax” and 1.2.2 “*Linum bienne*: Wild
 1013 Flax” of this thesis. Samples were collected from across Portugal, Spain, France, and the UK (Fig 3). A
 1014 full list of the individuals used is available under appendix 2.

1015

A:



B:



1016 Figure 3. A map of the origin of collection of cultivars (A) and wild (B) *Linum*. Each population is alphabetically ranked by
1017 their latitude, with the most Southern population ranked A. For cultivars, since latitude was only recorded by country level,
1018 numerous populations belonging to the same latitude are depicted per latitude level which are alphabetically ranked.

1019

1020 Out of the total of 47 populations, 28 were wild types (*L. bienne*) and 18 were cultivars (*L.*
1021 *usitatissimum*). Wild seeds were initially grown in glasshouses available at the Department of
1022 Biosciences in Durham University. The controlled glasshouse had 16:8 hours daylight to no light ratio
1023 and the minimum temperature was measured at 13°C during winter seasons, with a maximum
1024 temperature measured at 28°C during summer seasons. S1 Seeds from these were utilized for
1025 vernalization experiments. A collection of 473 individuals over three experimental designs were
1026 studied. The vernalization experiments were conducted twice in the duration of this study (2018-
1027 2019 and 2020-2021). During both time frames, two controlled conditions (vernalized and non-
1028 vernalized) were specified using controlled growth chambers. Out of 473 individuals, 62 individuals
1029 were vernalized in 2020-2021; 257 individuals were vernalized in 2018-2019; and 154 individuals
1030 were non-vernalized in 2018-2019. The experiments in 2018-2019 were replicated (2 vernalized and
1031 2 non-vernalized chambers). In comparison, there was only one vernalization chamber and one non-
1032 vernalized chamber under the 2020-2021 experiment, albeit the same conditions were applied
1033 between the two experimental time frames. For the 2020-2021 experiments, samples were gathered

1034 from a collection of S1 and S2 seeds previously grown under experimental conditions based on no
1035 vernalization or under controlled glasshouses available in the department of Bioscience of Durham
1036 University. In total, there were 47 germinating populations from both vernalized and non-vernalized
1037 experiments. After 40 days of treatment, plants were morphologically examined by their flowering
1038 time, overall height, and number of stems when first flowering. Morphological measures between
1039 these traits will be examined in chapter 5 of this thesis. The vernalization conditions (see below)
1040 were setup in two Weiss Gallenkamp growth chambers model numbers A3655 and A3658. The
1041 conditions were kept the same throughout the experiment. The number of days to flower (flower
1042 initiation) was a point of phenotypic interest and genetic expression (RNA materials) as a point of
1043 interest for gene expression studies. Therefore, collection of RNA materials also took place at the
1044 corresponding first flowering of every individual.


1045 Preliminary measures for the S0 population grown in the controlled glasshouse were made
1046 before sowing of S1 generation for measures regarding vernalization which are plant height, stem
1047 numbers and bud numbers. The plants were exposed to vernalization conditions as follows; after
1048 sowing, *4°C (0-hour lights) for 72 hours, 22°C for 10 days with 16:8 h light ratio, and 4°C for 40 days*
1049 *with 16:8 h light ratio*. After vernalization plants were kept at 24°C to 16:8 h light ratio indefinitely.
1050 The non-vernalized conditions were as follows; after sowing, *4°C (0-hour lights) for 72 hours, 22°C for*
1051 *10 days with 16:8 light ratio, 24°C for 40 days with 16:8 light ratio indefinitely*. Note the difference
1052 between the vernalization and the non-vernalized being the *4°C for 40 days cold treatment*. The
1053 lightings consist of numerous fluorescent tube lighting units, which were “Philips Master TL5 HE”.
1054 These light tubes are 14 Watts in power requirements for each tube and emits 4000 Kelvin light
1055 temperature in colour by specifications. Plants from these S1 generations under vernalization
1056 treatments were also collected for RNA materials, for the genetic analysis purposes of this study. The
1057 phenotypic traits measured were divided into two categories: vegetative (stem number and plant
1058 heights) and reproductive (first day to achieve first flowering (flowering initiation) and seed size),
1059 with particular interest in reproductive measures, due to the interest in flowering initiation. The
1060 phenotypic measures were measured at the time of first flowering of every individual (± 1 day).

1061 Leaf materials were collected 14 weeks after sowing, and no more than three days after
1062 vernalization treatments have occurred. Leaf samples were collected from each treatment during a
1063 2-day window period from 11am to 1pm to avoid expression differences due to time of collection.
1064 Ten to twelve leaves from the top 2cm of the longest shoot on each plant were collected inside a
1065 1.5ml Eppendorf tube for each sample and were each labelled, and flash frozen in liquid N² before
1066 storing in the -80°C freezer. Two vernalization experiments were conducted during this study under
1067 different years (2019/20 and 2020/21). However due to restrictions on caused by the Covid-19

1068 pandemic, the 2020/21 vernalization experiment was cut short. Due to this, differences between the
1069 two experiments were unavoidable. These differences were quantified by analysing comparative
1070 analysis of the two experiments. This can be found under the “Results” part of this chapter.

1071 When considering locality of a wild population, its environment is the climate of the local
1072 area in terms of longitude, latitude, and altitude of the places where these wild populations were
1073 collected. Local climate was analysed based on different climate variables such as precipitation,
1074 temperatures, and sunlight availability that were extracted from over a 30-year period. The data was
1075 retrieved from WorldClim database. Data were available for precipitation (mm), solar radiation
1076 (kJ/day*m²), average temperature (°C), minimum and maximum temperature (°C), vapour pressure
1077 (kPa), and wind speed (m/s). Climate variables for principle component analysis (PCA) were used at
1078 30arcsec resolution over seasonal months (June July August for summer, September October
1079 November for autumn, December January February for winter and March April May for spring). To
1080 include all 6 dimensions of climatic variables, they were loaded into a principle component analysis
1081 with a principle component one (PC1) level explaining 63.8% of climatic variables (data available in
1082 appendix 3). Loaded value also suggests that lower PC1 values are associated with colder
1083 temperature associated with more Northern latitude. These data were used in a linked study of
1084 flowering time study (Landoni B. et al., 2022). All the climate variables were loaded in a collective
1085 PCA, whereby climatic PC1 values for each population were processed as the final representative
1086 values for all climatic variables. This would make further downstream analysis, regarding climate
1087 variables easier to process.

1088 When plotting PC1 and latitude of origin for each of our population, there is a visible
1089 relationship between the two variables, suggesting that lower PC1 values reflects higher latitudes.
1090 This can be seen in the plot below:



1091 Figure 4. Scatterplot to show negative relationship between climate variable representative (PC1) and latitude.
1092 Lower PC1 values represent lower temperatures (°C) and solar radiation (kJ/day*m²).

1093 2.2.2 Relative Gene Expression: RNA Extractions

1094 Real-time quantitative PCR (rt-qPCR) is well suited for relative gene expression
1095 quantification (Livak and Schmittgen, 2001). For this, RNA is extracted and then converted back into
1096 copy DNA (cDNA) by using reverse transcriptase. This was done to an individual level, with no
1097 bulking over population. cDNA products were then subjected to polymerase chain reactions by
1098 designing primers specifically for loci of interest. The amount of cDNA target is quantified using dye
1099 markers that fluoresce when they bind DNA PCR product and comparing real-time amplicons with
1100 those of housekeeping genes from the same sample. The use of real-time polymerase chain reaction
1101 is a routine tool in molecular biology for the study of gene expression.

1102 Leaf tissues from vernalization experiments (as mentioned under section 2.2.1 “Samples and
1103 Experiment”) were first ground using a homogeniser. Promega’s ReliaPrep RNA Tissue Miniprep
1104 System was used to extract RNA and solutions were prepared using manufacturer’s instructions.
1105 After solutions were prepared, LBA+TG buffer were added into previously ground samples and were
1106 mixed using micro-pestles and tissue homogeniser until thoroughly mixed. Materials were lysated
1107 and processed following manufacturer’s protocol. After the centrifugation steps, RNA was washed
1108 according to manufacturer’s protocol with an additional second wash. Further centrifugation steps
1109 took place at 14,000g for 2 minutes after the second wash.

1110 After collection, samples were ready for elution. Samples were eluted according to the
1111 manufacturer’s instructions. Centrifugation steps were repeated using the same elutant to make
1112 sure all the available RNA were eluted out of the membrane. 1µL of elutant from each sample tube
1113 were used for RNA measurement under a Nanodrop ND1000. When satisfied with measurements,
1114 samples were stored in -80°C freezer to preserve RNA integrity. Satisfactory measures were decided
1115 at >20ng/µL of RNA with 260:230 ratio of >1.20 and 260:280 ratio of >1.60.

1116

1117 2.2.3 Relative Gene Expression: cDNA synthesis:

1118 For cDNA synthesis, Applied Biosystem’s High-Capacity cDNA Reverse Transcription Kit was
1119 used. For control, no template control (NTC) and no reverse transcriptase (NRT) samples were
1120 added. Each sample reaction was made to the same amount of RNA template in a total of 10µL
1121 volume. The amount of RNA template needed for the cDNA synthesis was calculated by
1122 standardizing all samples to the sample with the lowest amount of detectable RNA. This was done by
1123 calculating how much RNA could be extracted for a 10µL eluted RNA additive (to the mastermix),
1124 based on measures of RNA by spectrometry in ng/µL for specific samples. The maximum amount of
1125 RNA for the cDNA synthesis protocol was 2µg in 10µL final volume. Samples which had more

1126 detectable RNA were diluted down to the same concentration as the lowest concentrated sample.
 1127 This dilution was done using nuclease-free water accordingly. For the no template control, RNA was
 1128 replaced with nuclease free water. 10µL was added per NTC. For the NRT, 5µL of random RNA
 1129 sample with 6µL of nuclease free water were added.

1130 For the rest of the reactants, a master-mix were prepared in an Eppendorf tube per
 1131 manufacturer’s instructions. Master-mixes were vortexed for 10 seconds. After the addition of the
 1132 master-mix ,1µL of Multiscribe Reverse Transcriptase were added into each sample and the NTC
 1133 control. Samples were then centrifuged briefly and vortexed subsequently. After the addition of
 1134 both master mix and Transcriptase, a thermocycling step can take place. The thermocycling steps for
 1135 cDNA synthesis were as follows:

| | Step 1 | Step 2 | Step 3 | Step 4 |
|--------------|------------|-------------|-----------|--------|
| Temperature | 25°C | 37°C | 85°C | 4°C |
| Time Elapsed | 10 minutes | 120 minutes | 5 seconds | ∞ |

1136 Table 1. Thermocycling steps for cDNA synthesis.

1137 After the thermocycling steps were finished, samples were stored in a 4°C fridge for short
 1138 term (<72 Hours) and a -21°C freezer for long term storage(>72 Hours). The final Ct-value from the
 1139 later real time qPCR protocol would serve as quality control. If quality of RNA from the extractions
 1140 are satisfactory there was no need to measure cDNA output (Schmitz and Amasino, 2007).

1141

1142 2.2.4 Expression of Genes of interest and House Keeping genes:

1143 Housekeeping genes (HKGs) are required for maintenance of cells (Butte, Dzau and Glueck,
 1144 2001). These genes are usually expressed relatively constantly under different conditions, which
 1145 makes them suitable for comparison against the expression of other genes. Generally, HKGs
 1146 constitute an important component for rt-qPCR procedures. However, a study suggested that
 1147 expression of HKGs may change with cellular density of samples and that they could be affected by
 1148 experimental conditions (Greer *et al.*, 2010a). Therefore, use of multiple HKGs within the same
 1149 experiment can guard against these effects. *GAPDH* (Glyceraldehyde-3-phosphate dehydrogenase)
 1150 and *UBI* (Ubiquitin) represent a couple of the commonly used HKGs in plants and therefore used for
 1151 assessing different abiotic stress (Liang *et al.*, 2018a) (Carmona *et al.*, 2017). For *Linum*, the HKGs
 1152 which will be used for this experiment are *LuGAPDH* and *LuUBI2*. This reference genes for *Linum*
 1153 were developed by Huis *et al.* and are specifically useful for real time PCR (qRT-PCR) protocols (Huis,
 1154 Hawkins and Neutelings, 2010). Our study will be incorporating these HKGs, as were also found on
 1155 the study investigating photoperiodicity in *Linum* by Sun *et al* (Sun *et al.*, 2019).

1156 We hypothesized that there would be variation in gene expression for flowering time genes
1157 according to latitude and due to vernalization. We tested flowering time genes already studied in
1158 *Linum usitatissimum* by Sun et al (Sun *et al.*, 2019). In our study however, we will be testing the gene
1159 expression based on vernalization stimulus and will be controlling for other environmental stimulus
1160 such as light availability. Whilst the study by Sun *et al* (2019) had found significant differences in
1161 expression of flowering time genes for *L. usitatissimum* under different light-length treatments, they
1162 have not tested vernalization as an environmental factor. In addition to this, they have not tested
1163 the wild relatives. It would be of interest to look at the expression of floral integrator gene, *LuFT*,
1164 along with the other photoperiodicity genes when differences in vernalization treatments is applied.
1165 This is because *Linum* is a temperate plant found across the sub-temperate into the temperate
1166 regions, thus different population may have adapted differently to vernalization as a stimulus to
1167 flowering initiation.

1168 In addition, the lineage leading to *L. usitatissimum* and *L. bienne* as a genus have been
1169 understood to have undergone a polyploidy event 20-40mya (Sveinsson *et al.*, 2014a). This has
1170 repercussions as there potentially are multiple copies (paralogues) of the same genes within *Linum*.
1171 This would mean there may be a need to test multiple copies of the same gene to see an effect
1172 which could be from one or multiple set of these copies. Wild and cultivars were tested for seven
1173 paralogue-specific flowering time genes using a modified real-time qPCR protocol. These genes were
1174 *LuFT1*(*Flowering Locus T 1*), *LuFT2*(*Flowering Locus T 2*), *LuGI1.1* (*GIGANTEA 1.1*), *LuGI1.2* (*GIGANTEA*
1175 *1.2*), *LuGI2* (*GIGANTEA 2*), *LuCO1*(*CONSTANS 1*), and *LuCO2* (*CONSTANS2*). A full list of nucleobases
1176 codes used for each primer can be found in appendix 4 (Sun *et al.*, 2019).

1177

1178 2.2.5 Primer Testing

1179 The primers for the genes of interest were initially tested by performing PCR and analysing
1180 the products on an agarose gel (Jarman, Ward, and Elliott, 2002). Primers was tested using
1181 Promega's Go-Taq green master-mix; 2.5µl of each Forward and Reverse primers along with 2.5µl of
1182 *Linum* cDNA template for a 25µl total reaction volume. This would mean that the concentration of
1183 each primer is 1µM with < 250ng of cDNA template. We added 12.5µl of the Go-Taq green
1184 mastermix for a 1x concentration solution. The rest of the solution is made up with nuclease-free
1185 water up to 25µl. PCR mixes were treated in a thermal cycler using the following programme:
1186 Denaturation at 95°C for 3 minutes; Annealing with 35 cycles of 95°C for 45 seconds, 55°C for 30
1187 seconds, 72°C for 1 minute; and finally, an extension stage at 73°C for 5 minutes. Products were
1188 examined on a 2% TAE gel for PCR products. Products at <200bps with minimal amount of smearing

1189 observed were deemed acceptable. A decision was made to not use LuGI1.2 and LuGI2 primers
1190 based on inconsistencies with the gel such as smearing, and product sizes observed as > 200bps,
1191 leaving 5 primers (*LuFT1*, *LuFT2*, *LuCO1*, *LuCO2*, and *LuGI1.1* with the 2 housekeeping genes
1192 (*LuGAPDH* and *LuUBI2*).

1193

1194 2.2.6 Quantitative Real-Time Procedure

1195 For real-time quantitative PCR reactions, an Applied Biosystem model 7300 real-time
1196 thermal-cycler was used with StarLab's 96-Well PCR Plates (96-Well PCR Plate, Skirted, Low Profile,
1197 White – STARLAB, 2022).

1198 Diluted cDNA was prepared for each sample. Primers were ordered from Integrated DNA
1199 Technologies (IDT). Each primer was aliquoted and diluted to 10 μ M working solution. To perform a
1200 real time qPCR, GoTaq qPCR SYBR Green were used (Promega Corporation). A master mix of the
1201 reaction components were then prepared as follows:

| Component | Volume / reaction (μ l) | Final Concentration |
|----------------------------|------------------------------|---------------------|
| SYBR Green mix | 7.5 | 75% |
| Forward-Primer | 0.9 | 9% |
| Reverse-Primer | 0.9 | 9% |
| Nuclease free water | 0.7 | 7% |

1205 *Table 2. The components required for the real-time qPCR reaction.*

1206

1207 SYBR Green Mix (included in the Promega qPCR GoTaq RT-qPCR system), and Nuclease free water
1208 were mixed by considering three repetitions per sample and 2 controls (No reverse transcriptase
1209 (NRT) and No template control (NTC)). For 14 samples we had $(3 \times (14+2)) = 48 \times 2 = 96$. As there are
1210 possibilities for pipetting errors, 100 samples were considered for the SYBR and the water. SYBR and
1211 water were mixed in a 1.5ml Eppendorf tube and vortexed at 2000RPM for 30 seconds.

1212 The small volume of primers was often easier to be dispensed onto the sides of each well,
1213 this way loaded wells can be marked by the presence of the primer on the side to avoid
1214 contamination by other primers. After the primers were fully loaded, the plate was centrifuged
1215 down for 1 minute at 3000RPM at room temperature using the F2096 rotor in an Allegra X22
1216 refrigerated bench centrifuge. Making sure all the liquid has reached the bottom of the well, 8.2 μ L of
1217 the SYBR Green mix were loaded in with 5 μ L of each respective diluted cDNA samples following.
1218 Plates were sealed using an appropriate 96-well plate plastic sealer (we used Starlab's Self-Adhesive
1219 sealing films), making sure that each well was tightly secured with the sealer by pressing on each

1220 well after sealing. Another centrifugation step for 1 minute at 3000RPM were applied to the plate
1221 and samples were ready to be thermal cycled as follows:

| Stage | Temperature (°C) | Time (s) | Number of repeats |
|-------------------|------------------|----------|-------------------|
| Holding | 50 | 120 | 1 |
| Enzyme activation | 95 | 600 | 1 |
| Cycling I | 95 | 15 | 40 |
| Cycling II | 60 | 60 | 40 |

1222 Table 3. The cycling steps for the real-time qPCR.

1223
1224 SYBR green (Promega) was used to quantitatively assess amplified PCR product. SYBR green dye
1225 fluoresces at 497-520nm blue to green light when binding to double stranded DNA (ds-DNA). The
1226 SYBR dye fluorescence intensity can be used to quantify how much of each gene were amplified
1227 using specific primers (Zipper *et al.*, 2004). The ds-DNA will increase by each thermal cycle thus
1228 increasing their binding to SYBR until the amount of DNA material reaches the cycle-threshold (Ct)
1229 values which is the number of cycles it takes for the dye to be distinguishable to the background as it
1230 binds to the DNA material (Zipper *et al.*, 2004).

1231 The Ct-value reads were collected at end of stage cycling II, whereby the real-time machine
1232 measures the cycling threshold values against amplifications of the targeted cDNA expressed in the
1233 sample. After this, Ct-values were further analysed as described under following section 2.2.10
1234 “Relative Gene Expression: Quantification”.

1235 1236 2.2.7 Primer Efficiency

1237 A standard curve calculation was done for the Ct-values of the different dilutions to define
1238 efficiency of primers whereby the primer efficiency is tested by calculating the slope against the
1239 concentration of tested primers (Pfaffl, 2001). This test is essential, because calculation of relative
1240 gene expression is based on the delta-delta (difference) of the Ct-values between the HKGs and the
1241 genes of interest. The difference in Ct-values is influenced by how efficient a PCR product can react
1242 with the SYBR dye.

1243 Primer efficiency was tested by calculating the average Ct-values for runs of a sample that
1244 has been serially diluted and calculating the coefficient of Determination (R^2) values across sample
1245 dilutions. In theory, the more dilute the samples are, the more slowly the primers are going to
1246 amplify, and this will form a standard curve for fitting a model and performing R^2 calculation (Glantz,
1247 Slinker and Neilands, n.d.) using:

1249
$$R - squared = \frac{Sum\ of\ squares\ due\ to\ regression}{Total\ sum\ of\ squares}$$

1250

1251 In this case the dilutions of primer tested were as follows; undiluted, 1/20, 1/200, 1/2000.
1252 After applying *Log* to the sample quantity, the coefficient of determination (slope) was calculated.
1253 The efficiency (%) was then calculated using the following equation described by Ginzinger
1254 (Ginzinger D, 2002):

1255
$$Efficiency\ (\%) = \left(10^{\frac{-1}{Slope}} - 1 \right) \times 100$$

1256

1257 Efficiency values from the above equation are represented in percentage. However, for the
1258 relative gene expression calculations, the percentage values were converted into decimal values
1259 whereby a value of 2.00 will indicate 100% efficiency and 1.00 will indicate 0% efficiency. This
1260 conversion is essential for the final efficiency value input calculation in relation to ΔCt of each gene
1261 further downstream. Efficiency values were converted from % using the following equation
1262 (Ginzinger D, 2002):

1263
$$Efficiency\ Value = \frac{Efficiency\ \%}{100} + 1$$

1264 Three primer efficiency tests were performed using real-time qPCR procedures for each
1265 respective primer but only the third test values used further for down-stream calculations as these
1266 seemed the best results from several tests, based on the efficiency (%) values closer to 90-110%. The
1267 results from the primer tests are noted in appendix 5.

1268 These primer efficiency values represent values which were inputted as a correction when
1269 calculating relative gene expression (RGE) of each gene tested. The efficiency value was calculated
1270 for each of the tested primers as well as the tested HKGs according to the separate ΔCt (avg Ct
1271 values – actual) for each gene and for each separate real time run (i.e repeats). The final Primer
1272 efficiency correction were calculated using this formula:

1273
$$Final\ efficiency\ input = Efficiency\ value^{\Delta Ct}\ for\ each\ gene$$

1274 The measured efficiencies values were not optimal as efficiencies of some primers are <95% as
1275 recommended (Miranda and Steward, 2017). Ideally these primers would be re-designed and re-
1276 ordered, but with the RNA materials in some cases already extracted and time limit considerations, a
1277 decision was made to lower the threshold efficiency to 85% to include all the tested primers were
1278 above the threshold.

1279

1280 2.2.8 Real time qPCR of Vernalised and Non-vernalised Samples

1281 Regarding the real time qPCR experiment described above, a set of samples that had
1282 experienced both vernalization treatments were chosen to compare flowering time gene expression
1283 differences. The final set of samples included 14 wild and 14 cultivar pairs each with samples from
1284 the non-vernalization and vernalization experiment. Some additional samples could not be utilized
1285 due to low quality RNA. Subsequently, cDNA from 2 samples at a time was added to 96 well plates
1286 designed to test the 5 Flowering time genes along with the 2 HKGs.

1287 Primer-sample combinations tested totalled up to 14 per plate with 3 repetitions each. The
1288 primer sequences are available under appendix 4. These, with the addition of No Reverse
1289 Transcriptase (NRT) and No Template Control (NTC) made up a 96-well plate.

1290

1291 2.2.9 Clean-ups and quality controls:

1292 As a pre-cursor to calculating relative gene expression (RGE), the Ct-values of the
1293 Housekeeping (HK) genes were checked for quality control. This was of importance because the
1294 observed primer-efficiency values were not ideal (87.273% for *LuGAPDH* and 51.464% for *LuUBI2*).
1295 Individual Ct values were removed if either; Ct-values were either too high (>29 cycle-threshold
1296 values) or too far away from the other values (i.e >1 Ct variation between the triplicates) for
1297 *LuGADHP*; 51/144 Ct-values were either too far away from other Ct-values or too high. These steps
1298 addressed potential experimental errors such as poor qualities and low quantities of cDNA and
1299 pipetting errors, which will impact the real-time qPCR quantification process.

1300 When applying the delta-delta Ct ($2^{-\Delta\Delta Ct}$) method to the final relative gene expression values,
1301 all initial samples were included, except for one wild sample (Saf_10) for both Flowering Time genes.
1302 This sample was excluded because of the much bigger relative gene expression values that make it
1303 an extreme outlier. This could be due to experimental errors such as the quality of RNA.

1304

1305 2.2.10 Relative Gene Expression: Quantification

1306 RGE data were processed in Microsoft Excel. For quantification purposes, a template was
1307 built for primer efficiency using formula stated under section 2.2.7 “Primer Efficiency” part of this
1308 chapter. In addition to these, further calculations are required to calculate relative gene expression.
1309 The Ct results for each gene of interest (GOI) was added to a Microsoft Excel template and are
1310 specified to the same House Keeping Genes (*GAPDH*) Ct-values for each respective samples. To

1311 calculate relative gene expression, a difference (delta) of Ct values were calculated between a
1312 reference house-keeping gene and the gene of interest based on previous method described by
1313 *Livak et al* (Livak and Schmittgen, 2001a). There are two methods which are available to explore
1314 using one HKG or using two HKGs. Both are described below.

1315 The default formula for calculating relative gene expression differences is the delta-delta Ct
1316 ($2^{-\Delta\Delta Ct}$) method. This method uses only one HKG. This is where, for each gene and sample, average
1317 Ct-values for the gene of interest (GOI) and the housekeeping gene (HKG) are calculated. Differences
1318 (delta-Ct) between the gene of interest and the housekeeping gene Ct-values were calculated and
1319 then the delta-delta Ct ($\Delta\Delta Ct$) were calculated using a calibrator value (i.e., *delta-Ct – calibrator*
1320 *value*). The choice calibrator value differs depending on experimental design but in this experiment,
1321 the average value of all the non-vernalized treatment ΔCt for the gene of interest was used as the
1322 calibrator value for that specific gene. This is an important factor to the quantification of RGE, as the
1323 choice of calibrator will impact the final gene-fold value (Schmittgen and Livak, 2008).

1324 ΔCt for a specific gene, using one HKG was calculated using (Rao X *et al*, 2013):

$$1325 \quad \Delta Ct = Ct(a \text{ Gene of interest } (GOI)) - Ct(a \text{ reference gene } (HKG))$$

1326 $\Delta\Delta Ct$ are then able to be calculated using (Rao X *et al*, 2013):

$$1327 \quad \Delta\Delta Ct = \Delta Ct(a \text{ Gene of interest } (GOI)) - \Delta Ct(a \text{ reference gene } (HKG))$$

1328 The formula above can be applied to a specific GOI with a specific HKG which are constant for the
1329 entire experiment (i.e GAPDH only). Using this method, a fold-change of the target gene (either
1330 controlled (non-vernalized) or treated (vernalized) can be calculated, normalized, and related to a
1331 HKG using calibrator values as described above. The method was fully described and updated in a
1332 data analysis by Rao *et al* (Rao X *et al.*, 2013).

1333 Although the delta-delta Ct method is the most used, the use of multiple house-keeping
1334 genes is regarded as more reliable in giving a background HKG expression against which to calculate
1335 relative gene expression (Riedel *et al.*, 2014a) (Manoli *et al.*, 2012). The variation that may occur
1336 between the multiple different house-keeping genes is resolved by normalization steps (Huggett *et*
1337 *al.*, 2005). According to *Vandesompele et al*, 2002, the normalization is done by geometric averaging
1338 of the multiple house-keeping genes involved as controls (Vandesompele *et al.*, 2002; Riedel *et al.*,
1339 2014). The equation for using multiple reference genes is as follows:

$$1340 \quad \text{Relative Gene Expression} = \frac{(E_{GOI})^{\Delta Ct_{GOI}}}{GeoMean[(E_{REF})^{\Delta Ct_{REF}}]}$$

1341

1342 The delta-Ct (ΔCt) values for each of the tested samples were determined using the following
1343 equation:

$$1344 \quad \Delta Ct = \text{Control } Ct - \text{Treated } Ct$$

1345 The treated Ct values in this case will be the Ct values from the vernalized samples and the control
1346 from the non-vernalized. The PCR performed consisted of 2 reference genes (*LuGAPDH* and *LuUBI2*)
1347 with a gene of interest from one of our 5 chosen loci. Using the equations above and the primer
1348 efficiency values gained previously as formulated using methods described under section 2.2.5
1349 “Primer Testing”, it is possible to compare relative genetic expression between treated (vernalized)
1350 and control (non-vernalized) individuals in our *Linum* sample set.

1351

1352

1353 2.3 STATISTICAL ANALYSES FOR RELATIVE GENE EXPRESSION

1354 Relative Gene Expression (RGE) is calculated using the $2^{-\Delta\Delta Ct}$ formula as described in section
1355 2.2.10 “Relative Gene Expression: Quantification” above. For follow-up analysis, various packages
1356 were used in R (R Core Team, 2022) ran under the R graphical interface of Rstudio (Rstudio Team,
1357 2020). $\Delta\Delta Ct$ can be calculated by creating a template using the formula to calculate ΔCt (see section
1358 2.2.10) for each of the genes that were tested, including the HKG. The template was created in
1359 Microsoft Excel. the raw data were further analysed in R.

1360 For comparison between the different experiments that took place during different years
1361 (2018 and 2021), an F-test (Box G, 1953) was applied in R. For genetic expression analyses, templates
1362 for the calculation of Ct-values (see “Relative Gene Expression: Quantification”) were made and bar
1363 charts can be plotted in Microsoft Excel. After values for Relative Gene Expressions (RGE) were
1364 gathered for each respective locus, species and treatment, data could be modelled against
1365 environmental variables.

1366 RGE differences of vernalized and non-vernalized treatments, and wild or cultivar
1367 combinations was tested for each gene using a 2-sample, 2-tailed unequal variance t-test. This test
1368 compared fold-gene expression of the gene of interest between the two treatments (vernalized and
1369 non-vernalized). The tests were performed in Microsoft Excel. The formula used for the t-test is
1370 “=TTEST(array1,array2,2,3)”. RGE differences were visualised using R package ggplot including the
1371 command: “+stat_compare_means(label = “p.signif”, method = “t.test”, ref.group = “.all.”)”
1372 (appendix 6).

1373 The differences of flowering initiation between vernalized and non-vernalized individuals
1374 were correlated against latitude of origin ($^{\circ}$ N) using Pearson's coefficient of determination (r^2) in R.
1375 This analysis measured how correlated flowering initiation is to latitude in each data sets (wild and
1376 cultivated; Steel R. *et al.*, 1960). P-values and r^2 values explained by the models were visualised
1377 within scatter plots of the data, showing lines of regression using the R-package "ggplot2" (Wickham
1378 H., 2016). The full command used for drawing correlation coefficients for this analysis can be found
1379 under appendix 6. To further analyse these relationships using modelling, a Linear model analysis
1380 was able to run using R. The command can also be found under appendix 7. For modelling against
1381 environmental variables such as "latitude" and "climate" (described under the "Study and Methods"
1382 part of this chapter), a GLM (General Linear Modelling) approach was used. Various commands used
1383 for the GLM analysis in R can be found under appendix 7.

1384 2.4 RESULTS: FLOWERING TIME BETWEEN EXPERIMENTS FROM DIFFERENT
1385 YEARS

1386

1387 The vernalization results were divided into three different categories: “non-Vernalized
1388 2018”, “Vernalized 2018” and “Vernalized 2021”. Due to COVID-19 pandemic, vernalization
1389 experiment in 2021 had to be terminated early in March 2020. Consistencies between the different
1390 years couldn’t be kept the same and a bias towards earlier flowering plants can be observed in the
1391 2021 experiment as a result.

1392 Differences between the 2018 and the 2021 vernalization experiments was observed when
1393 looking at variation in number of days to flower. This was due to bias towards earlier flowering
1394 plants in the 2020-2021 replicate as the experiment was stopped prematurely (due to Covid-19
1395 restrictions), resulting on limited data collection for late flowering plants. Statistical comparison of
1396 the number of days to flower (flowering initiation) between the two vernalization experiments,
1397 under the different years found significant differences between “vernalized 2018” and “vernalized
1398 2021” experiments (*F-test*, $F=2.523$, $df=196$, $p<0.005$) (Figure 5A). When data were filtered to the
1399 same individuals present in both experiments, the difference between the two datasets remained,
1400 with biases for earlier flowering individuals expressed (*2-sample, 2 tailed t-test* $p<0.001$) (Figure
1401 5B). The significant variation between flowering initiation under the 2018 and 2021 vernalization
1402 experiment suggests that at least for phenotypic measures, data from the 2021 vernalization
1403 experiment should be excluded to avoid biases towards earlier flowering plants.

1404

1405

1406

1407

1408

1409

1410




Figure 5A. A boxplot showing flowering time differences between all the individuals tested across the 2018 non-vernalization, the 2018 vernalization and the 2021 vernalization experiment (*F-test*, $F=2.523$, $df=196$, $p<0.005$). Figure 5B. A boxplot representing flowering time of a subset of all the mutual individuals found in the two vernalization experiments, showing more significant variations (*2-sample, 2 tailed t-test* $p<0.001$). Boxplots suggests a bias for earlier flowering plants in the 2021 vernalization experiment, due to ending prematurely.

1411 2.4.1 Results: Vernalization and flowering time

1412 Data were processed without flowering initiation data from the 2021 vernalization
1413 experiment. The number of days it took for individuals to flower were tested against latitude for wild
1414 and cultivars (Figure 6). Under the 2018 vernalization experiment, dramatic reduction of number of
1415 days to flowering was observed in the wild *L. bienne* that were treated by vernalization. A positive
1416 correlation between the latitude of origin and the number of days to flower in the wild species were
1417 also observed. The cultivars showed a much smaller difference between treatments and a negative
1418 latitudinal correlation with number of days to flowering. This suggests that requirement of
1419 vernalization differs between the two species. *L. bienne* (wild) conveys a dramatic change in
1420 flowering initiation. This was observed on the more northern populations. In the most northern
1421 populations, vernalization reduced the number of days to first flowering by more than 100 days. The
1422 Northern wild species were more sensitive to vernalization, in terms of reducing the number of days
1423 required to flower. In contrast for *L. usitatissimum* (cultivar), vernalization only slightly increases the
1424 number of days it takes for individuals to flower, in comparison with non-vernalized individuals. In
1425 addition, linear modelling showed that for wild samples in non-vernalization, latitude has a
1426 significant influence on days to flowering (*Non-vernalized* $R=0.46$, $p<0.01$; *Vernalized* $R=0.48$
1427 $p<0.01$) (Figure 6). In contrast for cultivars, the requirement to vernalize is dramatically reduced,
1428 with correlation becoming negative between the number of days a plant takes to flower and
1429 latitude, however this was not statistically significant.




Figure 6. Scatterplots with regression line and linear models for flowering initiation to latitude. Note the differences because of latitude and vernalization in the number of days to flowering between wild and cultivated and samples. Reduction of more than 100 days to flowering initiation is observed in more Northern individuals of wild *Linum* whilst this was not observed in the cultivars. In contrast, the cultivars reveal an addition to the number of flowering due to the vernalization stimulus. In the wild number of days to flowering is positively correlated to latitude in both vernalization treatments ($p<0.05$). Shaded areas represent a 95% confidence interval in the dataset respectively.

1430

1431 **2.4.2 Results: Primer Efficiency Values**

1432 For primer efficiency, the desirable value would be 90-110% of efficiency. When we look at
1433 the efficiency percentage of the five flowering time loci however, only *LuFT1* and *LuCO1* fell into this
1434 category. Efficiency values for *LuUBI2* were very low. However, *LuGAPDH* showed close to ideal
1435 values at 87% efficiency. Sub-optimal primer efficiency may cause the representation of false fold-
1436 change; thus, this may affect the representation of the data in this study. Even with these sub-
1437 optimal primer efficiency value, time was of the essence during this study. It was deemed that any
1438 relationship which may be inferred in this study is taken with precautions to this sub-optimal primer
1439 efficiency, as efficiency values is significant to quantification of relative gene expression (Sreedharan
1440 S. et al., 2018).

1441

| Primer | Type | R² | Slope | Efficiency (%) | Converted value |
|----------------|-------------|----------------------|--------------|-----------------------|------------------------|
| <i>LuGAPDH</i> | HKG | 0.995 | -3.670 | 87.273 | 1.873 |
| <i>LuUBI2</i> | HKG | 0.903 | -5.546 | 51.464 | 1.515 |
| <i>LuGI1.1</i> | GOI | 0.333 | -0.559 | 6034.644 | 61.346 |
| <i>LuCO1</i> | GOI | 0.552 | -3.564 | 90.818 | 1.908 |
| <i>LuCO2</i> | GOI | 0.962 | -3.614 | 89.114 | 1.891 |
| <i>LuFT1</i> | GOI | 0.799 | -3.085 | 110.949 | 2.109 |
| <i>LuFT2</i> | GOI | 0.980 | -2.380 | 163.090 | 2.631 |

Table 4. Primer efficiency values for genes of interest (GOI) and housekeeping genes (HKG).

1443


1444 **2.4.3 Results: Comparison of RGE using one and multiple HKGs.**

1445 With the variation observed in the primer efficiency for this study, it is of interest to
1446 compare RGE results using one or multiple HKGs. Treatments are sorted into four groups: non-
1447 vernalized cultivars (NV-Cultivars), vernalized cultivars (V-Cultivars), non-vernalized wild (NV-Wild),
1448 and vernalized wild (V-Wild). A summary boxplot is shown below for the relative gene expression of
1449 the floral integrator *LuFT1*.

1450 Relative gene expression varies when using one or two HKGs. Using one HKG, the vernalized,
1451 cultivar treatment showed a significantly increased *LuFT1* expression (Figure 7) to the rest of the
1452 treatments. This reveals that using different methods and respective calculations could affect further
1453 analysis. Based on the primer efficiency test result that *LuUBI2* (one of the HKGs) have a very low
1454 primer efficiency (51.46%), it is thought that using *LuUBI2* may have skewed relative gene expression
1455 results. Therefore, for further analysis, only one HKG (*LuGAPDH*) was considered and the use of the

1456 delta-delta ct ($2^{-\Delta\Delta Ct}$) method would be further applied for the rest of the analysis regarding relative
 1457 gene expression.

1458



1460

1461 Figure 7. Boxplots comparing Relative Gene Expression for *LuFT1* using two (Figure 6A) or just one (Figure 6B) house-
 1462 keeping genes. In Figure 5B the relative gene expression of *LuFT1* is found to be significantly different (marked by *) to the
 1463 other treatments ($p < 0.05$, multiple pairwise t-test).

1464


1465 2.4.4 Results: Relative Gene Expression in Response to Vernalization

1466 Relative gene expression results were initially sorted into groups reflecting treatments and
 1467 species (non-vernalized-cultivated, vernalized-cultivated, non-vernalized-wild, and vernalized-wild).
 1468 Relative gene expression ($2^{-\Delta\Delta Ct}$) was compared across each of the treatments to see whether any
 1469 species/treatments show significant differences in relative gene expression respective of each gene
 1470 for the treatment. Boxplots can be illustrated with a Bonferroni-corrected multiple pairwise t-test to
 1471 reveal significance between treatments (Figure 8).

1472 Variation in expression was mostly seen in cultivars with a potential down-regulation in all
 1473 the tested genes when cultivars were vernalized (Figures 7 A-E). Less strong RGE differences were
 1474 seen in the wild samples. In the case of the comparisons between treatments, most of the
 1475 differences in the data were observed when treating cultivars to vernalization. This is interesting as
 1476 phenotypically; this difference wasn't to be expected. This is because flowering initiation in the
 1477 cultivars did not show faster flowering initiation in response to vernalization and instead showed
 1478 slightly slower flowering initiation (Figure 4). The multiple pairwise t-test however, was comparing
 1479 differences of relative gene expression when vernalized, against all wild and cultivar individuals


1480 which are perhaps behaving very differently in terms of expressing each of these genes and maybe
1481 independent of each other. Therefore, variation among individuals was explored in more detail.
1482 Comparisons between expression responses (relative gene expression) against individuals with
1483 different treatment were undergone using a paired t-test as in the next part of this chapter.

1 A



1485 Figure 8A. Relative Gene Expression (RGE) of samples and treatments for the locus *LuCO1*. The significance of multiple
1486 pairwise t-tests, adjusted with Bonferroni correction, are shown at the top of each boxplot. The labels "ns" means no-
1487 significant differences were observed between 4 observed treatments.


B




1488

1489

1490 Figure 8B. Relative Gene Expression (RGE) of samples and treatments for the locus *LuCO2*. The significance of multiple
 1491 pairwise t-tests, adjusted with Bonferroni correction, are shown at the top of each boxplot. "Vernalized cultivated"
 1492 ("NV_Cultivars") show a significant RGE variation when compared to other treatments ("**" means $p < 0.01$).
 1493




1494 Figure 8C. Relative Gene Expression (RGE) of samples and treatments for the locus *LuFT1*. The significance of multiple
 1495 pairwise t-tests, adjusted with Bonferroni correction, are shown at the top of each boxplot. "Vernalized cultivated" (V-
 1496 Cultivars) shows a significant RGE reduction ("*" means $p < 0.05$).
 1497



1498 Figure 8D. Relative Gene Expression (RGE) of samples and treatments for the locus *LuFT2*. The significance of multiple
 1499 pairwise t-tests, adjusted with Bonferroni correction, are shown at the top of each boxplot. "Vernalized cultivated" (V-
 1500 Cultivars) shows a significant RGE reduction ("*" means $p < 0.05$).
 1501

1502



1503 Figure 8E. Relative Gene Expression (RGE) of samples and treatments for the locus *LuGI1.1*. The significance of multiple
1504 pairwise t-tests, adjusted with Bonferroni correction, are shown at the top of each boxplot. "Vernalized cultivated" (V-
1505 Cultivars) shows a significant RGE reduction ("**" means $p < 0.01$).

1506

1507 Variation in expression was mostly seen in cultivars with a potential downregulation in all
1508 the tested genes when cultivars were vernalized. Less strong RGE differences were seen in the wild
1509 samples. In the case of the comparisons between treatments, most of the differences in the data
1510 were observed when treating cultivars to vernalization. This is interesting as phenotypically; this
1511 difference wasn't to be expected. This is because flowering initiation in the cultivars did not show
1512 faster flowering initiation in response to vernalization and instead showed slightly slower flowering
1513 initiation (Figure 6). The multiple pairwise t-test, however, was comparing differences of relative
1514 gene expression when vernalized, against all wild and cultivar individuals which are perhaps
1515 behaving very differently in terms of expressing each of these genes and maybe independent of each
1516 other. Therefore, variation among individuals was explored in more detail. Comparisons between
1517 expression responses (relative gene expression) against individuals with different treatment were
1518 undergone using a paired t-test as in the next part of this chapter.

1519


1520 2.4.5 *LuGI1.1* (*GIGANTEA*) Expression Response

1521 We tested fold-change differences (delta-delta ct) as a relative gene expression (RGE)
1522 measure to investigate whether differences in gene expression under vernalization treatment can be
1523 observed. To be more conservative, since we have two treatments compared with 5 genes, our
1524 alpha needs to be adjusted using Bonferroni's correction to 0.005 (0.05/10). Overall differences in
1525 *LuGI1.1* expression in cultivars was the closest to approaching significance, however, are not

1526 significant when considering Bonferroni's correction (*paired, 2-tailed t-test, p = 0.012 (p > α), t-stat =*
 1527 *3.043, t-Critical two-tail= 2.228*). The greatest differences were observed in cultivars EDE, BLE, PRI,
 1528 GIS, and MON (Figure 9).


1529 There was no significant difference expression of *LuGI1.1* between treatments in wild
 1530 samples (*paired, 2-tailed t-test, p = 0.407, t-stat = -0.864, t-Critical two-tail= 2.228*) (Figure 10).

1531 Expression varied more among samples with observations of both up and down-expression of
 1532 *LuGI1.1*. Samples Lla_A and Tor_4 increased *LuGI1.1* expression the most and vernalized sample
 1533 Tal_4 decreased expression the most.



1534 Figure 9. Bar charts showing *LuGI1.1* fold change between cultivars under non-vernalized and vernalized treatments
 1535 (*paired, 2-tailed t-test, p = 0.012 (p > 0.005)*). The cultivars showing the largest downregulation differences are shaded in
 1536 darker grey.

1537




1538 Figure 10. Bar charts showing *LuGI1.1* fold change between wild samples under non-vernalized and vernalized treatments.
 1539 Highlighted in the bar charts with darker grey and orange are samples which seem to decrease and increase expression of
 1540 *LuGI1.1* when vernalized, respectively.

1541

1542 **2.4.6 *LuCO1* and *LuCO2* (*CONSTANS*) Expression Response**

1543 The two genes *LuCO1* and *LuCO2* did not show a significant expression difference between
 1544 vernalization treatments for either wild or cultivated samples (Figure 11). However, the variation
 1545 within the data could still be observed.


1546 There were no significant differences in *LuCO1* expression change in cultivars (*paired, 2-*
 1547 *tailed t-test, p = 0.229, t-stat = 1.281, t-Critical two-tail= 2.228*). Most of the variation observed
 1548 occurred in sample LIR with decreased expression when vernalized. Expression of *LuCO1* was neither
 1549 significantly different between non-vernalized nor vernalized treatments of the wild samples (*paired,*
 1550 *2-tailed t-test, p = 0.310, t-stat = -1.070, t-Critical two-tail= 2.228*). Different samples showed
 1551 increased and decreased expression of *LuCO1*. As in the case of *LuGI1.1*, samples LLa_A and Tor_4,
 1552 and 19_30, increased expression after vernalization. Sample Tal_4, as well as Tym_30 and Saf_10
 1553 decreased expression after vernalization.



1554
 1555 Figure 11. Bar charts showing *LuCO1* fold change between cultivars (A) and wild types (B) under non-vernalized and
 1556 vernalized treatments. Highlighted in the bar charts with darker grey and orange are samples which seem to decrease and
 1557 increase expression when vernalized, respectively.

1558

1559 There were non-significant differences in *LuCO2* expression between the non-vernalized and
 1560 the vernalized treatments for cultivars (*paired, 2-tailed t-test, p = 0.132, t-stat = 1.640, t-Critical two-*
 1561 *tail= 2.228*), and wild samples (*paired, 2-tailed t-test, p = 0.222, t-stat = -1.301, t-Critical two-tail=*
 1562 *2.228*) (Figure 12). The trend was for *LuCO2* expression to decrease in cultivars during vernalization,
 1563 while the trend for wild samples was to increase *LuCO2* expression. This revealed that although
 1564 there were overall significant differences when comparing vernalized cultivated samples gene
 1565 expression for *LuCO2*, when testing between vernalized and non-vernalized cultivated samples only,
 1566 the expression of *LuCO2* shows no significant difference and results to vernalization having no
 1567 significant effect in the relative gene expression of *LuCO2*, for both cultivar and wild samples.
 1568



1569 Figure 12. Bar charts showing *LuCO2* fold change between cultivars (A) and wild types (B) under non-vernalized and
 1570 vernalized treatments. Highlighted in the bar charts with darker grey and orange are samples which seem to decrease and
 1571 increase expression when vernalized, respectively.


1572

1573

1574 2.4.7 *LuFT1* and *LuFT2* (FLOWERING LOCUS T) Expression Response

1575 The cultivars showed a trend of decreased expression of *LuFT1* when vernalized. This was
 1576 seen more in cultivar samples EDE and BLE, alike the *LuCO2* results. The overall change in gene

1577 expression was not significant (*paired, 2-tailed t-test, p = 0.093, t-stat = 1.856, t-Critical two-tail=*
 1578 *2.228*) (Figure 12A). Wild samples showed a mixed pattern with one potential anomaly for sample
 1579 Saf_10 (not shown) with much greater *LuFT1* expression than other wild samples across both
 1580 treatments. There was no significant expression change in *LuFT1* expression when wild samples were
 1581 vernalized (*paired, 2-tailed t-test, p = 0.669, t-stat = 0.440, t-Critical two-tail= 2.228*) (Figure 13B).
 1582




1583 Figure 13. Bar charts showing *LuFT1* fold change between cultivars (A) and wild types (B) under non-vernalized and
 1584 vernalized treatments. Highlighted in the bar charts with darker grey and orange are samples which seem to decrease and
 1585 increase expression when vernalized, respectively.

1586


1587 As was observed for *LuFT1*, cultivar samples EDE and BLE showed the most reduced *LuFT2*
 1588 expression after vernalization with non-significance on the fold change between the two treatments
 1589 for the entire cultivar sample set (*paired, 2-tailed t-test, p = 0.087, t-stat = 1.916, t-Critical two-tail=*
 1590 *2.262*) (Figure 14A). Among the wild samples, an exceptionally high downregulation in expression of
 1591 *LuFT2* (>100 fold difference, not shown) was observed sample Saf_10. This sample was treated as an
 1592 anomaly. Nearly the same trends as *LuFT1* were seen, with sample Tor_4 showing the most increase
 1593 in expression of *LuFT2*. However, the results between the two treatments for the wild sample set
 1594 were still not statistically significant altogether (*paired, 2-tailed t-test, p = 0.575, t-stat = 0.579, t-*

1595 *Critical two-tail= 2.228*) (Figure 14B). Relative gene expression was also tested between species in
 1596 only the vernalized samples using a paired, 2-tailed t-test. None of the tested flowering time genes
 1597 have shown significant expression difference between species when vernalized ($\alpha = >0.05$).

1598



1599



1600


1601 Figure 14. Bar charts showing *LuFT2* fold change between cultivars (A) and wild types (B) under non-vernalized and
 1602 vernalized treatments. Highlighted in the bar charts with darker grey and orange are samples which seem to decrease and
 1603 increase expression when vernalized, respectively.

1604

1605 2.4.8 Gene Expression and Flowering Time

1606 General linear models (GLM) were used to test whether the difference in RGE between
 1607 vernalized and non-vernalized wild and cultivated *Linum* were associated with the flowering time
 1608 initiation (Figure 15). In wild samples, the relative gene expression (RGE) differences of *LuFT1* and
 1609 *LuFT2* were significantly positively associated with days to flower after vernalization (*LuFT1* $t=6.359$
 1610 $p<0.001$; *LuFT2* $t=5.128$ $p<0.001$). The other tested loci, *LuCO1*, *LuCO2*, and *LuGI1.1* showed no
 1611 significance when tested for association with flowering time (*LuCO1* $t=1.510$ $p=0.162$; *LuCO2* $t=1.324$
 1612 $p=0.215$; *LuGI1.1* $t=1.766$ $p=0.108$).

1613 For the cultivars, it was found that *LuFT1* and *LuFT2* expression differences were not
 1614 significantly associated with flowering initiation (*LuFT1* $t=1.226$ $p=0.237$; *LuFT2* $t=1.732$ $p=0.122$).
 1615 Neither did *LuCO1* and *LuCO2* expression differences show significant associations with flowering
 1616 initiation (*LuCO1* $t=-1.575$ $p=0.150$; *LuCO2* $t=1.287$ $p=0.230$). Interestingly, *LuGI1.1* expression
 1617 differences showed a significant positive association with flowering initiation (*LuGI1.1* $t=2.932$
 1618 $p=0.016$).
 1619



1620 Figure 15. Scatterplots showing number of days to flower after vernalisation labelled as “Days_to_fl” in relation to relative
 1621 gene expression difference in wild samples and cultivars separately.


1622

1623 **2.4.9 Relative Gene Expression and Latitude**

1624 The earlier phenotypic analysis found that vernalized wild individuals dramatically reduce
1625 the number of days required to flower when compared to non-vernalized wild plants, with
1626 populations from more northerly latitude of origin showing vernalization sensitivity. Here we tested
1627 latitude against relative gene expression for the cultivars and wild samples (Figure 16).

1628 Latitude was positively associated with relative gene expression difference in the genes; *LuFT1*
1629 ($t=2.668$ $p=0.0257$); *LuFT2* ($t=3.443$ $p=0.0076$); and *LuGI1.1* ($t=2.320$ $p=0.0428$). In the cultivars, there
1630 was no correlation between any gene expression tested in this study and latitude.

1631




1632 Figure 16. Scatterplots showing latitude of origin (Lat) in relation to relative gene expression difference in wild samples and
1633 cultivars separately.

1634

1635 **2.4.10 Results: Relative Gene Expression and Climate**

1636 More Northerly location (higher latitude) suggests of colder, more wet, and windier
1637 climates. Wild samples were analysed for the relationship of flowering regulations genes expression
1638 differences and several climatic variables under that latitude. Climate of origin was summarised into
1639 principal components, and princliple component 1 (PC1) was then tested against relative gene
1640 expression differences as mentioned under section 2.2.1 “Samples and Experiment” of this chapter.



1641 Figure 17. Scatterplots showing climate variable PC1 (pc) in relation to relative gene expression difference of wild samples.

1642 Expression differences of *LuFT1* and *LuFT2* were significantly negatively associated with PC1.
1643 Lower PC1 values suggests colder climate (appendix 3) a summary of all the climatic variables (*LuFT1*
1644 $t=-2.905, p=0.017$; *LuFT2* $t=-3.812, p=0.004$). There was also a significant negative association
1645 between climate and *LuGI1.1* ($t=-3.180, p=0.009$). The correlation observed in the wild for *LuGI1.1*
1646 against PC1 is more significant than it is observed under latitude alone. This suggests that climatic
1647 variables represent more significant correlation to the RGE observed under *LuGI1.1* and that that
1648 locus may be more varied in the wild according to the local climate than the latitude alone.

1649

1650

1651 2.5 DISCUSSION AND CONCLUSION

1652

1653 In the 2018 vernalization experiment, vernalization reduced the number of days to flowering
1654 in the wild *L. bienne*, especially in more Northern populations. This suggests that vernalization is an
1655 important mechanism in *L. bienne* and there was variation in regulation of flowering time in the wild
1656 relatives based on their latitude of origin. In addition to this, the results for *L. usitatissimum* suggests
1657 that requirements for vernalization were different between wild and cultivated *Linum*. This
1658 difference could be due to artificial selection for faster flowering of cultivars as part of domestication
1659 in *L. usitatissimum*. In the literature, genetic and association mapping have identified hundreds of
1660 genes as targets of divergence due to domestication (Smýkal P. et al., 2018). Evidence of this has
1661 been previously observed through resequencing of candidate genes related to seed sizes and weight
1662 (Guo et al., 2020a). Vernalization insensitive flowering time could also be linked to productivity of
1663 *Linum* in temperate climates, and thus mechanism to vernalization is of interest in *Linum* breeding
1664 (Gutaker et al., 2019).

1665 Other studies concluded that temperature, photoperiod (day length), and light availability
1666 (amount of sun light during the day) can influence developmental rates in crop plants such as wheat
1667 and other annual crops (Craufurd and Wheeler, 2009; Cave et al., 2013; McMaster et al., 2008). In
1668 wild *L. bienne*, variation in environmental conditions at different latitudinal localities have selected
1669 for differences in flowering initiation due to vernalization. This variation was not observed in our
1670 cultivated *L. usitatissimum* collection, suggesting no sensitivity to vernalization. However, in the
1671 literature there are suggestion that winter type *L. usitatissimum* found in Texas, USA, are sensitive to
1672 vernalization (Darapuneni M. et al., 2014). They suggest that these winter types are different to
1673 varieties grown in more Northern areas because they are grown in Autumn due to higher spring and
1674 summer temperatures in Texas. We saw no evidence of sensitivity to vernalization in our Northern
1675 European Spring and Winter varieties of *L. usitatissimum*. We suggest that there is wider variation
1676 within *L. usitatissimum* for requirement to vernalize. Perhaps growing seasons of each variety can
1677 affect the sensitivity of *L. usitatissimum* variant to vernalization.

1678 Another result illustrated in this study is the difference of gene expression due to
1679 vernalization conditions. After considering correction for multiple testing and variation among
1680 individuals, there was no significant difference due to vernalization in the fold change of any of the
1681 tested genes in both wild and cultivar types. Sun et al., 2019 also did not find an expression
1682 difference for *LuGI1.1*, between their cultivar lines (Sun et al. 2019). In addition to this, vernalization
1683 conditions tested in this study were not significant for *LuGI1.1* gene expression either. There is little

1684 evidence that among the genes tested in this study, cultivated and vernalized individuals showed a
1685 significant difference in gene expression due to vernalization. This suggests that fold changes in all
1686 three flowering time genes expression were not significantly affected by vernalization. However, we
1687 have not tested fold-change between populations, as we did not have enough samples over
1688 population in our samples to compare for this. To test for population variation, more individuals
1689 over each population could be tested in the future.

1690 It is interesting to observe that in our wild *Linum*, there was a clear positive correlation
1691 between the RGE difference of both tested *LuFT* copies in relation to the number of days to flower,
1692 otherwise stated as “flowering initiation” in this thesis. These results are alike to previous
1693 photoperiodicity experiments of other cultivar types, whereby both *LuFT1* and *LuFT2* were shown to
1694 be associated with flowering initiation under photoperiodicity, but not the other flowering time
1695 genes tested (Sun et al, 2019). However, this relative gene expression difference was not observed in
1696 the cultivated *Linum* samples tested in this study. Instead, expression of *GIGANTEA (LuGI1.1)* was
1697 observed to be positively correlated to the number of days to flowering. There are suggestions that
1698 the flowering time gene *GIGANTEA* is involved in flowering and maturity development in
1699 heterozygous lines of the plant species *Glycine max* (Watanabe *et al.*, 2011). This becomes
1700 particularly interesting when considering the population genetics of a given study species. The
1701 illustration that the gene *GIGANTEA* is involved in flowering development of heterozygous *Glycine*
1702 *max* varieties, suggests that heterozygosity could become a variable to test against *GIGANTEA*
1703 expression under future studies involving cultivated *Linum*.

1704 The difference in *LuGI* expression was not observed in the wild *Linum* as was observed in the
1705 cultivars. The differences in relative gene expression found in the current study, reveal that there is a
1706 difference in gene regulation, perhaps due to different vernalization requirement between the wild
1707 and cultivated *Linum* and their population genetic variability. It is worth considering that flowering
1708 initiation is very different in the cultivated types when looking at the effects of vernalization to the
1709 number of days to flower between the two species. This study reveals that the expression of the
1710 flowering integrator gene *LuFT* have lost the ability to respond to vernalization in developed
1711 cultivars, and instead expression of another flowering time regulator gene may have taken its role in
1712 terms of initiating flowering. A cause could be changes in genetic expression of flowering regulators
1713 throughout developments of plants. Changes in flowering time gene roles and developmental
1714 switches from vegetative to reproductive stages of plants is not new. In the model *Arabidopsis*
1715 *thaliana*, it has been shown that flowering time regulator (*FT*) changes expression drastically in
1716 relation to developments (Blümel *et al.*, 2015). Upstream genes in the flowering time network, such
1717 as *CRY2* (Cryptochrome Circadian Regulator 2) and *LHY* (Late Elongated Hypocotyl) (Park M et al,

1718 2016) may influence the non-significant relative expression differences of *LuFT* to flowering initiation
1719 and may have a major role to play in terms of this loss of vernalization requirement in the cultivars.
1720 Determining how vernalization affects the flowering time gene network, especially in *Linum* cultivars
1721 would be useful for agriculture to find genes which are important for specific growing conditions and
1722 exploring the potential loss of certain genetic functions due to selective breeding and human
1723 cultivation.

1724 For most of the tested genes, vernalized, and non-vernalized expression changes were not
1725 significantly different in wild and cultivated samples. In the literature however, there are some
1726 studies in other non-model organisms, which reveal that *GIGANTEA* homologous genes were
1727 differently expressed under vernalization treatments. This was true in a study using Ryegrasses
1728 (Paina *et al.*, 2014), and white lupins (Rychel *et al.*, 2019). However, studies regard *GIGANTEA* to be a
1729 photoperiodic response gene. This comes with implication that *GIGANTEA* interacts with another
1730 photoperiodic gene, the *CONSTAN* (CO). However, some studies have illustrated that *GIGANTEA* may
1731 be independently mediating photoperiodic control of flowering (Jung *et al.*, 2007). Although, there is
1732 evidence the *GIGANTEA* may be variably expressed when vernalized in other non-model organisms,
1733 the experiment in this case was limited in the number of paired samples that experienced both
1734 vernalization and non-vernalization treatments in 2021 (9 wild and 8 cultivated). This is due to
1735 availability of DNA materials (only available for 2021 experiments) and restrictions caused by the
1736 Covid-19 pandemic. There were 62 germinating individuals in the vernalization 2021 chamber.
1737 However, only around 20 individuals germinated in the non-vernalization 2021 chamber. Some of
1738 these have not had enough time to produce leaf materials for RNA extraction. In this study, there
1739 were only 38 individuals which experienced vernalization in both 2018 and 2021 experiments for
1740 which there was both expression and phenotypic data. In the 2018 experiment (pre-Covid-19) there
1741 were 197 vernalized individuals. This gives an idea about the limitation of data available for genetic
1742 study under the 2021 experiment due to restrictions caused by the Covid-19 pandemic. However, as
1743 in Figure 4, flowering time results from the 2018 experiment revealed correlation in flowering time
1744 and latitude in wild relatives and in addition suggests the loss of vernalization in cultivars. In the
1745 future, time permitting, an experiment involving vernalization, looking at a larger set of samples,
1746 may in turn reveal a stronger difference between the gene fold-change in vernalized and non-
1747 vernalized individuals.

1748 Another aspect that could be of interest from the genetic point of view is epigenetic
1749 regulation. Epigenetics is related to the idea that genetic regulation may be affected by different
1750 environments, which can include temperature changes. This was studied in the model *Arabidopsis*,
1751 looking at genetic regulation of flowering time (Khan, Ai and Zhang, 2014) and epigenetic responses

1752 to heat (Liu *et al.*, 2015). In the current study, certain genes in the wild samples showed a mix of
1753 increased and decreased expression across treatments. For example, the Southern samples Lla_A
1754 and Tor_4 tended to show increased expression (upregulation) when vernalized, and some Northern
1755 samples such as Tym_30 showed decreased expression (downregulation) of multiple flowering
1756 initiation genes when vernalized. Across wild sample populations, there were differences in genetic
1757 regulation based on their locality, or in this case latitude for *LuFT1*, *LuFT2*, and *LuGI1.1*. This further
1758 adds to the point that different population may regulate flowering time genes differently based on
1759 adaptation to their local environment which often affect downstream processes such as biosynthesis
1760 of plant hormones leading to developmental differences (Jaakola and Hohtola, 2010). A study of the
1761 model *Arabidopsis* comparing differences in altitude reveals that there was variation in genetic
1762 regulation of flowering time regulators, for populations under different altitude. These include a
1763 tested flowering regulator in this study, the Flowering Locus T (*FT*), with addition of vernalization
1764 genes such as VERNILIZATION INSENSITIVE 3 (*VI3*). Regulations of flowering regulator have been
1765 revealed to regulate MADS protein downstream and this differed between low and high altitudes
1766 genotypes (Suter *et al.*, 2014). Test of more *Linum* samples in consideration of altitude of different
1767 populations could help verify these results for *Linum* in future studies.

1768 Regarding the cultivars, most of the expression differences were reduced expression
1769 following vernalization. Reduced expression of flowering time genes such as the *LuCO1* and *LuCO2*
1770 (*CONSTANS 1* and *2*) could be a response to the expression of downstream flowering meristem
1771 identity such as *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS (SOC)* to initiate flowering
1772 following vernalization as has been found in the model *Arabidopsis* (Sasaki *et al.*, 2017)(Li *et al.*,
1773 2015a)(Valentim *et al.*, 2015a). Therefore, further studies relating expression of homologues to *SOC*
1774 genes in *Linum* to expression changes of *CO* genes in response to vernalization could help confirm
1775 this vernalization pathway in *Linum*.

1776 Flowering time gene expression of *LuFT1*, *LuFT2*, and *LuGI1.1* were found to be associated
1777 with environmental variables related to latitude and climate of origin of wild samples. This result
1778 suggests that in the wild, phenological adaptation to the local environments could be due to
1779 different gene expression responses of these three flowering time genes. This is supported by the
1780 suggestion that *FT* was found to be a major locus contributing to local adaptation in flowering of
1781 perennial plants (Wang J. *et al.*, 2018). The strongest associations between genetic expression as
1782 well as phenotypic responses (flowering time) to the local environments was shown by both *LuFT*
1783 genes. This contrasts with the result for wild flowering time that was not significantly associated
1784 with *LuGI1.1* relative gene expression. However, latitude and environmental variables were
1785 associated with *LuGI1.1* relative gene expression. In addition, in the cultivar set, *LuGI1.1* was the only

1786 gene that showed an association with flowering time. It is of interest to look at genes that maybe
1787 influencing the expression of *LuGI1.1*. It may be that expression of those genes were more related to
1788 flowering time than *LuGI1.1* itself. For example, there are multiple variants of the gene *GI* within
1789 *Linum* that were not tested in this study. These genes could be tested as part of future to further
1790 explore the flowering time network in *Linum* and implications of environmental variables. In
1791 addition, other vernalization related genes have also been observed to be related to local
1792 adaptation. The *FRI* (*FRIGIDA*) gene are related to local adaptation to drought tolerance by
1793 controlling flowering time in *Arabidopsis thaliana* (Lovell J. et al., 2013; Tigano and Friesen., 2016). In
1794 future studies, *FRI* can be another gene of interest when suggesting local adaptation to vernalization
1795 responses in *Linum*. In conclusion, evidence of local adaptation can be inferred from the phenotypic
1796 variation observed in the wild *L. bienne* tested here. This is further supported by the variation in
1797 expression of *FT*, linked to correlation towards environmental variables.

1798 One issue in this study was the decision to use of only one HKG. The use of multiple HKGs is
1799 regarded as superior to using only one HKG for quantification purposes as it gives a better picture of
1800 background expression (Remans *et al.*, 2008) (Manoli *et al.*, 2012). However, in the case of this
1801 experiment, the efficiency of *LuUBI2*, one of the tested HKG primers was very low (51%). This means
1802 that using both genes may skew the results due to the less efficient HKG so the Ct-values for *LuUBI2*
1803 were dropped from the analysis to conserve for this. In addition, the efficiency of some of the genes
1804 of interest (GOI) could be improved as part of future research. Quantification of relative gene
1805 expression using two HKGs was also explored as part of the analysis, with the less efficient *LuUBI2*.
1806 The results showed a drastic difference in relative gene expression when using two HKGs, with no
1807 significant differences between the treatments, potentially skewing relative expression. Using two-
1808 HKG indicate that the genes *LuFT1* and *LuCO2* showed the greatest difference in expression between
1809 vernalized and non-vernalized (non-vernalized) samples, but these differences were not statistically
1810 significant (appendix 8). This result suggests that using the less efficient primer may skew our results.
1811 We suggest improving the efficiency of *LuUBI2* for future research looking on the relationship of
1812 flowering time gene expressions with vernalization, using multiple HKGs. This would strengthen our
1813 results suggesting *LuFT1*, *LuFT2*, and *LuGI1.1* associations with environmental variables as found
1814 using one HKG in this study. Until such a time, our results are to be treated cautiously as a signal for
1815 the effects of environmental variables in relative gene expressions of *LuFT1*, *LuFT2*, and *LuGI1.1*.

1816 The relationship between relative gene expressions (RGE) difference calculated with two
1817 HKGs was tested against latitude of origin of wild samples (appendix 9) and no significant
1818 relationships were found. These results contrast to the results found above, using one HKG and
1819 suggest that the low efficiency value of *LuUBI2* could have contributed to masking some significance

1820 seen with only one HKG. In the future, using two HKGs with a higher efficiency value are
1821 recommended to test the relationships between gene expression to latitude, and climatic variables.
1822 We reflected that using one HKG is not ideal as most recent relative gene expression studies
1823 recommends using multiple HKGs. However, limited results on the expression of the floral integrator
1824 *FT* using one HKG are suggestive. To robustly test the expression of *FT* in *Linum* under vernalization,
1825 observation of HKG with higher efficiencies are recommended for future research.

1826 To further this research, other important regulatory flowering time genes in the network
1827 could be tested such as *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS (SOC)* and as well as
1828 homologues of further downstream meristem identity genes found in the model *Arabidopsis* such as
1829 *LEAFY (LFY)*, *APETALA1 (AP1)*, *SEPALLATA3 (SEP3)* and *FRUITFULL (FUL)* (Schopf *et al.*, 1996) (Gregis
1830 *et al.*, 2009). Meristem development in *Linum* could be measured directly as a phenotypic
1831 comparison as has been done in other studies (Kayes and Clark, 1998) (Heisler and Jönsson, 2007).
1832 The complex network of flowering time genes could also imply that there is no single gene
1833 responsible for the phenotypic differences seen when both wild and cultivar type are vernalized. It
1834 may occur that several genetic regulations contribute, including the five major genes tested here.
1835 However, real-time primers that are specific to the two paralogues of each gene present in *Linum*
1836 would first need to be designed and tested, which is not a trivial task (Sun *et al.*). More genes which
1837 are specific to the vernalization pathway should be tested. Outside of the main floral pathway
1838 integrators, which *FT* (Flowering locus T) is a part of, this study looked only at previously designed
1839 flowering time genes which were related to photoperiodicity, such as *CO* and *GI*. It is interesting
1840 therefore, that the two *FT* loci showed significant relationships with latitude and climate of origin
1841 when observing the wild types. Vernalization-specific genes such as *FLC* (Flowering locus C) and *FRI*
1842 (FRIGIDA) have been shown to also play a role in the regulation network to suppress flowering locus
1843 T in the model *Arabidopsis* (Flowers *et al.*, 2009a). It would be of interest then to look at these
1844 vernalization pathway genes with appropriately designed primers specific for those genes in *Linum*
1845 as part of future studies.

1846 For further studies, it would also be of interest to measure traits such as seed yield. *Gutaker*
1847 *et al.*, have talked about potential link of flowering time to fruiting. Expression data for genes which
1848 may affect flowering time could be linked to whether seed yield from these measured individuals are
1849 affected by the changes of expression. Furthermore, variation in seed germination due to
1850 vernalization has been illustrated in the model *Arabidopsis* (Auge *et al.*, 2017). This may potentially
1851 link to productivity in *Linum* that may vary due to the requirement of vernalization and therefore
1852 affecting flowering time.

1853 CHAPTER 3: THE QUANTIFICATION OF POLLEN AND POLLEN TUBES TO REVEAL
1854 POLLEN GERMINATION VIABILITY IN *LINUM* UNDER DIFFERENT TEMPERATURE
1855 TREATMENTS.
1856


1857 Pollen is the vehicle in which the male gamete is found. It is considered as a
1858 microgametophyte (sperm-producing gametophyte) in plants, a powder-like substance where the
1859 male gamete is produced and transferred (Johnstone, 2001). Pollen grains needs to disperse and
1860 travel to the stigma to fertilize egg cells which are found in the ovule and are non-dispersal. Pollen
1861 will often encounter many ecological and environmental pressures as they disperse (Loveless &
1862 Hamrick., 1984). They are considered as one of the major contributors to the ability of flower to
1863 turn into seeds and produce progenies, a transformation from plant haploid gametophytes into
1864 diploid sporophytes (Pacini & Dolferus, 2016). Pollen is essential for seed production and thus
1865 agriculturally significant in crop plants such as *Linum*. As well, wild flowering plants often deploy
1866 different flowering strategies to optimise germination by pollen, and along with this, have evolved
1867 different germination mechanisms useful for artificial selections of male genotypes (Williams, 2012)
1868 (Tel Zur, et al., 2020). In this chapter we quantified pollen and pollen tube counts in *Linum*
1869 individuals treated under different temperature stress.

1870 Pollen must fertilize the ovary through the stigma-style-ovary system to form diploid
1871 sporophytes and form seeds (Pacini & Dolferus, 2016). The receptive ability of the stigma-style-ovary
1872 system is another aspect that could be observed with regards to the success of pollen germination.
1873 In this study however, we will focus on the pollen's ability to germinate and fertalize ovaries.
1874 Assessment of *in vivo* pollen germination success can be done by counting the number of pollen
1875 tubes in the style (Alonso et al., 2013; Williams & Reese, 2019). We decided to observe pollen
1876 initially because pollen is relatively easier to stain and study under confocal microscopy than the
1877 stigma. In a book published in 1998, Delph F, *et al*, reviewed variation in pollen growth, where there
1878 is an observation that pollen growth varies between species due to pollen/sperm competition
1879 whereby pollen traits such as, how quickly pollen reach the ovules, or how many pollen grains are
1880 transferred influence the success of germination. In addition, pollen competition in plants is
1881 described as gametophytic selection, potentially affected by environmental cues such as ambient
1882 temperature. This competition may cause pollen traits to differ in different flowering plants (Delph
1883 and Havens, 1998). In our *Linum* samples, we observed the viability of pollen and pollen tubes for
1884 the germination into seeds. We are particularly interested in the effect of temperature towards
1885 pollen viability under both wild and cultivated species.

1886

1887 3.1. Pollen Structure

1888 In terms of their physical appearance, pollen grains varies in size across species (2-200 μ m)
1889 and contains parts where the generative cells are stored, often with a tube nucleus, within an intine,
1890 exine, and an aperture pore whereby the gametes can be omitted from using a structure widely
1891 known as pollen tubes (Johnstone, 2001; Pacini E, 2008). Pollen grains are microscopic. The smallest
1892 pollen size in diameter is found in *Myosotis*, with a diameter of 2.4-5 μ M (Sporemex, 2022), this
1893 often makes microscopy an ideal tool for observing pollen structures. The structure of a typical
1894 pollen is illustrated as below (Unacademy, 2022):



1895 Figure 1. A diagram depicting the well-known structure of a mature pollen.

1896

1897 During germination, the tube nucleus part of the pollen expands into a pollen tube whereby the
1898 male gamete of the pollen can be transferred to the ovary. Pollen tube expands out of the pollen
1899 through the “aperture pore”. The number of aperture pores can vary between species. It is thought
1900 that increased number of apertures in angiosperm pollen grains can offer the species a selective
1901 advantage as it increases the number of germination sites and facilitates the chances of contact
1902 between the pollen tubes and the female ovule (Furness & Rudall, 2004).

1903 For germination, pollen often require active RNA synthesis. This was evident in the conifers,
1904 where RNA synthesis for pollen germination was studied for its dependence on transcription and
1905 translation (Breygina et al., 2021). When pollen is germinating, the tube nucleus will extend towards
1906 the aperture pore and will then form a pollen tube. These pollen tubes are organelles which can
1907 elongate and are responsible for the transfer of pollen male gametes into the stigma and down to
1908 reach the ovule (Adhikari et al., 2020). When pollen tubes reach ovules, male gametes within the
1909 tube nucleus enter the ovaries to form zygotes. The zygotes are the progenitor stem cells which can

1910 form embryos that later form into seeds (Kanday and Sundaresan, 2021). With this knowledge, it is
1911 then important to consider pollen viabilities in terms of their ability to form zygotes. This includes,
1912 amount of pollen, amount of pollen tube, and the rate which pollen tube reaches ovary. This can
1913 lead to factors influencing seed formation success (Iwazumi and Takahashi, 2012).

1914 Pollen viability is therefore of interest related to the breeding strategies of *Linum* as both
1915 wild and cultivated types are present under different environmental conditions. In *L. usitatissimum*,
1916 pollen can be used as a tool to measure gene flow using pollen as a mediator, which may in turn can
1917 detect different strategies and local adaptations related to the rate of pollen germination under
1918 different conditions (Jhala, et al., 2010). It is therefore of interest, to look at pollen germination rate
1919 and pollen viabilities when looking at local adaptations as well as strategies of breeding systems. In
1920 the literature, study species such as cotton (*Gossypium hirsutum*) shows a high tolerance to heat.
1921 Optimum temperature for pollen growth and pollen tube length in cotton have been observed to be
1922 above 32°C (KAKANI et al., 2005). Other observations in temperate cultivars such as apricots and
1923 sweet cherries (*Prunus spp.*) suggests that pollen germination and pollen tube length are higher in
1924 some cultivars treated with colder temperature (5°C), although varying within different lines (Pirlak.,
1925 2002). Previous observations of pollen tubes were made in another agriculturally significant plant,
1926 peach (*Prunus persica*), by Hedhly A. et al (Hedhly & Herrero., 2008). This study found that drastic
1927 increase in temperature had a negative effect on stigmatic reception of the pollen tubes, thus
1928 significantly slowing pollen tube growth. These results suggest that stigmatic cells' ability to sustain
1929 and adhere the pollen tube cells were reduced because of drastic temperature effect and the
1930 kinetics of pollen tube growth was also affected. This behaviour was also observed in the field
1931 (Hedhly & Herrero., 2008). A similar negative effect of temperature on pollen kinetics and stigmatic
1932 cell adhesion was also previously observed by the same group in cherries (Hedhly et al., 2004).

1933 It has been determined that pollen germination and pollen tube growth could be observed
1934 *in vitro* and semi *in vivo* for the model *Arabidopsis thaliana* (Dickinson et al., 2018), making them
1935 versatile for observations in the laboratory or the wild. In angiosperms, the dynamics between the
1936 pollen and its pollen tube formation right into the ovaries represents some of the most
1937 environmentally sensitive part of their sexual reproduction. Temperature is considered an important
1938 variable, currently studied under various environmental research because it is a "fragile" climatic
1939 variable, and temperature represents one of the main environmental variables which could affect
1940 the health of living organisms, especially in plants (Reynolds & Casterlin, 1980) (Nievola et al., 2017)
1941 (Qaderi et al., 2019). Pollen tube growth dynamics have been highlighted to potentially be controlled
1942 by Ca²⁺ signalling and sensitive to heat stress, making it vulnerable to changes to the environment
1943 (Johnson et al., 2019). It has been suggested that Ca²⁺ signalling acts as a signal transduction factor

1944 under heat stress conditions in the model *Arabidopsis thaliana* (Xu et al., 2022). This may affect
1945 pollen tube dynamics in *Linum*.

1946 This part of the chapter will test several growing temperatures to establish whether pollen
1947 germination is affected by using quantification and analysis of pollen structures. One of the interests
1948 in pollen observation is whether different temperature conditions impact the performance of pollen
1949 tubes to reach the ovule and form a gamete. To do this, pollen tube would need to be able to travel
1950 down the style into the ovule which contains the female egg cells. Whether temperature presents a
1951 significant stress for pollen tubes to be able to reach the ovule is of important agricultural interest in
1952 crops including *Linum*. A further interest is if there would be a difference in species and population
1953 level across cultivated *Linum* (*L. usitatissimum*) and its wild relatives (*L. bienne*). Another interest in
1954 terms of pollen viability is whether different climate of origin variables is associated with pollen
1955 viability. For the purposes of the study of this chapter, the vulnerability to changes in temperature
1956 are hypothesized to be of key importance for seed set in *Linum*. Pollen viabilities were previously
1957 tested in *L. usitatissimum* and were found to be little affected by heat stress. However, findings
1958 suggest the formation of bolls and seeding were negatively affected by heat stress (CROSS et al.,
1959 2003). Temperature effectson pollen have not been tested in *L. bienne*. Other cultivated species
1960 showed a negative effect on pollen viability and germination post high temperature exposure (Aloni
1961 et al., 2001).

1962 In this study, temperature as an environment variable was evaluated against pollen viability
1963 in the wild and cultivated *Linum* samples which were also used across other studies in this thesis.
1964 Observations were made of pollen viability at various stages. These were pollen count, pollen tubes
1965 count, and whether pollen tubes were able to reach the ovaries as a sign of pollen gamete being
1966 “successfully” transferred. Different temperatures were tested *in vivo* for pollen germination
1967 strategies comprising heat and cold treatments as well as a control typical temperature treatment.
1968 The main hypothesis assessed in this chapter was that pollen viability is affected by temperature as
1969 an environmental variable in wild and cultivated *Linum*. We also expect wild and cultivated flax to
1970 show variability within their pollen viabilities under different treatments. With local adaptation in
1971 mind, we expect differences in pollen responses to cold and heat treatments between populations
1972 originating in warmer and colder climates. This may be presented as correlations between pollen
1973 viability and local climate variables. More Northern plants are expected to perform better under
1974 colder temperatures in terms of pollen viability and vice versa for Southern plants.

1975

1976

1977 3.2 STUDY MODELS AND METHODS

1978 Under this study, 51 individuals representing 18 wild populations, and 14 cultivar varieties
1979 were observed (appendix 10). These individuals were harvested from the vernalization experiments
1980 and so constitutes for S1 (first selfing generation). Fully grown plants were able to be used for pollen
1981 observations in their subsequent flowerings. The sample plants were tested using different
1982 controlled environments inside a Weiss Gallenkamp controlled chamber: models A3655 (Arctic) and
1983 A3658 (Tropical). Each of the chambers were connected to a main computer and were equipped
1984 with numerous fluorescent tube lighting units, which were “Philips Master TL5 HE”. These light tubes
1985 are 14 Watts in power requirements for each tube and emits 4000 Kelvin light temperature in colour
1986 by specifications. The artic chambers were used to provide colder temperatures while the tropical
1987 chambers were used to provide the warmer treatments. During the growing stages, plants were
1988 grown in a pot with F2S compost (Levington Advance Seed and Modular + Sand) combined with
1989 dried rice husks (3:1 ratio) for aeration purposes. The same 750ml black square plastic pots were
1990 used for each sampled individual. In each pot, five seeds from the same maternal individual were
1991 sown. Plants were grown under a controlled glasshouse condition (16:8 daylight ratio, 25°C daytime
1992 temperature and 13°C night-time temperature) until first flowering was observed.

1993 For the different temperature treatments, individuals were randomized under controlled
1994 glass-house conditions. Plants were grown until first flowering at 25°C during daytime before
1995 treatments. Plants were then treated with either a heat (+5°C) or a cold (-5°C) treatment under the
1996 corresponding tropical or arctic chambers as the first flower were observed. Plants were treated for
1997 at least 72 hours before data were collected, to allow for acclimatation. Post acclimatation, flowers
1998 were collected at an “open stage”. As described in Schewe *et al* 2011, flowers at an “open stage” are
1999 categorised as flower at anthesis, showing an opened 5-part whorl where the stamens are found
2000 with the stigma (stage 11 in Schewe *et al*, 2011) (Schewe, et al., 2011). Aniline blue staining protocol
2001 was utilized along with a confocal microscopy. Full protocol and information of the Aniline blue are
2002 available under section “3.2.3 Pollen Observation” .

2003 Whole flowers were harvested per individual and preserved in a 70% ethanol (70%EtOH)
2004 solution, inside separate 1.5ml Eppendorf tubes. Floweres were preserved at 4°C. Observations
2005 found 70%EtOH solutions to be adequate for floral and pollen preservation after 24-hour periods
2006 (appendix 11). Flowers were submerged in at least 1ml of 70% EtOH before storing in a 4°C fridge.
2007 Before microscopy, petals and anthers were removed. This resulted in the boll and the stigma
2008 exposure. Some errors in pollen measures are expected as pollens may or may not be washed out
2009 during preservation stages, but the same procedures were applied to all samples under all
2010 treatments. Counts of pollen, pollen tube and pollen reaching ovary were recorded over several

2011 weeks of flowering, using a combination of pollen staining and confocal microscopy techniques
 2012 (described below). Between the different temperature treatments, samples were randomly swapped
 2013 over a 72 hour rotation period with varying number of flowers at “open stage” able to be harvested
 2014 per-observation.

2015

2016 3.2.1 Pollen Viability *In Vitro*

2017 Studies have previously shown several ways to test pollen viability. These often consist of
 2018 staining by a fluorescent dye and assessing the presence of pollen and pollen tubes under a confocal
 2019 microscope, as performed previously in *Panicum* (Ge et al., 2011). Optimization of pollen
 2020 fluorescence microscopy is also of interest in the model *Arabidopsis* (Bou Daher et al., 2008) and
 2021 thus this study involving the non-model *Linum* will explore staining methods for fluorescent
 2022 microscopy.

2023 As a precursor to quantification, tests were done using *in vitro* germination under heat stress
 2024 which uses different germination media to *in vitro* germinate pollen from different temperature
 2025 treatments. At the aim was to find out whether *in vitro* germination using germination medium was
 2026 possible in *Linum* to determine heat stress (Rodriguez-Enriquez et al., 2012). In the model
 2027 *Arabidopsis thaliana*, it is suggested that *in vitro* germination is possible, with a varied germination
 2028 levels at different treatments, however pollen germination and tube growth were dependent on
 2029 pollen density in both liquid and solid medium (Boavida & McCormick, 2007). Recent past studies
 2030 have tested the effect of germination media with extra sucrose, and boric acid for *in vitro* pollen
 2031 tube germination in the non-model Chinese fir (Fragallah et al., 2019). Then, *in vitro* methods were
 2032 used to germinate different pollens under different environmental conditions. We assessed different
 2033 solutions in *Linum in vitro* germination. Different germination solutions were made using Sucrose as
 2034 the main sugar (Table 1).

| Components | Solution 1 | Solution 2 | Solution 3 |
|------------------------|------------|------------|------------|
| 12% (w/v) Sucrose | 50% | 40% | 60% |
| 5mM CaCl ₂ | 20% | 20% | 20% |
| 0.01% (w/v) Boric Acid | 15% | 20% | 10% |
| 1mM MgSO ₄ | 15% | 20% | 10% |

2035 Table 1. A table of the three different *in vitro* germination solution which were tested in the laboratory.

2036 The pollens were firstly place within the different solutions for at least 3 hours before observation.
 2037 To assess germination success, a control was also provided with just nuclease free water as the
 2038 solution. 20% Toluidine blue was used to dye the pollens after the treatments and pollens were
 2039 observed under a Leica DM500 microscope with a camera model MC190HD connected to a
 2040 computer running Leica’s LAS software. Results for the *in vitro* germination were not very convincing

2041 (see the results part of this chapter). This result would mean that *in vitro* germination procedures in
2042 *Linum* pollen needed progress. Therefore, it was decided that pollen would need to germinate *in*
2043 *vivo* on flower pistils, with measures at different pollen maturity stages to measure heat stress under
2044 different temperature conditions.

2045 For *in vitro* germination attempts, three solutions were prepared according to Table 1 and
2046 was adjusted to pH of between seven and eight. Three to five drops were put onto a glass
2047 microscope slide with a Pasteur pipette. Wild *Linum* flowers from control 25 C conditions that were
2048 still not opened (to avoid *in vivo* germination) were collected by cutting from the stem and all petals
2049 were carefully removed using forceps to expose the stigmas, and anthers. These parts were then
2050 dipped onto the solution on the microscope slides several times to release pollen into the solution.
2051 At least 3 Hours have passed to allow germination before the staining procedure was undertaken.
2052 For the stain, 20% Toluidine blue were prepared within a mixture of 50% nuclease free water and
2053 50% glacial acetic acid. Two drops of the dye solution were added to the slide and the pollen
2054 samples. Observation took place using a Leica DM500 microscope fitted with MC19HD camera,
2055 connected to a computer able to run Leica's LAS software. In this study 30×10 and 50×10
2056 magnifications were used for the pollen *in vitro* germination observations.

2057

2058 3.2.2 Pollen Treatments

2059 For the different temperature treatments, individuals were randomised under a controlled
2060 glass-house conditions before measurements under confocal microscopy. Plants were initially grown
2061 at 25°C during daytime. Plants were then treated with either a heat (+5°C) or a cold (-5°C) treatment
2062 under the corresponding tropical or arctic chambers as the first flower were observed. Plants were
2063 treated for at least 72 hours before further progress to allow for acclimatation. Post acclimatation,
2064 flowers were collected at an "open stage.

2065 Aniline blue staining protocol was deployed along with a confocal microscope for visual aid.
2066 Full protocol and information of the Aniline blue are available under the "3.2.3 Pollen Observation"
2067 part found below in this chapter.

2068 As described in *Schewe et al*, flower at an "open stage" are categorised as flower at anthesis,
2069 showing an opened 5-part whorl where the stamens are found with the stigma (stage 11) (Schewe,
2070 et al., 2011). Whole flowers were harvested and preserved with 70% EtOH inside separate 1.5ml
2071 Eppendorf tubes for each line. Two solutions were evaluated for pollen preservation. One solution is
2072 Formaldehyde-acetic acid alcohol (50ml of >95% ethyl alcohol, 2.5 ml of glacial acetic acid, 5.5 ml of
2073 formaldehyde, and 42 ml of dH₂O) (FAA), and the other solution was 70% ethanol with dH₂O

2074 (70%EtOH). Observations found both solutions to be adequate for floral and pollen preservation
2075 after 24-hour periods (appendix 12). After collection, flowers were submerged in at least 1ml of 70%
2076 EtOH inside a 1.5ml Eppendorf tube for preservation. Petals were removed as well as anthers which
2077 results in the boll and the stigma exposure. Some errors in pollen measures are expected as pollens
2078 may or may not be washed out during preservation stages, but the same treatments were applied to
2079 all samples under all treatments. Measures of pollen and pollen tube counts have been accumulated
2080 over several weeks. Between the different temperature treatments, individuals were randomly
2081 swapped, and different flowers were also measured per treatment.

2082 3.2.3 Pollen and Pollen Tube Observation

2083 For the purposes of pollen and tubes observations, a modified protocol based on the
2084 protocol by Lu Yongxian (Lu Y, 2011) were used. The fluorescent staining dye applied into stigmas to
2085 observe pollen number and observe pollen tubes was 0.5% (w/v) Aniline blue in 0.1M KH_2PO_4 .
2086 Glycerol was recommended in some fluorescent dye protocols, but it was not used in this study as
2087 fluorescence dye were observed better in non-glycerol solutions. Optimisation of aniline blue
2088 dilution was of interest, as too dilute will limit penetration into the pollen cells and too concentrated
2089 will act as a background noise, which may interfere with the images surveyed (Herburger &
2090 Holzinger, 2016). Preliminary optimisation found that 0.5% (w/v) Aniline blue in Potassium
2091 Phosphate (KH_2PO_4) provided images that were useful for pollen and pollen tubes observations
2092 (Appendix 13).

2093 Bolls and stigma attached were examined onto a 25mm x 75mm glass microscope slide with
2094 a thickness of $\leq 1.2\text{mm}$. 2-3 drops of 0.5% aniline blue were sufficient for dyeing purposes. In theory
2095 aniline blue can penetrate callose better than other cells (Herburger & Holzinger, 2016), and
2096 therefore more sensitive towards pollens and pollen tubes. This made it easier to distinguish the two
2097 pollen tissues from other cell types and to quantify them. Samples were covered using 22mm x
2098 22mm cover slip with a thickness of 0.1mm to allow for optimal viewing under a confocal
2099 microscope. Samples were stored at room temperature for >3 hours prior to viewing to allow for the
2100 dye to penetrate the callous cells. The observations were also done in a dark room with as little light
2101 as possible.


2102 A Zeiss model 880 Confocal Microscope was used to observe fluorescence and presence of
2103 pollen and pollen tube structures with fluorescence absorbance between 400-500nm for aniline blue
2104 absorption (Yang et al., 2007) and images were further processed by Zeiss' own software (ZEN Black
2105 edition) to optimise image settings. Zeiss' 10 \times magnification and Zeiss' 20 \times magnification eye pieces
2106 all combined with a 10 \times eyepiece was used to observe pollen and pollen tubes. Different focal layers

2107 were examined and in cases where pollens could not be observed under one layer of focus, a 3D
2108 image with appropriate number of Z-stacks was acquired to scan through all axes.

2109

2110 3.2.4 Pollen Counts

2111 *Linum* pollen structures were circular vessel like grains and have high callose content and
2112 therefore would be highlighted more intensively by the aniline blue dye than surrounding cells. We
2113 observed no differences in structure for the wild and cultivar species of *Linum* observed for this
2114 study. The pollen was expected to be found adhered around the stigma. Pollen tube structure were
2115 identified as thread like structures coming out of the pollen tube and penetrating the stigma, often
2116 in a straight line down to the ovary (Figure 2, next page).



2117

2118 Figure 2. A. 10×20 magnified stigma, fluoresced with Aniline blue illustrating an absence of pollen, B. the presence of
 2119 mature pollens on the stigmas, conveyed as blue circular shapes C. more fluorescent callose makes pollen more visible in
 2120 different colour ranges. D. A 10×20 magnified 3D depiction of stigma and pollens. E. A 10×20 magnified, and labelled image
 2121 of pollen structures observed.

2122 Absence of pollen and pollen tube structures can be observed in some cases under fluorescent
2123 microscopy (Figure 2A). In other cases, presence of pollen and pollen tubes at the stigma were also
2124 be observed (Figure 2B-E). Pollen were rounded in shape and fluoresce brighter than the stigma
2125 cells. The pollen tubes were string-like structures growing out of pollen into the stigma (Figure 2E).

2126 Images were further processed in Zeiss' Zen Black software to apply colouration to highlight
2127 fluorescence absorption at 400-500nm wavelength and increase contrasts between the pollen and
2128 pollen tube cells and surrounding stigma/style cells. Callose in pollen and pollen tubes can be seen
2129 as more fluorescent (Figure 2C). A 3-Dimensional observation can also be made using the Z-stacks
2130 function of Zeiss' Zen Black software (Figure 2D). This feature enabled layers to be taken stack by
2131 stack under a new Z dimension, thus building a 3-dimensional image. This tool represents an aid, as
2132 pollen and pollen tubes may have not been fully observed on a 2-dimensional image. After
2133 optimisation for each slide, sample images were captured, and pollen and pollen tubes were
2134 manually counted to build a dataset.


2135

2136 3.2.5 Pollen Tube Counts

2137 Observation was made for three different temperature treatments to observe whether
2138 pollen tubes were able to reach the ovule at the open flower stage. Aniline blue staining and
2139 confocal microscopy methods were used as before, but for this observation, we observed pollen
2140 tubes inside pistils and stigma cells. Colouration of the aniline blue were referred to "cyan" and
2141 contrast were increased with using Zeiss' Zen Black software. For stigma and styles with little pollen
2142 tube growth, wavelength was contrasted from fluorescent green to blue, highlighting dyed cells
2143 fluorescing green. The processed images could then be observed to see whether pollen tubes were
2144 able to reach down to the bottom of the style and reach the ovule. This was observed for each of the
2145 temperature treatments and recorded in binary, 0 representing pollen tube not reaching ovary and 1
2146 representing pollen tube reaching ovary for every individual tested. The ovary as well as the pollen
2147 tube fluoresce when dyed with the Aniline blue. An observation of a pollen tube structure going into
2148 the ovary at the bottom of the pistils would suffice the question of whether the pollen tube have
2149 managed to reach the ovary or not. The resulting difference in images with pollen tubes and without
2150 pollen tubes could be observed on the Figure below:

2151

2152



2153 Figure 3. A. 10×20 magnification of a flower pistil with observed pollen tube growth fluorescence highlighted in cyan. B.
2154 10×20 magnification of fluoresced ovary cells highlighted in “fluorescent green”, depicting no pollen tubes going into the
2155 ovary.

2156

2157 Figure 3 compares 10×20 microscopic images between individuals with pollen tubes growing
2158 through the pistils and ovaries, and individuals with no observable pollen tube in the ovaries. Pollen
2159 tubes growth were highlighted in cyan when dyed with aniline blue (Figure 3A). Long thread like
2160 structure could be seen going through the pistils into the ovary parts of the flower. Both pollen tube
2161 cells and ovary cells were highlighted and dyed in aniline blue as observed here. Absence of ovary
2162 cells can be observed just after the end of the style, which were highlighted in fluorescent green
2163 (Figure 3B). This reveals that for this individual, no pollen tube was observed going into the ovary
2164 structure and therefore pollen tube did not successfully reach the ovary. As an additional measure,
2165 this proportion of pollen forming pollen tubes was also calculated for further analysis.

2166

2167

2168 3.3 ANALYSIS

2169 ANOVA (Girden ER, 1992), linear modelling (Yan and Su, 2009) and tests for binomial models
2170 for this study were undertaken using the statistical programme R (R Core Team, 2022). The graphical
2171 interface used to run R was RStudio (Rstudio Team, 2020). Raw data for pollen counts, pollen tube
2172 counts as well as whether pollen tubes were able to reach the ovary for each of the treatment were
2173 firstly processed and organised in Microsoft Excel and converted to a tab-delimited CSV format for R
2174 to read. Raw data exist in both binomial and count format. Each of the linear modelling stages need
2175 to take account the data format in such cases.

2176

2177 3.3.1 Analysis: Pollen and Pollen Tube Counts

2178 Analysis was performed to determine whether pollen and pollen tube counts for treatments
2179 differ significantly. First, the data was checked for normal distribution with Shapiro-Wilk's normality
2180 tests (SHAPIRO & WILK, 1965). This test uses a modified analysis of variance, to calculate a *p-value*. A
2181 histogram of the data counts was also observed to show whether the data is normally distributed.

2182 General linear modelling (GLM) was utilized to test the effect of changing temperature on
2183 the number of pollen counts (McNeil *et al*, 1996). For non-fractional count data, Poisson's regression
2184 was used (Hayat & Higgins, 2014) with "family=poisson()" in the GLM command under R. A Poisson
2185 Regression model is a Generalized Linear Model (GLM) that is used to model count data (Hayat &
2186 Higgins, 2014). The output Y (count) is a value that follows the Poisson distribution. The model
2187 assumes logarithm of expected values (mean) that can be modeled into a linear form by some
2188 unknown parameters. For a post-hoc analysis comparing pairs of treatments, a Tukey's adjustment
2189 was added to the general linear model by using the R package "Multcomp" (Hothorn et al., 2008).
2190 The full command is available under appendix 14. Data were summarized using a bar chart for the
2191 pollen count against different treatments with the "barplot" command in R. For interactions
2192 between treatment and species, an R function "interaction()" can be used to interpret interactions
2193 between two categorical variables (Chambers and Hastie, 1992) to be integrated into the linear
2194 modelling.

2195

2196 3.3.2 Analysis: Proportion of Pollen Tubes

2197 The number of pollen tubes relative to the total number of pollen counts were tested for
2198 interaction against the different treatments and between the two wild and cultivated species within
2199 the *Linum* samples. Data was modelled using general linear model with a quasibinomial function

2200 using the R package “lmerTest” (Kuznetsova et al., 2017). A ‘logit’ link was added using the command
2201 “(link = 'logit')” as an addition to the quasibinomial family specification which attempts to describe
2202 additional variance in the data that cannot be explained by a Binomial distribution alone. After the
2203 linear modelling, analysis of variance (ANOVA) for the different variables (Girden ER, 1992) was
2204 applied to the model in R using the command “anova(model,test=F)”. Interaction between
2205 treatments and species was measured using the “interaction()” command of R. The Tukey’s contrast
2206 treatments were added to the model using the R package multcomp (Hothorn et al., 2008), using the
2207 command: “model<-glht('modelprefix',mcp(interact="Tukey"))”. The full model commands can be
2208 found under appendix 14.

2209 To evaluate whether pollen tube was able to reach the ovary, a binomial “yes or no” test
2210 was utilized. To test for this, a general linear model with a binomial adjustment was run in R with the
2211 function “family=binomial(link='logit')” added to the general linear model command (Gelman, A. and
2212 Hill, J., 2006).

2213

2214 3.3.3 Analysis: Population of Origin Variables

2215 To consider environmental effect, we gathered geographical data for the wild *L. bienne*.
2216 Climatic variables from the origin of the wild individuals were gathered from WorldClim database in
2217 *Chapter 2* of this thesis. Latitudinal and climatic data was only reliably available for the wild *L. bienne*
2218 samples. This is because location data for our cultivars was not precise. Therefore, for the analysis
2219 against local variables, only wild *L. bienne* individuals were considered. Climatic variables were
2220 processed into summary principle components (PCs) as was described previously in chapter 2 (see
2221 section 2.2.1 “Samples and Experiment”). Latitude and climatic variables were used to further
2222 describe temperature variables of the local areas which the wild *L. bienne* were collected from. More
2223 Northern origin plants are expected to perform better under colder temperatures in terms of pollen
2224 viability and vice versa for Southern origin plants. The loadings of the principal component analysis
2225 are as in appendix 3. The temperature loadings of climate PCs will be of particular interest in this
2226 chapter. Since the temperature loadings were positive, a higher PC1 value indicated a higher
2227 temperature. For example, the PC1 value for the Southern population ‘3’ is 5.367 while the PC1
2228 value for the Northern population ‘Tym’ is -6.616. The average temperature loading for population
2229 ‘3’ collected from the WorldClim database is at 12.07°C in winter (December, January, February)
2230 while for population ‘Tym’ it is 5.02°C. For the summer (June, July August) the average temperature
2231 for population ‘3’ is 22.6°C and for population ‘Tym’ it is 15.05°C. A linear model with a Pearson’s
2232 correlation coefficient can be used to test whether latitude of locality and climatic variables affects

2233 either pollen count, pollen tube counts and or the proportion of pollen able to reach the ovary. A
2234 scatterplot of relationships was generated using the “ggscatter” command of the package ggplot2
2235 built for R (Villanueva & Chen, 2019).

2236

2237


2238 **3.4 RESULTS**

2239

2240 **3.4.1 Results: *In vitro* pollen germination**

2241 The processed microscopic images reveal that in all cases of the *in vitro* solutions 1-3, the
2242 pollen structure burst. We found that, the *in vitro* treatments of the *Linum* pollens were not up to
2243 standard and resulted in pollen materials bursting. Comparison between pollen treated in nuclease
2244 free water and the germination solution 1 can be seen (Figure 4A-C). A 500x magnification of one of
2245 the sites of the aperture pore whereby bursting occurred were also observed (Figure 4C). This
2246 bursting was observed using all germination solutions.

2247



2252 Figure 4. A. 10×50 cropped magnification of a mature pollen dyes with 20% Touludine blue, observed in nuclease free
2253 water and no added germination solutions. B. A 10×20 magnified image of a 20% Touludine blue dyed pollen cell which is
2254 undergoing bursting from all the aperture pore sites due to the germination solutions. C. 10×50 magnification of one of the
2255 aperture pore sites of a recently burst pollen.

2256


2257 **3.4.2 Results: Test for Normality**

2258 As part of the analysis against raw pollen/pollen tube counts, tests were done to reveal
2259 whether the data is normally or abnormally distributed, to inform the statistical analysis which could
2260 be used against the data. For normality, a histogram of the raw data distribution should peak around
2261 the average with decreasing number of raw counts either side of the data to form what is widely
2262 known as a bell curve when a line of best fit is applied to the histogram. A histogram of the raw data
2263 was able to be illustrated with a non-normal distribution (Pollen count, % of pollen tubes) (Figure 5).

2264 A Shapiro-Wilks test was applied to the raw data to statistically quantify normality revealed
2265 that the data is not normally distributed and therefore adjustments were made to statistical analysis
2266 tests to take this non-normality into account by using the appropriate Poisson or quasibinomial
2267 distribution family (pollen counts: $W = 0.89365$, $p\text{-value} = 4.099e-14$ and % pollen tubes $W =$
2268 0.77881 , $p\text{-value} < 2.2e-16$).

2269

2270




2276 Figure 5. Histograms and density curves for the data distribution of both pollen tube % and pollen count showing a non-
2277 normal distribution.

2278

2279 3.4.3 Results: Pollen and Pollen Tubes Count

2280 As a matter of ease, the treatments were all respectively known as 20°C (Cold), 25°C
 2281 (Glasshouse), and 30°C (Heat) from here on. Summary of all the data were illustrated in a barchart
 2282 (Figure 6). Both treatments (heat and cold), revealed a significant reduction on the number of pollen
 2283 count (Figure 6 and Table 2). Only the heat treatments suggested significant difference in terms of
 2284 tube count (Figure 6 and Table 3).



2285

2286 Figure 6. A bar chart of the overall average pollen data for the different heat treatments. Glasshouse treatments are
2287 highlighted in blue, heated treatment in orange and cold treatment in grey.

2288

2289 Pollen count of each treatment were statistically compared to check for effects on the
 2290 number of pollens due to temperatures according to the hypothesis that the temperature affects the
 2291 number of viable pollens within both wild and cultivated *Linum*. There was variation in the average
 2292 pollen count when comparing the three treatments (Figure 6). To statistically test for this variation,
 2293 we ran an ANOVA with a Tukey’s post-hoc for multiple comparisons and we illustrated a model using
 2294 a Poisson’s regression model formula of “Pollen\$Count ~ Pollen\$Treatment” for the data comprising
 2295 all the samples within this study. The analyses suggested that there was significant decrease
 2296 between the pollen counts in the glasshouse (25°C) compared with both the cold (20°C) and heat
 2297 (30°C) treatments (Table 2 below). This suggests that the treatments have a significant effect on the
 2298 number of pollens within the samples tested in this study.

2299

Linear Hypotheses:

| | <i>Estimate</i> | Estimate | Std. Error | z value | Pr(> z) | |
|------------------------|-----------------|-----------------|-----------------|---------------|------------------|----------|
| 25C vs 20C == 0 | | 0.104605 | 0.01972 | 5.305 | <1e-05 | * |
| 30C vs 20C == 0 | | 0.005315 | 0.021423 | 0.248 | 0.967 | |
| 30C vs 25C == 0 | | -0.09929 | 0.019749 | -5.028 | <1e-05 | * |

2303

2304 Table 2. A linear model with Tukey’s correction for pollen count summary between the different temperature treatments.
 2305 Significance ($p < 0.05$) was showing for a linear hypothesis between the 25°C – 20°C and the 30°C – 25°C treatments.

2306

2307 To see whether the same could be seen in the number of pollen tubes, average pollen tube
 2308 count was also tested for each treatment (Figure 6). When statistically compared, the number of
 2309 pollen tube on average between the treatments decreased with both cold and heat treatments. Just
 2310 as were seen in the pollen count dataset. In addition, the number of pollen tubes seem to be
 2311 significantly reduced ($p < 0.05$) for the heat (30°C) and cold (20°C) treatments (see Table 3).

2312

Linear Hypotheses:

| | Estimate | Std. Error | z value | Pr(> z) | |
|------------------------|-----------------|----------------|----------------|------------------|----------|
| 25C vs 20C == 0 | 0.3805 | 0.04108 | 9.263 | <1e-08 | * |
| 30C vs 20C == 0 | -0.31057 | 0.05121 | -6.065 | <1e-08 | * |
| 30C vs 25C == 0 | -0.69108 | 0.04588 | -15.064 | <1e-08 | * |

2313 Table 3. A linear hypothesis with a Tukey’s adjustment for the number of pollen tubes observed between the different
 2314 treatments. The test shows significant differences between all the treatments ($p < 0.05$).

2315

2316 We observed variation in pollen tube count and pollen count when looking at different
 2317 temperature treatments. However, in relation to pollen count we wanted to see if there is variation
 2318 in the proportion of pollen tubes able to be formed under our temperature treatments. To observe
 2319 whether the proportion of pollen tubes to pollen count is substantially affected by the temperature
 2320 treatments, the average proportions of pollen tubes formed against pollen for each temperature
 2321 treatments were calculated and modelled. The results suggests that against the glasshouse (25°C)
 2322 treatments, the cold (20°C) treatments showed a decrease in the proportion of pollen forming pollen
 2323 tubes but that the difference was not significant (Table 4). Just as the cold (20°C) treatments, the
 2324 heat (30°C) treatments showed a decrease in the proportion of pollen tube forming against the
 2325 pollen count (Table 4). This decrease in proportion, however, was shown to be significantly different
 2326 to the glasshouse (25°C) treatments when modelled against each other ($p = <0.001$) (Table 4). This
 2327 suggest that warmer treatments could potentially affect proportion of pollen tube able to be formed
 2328 in relation to the pollen count. Below is a linear hypothesis summary (with Tukey’s modification) for
 2329 the proportion of pollen tube formed against the number of pollens for each treatment.

2330

Linear Hypotheses:

| | Estimate | Std. Error | z value | Pr(> z) |
|-------------------------------|----------------|---------------|---------------|--------------------|
| <i>25C vs 20C == 0</i> | 0.2759 | 0.1184 | 2.331 | 0.0511 |
| <i>30C vs 20C == 0</i> | -0.3159 | 0.1442 | -2.19 | 0.072 |
| <i>30C vs 25C == 0</i> | -0.5918 | 0.1298 | -4.561 | <0.001 * |

2331 Table 4. Linear hypothesis with a Tukey’s adjustment for the proportion of pollen tube forming, showing significance
 2332 between 30°C and 25°C treatments ($p < 0.05$).

2333 To summarize the linear hypothesis significance for section 3.4.3 “Results: Pollen and Pollen
 2334 Tubes Count”, we illustrated the following Table:

| Treatment | Pollen Count | Pollen Tube Counts | Proportion of Pollen tubes formed |
|-------------------|-----------------|--------------------|-----------------------------------|
| 25C vs 20C | SIGNIFICANT | SIGNIFICANT | NOT SIGNIFICANT |
| 30C vs 20C | NOT SIGNIFICANT | SIGNIFICANT | NOT SIGNIFICANT |
| 30C vs 25C | SIGNIFICANT | SIGNIFICANT | SIGNIFICANT |

2335 Table 5. Summary of linear hypothesis tests significance between different temperature treatments and counts as per
 2336 section 3.4.3 “Results: Pollen and Pollen Tubes Count”.


2337

2338 **3.4.4 Results: Pollen Tube Reaching the Ovary**

2339 To test a model of whether any of the three treatments were significantly different in the
 2340 number of pollen tubes able to reach the ovary, a binomial general linear model (GLM) was
 2341 performed for “yes or no”, binomial data under the different treatments. The model was
 2342 summarised using a Tukey’s adjustment for abnormally distributed data (see section “3.3.2 Analysis:
 2343 Proportion of Pollen Tubes”). The results reveal that the percentage of pollen reaching the ovary is
 2344 significantly different in the 30°C treatments (see Figure 7 and table 6).

2345 The results from the general linear model suggests that between the glasshouse (25°C) and
 2346 the cold (20°C) the difference in the proportion of pollen reaching the ovary was not significantly
 2347 different ($p = 0.85$). This being the case, significant difference in the number of pollen tube reaching
 2348 ovaries could be observed between the heated (30°C) treatment and both the glasshouse (25°C) and
 2349 cold (20°C) treatments ($p < 0.001$ for both) (see table 6).

2350



2351 Figure 7. Bar chart showing the proportion of pollen reaching ovary according to the different temperature treatments.

2352

Linear Hypotheses:

| <i>Treatments</i> | Estimate | Std. Error | z value | Pr(> z) | |
|-------------------|----------------|--------------|---------------|-------------------|----------|
| <i>25C vs 20C</i> | 0.2126 | 0.399 | 0.533 | 0.85457 | |
| 30C vs 20C | -1.3132 | 0.357 | -3.678 | 0.00067 | * |
| 30C vs 25C | -1.5258 | 0.346 | -4.41 | < 1e-04 | * |

2353

2354 Table 6. General linear model with Tukey’s adjustment for the proportion of pollen tubes reaching ovaries. The test shows
 2355 significant differences between the temperature treatments of 30°C – 20°C and 30°C – 25°C ($p < 0.05$).

2356


2357 **3.4.5 Results: Pollen Tube Proportion Over Pollen Count for Different *Linum* Species**

2358 Additionally, we tested pollen tube proportion and pollen count to observe variation between
2359 species. Initial ANOVA revealed that when taking into consideration the interaction between
2360 treatment and species, the pollen tube proportion is different ($p = 0.0007$ ($p < 0.05$)). The proportion
2361 of pollen tube were linearly modelled with logistic distribution against the interaction between
2362 treatment and species. Data for either wild and cultivated species were separated and analysed with
2363 a linear model hypothesis separately. Bar plots illustrates a reduction in the proportion of pollen
2364 forming pollen tubes when heat treated for both *L. bienne* and *L. usitatissimum* (Figure 8 A and 8B).
2365 The proportion of pollen tube formed was found to be highest in the glasshouse (25°C) treated for the
2366 cultivars *L. usitatissimum*. This was reduced when treated under the cold (20°C) treatment, but not as
2367 much as when heat (30°C) treated. This wasn't the case when looking at *L. bienne*. When cold (20°C)
2368 treated the proportion of pollen tube formed in *L. bienne* did not decrease.

2369 We inferred a linear model summary when considering each species for treatments against each
2370 other. This was done with a Tukey's post-hoc correction (tables 6A and 6B). The proportion of pollen
2371 tubes saw the largest difference when looking at the heat (30°C) treated *L. usitatissimum* against the
2372 glasshouse (25°C) treated *L. usitatissimum* ($p < 0.001$). This was followed by the glasshouse (25°C)
2373 treated *L. usitatissimum* and the cold (20°C) treated *L. usitatissimum* ($p > 0.001$). *L. bienne* showed a
2374 statistically non-significant difference between the glasshouse and the cold/heat treatments when
2375 modelled (25°C – 20°C Treatments $p = 0.4556$, 25°C – 30°C Treatments $p = 0.4244$). This illustrates that
2376 the proportion of pollen able to form pollen tube is significantly affected by changes in temperature
2377 in the cultivar *L. usitatissimum* than in their wild relative.

2378


2379 **A – *L. bienne***



2380

2381 **B – *L. usitatissimum***

2382



2383 Figure 8. Bar chart to represent the proportion of pollen tube forming against temperature treatment for the wild relatives

2384 *L. bienne* (8A) and the cultivar *L. usitatissimum* (8B).

2385

2386 **A– *L. bienne* Summary**

Linear Hypotheses:

| <i>Treatments</i> | Estimate | Std. Error | z value | Pr(> z) |
|-------------------|----------|------------|---------|----------|
| <i>25C vs 20C</i> | -0.158 | 0.1323 | -1.194 | 0.4556 |
| <i>30C vs 20C</i> | -0.3307 | 0.1517 | -2.179 | 0.0743 |
| <i>30C vs 25C</i> | -0.1727 | 0.1385 | -1.247 | 0.4244 |

2387 **B – *L. usitatissimum* Summary**

Linear Hypotheses:

| <i>Treatments</i> | Estimate | Std. Error | z value | Pr(> z) |
|--------------------------|----------------|---------------|---------------|---------------------|
| <i>25C vs 20C</i> | 0.6084 | 0.1892 | 3.217 | 0.00372 * |
| <i>30C vs 20C</i> | -0.346 | 0.2434 | -1.422 | 0.32649 |
| <i>30C vs 25C</i> | -0.9544 | 0.2178 | -4.382 | < 0.001 * |

2388

2389 Table 7 A and B. Summary tables of the linear model with Tukey’s adjustments for the proportion of pollen tube formation

2390 between each *L. bienne* (A) and *L. usitatissimum* (B). No significant differences were observed between the different

2391 treatments for *L. bienne*. There were significant differences between the 25°C – 20°C and the 30°C – 25°C treatments for the

2392 *L. usitatissimum* ($p < 0.05$).

2393

2394 To summarize the linear hypothesis for section 3.4.4 Results: “Pollen Tube Reaching the
 2395 Ovary” and section 3.4.5 “Results: Pollen Tube Proportion Over Pollen Count for Different *Linum*
 2396 Species”, a Table is illustrated in below (Table 8):

| Treatments | proportion of pollen reaching ovary (<i>L. bienne</i>) | proportion of pollen tube forming (<i>L. bienne</i>) | proportion of pollen tube forming (<i>L. usitatissimum</i>) |
|------------|--|--|---|
| 25C vs 20C | NOT SIGNIFICANT | NOT SIGNIFICANT | SIGNIFICANT |
| 30C vs 20C | SIGNIFICANT | NOT SIGNIFICANT | NOT SIGNIFICANT |
| 30C vs 25C | SIGNIFICANT | NOT SIGNIFICANT | SIGNIFICANT |

2397 Table 8. Summary of linear hypothesis tests significance between different temperature treatments, ability of pollen
 2398 reaching ovary and proportion of pollen tube forming for each species.

2399


2400 3.4.6 Results: Propotion of Pollen Tubes Reaching Ovary for Different *Linum* Species

2401 We evaluate the ability of pollen tube reaching the ovary for the different species. Evaluation
 2402 of whether pollen tube was able to reach the ovary was applied as described under section “3.3.2
 2403 Analysis: Proportion of Pollen Tubes” of this chapter. An initial Anova, revealed that the interaction
 2404 between species and the treatments together were not significant (Treatment/Species $p = 0.2684$).
 2405 The bar plot below (see Figures 9A) reveals a reduction in the proportion of *L. bienne* pollen able to
 2406 reach the ovaries, under the heat (30°C) treatment.

2407 Summaries of the general linear models reveals that for both wild and cultivated species, the
 2408 proportion of individuals with pollen tube able to reach the ovary is significantly affected by the heat
 2409 treatments (30°C) (Tables 9A (*L. bienne*) and 9B (*L. usitatissimum*)). This was significant when
 2410 compared against the cold (20°C) treatment (for *L. bienne* $p < 0.00179$, for *L. usitatissimum*
 2411 $p = 0.00195$), and the glasshouse (25°C) treatments (for both species $p < 0.001$ ($< \alpha$ of 0.05)). The
 2412 result also shows some reduction on the proportion of individuals with pollen tube able to reach the
 2413 ovary when observing the glasshouse and the cold treatments. However, this difference isn’t
 2414 statistically significant ($p > 0.05$ in both cases).


2415

2416 **A – *L. bienne***



2417

2418 **B – *L. usitatissimum***



2419 Figure 9 . Bar chart for the proportion of pollen tube reaching the ovaries between different treatment for *L. bienne* (A) and
2420 *L. usitatissimum* (B).

2421

2422

2423

2424

2425 **A – *L. bienne* Summary**

Linear Hypotheses:

| <i>Treatments</i> | Estimate | Std. Error | z value | Pr(> z) |
|--------------------------|----------------|---------------|---------------|---------------------|
| <i>25C vs 20C</i> | 0.2034 | 0.6393 | 0.318 | 0.94515 |
| <i>30C vs 20C</i> | -1.7599 | 0.5135 | -3.427 | 0.00179 * |
| <i>30C vs 25C</i> | -1.9633 | 0.5445 | -3.605 | < 0.001 * |

2426

2427 **B – *L. usitatissimum* Summary**

Linear Hypotheses:

| <i>Treatments</i> | Estimate | Std. Error | z value | Pr(> z) |
|--------------------------|----------------|---------------|---------------|---------------------|
| <i>25C vs 20C</i> | 0.2034 | 0.6393 | 0.318 | 0.94515 |
| <i>30C vs 20C</i> | -1.7599 | 0.5135 | -3.427 | 0.00195 * |
| <i>30C vs 25C</i> | -1.9633 | 0.5445 | -3.605 | < 0.001 * |

2428 Table 9A and 9B. Summary tables of the general linear model with Tukey’s adjustment for the proportion of pollen tubes
 2429 reaching the ovary between the different treatments for the different species. In both species the test shows a significantly
 2430 reduced proportion of pollen reaching the ovary ($p < 0.05$).

2431

2432 To summarize our results for section 3.4.6 “Results: Propotion of Pollen Tubes Reaching
 2433 Ovary for Different Linum Species”, a Table is illustrated below (Table 10):

| Treatments | proportion of pollen tube reaching the ovaries (<i>L. bienne</i>) | proportion of pollen tube reaching the ovaries (<i>L. usitatissimum</i>) |
|-------------------|--|---|
| 25C vs 20C | NOT SIGNIFICANT | NOT SIGNIFICANT |
| 30C vs 20C | SIGNIFICANT | SIGNIFICANT |
| 30C vs 25C | SIGNIFICANT | SIGNIFICANT |

2434 Table 10. Summary of linear hypothesis tests significance between different temperature treatments and proportion of
 2435 pollen tube reaching the ovary for the two species.

2436

2437 **3.4.7 Results: Pollen and Latitude of Origin**

2438 For the wild *L. bienne*, there was interest in looking at whether locality of origin of the wild
 2439 plant had any roles to play on either the number of pollen count, the number of pollen tube observed
 2440 and or the proportion of pollen tubes reaching the ovary. This is of interest, as this would add to the
 2441 argument that local adaptation is present within our samples. Using Person’s correlation and a linear



2442 modelling (represented by the blue line in Figure 10A), pollen count had no correlation with the
2443 latitude of locality of the different wild *L. bienne* samples within the study.


2444 The number of pollen count in the glasshouse and heat-treated plants seems to increase, the
2445 more northern an individual is localized (For glasshouse ($R=0.024$, $p=0.86$), for heat ($R=0.1$ $p=0.45$)), but
2446 none of the correlations were significant (*all p values = >0.05*).

2447 Both glasshouse and cold treated individuals showed an increase in the number of observed
2448 pollen tubes, the more northern they are localized, but only the cold treatment was significantly
2449 correlated ($R=0.24$, $p=0.085$ (glasshouse), $R=0.44$ $p=0.001$ ($p<0.05$)) (Figure 10B). This illustrates
2450 that, for our *L. bienne* individuals, northern individuals are more likely to form more pollen tube
2451 under cooler temperatures. When observing the heat treatment, it can be observed that correlation
2452 is opposite to that of the glasshouse and cold treatments ($R=-0.16$ $p=0.23$), all be it not significantly
2453 correlated. This change in correlation trend, shows some signs of the Northern individuals in this
2454 sample test, being more sensitive in their pollen tube formation to increases in temperature. This
2455 illustrates that the more Northern an individual within our *L. bienne* sample is localized, the more
2456 likely that they would favour cooler temperature for pollen tube formation.

2457 In Figure 10C, we plotted the proportion of pollen tube observed over the pollen count (in
2458 percentage) and against latitude. The trend illustrates the positive correlation in the cold treatment,
2459 of pollen tube amount against latitude is even more significant when considering the proportion of
2460 pollen tube able to form against the pollen count per individuals ($R=0.58$, $p<0.0001$). This
2461 strengthens the point that the more Northern an individual is, the more able they are to form pollen
2462 tubes under the cold treatments.

2463





2466

2467 Figure 10. Pearson's correlations between pollen counts (10A) and pollen tube counts (10B) and pollen tube to pollen
 2468 count proportion (10C) in relation to the latitude of localities. The only significant correlation seen was between pollen
 2469 tube count and latitude of localities as seen in Figure 10B. Pollen tube counts under cold treatment ($R=0.44, p=0.001$). Fig
 2470 10C. Represents the proportion of pollen tube (%) to pollen count against latitude of individuals representing correlation
 2471 under cold treatment ($R=0.58, p=0.000006$).

2472 3.4.8 Results: Pollen and Local Climate

2473 With the latitude of localities showing significance on the ability of pollen to form pollen
2474 tubes under the cold treatment, it is possible to link pollen viability with climatic variables, especially
2475 variables such as average temperature. We suggest that more Northern individuals are varying
2476 significantly in terms of pollen tube formations to the Southern individuals. We used summary PC for
2477 climate variables from chapter 2 to compare against pollen variables and climate on our wild
2478 samples.


2479 The number of pollen counts observed (Figure 11A) illustrates that there was no significant
2480 correlation between the pollen count and the principle component 1 (pc1) of the climatic variables
2481 ($p > 0.05$ for all temperature treatments). This agrees with the findings for pollen count against
2482 latitude (see above section) whereby pollen count doesn't correlate with latitude of locality. The
2483 number of pollen tubes, reveals a difference in trend against the cold and glasshouse treatments
2484 compared with the heat treatment. With higher levels of PC1 observed, the less pollen tube is
2485 observed within the cold and the glasshouse treatment while under the heat treatment, the higher
2486 the level of PC1 for climatic variables the more pollen tube observations were made. However, just
2487 like under the findings against the latitude model, only the cold treatment showed a significant
2488 decrease in the number of pollen tubes ($R = -0.41$ $p = 0.0026$).

2489 More significant correlations between pollen tube proportions and climatic variables can
2490 also be summarised by PC1 (Figure 11C). The correlations reveal the proportion of pollen tube
2491 decreases more significantly with an increase in the PC1 value under the cold treatment ($R = -0.54$,
2492 $p < 0.0005$). However, in contrast with the pollen tube count alone; when considering pollen tube
2493 formation in relation to the pollen count, the glasshouse treatment also shows a significant decrease
2494 in proportion in the event of an increase in PC1 value, a trend which was not significant under pollen
2495 tube count only. This reveals that proportion (in percentage) of pollen tube able to form were
2496 significantly correlated with climatic variables in the form of PC1 under both cold and glasshouse
2497 treatments. There was still a positive trend when considering the heat treatment. However, this
2498 remained insignificant ($R = 0.12$, $p = 0.42$).


2499 A higher level of PC1 reveals a loading of warmer average temperatures for the local areas
2500 where the population were collected from in the wild (see section "3.3.3 Analysis: Population of
2501 Origin Variables"). Therefore, the significant decrease in the number of pollen tube count against a
2502 higher PC1 value reveals that at least in the cold treatment, the number of pollen tube able to be
2503 formed in our wild samples were significantly correlated with the higher temperature of their local
2504 climate. This result illustrates that under cold temperature treatments, individuals that are localised

2505 to a warmer climate are less able to form pollen tubes than that of individuals originating from
2506 colder climate.

2507



2508



2509

2510

Figure 11. Pearson's correlations and linear modelling between pollen counts (A) and pollen tube counts (B) and climate PC1 values. The only significant correlation seen was between pollen tube count and PC1 values at cold temperature as seen in Fig 17 B ($R = -0.41, p = 0.0026, < \alpha$ of 0.05). Pearson's correlations and linear modelling between pollen tube proportion (in percentage) to the count against the PC1 climatic variable were also illustrated (Figure 11C).

2514

2515

2516 3.5 DISCUSSION AND CONCLUSION

2517 For pollen germination, *in vivo* germination worked better than *in vitro* germination. This is
2518 because the solutions and conditions that were tested under this study did not provide an ideal *in*
2519 *vitro* condition for *Linum* pollen germination. This caused the pollen's content to "burst out" of the
2520 pollen exine and thus making pollen germination impossible. This "bursting" was perhaps due to the
2521 osmotic pressure applied by the solutions. Some studies have found that "bursting" of pollen could be
2522 possible due to mechanisms of osmosensory regulations (Shachar-Hill et al., 2013). Other research in
2523 Tobacco (*Nicotiana tabacum*) pollen, revealed that pollen cell volume changed "rapidly" in response
2524 to extracellular osmotic potential (Zonia & Munnik, 2004). Another study found that swelling and
2525 bursting of pollen grains was caused by the effect of the environment in protein production and
2526 protein infolding to the otherwise stiff exines of pollens (Božič & Šiber, 2022). Another study revealed
2527 that pollen is fragile to water status and associated processes once they leave the environment of the
2528 anthers (Firon et al., 2012). Future *Linum* studies could explore how osmotic pressure of different *in*
2529 *vitro* germination solutions could affect the integrity of *L. bienne* pollen.

2530 In this study, the pollen count was reduced by the different temperature treatments (cold and
2531 heated) relative to standard growing temperature of 25°C. In the overall data, averages of measures
2532 such as the pollen count, % of pollen forming tubes and the pollen tube counts all revealed a reduction
2533 in the heat and cold treatments relative to standard glasshouse treatment (Figure 6). In tomato
2534 (*Lycopersicon esculentum*) it has been previously shown that the effect of lower temperatures had no
2535 significance reduction in the counts of pollens observed, and even in cases increased pollen tube
2536 proportion (Peet & Bartholemew., 1996). In contrast, a temperate species borage (*Borago officinalis*)
2537 showed reduction in pollen grains when treated with heat treatment (Descamps et al., 2021). This
2538 species, is heavily reliant on insect pollinators. This is not the case for *L. usitatissimum* which is known
2539 to be a better self-pollinator (Williams et al., 1990). Perhaps, differences in breeding strategies and
2540 environmental variables could lead to different sensitivity in pollen production for different species,
2541 since selfing species doesn't necessarily interact with other organisms to germinate, making it less
2542 desirable to produce as much pollen as possible for success of pollination due to reliance on
2543 environmental pollinators. Considering the number of pollen tubes, this measure was always
2544 significantly different between the different treatments (Table 3). In all cases, there was a reduction
2545 of the number of pollen tubes observed in the cold and heated treatment. This may suggest that in
2546 our samples, there was an optimal temperature in which the number of pollen tube observed was
2547 maximized. In a study involving several *Rosaceae* species, it was suggested that pollen germination
2548 was optimized differently for different species at temperatures between 15°C to 30°C (Beltrán et al.,
2549 2019). However, when looking at the different in average pollen tube counts between the cold and

2550 the heated treatments, this difference was also significant. This suggests perhaps pollen tube
2551 formation was better under cold stress than it was under heat stress. One example which observed a
2552 better pollen tube growth under lower temperature was one that was conducted with *Citrus* (Montalt
2553 et al., 2019).

2554 When the number of pollen tubes was combined with the pollen counts to consider a
2555 proportion of pollen tube measure, the proportion of pollen tube decreased significantly with the heat
2556 treatment (Table 4). This illustrates that in addition to the decreasing number of pollen counts, under
2557 increased temperatures, the proportion of pollen able to form pollen tubes also decreased when
2558 temperatures were increased. For the colder temperature treatment however, there was a decrease
2559 in the proportion of pollen tube number, but this wasn't significant. This reveals that pollen tube
2560 formation in *Linum* is more sensitive to an increase in temperature rather than a decrease in the
2561 temperature. There seems to be some agreement in other temperate species. In cultivars of cherry
2562 (*Prunus avium* L.), pollen germination reduced while pollen tube growth accelerated when there was
2563 an increase in temperature. Therefore, pollen kinetics seemed to be affected with higher
2564 temperatures (Hedhly et al., 2004). Other studies also point to changes in stigma reception ability to
2565 the pollen tubes. In a study of the cultivated peach (*Prunus persica* L.), stigmatic receptivity was
2566 affected by temperature. However, accelerated pollen tube growth was observed, which the study
2567 explained as the opposite effect of temperatures on the male and female side in peach (Vuletin Selak
2568 et al., 2013). Another study of pollen performance in olives (*Olea europaea* L.) found that temperature
2569 affected pollen tube viability and pollen tube growth was reduced under increased temperatures. The
2570 study also suggests that temperature and genotype interaction was significant for pollen performance
2571 in olives (Vuletin Selak et al., 2013). There are various suggestions when it comes to pollen tube growth
2572 and temperatures in the literature. For different species and cultivars, it maybe that pollen tube
2573 growth is affected differently under different temperature treatments. This was evident when looking
2574 at species level (wild vs cultivars) in our study. Our result suggests that in cultivated *L. usitatissimum*,
2575 the proportion of pollen tubes was significantly affected by increasing and decreasing temperature
2576 treatments relative to the glasshouse treatments. It seems that the optimal temperature for the
2577 cultivar in our samples was the glasshouse treatment at 25°C daytime and 13°C nighttime
2578 temperature. In the wild *L. bienne* samples however, the greatest pollen tube proportions were
2579 observed under the cold treatment Although, this was not significantly different to the other
2580 temperature treatments for *L. bienne*, it showed a trend towards a lower temperature optimum to
2581 that of their cultivar relatives.

2582 When looking at the proportion of pollen tubes for each species, it was evident that cultivated
2583 *L. usitatissimum* was more sensitive to temperature than the wild relative *L. bienne*. This was evident

2584 because under the different temperature treatments, pollen tube proportions were significantly
2585 smaller in both cold and heat treatments with the cultivars. There was a more significant reduction in
2586 the proportion of pollen tubes than their wild ancestors. The effects of higher temperature stress
2587 negatively affected pollen tube formation more so than lower temperatures, especially in the
2588 cultivars. This effect of higher temperature stress on pollen viability is not new in plants. In chickpea
2589 (*Cicer arietinum*) for example, the pollen grains have been shown to be more sensitive to heat stress
2590 than their stigma counterparts (Devasirvatham et al., 2012). The fact that the crop wild relative *L.*
2591 *bienne* showed a potentially more resilient trait to rising temperature could be of interest in crop
2592 development. This importance has been highlighted in durum wheat (*Triticum*) before, which shows
2593 wild relatives yielding more grain under temperature stress when compared to the cultivar relatives
2594 (El Haddad et al., 2020). The fact that lower temperature affects pollen tube formation has been
2595 observed in other plants. Studies using pear also suggests disruption in pollen tube formation. They
2596 suggest that this is mediated by mitochondrial metabolic dysfunctions (Gao et al., 2014). The
2597 CBF/DREB1 proteins have also been identified to control expression of cold-induced genes
2598 (Thomashow, 1999). In Barley, novel alleles to frost resistance have also been identified as *FR-H2*
2599 amongst others (Sallam et al., 2021). There is potential that these cold resistance genes identified in
2600 other study models may play a role in higher tolerance to cold temperature observed in *L. bienne*
2601 pollen tube formation ability. In *Linum* this phenotypic difference in temperature stress pollen tube
2602 sensitivity may represent more diverse genetic resources for the improvement of crop heat stress
2603 resilience, as was reviewed in various wild relatives for crop improvements (Dempewolf et al., 2017).

2604 In terms of the ability of pollen tube to reach the ovary, the heat treatments revealed a
2605 significant reduction of pollen tubes that were able to reach the ovary. This was not the case in the
2606 cold temperature treatments. This suggests that an increase in temperature had a significantly
2607 negative effect on the ability of *Linum* pollen tubes to reach the ovary (Table 5). This result illustrated
2608 that *Linum* pollen's abilities to reach ovaries are more sensitive to an increase in temperature rather
2609 than a decrease. In a study with wheat (*Triticum aestivum* L.) pollen tube growth into the ovary was
2610 reduced due to an increase in temperature. It was suggested that this was because heat causes
2611 abnormal conditions for the ovary (Saini et al., 1983). In peach (*Prunus persica* L.), stigmatic receptivity
2612 to pollen tube was negatively affected by a rise in temperature (Hedhly et al., 2005). Suggestions in
2613 the literature often reveal negative effects of temperature increase on the female parts such as stigma
2614 and ovary that results in the observed reduced ability for pollen to reach the ovary. In this study
2615 however, we did not look at stigmatic and/or ovary 98arlierpment. In line with what is seen with the
2616 proportion of pollen tube reaching ovaries in this study, it would be of further interest to see how this

2617 negative response to the increase in temperature correlates to stigmatic and/or ovary development
2618 in our *Linum* samples.

2619 Furthermore, when analysed separately, wild *L. bienne* showed a more significant reduction
2620 in the number of pollen tube reaching the ovaries than those of their cultivar relatives. This trend was
2621 not expected since the proportion of pollen tubes to pollen count is significantly more reduced in the
2622 cultivar *L. usitatissimum* compared to the wild *L. bienne*. In the proportion of pollen tube able to reach
2623 ovary/species the wild *L. bienne* seems to have responded more strongly than their cultivar relatives.
2624 There is a suggestion here that even though wild pollen tube to pollen count proportion were higher
2625 than the cultivars, their ability to reach the pollen was more sensitive towards warmer temperature.
2626 Perhaps the pollen's female counterparts such as the stigma and the ovary were more sensitive to
2627 temperature changes in wild relatives (Fábián et al., 2019). A similar observation was also made in
2628 peach whereby stigmas tend to lose their capacity to support pollen tubes (Hedhly A. et al., 2005).
2629 Hedhly A. et al., 2005 also suggests that there are contrasting effect of temperature on the male and
2630 female parts of the flower. It maybe the case that, the amount of pollen tubes is affected by the ability
2631 of female counterparts, such as the stigma, to sustain pollen tubes at different temperatures. This
2632 aspect is yet to be explored in *Linum*. In this study, we showed that the heat (30°C) treatments cause
2633 a significant reduction in the number of *Linum* pollen tubes, and pollen tubes reaching the ovaries.
2634 However, this can be due to the sensitivity of female counterparts to rising temperature, as is evident
2635 in other species (Pan et al., 2018; Fábián et al., 2019). In future studies, it would be interesting to
2636 observe the effects of temperature on the growth rate of *Linum* pollen tube inside the stigma. This
2637 can be done using live images from confocal fluorescence microscopy on *in vitro* germination studies.

2638 A major aim in this thesis was to look at local adaptations, that was done here by a comparison
2639 between pollen measures and latitude of locality and climatic variables. We were interested in
2640 whether there was any correlation between the pollen count, pollen tube count and/or the proportion
2641 of pollen tube forming against local geographic and climatic variables. None of the different
2642 treatments shows a correlation in latitudinal and climatic variables when considering pollen counts.
2643 This suggests that the number of pollen in our sample set is neither positively nor negatively correlated
2644 with local climates. Other study observing percentages of pollen across the Mediterranean region
2645 found ecological trends in not only pollen data but also other plant traits to climatic gradients across
2646 the region (Barboni et al., 2004). We expected to see a trend in *Linum* pollen count with climatic
2647 variables, however, we didn't observe a significant trend in our sample set. Although this was the case
2648 for our study, there was some evidence of a change in direction of correlation trend in the number of
2649 pollens observed and local climatic variables, when looking at the different temperature treatments.
2650 This correlation was not statistically significant but perhaps with a larger sample set and from a more

2651 varied local environment, the trend would be more significant and more obvious, and a larger sample
2652 set would help in statistical power as well. Another study suggests that ‘fresh’ pollen should be used
2653 for observation studies as pollen conservation may impact germination abilities of pollen (Beltrán et
2654 al., 2019). Although laborious, it would be of interest to use ‘fresh’ pollen instead of a preservation
2655 method, as was done in this study. The number of pollen observed under different climatic variables
2656 is important as this may influence the possibility of gene flow between different populations and
2657 species. As *Linum* is a self-pollinating species, it is perhaps plausible that environmental segregation
2658 of populations by pollen production could influence gene flow dynamics as well as pollen/pistil
2659 incompatibility as was shown post-pollination in *Polemoniaceae* (Ruane & Donohue, 2007).

2660 When observing pollen tube count however, there was a significant association when looking
2661 at the cold treatments. This observation was true for both the latitudinal and climatic variables. Even
2662 within our limited sample sets, there was a latitudinal trend that, the more northern our sample was
2663 collected from, the more pollen tube it can produce under colder temperatures. The correlation with
2664 climatic variables also agrees with this observation. Local climate may, in the case of wild *L. bienne*,
2665 influence pollen viability, not just in terms of temperature, but also other climatic viables which may
2666 have played a role. In oak, pollen viability and sunlight availability in the local areas possibly influence
2667 pollen mediated gene flow between populations (Schueler et al., 2004).

2668 In addition, more northern wild population showed potential local adaptation with higher proportions
2669 of pollen tube forming observed in the cold treatments (Table 6A). This suggests that wild *L. bienne*
2670 are perhaps more adapted to colder temperature. There is potential for GWAS study here for Northern
2671 wild *L. bienne* to identify novel genes for cold tolerance. This can have implication towards
2672 improvement of cultivars. There was also a difference in the trend with heat treatments. When
2673 heated, the correlation changed direction between the latitude of localities and climatic variables and
2674 pollen and pollen tube counts. This was not significant but could be tested with a larger wild individual
2675 from a wider range of local variables in the future.

2676 In conclusion, different temperatures affected male reproductive function in terms of pollen
2677 number, pollen tube number, proportion of pollen tubes forming against pollen count, and proportion
2678 of pollen tubes reaching the ovary. In all cases there seemed to be an adverse effect of increases of
2679 temperature, be it pollen, and pollen counts and the amount of pollen tubes able to reach the ovary.
2680 Therefore, in our samples, an increase in temperature lessened pollen viabilities for *Linum*. There was
2681 also evidence that in the colder temperature treatments, pollen viability is correlated with latitude of
2682 localities and climatic variable. Local adaptation in pollen viability may have resulted in the differences
2683 seen. Additionally, we would like to observe the female counterpart of these *Linum* samples do in
2684 relation to the pollen viability trend that was seen in this study. Further study on the reception of

2685 female structures, such as the the ovary may reveal that, in line with the trend seen in pollen viabilities
2686 in *Linum*, the female counterparts also play a major part. Illustration in Sorghum have observed
2687 deterioration of ovary under heat stress (Chilawal A. et al., 2020). There is also interest in looking at
2688 further colder treatments for the wild population to see whether there is optimal temperature for
2689 pollen viability and to generate results for more samples from more diverse latitudes. The differences
2690 seen here are evidence of local adaptation in the wild *Linum* species. More northern individuals seem
2691 to be more locally adapted to colder temperatures and could be more adversely affected by an
2692 increase in temperature in terms of their ability to form pollen tube and the ability for pollen tube to
2693 reach ovaries.

2694

2695 CHAPTER 4: POPULATION STRUCTURE AND DIVERGENCE WITHIN LINUM
2696 SAMPLES ORIGINATING ACROSS WESTERN EUROPE LATITUDE RANGE.

2697

2698 There is ample evidence that climate change has the potential to drive distributions of
2699 population within both plants and animals, often having significant changes to ecosystems (Hampe
2700 & Petit, 2005; Mori et al., 2022). When looking at distribution of wild populations, it is with interest
2701 that latitudinal and longitudinal ranges for local populations could return insights into genetic
2702 diversity and conservation ecology. Genetic diversity is important in living organisms for adaptation
2703 to changing environments. Thus, conservation of species often depends on genetic diversity in each
2704 population (Alcala et al., 2013). In the wild, events such as genetic bottlenecks, genetic
2705 insertion/deletion, selection, and genetic duplication are widely known to shape patterns of genetic
2706 diversity in each population (Alcala et al., 2013). The central-marginal hypothesis predicts that at
2707 range margins there is decline in genetic diversity but an increase in differentiation towards
2708 speciation, due to variation which are caused by events such as genetic drift and gene flow (Langin et
2709 al., 2017).

2710 *Linum* is a plant genus with more than 150 species (Muravenko et al., 2010). There is a wider
2711 interest establishing the extent of genetic diversity and structure within different populations in wild
2712 *Linum* populations. A wide interest in plant research is looking at genetic implication of different
2713 phenotypes within plants, especially for crop development. In crop development, genetic analysis
2714 tool such as quantitative trait locus (QTL) analysis are often used to pinpoint loci which are
2715 responsible for a desired trait within a plant type (Asíns M., 2002). Once desired traits are identified,
2716 further cultivar developments could be approached with marker assisted selection. However, in
2717 these types of studies a whole genome sequence is often required. More recently new sequencing
2718 approaches are being developed which enables genetic analysis studies to be done relatively faster
2719 and cheaper (Behjati & Tarpey., 2013). Studies involving genetic marker systems to genotype wild
2720 and cultivated types were able to identify genetic structuring to further reveal diversity and
2721 consequences of selective breeding in the cultivars without the need of a whole genome (Bacilieri et
2722 al., 2013). This can be achieved using sequence repeats such as simple sequence
2723 repeats/microsatellites (SSRs). SSRs are sequences which are ubiquitous in a specific eukaryotic
2724 genome and are amongst the most common genetic markers to be developed (Goldwin et al., 2005).
2725 SSRs were developed along with PCR procedures to amplify sequences, often looking at genetic
2726 diversity within set populations, and between plant types (Grapin et al., 2005). Studies looking at
2727 non-model species have been able to identify reduction of genetic diversity due to hybridization and

2728 genetic bottlenecks events which revealed consequences of cultivation of the species (Guan et al.,
2729 2021). This is also of interest in terms of molecular ecology. Genetic analysis has previously
2730 identified localization and invasion histories of non-model plants (Hernández et al., 2019). To look at
2731 differences across population, it is of interest to establish a structure and diversity between the local
2732 population.

2733 In *Linum*, previous research used Single Sequence Repeats (SSR) techniques. In the research,
2734 SSRs were developed for 34 Turkish wild flax (*Linum bienne*) accessions and accessions from
2735 different cultivated flax (*Linum usitatissimum*) lines. The research found variation in polymorphisms
2736 between the accessions and clustering of 493 individuals according to their respective types,
2737 suggesting wild flax are more closely related to the deshiscent type. Along with this, the research
2738 also suggests genetic distancing among the wild types are significantly related to their geographical
2739 distances as well as their elevation (Uysal et al., 2010). The research revealed potential for marker
2740 studies using *Linum* types to understand genetic structuring, domestication, and genetic diversity of
2741 *Linum*. For this purpose, short reads within the whole sequence were used as a genetic marker for
2742 different individuals. This is substantially easier than obtaining a whole genome, as a whole genome
2743 for the wild *Linum* species is not necessary for observation of population structure based on these
2744 markers. This being the case for the wild species, a chromosomal-level whole genome is readily
2745 available for the cultivar relatives *L. usitatissimum* (Sa et al., 2021). In this chapter, we use ddRAD
2746 sequencing to investigate whether there are structuring, diversity and or divergence within our
2747 *Linum* samples.

2748 We have a collection of wild *L. bienne* and several oil and fibre cultivar variety of *L.*
2749 *usitatissimum* originating from Western Europe. Within the *Linum* samples available under this
2750 project, there were a variety of individual plant populations that originated from different places
2751 across the latitude of Western Europe. We hypothesize that there is genetic variation due to
2752 localization within different environments. This could be implied by structuring and genetic diversity
2753 measures which may have taken place across time as different population form and localize to their
2754 local environment. When looking at the collection of *Linum* samples from across western Europe,
2755 genetic structure would be revealed as the genetic distance between the different *Linum*
2756 populations as well as between the wild and cultivar types. This would help in explaining their
2757 population genetics and as well as their breeding systems. There will be an interest in answering the
2758 question of “can we use genetics to interpret population structures and breeding systems in
2759 *Linum*?”.

2760 Our *Linum* samples consisted of 121 different individuals which made up 15 wild population
2761 (*Linum bienne*) from across Western Europe and 12 Cultivars (*Linum usitatissimum*). In the 15
2762 different wild populations, there were 109 individuals, and, in the cultivars, there is only 1 individual
2763 per cultivar line, which we have classified under one “cultivar” population for this study. The wild
2764 populations were collected respectively from the southern Spanish through to northern UK regions
2765 whilst cultivars were gained from IPK World Collection (Figure 7). This provides a wide latitudinal
2766 range for the temperate plants. When looking at wild types regarding their population structure, this
2767 was more likely affected by localization by the different population under the different latitudinal
2768 range. Groupings between the different populations are expected whereby groups can be
2769 distinguished by their local population. Population structures are inferred by determination of
2770 genetically related clustering (groups) observed without prior knowledge of the populations (Odong
2771 et al., 2011). The cultivars were expected to group together while the wild populations were
2772 expected to be grouped based on their region of locality. There is an expectation that populations
2773 within the sample sets group between northern and southern European populations. This would
2774 agree with morphological differences found under the vernalization experiment in this thesis (Part
2775 “2.4.1 Results: Vernalization and flowering time”). We also examined ancestry, to determine wild
2776 populations relationships to the set of cultivars within the experiment.

2777 For the purposes of sequencing genetic markers, a double-digest Restriction-site Associated
2778 DNA Sequencing (ddRADSeq) protocol was optimized to enable a digestion of DNA which could be
2779 done with multiple restriction enzymes. The use of a second restriction enzyme allows for more
2780 precise and consistent cutting of the DNA as well as provide more combination of previously
2781 multiplexed samples, allowing for a more homologous indexing of many individuals (Peterson B. et
2782 al., 2012). This enabled the large-scale generation of short read sequences without the need to
2783 sequence a whole genome. DdRADSeq in turn enabled genomic studies in models without whole
2784 genome databases and non-model organisms alike (Arnold et al., 2013) (Peterson B. et al., 2012). A
2785 previous SNP comparison was used to estimate genetic diversity and population structure in rice
2786 (*Oryza sativa*) (Singh et al., 2013). SNPs are thought to have numerous advantages over SSRs. These
2787 include a more precise estimate of population diversity and the ability to consider local adaptation
2788 through identification of groups by clustering methods (Zimmerman., 2020).

2789 Illustration of population structure, genetic diversity, and selection analyses were carried out
2790 through the data gained under the ddRADSeq protocol. Genetic information regarding the *Linum*
2791 population was gained in a relatively cost effective and time effective way. We hypothesize that
2792 genetic structuring is visible within our *Linum* samples as a signal of local adaptation. In addition we
2793 also investigates whether there are other genetic signal such as genetic diversity and heterozygosity

2794 in our samples for the implication of genetic distinction within our *Linum* populations. The
2795 expectation in the *Linum* sample, is that individuals would be grouped based on their population and
2796 location, and that grouping between the northern and southern populations would match what was
2797 previously observed in previous chapters which have examined the *Linum* samples morphologically.
2798 This would reveal that, structurally, northern, and southern populations in the *Linum* collection are
2799 genetically distant to each other. Further analysis could explore genetic traits such as heterozygosity
2800 and whether any population is genetically more diverse than other populations in the sample set.
2801 This would highlight potential breeding system differences in our *Linum* samples.

2802

2803

2804 4.2 STUDY AND METHODS

2805

2806 4.2.1 Modified RAD Seq Protocol

2807 For this chapter ddRADSeq protocol was used along with some modifications to the original
2808 protocol described by Peterson et al (2012). DNA was obtained through a modified DNA extraction
2809 protocol. The DNA extraction protocol was based on a method described by Doyle & Doyle (Doyle &
2810 Doyle 1987). After DNA extraction, a digestion procedure was carried out using two enzymes (we
2811 used MseI and PstI in this study) whereby the digestion was used to cut DNA restrictively into small
2812 sequences around 250-500bp long, further size selection (250-500bp) occurs during the Pipin stage
2813 of the protocol, after the ligation stages. DNA quality and quantities were observed through several
2814 quantification methods which includes gel electrophoresis and spectrophotometry. After satisfaction
2815 with DNA quality and quantity, libraries were built with sequences that are depicted as the Figure
2816 below.




Figure 1. An illustration of the ligated DNA material per sample which are both adapted and barcoded.

2817

2818 The final product of the DNA sequence after the ligation and barcoding procedures can be
2819 illustrated (Figure 1). The protocol used in this experiment was based on a ddRADSeq protocol
2820 previously described by Peterson *et al* (Peterson et al., 2012). Population structure analyses were
2821 subsequently performed to group individual samples based on their genetic distance relative to each
2822 other.

2823

2824 4.2.2 DNA Extraction Procedures

2825 For RAD-Sequencing purposes, we required young leaf materials for optimal DNA quality.
2826 Wild S0 seeds were grown in the glasshouse (16:8 day/night ratio and day/night temperatures at
2827 25/13°C) for optimal growth condition. 121 unique individuals representing the widest possible
2828 latitudinal ranges (from 31.791° South to 53.352° North) were chosen for this purpose, covering
2829 temperate regions of western Europe. The samples have a latitudinal range which covers northern
2830 UK down to the Southern coast of Spain. These samples include wild and cultivar species with the

2831 wild species collected directly from the field. A modified CTAB DNA extraction protocol was used
2832 from methods described by Soltis lab (Doyle, 1987).

2833 We modified the main CTAB washing buffer for DNA extractions. The presence of salt and
2834 polysaccharides could potentially interfere with downstream processes. To reduce this interference,
2835 suggestions were adapted from Clarke *et al* (Clarke J., 2009). The final modifications to the samples
2836 were made as follows:

| Reagent | Amount (For 100mL) | Final concentration |
|---|--------------------|---------------------|
| 10% CTAB in dH ₂ O | 30mL | 3% |
| 5M NaCl in dH ₂ O | 28mL | 28% |
| 0.5 M EDTA (pH 8.0) | 4mL | 4% |
| 1 M Tris-HCl (pH 8.0) | 10mL | 10% |
| Polyvinylpyrrolidone (PVP) (MW 40 kDa) | 3g | 3% |
| β-Mercaptoethanol | 0.2mL | 0.2% |
| dH ₂ O | 24.8mL | 24.8% |

2837 Table 1. Reagents for the CTAB extraction buffer

2838 With the CTAB buffer made, the protocol were carried out as described by Doyle (1987).

2839 4.2.3 The Modified CTAB protocol

2840 In the laboratory, dry baths were pre-heated to 60°C and RNase were removed and ice-
2841 thawed from -21°C storage. In the glasshouse, 10-12 young, 'green' leaves were collected per
2842 individual. Samples were individually homogenised. Alternative homogenising methods were also
2843 initially used, using 3mM metal beads, a pinch of sand and a Tissue-Lyser II. However, access to
2844 Tissue-Lyser II were limited due to Covid-19 restrictions and most samples were ground using a
2845 tissue homogeniser. Samples were ground as finely as possible inside Eppendorf tubes and flash
2846 frozen in liquid Nitrogen until all samples were ready.

2847 Per sample, DNA was cleaned as per Soltis lab protocol (Doyle, 1987). Samples were
2848 homogenised further with a vortexer until ground tissues are thoroughly mixed with the CTAB and
2849 RNase A solution. Samples were then placed on a 60°C pre-heated dry bath for at least 60 minutes
2850 (modified to allow time for optimal DNA lysis). Samples were then thawed to chill on ice for 3
2851 minutes before further processes.

2852 Samples were further washed using phenol-chloroform solutions as per protocol. To wash
2853 away any potential phenol after the wash, 200µL of chloroform isoamyl were added. Samples were
2854 centrifuged at 14,000RPM for 10 minutes before carefully removing the aqueous layer into newly
2855 labelled Eppendorf tubes. Na Acetate were then added as per protocol. Salt was washed away at
2856 room temperature with isopropanol as per protocol. Centrifugation at ≥14,000 RPM for 15 minutes

2857 were applied after the addition of isopropanol. A pellet at the bottom of the tube were often
2858 observed after the last centrifugation step, although not all the time. Supernatant was subsequently
2859 removed from the tubes. Into these sample tubes, 500 μ L of “Ice cold” 70% ethanol were added to
2860 wash the DNA pellets. Tubes were then centrifuged at \geq 14,000 RPM for 15 minutes at 4°C and
2861 ethanol taken out. Samples were air dried (preferably under a laminar flow hood) for around 10
2862 minutes, until samples were dry. Dried samples were then re-suspended in 50 μ L of nuclease free
2863 water as elutant. Samples were then stored overnight in a 4°C fridge before measurements of both
2864 quality and quantity.

2865

2866 4.2.4 DNA Quality and Quantity

2867 We used a NanoDrop ND1000 to quantify 1 μ L of samples. Sample quantity and quality were
2868 measured as DNA concentration (ng/ μ L) and absorbance ratios at 260/230nm and 260/280nm. A
2869 more accurate Qubit method were also explored using Invitrogen’s Qubit 2.0 fluorometer. We used
2870 Promega’s QuantiFluor ds DNA System (QuantiFluor® dsDNA System, Promega Corporation), DNA
2871 samples were able to be quantified more precisely using this protocol. In most cases, DNA were also
2872 measured using gel electrophoresis.

2873 For further quality measures, 0.5 \times TBE gel electrophoresis were used. Biorad’s DNA Mini Gel
2874 tank, with a power supply unit were used to run these gels. 2 μ L of New England Biolab’s DNA purple
2875 loading dye and 3 μ L of nuclease free water were dispensed and lined up per number of loadings
2876 required. A 1Kb DNA HyperLadder (Bioline) were used to measure the size and quantity of DNA. Gels
2877 were electrophoresed and observed using a UVIDOC system.

2878

2879 4.2.5 Double-Digest RAD Sequencing

2880 After satisfactory DNA materials were acquired (>20ng/ μ l), we processed the DNA through a
2881 modified ddRADSeq protocol described by Peteron et al 2012 (Peterson B. et al., 2012). Samples
2882 were digested and ligated to specific adapters and barcodes. This would be proceeded by pooling
2883 and cleaning stages of the products before further size selection and sequencing.

2884

2885 4.2.5i RAD – Digestion:

2886 To fragment extracted DNA materials, enzymes were used to digest said DNA materials. There are
2887 several different digestion enzymes available to use, each with their own specifications. We tested
2888 three enzymes to digest DNA materials, namely, *Mse1*, *Pst1*, *Sbf1* (manufactured by New England

2889 Biolabs). Two different combinations were tested: *Mse1/Pst1* and *Mse1/Sbf1* with the digestion
2890 protocol using NEB's buffer 2.1. 500ng genomic DNA and 5µL of buffer 2.1 was used as
2891 recommended by the supplier for optimum enzyme efficiency. We prepared a 30µL reaction. A
2892 master-mix of the buffer and enzymes were prepared for every sample. For incubation, and
2893 inactivation purposes, a Prime Techne thermal cycler was used. A 30µL digestion reaction were
2894 prepared as per manufacturer's protocol.

2895 4.2.5ii RAD – Anneal adapter and Ligation

2896 Digested DNA was subsequently ligated by using a ligase enzyme (manufactured by New
2897 England Biolabs). During ligation, digested DNA was adapted and barcoded specifically. To do this,
2898 adapter oligos were first annealed together per manufacturer's protocol.

2899 After mixing briefly adapters were annealed using a protocol recommended by Eurofin
2900 Operon. In a thermo-cycler, adapters were heated to 95°C for 2 minutes and ramp cooled to 25°C
2901 over a period of 45 minutes. Annealed adapters were then stored in a -21°C freezer for long term
2902 and in a 4°C fridge for short term (<72 Hours). The process was repeated for every adapter available.

2903 Samples and adapters were now ready for ligation stage. In the experiment, one forward and one
2904 reverse working stock adapters were combined with fragmented sample DNA and ligated to barcode
2905 the samples. New England Biolab's T4 ligase enzyme at 400U/µL concentration and a 10× T4 ligase
2906 buffer were used for this. The samples were loaded according to the 30µL final reaction volume from
2907 the digestion steps. The following tables describes each component for ligation. Samples were
2908 incubated at 23°C for 30 minutes and then heat inactivated at 65°C for 10 minutes. Samples were
2909 then cooled down at 2°C per 90 seconds until they reached 23°C.

2910

2911 4.2.5iii RAD – Pooling and Cleaning Ligated products

2912 To pool and clean ligation products from primers, SeraMag magnetic cleaning beads were
2913 used. This step required all samples to be pooled into a single container. Total volume of the pooled
2914 samples was calculated as the volume of leftover ligation product after PCR (PCR product) × total
2915 amount of samples. 1.5mL Eppendorfs were used for pooling and cleaning. For each tube, the
2916 volume of pooled ligated product would be no more than 300µL, to be cleaned separately.

2917 Before using the SeraMag beads, working solutions were prepared per manufacturer's
2918 instructions. 1mL of the working solution were then transferred into a 1.5mL Eppendorf tube and
2919 further processed as per manufacturer's instructions.

2920 After the final magnetic stage, tube with the magnetic beads and 1mL TE were put set in a
2921 rack (non-magnetic) and PEG-8000 and Tween 20 solution was added per manufacturer's
2922 instructions. When the solution is mixed, 1mL of the previously prepared 1mL TE + SeraMag beads
2923 were pipetted into the conical flask. Conical flask was then filled up to 50mL volume with MilliQ
2924 water and mixed gently until the beads are evenly spread across the solution. The solution were
2925 then transferred into a 50mL Falcon tube, wrapped in aluminium foil, and stored in a 4°C fridge
2926 ready for use.

2927 Before using on ligated products, the cleaning beads were tested against a 100-1Kb DNA
2928 ladder (BioLine's HyperLadder). After confirmation against DNA Ladder (different ratio of beads
2929 should cut at different fragment sizes), the prepared Sera-Mag beads were then ready to be used for
2930 the ligated products. To do this, clarification needed to be made about the ratio of beads to be used
2931 for the fragment size which are of interest. In the case of this experiment, 250-500bp regions were
2932 of interest. A 1.5× SeraMag to ligated product volume ratio were used.

2933 After the addition of the beads working solution, solution was incubated at room
2934 temperature for 5 minutes. After this, samples were placed back into the magnetic stand and
2935 processed as per manufacturer's protocol. The produced DNA materials were then washed and dried
2936 per manufacturer's instructions. This wash step was repeated twice. When the second wash
2937 finished, beads were placed on a 37°C heat block until beads were dry. Rehydration took place with
2938 20µL of nuclease free water as elutants. When the beads were mixed in the water, samples were
2939 placed back in the magnetic stand until all beads were pulled towards the magnet. When all beads
2940 were pulled towards the magnet, the supernatant were then extracted and transferred to a new
2941 1.5mL Eppendorf tubes or into the next 300µL pooled tube until all were cleaned. Cleaned samples
2942 were then quantified using a NanoDrop ND1000 spectrometer.

2943

2944 4.2.5iv RAD – PCR Amplification to Generate Illumina Sequencing Libraries

2945 High-fidelity PCRs were undergone with all the samples for ligation verification and
2946 generation of sequencing libraries for Illumina. ThermoFisher's Phusion High Fidelity DNA
2947 Polymerase were used for this protocol with P1 and P2 adapter oligos as the primers. A total volume
2948 of 20µL reactions were prepared as per manufacturer's instructions. For this protocol, 20ng of input
2949 DNA were used. The input DNA into the digestion product was 500ng. 20ng in volume of DNA were
2950 calculated as follows:

$$2951 \text{ Input DNA} = \frac{500 (\text{digestion DNA input})}{20(\text{DNA required for PCR})} = 25$$

2952
$$\text{Final Volume} = \frac{37 \text{ (Final digestion volume)}}{25 \text{ (From input DNA)}} = 1.5\mu\text{L}$$

2953 The input DNA from ligation were calculated as 1.5µL. The samples were then run on the 111arlier-
2954 cycler for 12 cycles of the following programme: *98°C for 2 minutes, 98°C for 10 seconds, 65°C for 30*
2955 *seconds, 72°C for 30 seconds and a final 72°C for 10 minutes.* After the cycles were done samples
2956 were held at 4°C. A 1×TBE gel electrophoresis was performed to check the presence of products
2957 against a DNA ladder.

2958

2959 4.2.5v RAD – Pippin size selection Preparation

2960 To construct a ddradseq library of 250-500bp fragments, a Pippin Prep facility was used
2961 (Sage science). A 1.5% Agarose gel cassette (with marker L) was used for Pippin preparation using
2962 manufacturer’s protocol. The instrument was calibrated prior to the run as per manufacturer’s
2963 protocol.

2964 To run the Pippin size selection, the cassette was sealed with the provided seal. The
2965 automatic test was run and the current measured in each elution channel at room temperature. A
2966 size selection protocol was then manually inputted or selected from the “Protocol editor” tab with
2967 sample ID provided. Under the “Cassette” input, an appropriate cassette file was loaded and under
2968 the “Reference Lane” input, the marker well was inputted and applied to all lanes. The pippin was
2969 subsequently run for each sample well to select at 250-500bp for the purpose of this experiment.
2970 When the Pippin prep finished running, 30µL of sample was collected from the elution module. An
2971 Agilent Tapestation 2200 facility was used to identify and quantify the region of interest, with a High
2972 SensitivityD1000 ScreenTape.

2973

2974 4.2.5vi RAD – qPCR Quantification

2975 For a final quantification, a quantitative Polymerase Chain Reaction (qPCR) procedure was
2976 explored to quantify DNA materials. In this part of the quantification a BioRad CFX96 Real-Time PCR
2977 System was used. For this purpose, 4 dilutions of library materials were explored. The dilutions were
2978 made using previous pooling from section “4.2.5iii RAD – Pooling and Cleaning Ligated products” of
2979 the RAD methods, each with the appropriate amount for the pooled DNA in 10mM Tris-HCL solution
2980 as required per dilution. All dilutions were repeated three times. No template controls (NTC) were
2981 used as a control for contamination in the qPCR runs. There were 4 standard dilutions used as a
2982 standard in the Bio-Rad system for quantification purposes for each of the dilutions. In total there
2983 was 48 wells for the samples and dilutions, 3 NTCs, and 4×3 wells for the standards. This makes up to

2984 63 wells for the run. For the standards, we used 4 concentrations (10pMol, 1pMol, 0.1pMol, and
2985 0.01pMol).

2986 For qPCR purposes, a KAPA Library Quantification Kit (Roche Molecular Systems) were used. When
2987 reagents were fully made, a 96-well qPCR plate were designed. Standards and samples were loaded
2988 as per manufacturer's instructions. An appropriate cycle programme was then set up and ran.

2989 After the protocol had finished running, a "Quantification CQ result" can be seen. CQ stands
2990 for quantification cycles, in other qPCR machines these are also specified as cycle-threshold. We
2991 used starting quantity (SQ) values to calculate the final concentration for each dilution factor. For
2992 example, for a dilution factor of 1:2000, the SQ-value were multiplied by 2000. Size corrections were
2993 able to be calculated by multiplying the average fragment length (250-500bp) by the standards
2994 (452bp). Units were able to be converted from picomolar to nanomolar by multiplying by 1000 and
2995 subsequently nanomolarity to a concentration unit (ng/ μ L). Using the delta-CQ values, consistencies
2996 between the triplicates were checked before calculations of concentrations.

2997

2998 4.2.5vii RAD – Sequencing

2999 When satisfied with quality checks, an Illumina 2500 Hi-Seq sequencer was used at Durham
3000 University's DBS Genomics facility to sequence the ddradseq library. An appropriate amount of
3001 pooled library was calculated for sequencing. All the DNA library poolings from the previous steps
3002 were combined into 1.5mL Eppendorf tube. Illumina Hi-Seq 2500 requires certain amount of
3003 genomic DNA to be read successfully. DBS Genomics at Durham University's Biosciences department
3004 recommends at least 10ng/10 μ L concentration of processed DNA library to process the samples
3005 through the sequencer. After quality check measures were made and 2 \times 10 μ L tubes of sample were
3006 sent through the Illumina Hi-seq 2500 sequencer measuring at >10ng/ μ L each. An Eppendorf
3007 centrifugal vacuum concentrator were used as required to combine and concentrate pooled libraries
3008 to achieve a 10ng/ μ L threshold required for sequencing. After the sequencing process, two raw
3009 library files which needed further demultiplexing and mapping were outputted. The raw sequences
3010 consist of a forward and a reverse sequence for the pooled library which needed to be processed to
3011 imply information. We repeated the sequencing twice to gain a better-quality library read.

3012 4.3 ANALYSIS: PROCESSING RADSEQ

3013

3014 4.3.1 Analysis: Post RAD-Sequencing

3015 To make sure of homogeneity, we used the same pooled library from the same ddRADSeq
3016 run. The result of two sequencing processes were four raw sequences from the same pooling (two
3017 forward and two reverse sequences), for further analyses, these sequences were merged to provide
3018 one raw-forward sequence and one raw-reverse sequence. This required raw reads to be processed
3019 bioinformatically.

3020 High Performance Computing (HPC) facilities were available for access at Durham University.
3021 The interface for the HPC facility is provided by a Linux cluster using several Intel processors. We
3022 used “Bourne-Again Shell” (Bash) to operate the programmes within the HPC facility (Ramey, 2022).
3023 The HPC facility were also remotely accessed through internet connection. Bash environments and
3024 FTP protocols were loaded using SSH clients such as PuTTY (Tatham, 2022) and MobaXTerm
3025 (Mobaxterm., 2022).

3026

3027 4.3.2 Analysis: Demultiplexing Raw Reads

3028 Because the sequence was barcoded and adapted specifically during the ligation stages of
3029 the DDRADSeq protocol, the raw sequences were able to be demultiplexed and sorted based on
3030 their specific barcodes (Croissac et al., 2016). This enabled downstream processes to separate out
3031 the sequence based on the unique sample barcodes which were ligated during the ddRADSeq
3032 protocol. The demultiplexing protocol can be found under a pipeline prepared for *STACKS v2.61*
3033 (Catchen et al., 2013). The demultiplexing protocol were executed using the command
3034 “*process_radtags*” under *STACKS v2.61*. The two adapter sequences as found in appendix 15, were
3035 also specified in the command. The full barcode index is also provided in appendix 16. The
3036 programme was optioned to filter the data for uncalled bases, using built in default parameters. This
3037 is done by inputting “-c” with the “*process_radtags*” commands, it was also programmed to discard
3038 reads of low quality by inputting “-q”, and to rescue barcodes and RAD-Tag cutsites by inputting “-r”
3039 in the command line. The barcode option of the programme was specified to read barcodes which
3040 are in line with the sequence and occurs in paired end reads. The number of allowed mismatches
3041 when rescuing barcodes were set to two. The full command is listed under (appendix 17). After
3042 demultiplexing, *STACKS* were able to output 580,520,687 reads of which 22,639,406 were reads
3043 containing adapters and 814,583 were reads considered as “low quality”. After de-multiplexing with

3044 default parameter filters, 115/121 individuals were kept, 6 individuals were filtered out due to either
3045 excessive number of uncalled bases or low read qualities.

3046

3047 4.3.3 Analysis: Mapping to a Reference Genome

3048 Post-demultiplexing, samples were able to be mapped either without a reference genome
3049 (*de_novo*) or with a reference genome. We were interested in GWAS analysis, whereby a reference
3050 genome was essential. The sample set in this case was inclusive of cultivars (*L. usitatissimum*), and
3051 the wild pale flax (*L. bienne*) individuals which are widely regarded this wild species whereby the
3052 cultivars were domesticated from (Allaby et al., 2005). These are amongst the justification for opting
3053 to map to a whole genome which was already available for the cultivars of flax but not for the wild
3054 type.

3055 The Genome for the cultivar is publicly available and are assembled down to chromosomal
3056 levels, making it suitable to be mapped to (Sa et al., 2021). Mapping to a reference genome were
3057 undergone using the `ref_map.pl` programme of the STACKS v2.61 software (Catchen et al., 2013).
3058 The programme calls on each of the STACKS components noting “gstacks” as the component which
3059 align the positions for the RADSeq reads and calls SNPs in each of the sample based on the
3060 alignments. The programme expects the data to have been previously aligned with the genome
3061 using a separate aligner. For this purpose, BWA_MEM algorithm of BWA v0.7 was used to align the
3062 RADSeq reads to the cultivar genome (Li & Durbin., 2009). The outputs were pipelined to Samtools
3063 v1.15 to be converted to .BAM files which are the input file type required for `ref_map.pl` to work.
3064 BWA and Samtools were piped under one command. The command can be found under (appendix
3065 18). After obtaining .BAM files for each of the samples, `ref_map.pl` can be executed using the
3066 “`ref_map.pl`” command. The optional command to execute the “populations” programme of STACKS
3067 were also piped into the `ref_map.pl` command. This is a filter to accept a minimum percentage of
3068 individuals in a population to process a locus. This was set to 80% by using the command “-r 0.8”.
3069 The populations command also allows further file output options such as the PLINK and VCF formats
3070 to be processed further downstream with a population genetic specific software. A population map
3071 was also specified to map which samples belongs to which population as well as to map a
3072 geographical region specific to that population. This is useful for further analysis downstream. A
3073 population map file for the *Linum* samples is listed as in appendix 19. The full command for the
3074 `ref_map.pl` pipeline can be found under (appendix 20). Post `ref_map.pl` run, several outputs will be
3075 of interest. The first if the VCF output which can be used for analysis quality of the reads such as the

3076 read depths, allele frequency and PGD-Spider (Lischer and Excoffier, 2012), PLINK (Purcell et al.,
3077 2007) or vcftools (Danecek et al., 2011) was used to convert the SNP data into PED and MAP.

3078

3079 4.3.4 Analysis: Processing Mapped Reads

3080 To process the mapped reads an initial step was to convert .PED and .MAP output files into
3081 .RAW and additional .BED files for read inputs into further downstream processes. To do this, PLINK
3082 v1.90 was able to be used (Purcell et al., 2007). PLINK was able to specify .PED and .MAP files in a
3083 directory given the same name. The additional command "--recodeA" is a data management options
3084 whereby both .PED and .MAP files were able to be processed into a single .RAW data file which
3085 includes formatting that are useful for further population genetics analysis. For further analysis a
3086 .BED file format were also required. This is obtained through the "--make-bed" addition to the PLINK
3087 command. There were numerous (>1) chromosomes in the .MAP files. This means that PLINK needs
3088 to acknowledge and allow for this in the output .RAW file. To do this the "--allow-extra-chr" option
3089 for PLINK were inputted into the command line for both make-bed and .RAW file options. The full
3090 command for PLINK conversions can be found under appendix 21. After both .RAW and .BED files
3091 were constructed, further population analysis could take place. In this case downstream analysis
3092 would be carried out using a programme written in "R". R was run in Durham University's HPC server
3093 for the purposes of this analysis.

3094

3095 4.3.5 Analysis: Processing Data in SambaR

3096 To further the analysis of the mapped reads, output .RAW and .BED files were further
3097 processed through using a programme constructed under "R". The programme is called "SambaR"
3098 (Snp 115ar1 Management and Basic Analyses in R) and is available for open use with no requirement
3099 for a licence. SambaR functions to integrate numerous R packages such as "Adegenet", "poppr",
3100 "FactoMineR", et cetera. SambaR functions as "collections of functions which increase the power of
3101 existing R tools for population-genetic analyses" (De Jong et al., 2021). This convenience makes
3102 SambaR an ideal tool to save time with running population genetics analysis, which can include
3103 population structure, diversity, and selection analysis. SambaR were run in the "R" environment;
3104 therefore, the statistical tool R is essential for further downstream processing. For the analysis of the
3105 data within this thesis, R (version 4.1.3) were loaded from the HPC server into the environment.
3106 SambaR could be manually downloaded using from the official Github page
3107 (<https://github.com/mennodejong1986/SambaR>) and loaded into R manually. Alternatively, it could
3108 be loaded into R using the command

3109 `“source(“https://raw.githubusercontent.com/mennodejong1986/SambaR/master/SAMBAR_v1.07.tx`
3110 `t”)”`. This command executes a source code from the Github online server where the source code for
3111 SambaR can be found. Further downstream SambaR commands worked after the source codes were
3112 executed. R packages were able to be automatically checked and added through the `“getpackages()”`
3113 command. This command outputted a .txt file whereby each packages required could be checked as
3114 been successfully loaded or not. In the case of this analysis, all recommended packages were able to
3115 be loaded in addition to the essential packages to run the SambaR process. Data can be manipulated
3116 easily in SambaR using several pre-loaded R packages.

3117 SambaR also accepts a geographical input file. This is useful for implications of the
3118 geographical range within our samples. From collection, the samples within the pooled library
3119 should have a latitudinal range across Western Europe. This provides latitudinal cline for structure
3120 analysis. SambaR uses the function `‘getMap’` of the R package `rworldmap-1.3.6` (South, 2011). The
3121 addition of pie charts was included for mapping in SambaR, using the function `‘add.pie’` of the R
3122 package `mapplots-1.5.1` (Gerritsen, 2018). An input .txt file, consisting for 3 tab-separated columns
3123 were created using the geographical information of the different population observed within the
3124 sample. The information for this file is available in appendix 22. As an addition to SambaR’s
3125 `“importdata()”` command, the option `“geofile=fileprefix”` were included.

3126 For different analysis, data were processed differently. For genetic structure analysis it is
3127 better to have as many individuals and populations as possible to infer better structuring between
3128 the populations in the collections. However, for population diversity and differentiation analysis, we
3129 determined that it was better to include as many SNPs as possible. Therefore, filters that enables to
3130 retain as many SNPs as possible were desirable during population diversity and population
3131 differentiation analysis. This may in turn reduce the number of individuals in the samples, thus,
3132 number of populations may be reduced under the filters for population diversity and population
3133 differentiation. For recommendations, SambaR recommends not using `snpmis=>0.05` for population
3134 structure analysis while for diversity and selection analysis, SambaR recommends higher `snpmis`
3135 parameters.


3136 4.3.6 Analysis: Data Management in SambaR

3137 SNP data management and analyses were performed in R-4.1.2 (R Core Team, 2022) using
3138 wrapper functions of the R package SambaR (Github page:
3139 <https://github.com/mennodejong1986/SambaR>). The data was then imported into R and stored in a
3140 Genlight object using the function `‘read.PLINK’` of the R package `adegenet-2.1.5` (Jombart, 2008;
3141 Jombart and Ahmed, 2011). For the purposes of filtering, SambaR filter data per population and with

3142 much less individual/population, the data would be subject to stricter filtering procedure. The
3143 dataset for this study only consists of 115 individuals with 26 populations including eleven cultivars.
3144 Populations with only 1 individual will later be excluded from the population analysis. In the cultivar
3145 varieties, only one individual/population occurred and for the purposes of this analysis, they were
3146 put under one “cultivar (CUL)” population to see if any structural implication can be made against
3147 the wild *L. bienne*. In addition, having one individual per population is not recommended for
3148 population genetic analyses. Not only would it make the filtering/population impossible but also, for
3149 practical reasons, SambaR doesn’t allow populations which are only represented by one individual.
3150 This is because the mean difference between sequences (Nucleotide diversity) are estimated by
3151 mean sequence differences between individuals (Innan et al., 1999). For estimation of nucleotide
3152 diversity, population genetic analysis programmes often estimate by averaging the estimated
3153 number of nucleotide changes over all the samples (Innan et al., 1999). If a population only happen
3154 to have one individual, the mean difference is essentially only comparable to itself. For example, in
3155 heterozygous analyses, the nucleotide diversity would essentially just be the heterozygosity of the
3156 one individual as opposed to diversity amongst a population. As a result of this, cultivars would have
3157 to be merged and compared against the wild population. We included the cultivars to observe their
3158 genetic distance to the wild populations across Europe as a species. For initial data filtering option,
3159 all the populations were merged into one population using the “mergepop” command of SambaR.
3160 This was done after the data were imported. As a matter of downstream analysis, the output filtered
3161 data can be extracted into PLINK type format so that we can manually input the population prefix
3162 back in the dataset for further analyses.

3163 After merging, the data was filtered using the function ‘filterdata’ of the R package SambaR.
3164 For population structure analysis purposes, the data were filtered with the following parameters:
3165 *indmiss=0.7, snpmiss=0.05, min_mac=2, dohefilter=TRUE, snpdepthfilter=TRUE, and*
3166 *min_spacing=500*. After the filtering options, 97 out of 115 individuals (97-97 per population) were
3167 retained. For genetic diversity and differentiation analysis, the data was filtered with *indmiss=0.7,*
3168 *snpmiss=0.2, min_mac=2, dohefilter=TRUE and min_spacing=500*. After filtering 94 out of 115
3169 individuals (94-94 per population) were retained. After filtering 56328 out of 178847 SNPs were
3170 retained. Thinning (removal of missing data) reduced the dataset further to 2100 SNPs. The
3171 proportion of missing data per population can be summarised using Figure 2 below. Note that
3172 populations with proportion of missing data >0.07 were filtered out for genetic structure and
3173 diversity/differentiation analysis.

3174



3175 Figure 2. Boxplot to show the proportion of missing data after the filtering procedure over populations.

3176

3177 The “dohefilter” option in the “filterdata()” command refers to filtering of function which
3178 remove SNPs with heterozygosity levels which are potentially indicative of paralogs. Paralogs are
3179 genes that are present in a particular organism which are related to each other through gene
3180 duplication events (Koonin., 2005). Paralogous genes are often exacerbated in plant genetics
3181 because the events of gene duplications such as polyploidy is more prevalent in plants (Mastretta-
3182 Yanes et al., 2014). Paralogous genes have also been found to bias population genetic estimates,
3183 which will affect downstream population genetic type of analyses (Verdu et al., 2016). Since *Linum*
3184 was found to have undergone polyploidy, the identification of paralogs due to heterozygosity may
3185 be an issue for further population genetic type analysis (Sveinsson et al., 2014). The identification of
3186 paralogs was able to be inferred from locus specific heterozygosity against the locus specific minor
3187 allele frequency. The plot below (Figure 3) indicates these locus specific frequencies and reveals the
3188 amount of data which SambaR were able to filter out, due to the identification of paralogs: (The SNP
3189 dataset in red were subsequently removed after the “dohefilter” option was set to “TRUE”).

3190

3191
3192
3193
3194
3195
3196
3197
3198
3199
3200
3201
3202
3203
3204
3205
3206
3207
3208
3209
3210
3211
3212
3213
3214
3215
3216
3217
3218
3219
3220
3221




Figure 3. A scatterplot to reveal Sambar’s identification and filtering of paralogs.

In this case, SambaR’s default parameters for removing the paralogs were only removing a small proportion of heterozygosity (highlighted in red in Figure 3). This reveals that not all paralogs were potentially removed with SambaR’s “*dohefilter*” option. This was perhaps due to SambaR’s conservative use of Hardy Weinberg’s Equilibrium (HWE). HWE makes assumptions that in the populations, there are: no mutation, random mating, no gene flow, infinite population size, and no selection. With the knowledge that *Linum* are highly selfing (see previous chapters), the “*dohefilter*” option of SambaR may not be optimised for removing paralogs in *Linum* sequences (Jahnke & Etterson., 2019). Heterozygosity needs to be manually addressed from the input files before processing in SambaR. SambaR’s “*filterdata*” function filters individuals based on their proportion of missing datapoints considering all SNPs. As an addition to heterozygosity filter, SambaR were also instructed to filter SNPs based on SNPs depth (“*snpdepthfilter=TRUE*” command), This command filters SNP which has a high read depth. Subsequently, the function filters SNPs based on their proportion of missing datapoints considering retained individuals only. The filtering of individuals could return different estimates of missing data and, as a result, different numbers of retained SNPs and individuals occurs. In population structure filtering (i.e. more individuals and less SNPs), the GC-content of the retained dataset equalled to 0.53 and the ‘transversion vs transition’-ratio equalled 0.59. For genetic diversity and selection analysis (i.e. less individuals and more SNPs), the GC-content of the retained dataset equalled 0.53 and the ‘transversion vs transition’-ratio equalled 0.57. In the filtering for diversity and selection, linkage disequilibrium (LD) estimates were able to be calculated using *PLINK* (*-genome -r2 -ld-window-kb 1000000 -ld-window -r2 0*). LD can infer trait-associated region(s) of a genome which may be of interest to further studies. However, for short-read sequences LD estimates may be redundant. This is because most of the reads do not cover all

3222 sites and consequently restricts LD to loci which are potentially very close (Maruki & Lynch., 2014)
3223 (Bilton et al., 2018).

3224

3225 4.3.7 Analysis: Genetic Analysis in SambaR

3226 For the purposes of Structure analysis, several analyses took place within SambaR. The
3227 analysis is often based on *Nei's genetic distance* between populations (Nei, 1972). This includes
3228 Correspondence analyses (CA), Principal coordinate analyses (PcoA), Principal component analyses
3229 (PCA), DAPC analyses using principles such as Landscape and Ecological Association (LEA) tests
3230 (Frichot & Francois., 2015). Other genetic distance related analyses were also illustrated in SambaR.
3231 For Correspondence Analyses (CA), analysis was performed using the function "*dudi.coa*" of the R
3232 package *ade4*-1.7.19 (Dray and Dufour, 2007; Bougeard and Dray, 2018). Data was imputed per
3233 SNP/individual by calculating genotype probabilities from population specific minor allele
3234 frequencies. Principal coordinate analyses (PcoA) were performed using the function "*pcoa*" of the R
3235 package *ape*-5.6.2 (Paradis and Schliep, 2018). This occurred on distance matrices containing 3
3236 different measures of genetic distance, with *Nei's genetic distance*, calculated with the function
3237 "*stamppNeisD*" of the R package *StAMPP*-1.6.3 (Pembleton et al., 2013), Hamming's genetic
3238 distance, calculated with the function "*bitwise.dist*" of the R package *poppr*-2.9.3 (Kamvar et al.,
3239 2014), and pi (pairwise sequence dissimilarity), calculated with the function "*calcp*" of the R package
3240 SambaR. The principal component analyses (PCA) were performed using the function "*snpgdsPCA*"
3241 of the R package *SNPRelate*-1.28.0 (Zheng et al., 2012). DAPC analyses were performed using the
3242 function "*dapc*" of the R package *adegenet*-2.1.5 (Jombart, 2008; Jombart and Ahmed, 2011), both
3243 with and without prior population assignment. Multi-dimensional scaling (MDS) was performed
3244 using the function "*cmdscale*" (metric MDS) of the R package *stats*-4.1.3 (R Core Team, 2022) and
3245 the function "*isoMDS*" (non-metric MDS) of the R package *MASS*-7.3.57 (Venables and Ripley, 2002),
3246 on a Euclidean distance matrix generated with the function "*dist*" of the R package *stats*-4.1.3 (R
3247 Core Team, 2022). Neighbourhood joining (NJ) clustering was performed using the function "*NJ*" of
3248 the R package *phangorn*-2.8.1 (Schliep, 2011), using as input a Hamming's genetic distance matrix
3249 between individuals, calculated with the function "*bitwise.dist*" of the R package *poppr*-2.9.3
3250 (Kamvar et al. 2014). Bayesian population assignment (BPA) probabilities were calculated and
3251 plotted using the functions 'assign2pop' and 'plotassign2pop' of the R package SambaR. The optimal
3252 number of clusters (K) was determined using the elbow method on cross-entropy scores generated
3253 by the 'snmf' function, with the assumption that the startpoint of a plateau represents the optimal
3254 K.

3255 Divergence measures were measured using locus specific Fst estimates (according to Wright
3256 (1943), Nei (1977), and Cockerham and Weir (1987) (for all pairwise population comparisons) were
3257 subsequently calculated with the functions 'runWrightFst', 'locusNeiFst', and 'locusWCFst' of the R
3258 package SambaR. HWE, (2D) folded site frequency spectra (SFS), Tajima's D and genome wide
3259 heterozygosity analyses were able to be executed using the function 'calcdiversity' of the R package
3260 SambaR.

3261 Genome wide heterozygosity (genomeHe) was calculated in SambaR for each sample using
3262 the formula: $genome\ He = (He_seg * N_seg) / N_total$, in which: N_seg = the quantity of sites
3263 segregating within the population to which the tested individual belonged. He_seg = the proportion
3264 of Heterozygous sites within the investigated individual for those segregating sites. N_total = the
3265 total length of sequenced sites (polymorphic as well as monomorphic) which were able to pass the
3266 filter settings.

3267 Geographical maps were generated with the function 'getMap' of the R package rworldmap-
3268 1.3.6 (South, 2011). Piecharts were added using the function 'add.pie' of the R package mapplots-
3269 1.5.1 (Gerritsen, 2018). Admixture coefficients were implied with the functions 'snmf' and 'Q' of the
3270 R package LEA-3.8.0 (Frichot and Francois, 2014). Alpha was set to 10, tolerance to 0.00001, and
3271 number of iterations to 200. Ancestry coefficients were calculated with the software Admixture-1.3
3272 (Alexander et al., 2009) and illustrated using the 'plotstructure'-function of SambaR.

3273

3274 4.3.8 Analysis: Ancestry Coefficients


3275 As part of our population structure analysis, individual ancestry coefficients were able to be
3276 inferred using Landscape and Ecological Association (LEA) test. This was done using the package LEA
3277 under R (Frichot & Francois., 2015). These tests analysed population structure based on selection on
3278 a whole genome level. LEA applies landscape genomic data and identification of allele frequencies
3279 that illustrates genetic association with ecological associations. The LEA package derives adaptive
3280 alleles from large data sets, often referring to previous ecological association which is implied as
3281 ancestry coefficients (Frichot & Francois., 2015).

3282

3283 4.3.9 Analysis: Using different Genome reference

3284 There are multiple whole genomes now publicly available for *L. usitatissimum* and none for
3285 *L. bienne*. For this study we mapped to the *L. usitatissimum* whole genomes. They utilized different
3286 sequencing tools and have different read depths. For the purposes of this analysis, these differences

3287 between the reference genome could translate into differences in *snpmiss* and *indmiss* parameters
 3288 in the SambaR input. These differences translate into the number of SNPs and individuals able to be
 3289 analysed. For the purposes of comparison, we used another *L. usitatissimum* (Atlant variant) genome
 3290 sequenced by Nanopore and Illumina sequencing and contain another cultivar type when compared
 3291 to the CDC Bethune genome used for previous analyses in this thesis. The cultivar type used for this
 3292 whole-genome sequencing was found to have low variability of morphological and anatomical
 3293 characteristics under stress conditions, suggesting a variant that is adept under stress conditions.
 3294 After sequencing they found 8.4 Gb of sequence data with N50 of 12kbp and a read coverage of 23×,
 3295 and Illumina read coverage of 30× (Dmitriev et al., 2021). The CDC Bethune (v2) contained a
 3296 summary of 21.80 Gb HiFi reads generated with N50 of ≥12kbp (Sa et al., 2021). The difference in
 3297 initial data quality can be summarised in the below data quality plot when compared against our
 3298 short reads:




3307 Figure 4. Sambar's data quality plots for the CDC Bethune genome (4A) and the Atlant genome (4B).

3308

3309 The plot above illustrates the data quality when short reads are aligned to either the CDC
 3310 Bethune genome (Figure 4A) or the Atlant genome (Figure 4B). This quality plot revealed that
 3311 retained individuals for CDC Bethune genome was around the same as the Atlant whole genome
 3312 (120 individuals). A *snpmiss* parameter of 0.05 and *indmiss* parameter of 0.7 for example, will return
 3313 ~100 individuals with ~10k retained SNPs (before thinning). When aligning to the Atlant genome, this
 3314 number was reduced by 10. This reveals that at least for these two different genomes (both
 3315 different in terms of sequencing methods and cultivar type), that alignment of the short read
 3316 sequences to different reference genome have some effects in terms of the quality of retained SNPs

3317 and retained individuals after filtering in SambaR. The CDC Bethune genome reveals a higher
3318 preference based on the number of individual samples kept.

3319 One of the most important qualities when looking at the quality after filtering, is the amount
3320 of heterozygosity in the data after filtering. For this, two plots can be shown to illustrate linear
3321 relationship between the proportion of Heterozygosity in all sites and segregating sites. Using the
3322 same filtering parameters, two plots were illustrated with to compare between the different
3323 reference the short reads were aligned to.



3326
3327
3328
3329
3330

3331 Figure 5. Plots for heterozygosity after filtering short reads with the CDC Bethune genome (5A) and the Atlant
3332 genome (5B). Each dots illustrates different individuals differentiated by population (colours).


3333

3334 Figure 5A is the linear plot for Heterozygosity (He) proportion on all sites and segregating
3335 sites for the data aligned to the CDC Bethune reference genome. The Figure 5B is the same plot for
3336 the data aligned to the Atlant reference genome. This revealed that aligning the short reads to either
3337 genome will result in significant amounts of Heterozygosity. There were differences in the rankings
3338 of the individuals within the population, noted by the differences in colour labelling. This reveals that
3339 proportion of heterozygosity remains mostly the same even using different reference genome. This
3340 potentially convey those heterozygous reads in our short reads are real heterozygous reads or
3341 perhaps reads in the genome due to the polyploidic nature of *Linum*. However, it is interesting how
3342 the rankings of these heterozygosity have changed within populations in our short reads. For the
3343 purposes of structure, diversity and differentiation analysis in the populations sampled here, we
3344 aligned to the more commonly used CDC Bethune genome.

3345

3346 **4.3.10 Analysis: Removing Heterozygosity**

3347 As was seen in Figure 5A, all sites heterozygosity ranges from 0.000 to 0.20 within the
3348 samples and under different reference genomes. This was potentially an issue as the presence of
3349 heterozygosity might present an issue with paralogs which is especially of high importance to
3350 organisms which have previously undergone whole-genome duplication events such as polyploidy
3351 (McKinney et al., 2016). This is also revealed in the “locus specific heterozygosity” in the plot used
3352 for identification of paralogs in SambaR’s “dohefilter” function. There were potential paralogs due to
3353 these heterozygousities. SambaR’s filtering option may not remove all paralogs (Figure 6). In
3354 addition, we expected *Linum* to be highly selfing and polyploidy. As a result of this, explored options
3355 to filter for heterozygosity.



3363 Figure 6. Scatterplots to reveal SambaR’s identification and filtering of paralogs for the default “dohefilter”
3364 option, revealing potential paralogs not filtered by SambaR (in black).

3365

3366 An option to avoid paralogs is to remove heterozygous alleles altogether, however the
3367 consequence of removing 100% heterozygosity is that further analysis such as the population
3368 diversity analysis will be made redundant due to the lack of heterozygous alleles (Chapman et al.,
3369 2009). In SambaR this will return an error whereby SambaR will specify that all samples are
3370 homozygous and therefore a population diversity measure is made redundant. Measures of
3371 heterozygosity, such as multi-locus heterozygosity (MLH) is useful for predicting whether
3372 populations are inbreeding or outcrossing in terms of their breeding strategies (Jensen et al., 2007).
3373 It is then preferable if the heterozygosity in the data were not entirely removed. To do this there is a
3374 requirement to specifically filter out a certain number of heterozygous reads. This can be done by
3375 removing SNPs and/or loci locations which have a higher than the threshold amount of
3376 heterozygosity, manually.

3377

3378 4.3.11 Analysis: Filtering for Heterozygous Alleles

3379 To avoid the redundancy of a population diversity analysis, a specific genotype filter
3380 measure to exclude locations with exceptionally higher heterozygosity were applied to the original
3381 population output of STACKS. To do this, the original VCF output file were first converted from multi-
3382 allelic to biallelic genotype/allele reads. This was done using the function “*bcftools norm -m*” under
3383 *bcftools* V1.10. Formatting of alternative alleles were done using the command “*bcftools view -e*
3384 *“FORMAT/AD[:1]<2 && INFO/AD[1]<5”*” and extraction of genotype per sample is done by using the
3385 command “*bcftools query -f “%CHROM %POS[\t%GT]\n”*”. Formatting of alleles and extraction of
3386 genotypes were done in *bcftools* V1.10 (Li., 2011). Once genotyped and formatted to biallelic reads,
3387 the file were processed in R using genotyping to mark heterozygous allele as “1” and homozygous
3388 allele as “0”. Once heterozygous allele is marked as “1”, a filter was applied whereby loci location
3389 with more than extreme heterozygosity can be identified and outputted. We decided to filter only
3390 for the top 1% heterozygous alleles to avoid missing important information from these potentially
3391 true heterozygous alleles. Filtering for heterozygosity can be done using the addition of
3392 “*which(means[,q]>quantile(means,0.99))*” to the initial allele output in R. This will remove top 1%
3393 of loci with excessive “1s” or heterozygous alleles. R was then asked to output a “.txt” file whereby it
3394 listed the loci position which can then be filtered out in *vcftools*. The full command lines for this
3395 process can be found under appendix 23.

3396 We used *vcftools*’ “*--exclude-positions*” command in *Vcftools* for filtering against the output
3397 heterozygosity file (Danecek et al., 2011). First, the “.txt” file was made sure to be tab-delimited
3398 and was converted into a “Unix .txt” format using the Linux command “*awk ‘{sub(“\r\$”, “”); print }’*
3399 *winfileinput.txt > unixfileoutput.txt*”. These “.txt” formatting options were essential so that *vcftools*
3400 could read the loci positions without errors under *Linux* commands. In the same command, the file
3401 can be recoded using the “*--recode*” command of *vcftools* and can be pipelined to output a .VCF file
3402 output whereby exceptionally heterozygous loci location was filtered out. The full command for this
3403 process can be found under appendix 24. This would keep out loci with exceptionally high
3404 heterozygosity which may represent paralogous alleles while preserving an amount of
3405 heterozygosity which potentially represent outcrossing populations within our sample sets.

3406

3407 4.3.12 De novo mapping

3408 Mapping to *L. usitatissimum* was preferred in this analysis, to conserve the number of
3409 individuals and populations able to be analysed. However, using *L. usitatissimum* whole genome as a

3410 reference for population samples of *L. bienne* may not be optimal for SNP calling during further
3411 bioinformatic analysis, this may have implication on retained SNPs. We investigated how mapping
3412 “de novo” will impact the results gathered from mapping to the *L. usitatissimum* whole genome.
3413 After filtering for missing SNPs and individuals in SambaR, we found only 68 individuals were viable
3414 for further analysis when mapped de novo. After filtering and thinning, we found a total of 3029
3415 kept SNPs. The number of individuals kept for de novo mapping were significantly less than that
3416 from the option of using *L. usitatissimum* whole genome as a reference, where we kept 100
3417 individuals. We also lost potentially significant results because population 12 (Mediterranean
3418 population) were not kept from de novo mapping. Despite the loss of individuals using de novo
3419 mapping, we kept a significantly higher number of SNPs than mapping to the *L. usitatissimum* whole
3420 genome (kept only 800 SNPs with *L. usitatissimum* reference). This suggests that potential SNPs may
3421 have been lost due to mapping to *L. usitatissimum* whole genome. However, in the interest of
3422 keeping more individuals for population structure inference, we mapped to the *L. usitatissimum*
3423 reference. For future repeats of this population analysis, we suggest the use of *L. bienne* whole
3424 genome when they become publically available and/or including many more individual/population
3425 samples in the ddRADSeq protocol before mapping de novo. This may result in increases in the
3426 number of individuals retained after de novo mapping. We also suggest using more population from
3427 the Mediterranean region of Europe to observe for more variation from this region.

3428

3429

3430

3431 **4.4 RESULTS: IMPLICATIONS OF POPULATION ANALYSIS**

3432


3433 The final heterozygous-filtered data contained 866 SNPs (after filtering and thinning). After
 3434 the filtering procedure, SambaR retained 100 individuals that makes up to 15 populations with
 3435 retained grouping as Northern ($> 45^\circ$ North) and Southern ($\leq 45^\circ$ South) (Table 2), as described in
 3436 Landoni et al (2022). We expected to see structuring between Northern and Southern populations,
 3437 supporting those reciprocal responses to traits seen in Landoni et al (2022).

| Initial Population Name | Latitude | Group | Final Population Prefix |
|-------------------------|----------------|-----------|-------------------------|
| Cul | Not Applicable | Cultivars | CUL |
| 3 | 36.03633 | Southern | A-3 |
| 1 | 36.80044 | Southern | B-1 |
| 10 | 37.88211 | Southern | C-10 |
| 6 | 37.93551 | Southern | D-6 |
| 12 | 42.31008 | Southern | E-12 |
| 13 | 43.02902 | Southern | F-13 |
| Lla | 43.40738 | Southern | G-Lla |
| Vil | 45.09393 | Northern | H-Vil |
| Tal | 47.6997 | Northern | I-Tal |
| Mat | 48.35697 | Northern | J-Mat |
| Dor | 50.6 | Northern | K-Dor |
| low2 | 50.68183 | Northern | L-low2 |
| Tym | 53.30307 | Northern | M-Tym |
| Sut | 53.35291 | Northern | N-Sut |

3438 Table 2. List of output populations and their details of locality after the final filtering and thinning options of
 3439 SambaR. Populations which are found $\leq 45^\circ$ South were grouped as a more Southern population than those
 3440 $> 45^\circ$ in latitude. Apart from the cultivars (CUL), the populations were alphabetically ordered by latitudes with
 3441 the most Southern population first. Populations were specified and grouped as per previous reciprocal results
 3442 under Landoni et al (Landoni et al., 2022).

3443 In the following results the prefix of the population would be in the format of “Latitude (in
 3444 ascending order)-Population”. For example, for individuals from the most Southern population 3 the
 3445 label would be “A-3” and the most Northern population Sut the labels would be “N-Sut”. For Figures
 3446 depicting individual comparison in the sample (such as genetic distance trees), the data will be
 3447 formatted by “population_individual”, for example individual 10 belonging to the population 12 will
 3448 be formatted as “12_10”. The full individual and population details is available in appendix 25.

3449 Our samples represent a wide range of latitude within Western Europe, representing Northern and
 3450 Southern groups as was observed in Figure 7. In this results section we analysed the *Linum* samples
 3451 as stated above in terms of their genetic diversity, structure, and divergence.



3452

3453 Figure 7. A colour map revealing the geography of the samples after passing filters and thinning with reference
 3454 to their colours, population, and regions on the labels. For our analyses, Southern population are the wild
 3455 Spanish population and Northern population are wild population found in France and the UK. This grouping
 3456 aligns with previous reciprocal study groupings (Landoni et al., 2022)

3457

3458 4.4.1 Results: Genetic Diversity

3459 4.4.1 i Genome-wide Diversity

3460 We aimed to see population differentiation as well as to observe potential differences in
 3461 breeding strategies. We looked at genome wide diversity to observe these. We expected to see a
 3462 difference in diversity between Northern (> 45° North) and Southern (≤45° South) wild populations in
 3463 our samples.


3464 The bar plot reveals a lower proportion of segregating sites in most of the Northern
 3465 populations (Figure 8A). French populations such as “I-Tal” and “H-Vil” seems to reveal the lowest
 3466 proportion of segregating sites (0.0462 and 0.1039) along with the Northern UK populations of “L-
 3467 low2” and “M-Tym” (0.0727 and 0.1016). In exception of this is the French population of “J-Mat”,
 3468 which seems to have a high proportion of segregating sites. In contrast to this low segregating site
 3469 proportion in most of the Northern populations, the more Southern populations have a higher
 3470 proportion of segregating sites (>0.2). The highest proportion of segregating sites is observed in the

3471 Mediterranean population “E-12” (0.3915). These observations may suggest that the Southern
 3472 populations are more likely to contain genes which are not conserved and are therefore potentially
 3473 more outcrossing in terms of their breeding strategy when compared to their Northern relatives.

3474 In addition to this, the proportion of heterozygous sites can also suggest potential diversity
 3475 within populations. When looking at heterozygosity within segregating sites only (Figure 8B), there
 3476 wasn’t much of a pattern to be observed. The proportion of heterozygote alleles for all sites,
 3477 however, suggests that most of the latter, Northern population are less heterozygous. This suggests
 3478 that perhaps the Southern populations are employing a strategically different breeding system. We
 3479 expected the level of heterozygosity to not differentiate as much between the Southern and
 3480 Northern population should they all be “selfing” populations. It may be the case that the more
 3481 Southern populations are more outcrossing in breeding strategy than those of the more Northern
 3482 populations in our sample lists.

3483

3484



3485 Figure 8. Summaries of nucleotide diversity measures for our *Linum* samples. 8A. Bar chart to show the
 3486 proportion of segregating sites within each population levels in our samples based on regions. 8B. Box plots to
 3487 show the proportion of heterozygous alleles within the segregating sites, within each population in our


3488 samples based on regions. 8C. Box plots to show the proportion of heterozygous alleles for all sites within the
3489 genome. 8D. A scatterplot to show the nucleotide diversity or pi (%) against the genome wide heterozygosity
3490 (%) for all individuals colour coded by population. The colours on every part of this Figure corresponds to
3491 population list as inferred in the Figure.

3492

3493 4.4.1ii Multi-Locus Heterozygosity

3494 A measure of multi-locus heterozygosity can also be observed with a boxplot (Figure 9). In
3495 the box plot, an observation of multi-locus heterozygosity was illustrated. In the x-axis, the
3496 populations were lined alphabetically according to their latitude, with the most Southern
3497 populations to the left. Southern populations such as population B-1, C-10, E-12, F-13, and D-6 were
3498 seen to have a more diverse multi-locus heterozygosity, agreeing with the sequence heterozygosity
3499 as observed in previous section. When looking at the Northern population, the box plots are forming
3500 closer to 0. This infers less multi-locus heterozygosity in the more Northern individuals. Interestingly,
3501 population 3, which is a Southern population region-wise, is trending more with the Northern
3502 population and the Northern population “K-Dor” seems to have a more diverse multi-locus
3503 heterozygosity. Less multi-locus heterozygosity in a population is thought to illustrate more
3504 inbreeding within that population and less outcrossing. It could be that the breeding strategy of
3505 population 3 is genetically influenced by the more Northern population and are therefore more
3506 inbred than the rest of the Southern population and the opposite is true for some Northern
3507 populations such as “K-Dor”.

3508



3509 Figure 9. A boxplot to reveal multi-locus heterozygosity at a population level.

3510

3511 **4.4.1iii Numberof private alleles**

3512 A measure of the number of private alleles can suggest population divergence as higher
3513 mutation rate may be implied (Szpiech & Rosenberg, 2011). We reveal a higher number of private
3514 alleles for our cultivars than most of the wild populations observed within this study except
3515 population E-12 (Figure 10).

3516 Our analysis suggests that most of our wild population have around 228 – 246 out of 866
3517 SNPs which are private alleles. Our cultivars have more, at 269 SNPs showing as private alleles. The
3518 Mediteranean population E-12 suggests an even higher number of SNPs showing private alleles at
3519 374 SNPs. This suggests more discinction in this population, which suggests population divergence
3520 for the Mediterranean population.




Figure 10. Bar chart to illustrate the number of private alleles over population. # SNPs = 0.866k, number of private allels are in thousands (k).

3521


3522 **4.4.1iv Variation in Segregating Sites (Tajima's D)**

3523 One way to look at population divergence is to examine variation within segregating sites for
3524 the population. This will infer whether selection have occurred which have removed variation or
3525 there is potential selection in which variation is maintained within population. To look at this, we can
3526 look at the populations and how many sites are variable and how identical individuals are within the

3527 population. Tajima's D values can be interpreted for our populations, whereby a measure of pi and
 3528 pi related to the number of variable sites relative to the number of sequences can be normalized as
 3529 Tajima's D (Korneliussen T. et al., 2013). Negative Tajima's D will suggest that there is selection
 3530 removing variation within population and population is recently expanded. Positive Tajima's D will
 3531 suggest that there is selection maintaining variation and populations are not expanding.

3532 Tajima's D can be illustrated through a bar chart for every population within our samples
 3533 (Figure 11). It is with confidence that none of the population observed within our samples results in
 3534 positive Tajima's D. This reveals that there are selection removing variation within all *Linum* samples
 3535 in this study. This was however more observed in two populations (populations 12 and 6). There is a
 3536 suggestion here that based on the more negative Tajima's D, Southern populations are more
 3537 recently expanded than the Northern populations. In addition to this, summary of Tajima's D
 3538 estimation was able to be obtained from SambaR (see appendix 28) and a subsequent t-test were
 3539 able to be implied between the North and South wild populations. The t-test revealed that Tajima's
 3540 D estimation for Northern and Southern populations were significantly different ($P(T \leq t) \text{ two-tail} =$
 3541 $0.023 (<0.05)$).

3542



3543 Figure 11. A bar chart to show Tajima's D statistics for the different populations.

3544

3545 4.4.2 Results: Genetic Structure

3546 When the data was fully prepared with potential paralogs removed (top 1% most
3547 heterozygous loci removed), it is with further confidence that we can carry out structure analysis
3548 without paralogous alleles. To reveal population structure, we constructed a Ward D neighbour
3549 joining tree. When genotyped heterozygous filters were applied, the tree suggests genetic
3550 structuring between the different populations (Figure 12).


3551 The Ward.D neighbour joining tree reveals that there are three main population clusters.
3552 The tree reveals that most of the cultivars are grouped together (Figure 12). The Southern
3553 population were grouped further away from the Northern and Cultivar individuals in our sample set.
3554 In terms of the wild, there is an observation of two groups whereby Southern population were
3555 grouping as genetically closer to each other and the Northern population grouping on together. This
3556 reveals that, genetically, in terms of their population the two Southern and Northern populations
3557 are potentially genetically distanced to each other. The Southern group are also more genetically
3558 distanced to the cultivars and may have a higher allelic richness. The cultivar “Suz” grouped with the
3559 Northern cluster. In this cluster, there also occurred some Southern individuals such as the Southern
3560 population “3”. This could be due to unexpected history of migration in population A-3. We suspect
3561 that this is due to human errors such as seed labelling and labelling in the laboratory that may have
3562 led to these individuals not grouping to their clusters as expected. Future studies could examine
3563 structuring between more individuals from these population that may infer migration or
3564 hybridisation in population A-3.

3565 In other observations, the population highlighted in red circle are the only “Mediterranean”
3566 population observed within our samples (population E-12). On the heterozygous-filtered, unrooted
3567 tree, this group seems to be structurally different to the rest of the samples. They do not group with
3568 either Northern or Southern wild populations. This Mediterranean population seems to have a
3569 unique genetic structuring to the rest of the wild groups as well as the cultivars observed in this
3570 study. This reveals that in the samples contained within our RADSeq analysis, population E-12 is
3571 potentially unique to the rest of the wild populations found in the West, which are closely grouped
3572 to either the North or South cline of the population.

3573 In summary, the genetic network reveals that within the Western Europe wild *Linum*
3574 populations, there were observations of genetic structuring, generally between populations
3575 originating in the North and South of Western Europe. There are some individuals that are out of
3576 place from their expected cluster, which could indicate potential gene flow or human errors within

3577 the data. The discovery of population E-12 being structurally unique to either North or South
 3578 populations is an indication that the more Eastern, Mediterranean populations are structurally
 3579 different to the rest of the wild populations in the West.

parsimony score per site: 1.94; -log(likelihood): -10755



3580 Figure 12. A Ward.D (Ward's minimum variance method) Neighbour-joining tree for the illustration of genetic
 3581 distance on an individual level. The genetic tree reveals structuring in our samples with Northern and Southern
 3582 main clines for the wild. Cultivars are revealed to be more closely related to the Northern cline.

3583


3584

3585 4.4.2i Principle Coordinate Analysis (PcoA)

3586 When looking at genetic structure, dissimilarity between the individuals based on Nei's
3587 genetic distance can also be observed. A principal coordinate analysis (PcoA) was conducted, with
3588 PC1 (representing 46.7% variance) and PC2 (representing 29% variance) plotted to examine genetic
3589 similarity (Figure 12). Another multivariate statistical technique that will go hand to hand with the
3590 PcoA analysis is the more descriptive Correspondence analysis (CA), whereby another plot can be
3591 illustrated in appendix 26. There are observations here of clusters for both PcoA and CA analysis,
3592 highlighted in red boxes. Southern and Northern populations are mostly clustering with what seem
3593 like a cline between the two clusters highlighted (Figure 13 and Appendix 26). In addition,
3594 population E-12 seem to be clustering separately in what we have suggested as "Mediterranean
3595 population".

3596 Some Southern individuals from populations such as F-13 and A-3 can be observed clustering
3597 closer to the Northern cluster, suggesting potential gene flow between Northern and Southern
3598 populations. Some Northern populations such as the French population "Mat", can also be observed
3599 to have an individual outside of the cluster, closer to the Southern populations. The Spanish
3600 population of "Lla" are clustering with their relatives in the North. This suggests that this population
3601 may have implication of gene flow through past genetic events.

3602 Another point of interest is the separation of the population E-12 which we suggest as being
3603 the "Mediterranean population". In this analysis, this was the only "Mediterranean population" able
3604 to be analysed due to DNA material availability. They seem to be structuring differently to both
3605 Northern and Southern groups of the Western European *L. bienne* samples in this study (Figure 13
3606 and Appendix 26).



3607 Figure 13. A PcoA scatter plot with PC1=47.3% of total variance and PC2=34.9% of total variance. The red
 3608 boxes highlight potential clustering of populations. The plot agrees with the structuring suggested by the
 3609 genetic distance tree, in addition, a few intermediate individuals between the Northern and Southern
 3610 populations were also suggested, forming a potential cline between the Southern and Northern population.

3611

3612


3613 4.4.2ii Sequence Dissimilarity

3614 One way to illustrate genetic distance is to look at the proportion of dissimilarities between
3615 individual sequences or otherwise known as π . This is the average pairwise difference between
3616 individuals. A sequence dissimilarity measure (π) was illustrated as a matrix to illustrate
3617 dissimilarities between individuals and within their sequences, with Southern populations showing
3618 more dissimilarities (Figure 14).

3619 In terms of genetic structuring, there is an observation that most of the Southern
3620 populations are showing a degree of sequence dissimilarities when compared to the Northern
3621 populations (Figure 14). The sequence dissimilarity measures agree with the finding observed with
3622 the genetic tree, that the Southern and Northern populations are genetically distanced. Most of the
3623 Northern individuals are showing little signs of dissimilarity between themselves. For example,
3624 sequence dissimilarity between individuals in the Northern population “N-Sut” and “I-Tal” is close to
3625 0. Dissimilarities are observed to occur on the more Southern population with our *Linum* sample set
3626 (Figure 14).

3627 The sequence dissimilarity (π) matrix (Figure 14) also reveals that population E-12 (pink strip
3628 on the matrix) has the highest level of sequence dissimilarity (π) comparatively between all the
3629 other individuals analysed in this study. This is highlighted by the more intense shading of yellow
3630 when comparing individuals belonging to population E-12 to the rest of the individuals in the sample
3631 set (revealing π values between 0.385 to 0.44). What’s more interesting is that this dissimilarity is
3632 even more than the observed difference for cultivated and wild individuals between in our samples,
3633 when genotyping for highly heterozygous genes were implied to the data.

Sequence dissimilarity (π)




3634

3635 Figure 14. A sequence dissimilarity (π) matrix for individuals summarised into populations. The colour chart on
 3636 the left indicates the population each colour strip is representing in the matrix.

3637

3638 Similar observations were also seen in Nei's genetic distance. The higher the Nei's genetic
 3639 distance is the more distinct the sequence is when compared to the rest of the samples. A matrix
 3640 can be illustrated to observe Nei's genetic distance (Figure 15). One major observation in this matrix
 3641 is the higher number of Nei's genetic distance highlighted in shades of orange for the Southern
 3642 population (B-1, C-10, E-12, F-13, A-3 and D-6) when compared to the cultivars. This number was
 3643 observed to be lower for the Northern individuals. This further supports the findings that cultivars are
 3644 more genetically related with our Northern wild population as seen in our previous analyses.
 3645 Population A-3 is an exception from this trend seen with the Northern and Southern population. This
 3646 is perhaps a signal of genetic flow with the Southern population. Population A-3 is perhaps more
 3647 genetically derived from the more Northern population observed in our sample set, making them
 3648 more of a "Northern" population in terms of their genetic make-ups. A significantly higher Nei's
 3649 genetic distance was also observed with population "E-12" when compared with the rest of the
 3650 samples (>0.4), which can be inferred as Nei's genetic distance revealing more sequence distinction
 3651 on population 12 when compared with other populations. To put to perspective, the next highest

3652 Nei's genetic distance would be at 0.201 between the cultivated population and population 13. In
 3653 the literature, a Nei's genetic distance of more than 0.250 is considered as significant genetic
 3654 distance between populations observed (Nei M., 1972; Wright S, 1978). This finding suggests that
 3655 the Mediterranean population E-12 is considered as genetically more distanced to the rest of the
 3656 samples observed within this study.



3657 Figure 15. A Nei's genetic matrix between the populations observed in this sample. The matrix reveals a higher
 3658 genetic distance between the Northern and Southern populations. Darker orange shades reveals a more
 3659 significant genetic differences.

3660

3661 **4.4.2iii DAPC Analysis**


3662 As an addition, we used a multivariate method to illustrate population structure. One
 3663 method is the Discriminant Analysis of Principal Component (DAPC) (Miller et al., 2020). The DAPC is
 3664 calculated in SambaR by scoring the number of retained Principle Components (PCs), interpolation of
 3665 this scoring and scoring cumulative variance explained by the Principle Component Analysis (PCA)

3666 (De jong M. et al., 2021). The summary DAPC results for our genotype filtered data can be found
3667 under Appendix 27. A geographical map can be inferred using DAPC whereby population clustering
3668 can be illustrated under a population based on their latitude and longitude of origin (Figure 16).

3669 SambaR outputs pie charts on a map based on our population's locality to illustrate DAPC
3670 scoring (Figure 16). The Figure is separated by the number of clustering per population or otherwise
3671 known as 'K'. The cultivar individual "Mar" was excluded from this analysis as this was an exclusive
3672 Canadian population in the sample and we are interested only on Western European Latitudes for
3673 this study. From the Figure, we can observe groupings based on a DAPC analysis. When $K=2$ there
3674 was already some grouping of the Southern population. However, the cultivars are not distinctive
3675 yet. When $K=3$, the population 'E-12', highlighted in red became distinct. When $K=4$ populations
3676 showed further grouping whereby the four groups (Northern, Southern, Cultivars, Mediterranean)
3677 were observed as can be implied by the genetic distance tree. Here, the cultivars are seen as
3678 grouping in yellow, Northern populations in 'colourless', Southern population in brown and the
3679 population 'E-12' in red. Higher K 's can also be inferred whereby some of the individuals from
3680 Southern populations are grouping closer to the Northern populations as inferred in $K=5$. When $K=6$
3681 the Southern populations are further split, separating away from each other. When $K=4$, it can be
3682 inferred that some cultivars populations such as those in France and Netherlands are grouping with
3683 the Northern populations. The Netherlands population is the cultivar population "Suz", which if we
3684 refer to the genetic distance tree, is also grouping with the Northern clines, suggesting genetic
3685 relationship with Northern population and perhaps cultivation from these populations.

3686

3687




3688

3689 Figure 16. A geographical map of all the populations conveying genetic differences based on DAPC analysis
3690 and separated by K=2-6. Groupings was observed better when K=4. When K=4 Southern wild individuals were
3691 grouping in brown, Northern wild individuals colourless and the Mediterranean population in red. The cultivars
3692 were observed to be more spread out and are marked with a red dot in the middle of the respective pie charts
3693 under K=4.

3694 4.4.2iv Landscape and Ecological Association

3695 As part of the population structure analysis, a Landscape and Ecological Association analysis
3696 (LEA) were utilised whereby ancestry coefficients can be implied between populations. Number of
3697 sub-populations (K) was described at 1-6. A Cross-entropy criterion can be used to determine the
3698 best run for a fixed value of K. The plot below illustrates the cross-entropy criterion for our dataset.
3699 The lower the cross-entropy value the better prediction capability a K value has.


3700 The Cross-entropy criterion plot reveals that for our dataset, at higher K-values the minimal
3701 cross-entropy was the lowest (Figure 17). This starts to level off when K=4. This levelling off minimal
3702 cross-entropy suggests that the optimal number of populations of K would be equal to 4 for further
3703 LEA analysis. An LEA bar plot with ancestry coefficients can be constructed for K=1-6 to imply any
3704 admixture within our dataset.



3705 Figure 17. Cross-entropy criterion graph illustrating the best number of population (K) is K=4. When the trend
3706 levels off.

3707 We further illustrated genetic relationships between our sample using an LEA barplot
3708 revealing some admixture within the Southern and Northern populations when $K \geq 4$ (Figure 14).
3709 Admixture is suggested by the presence of sequences/SNPs from multiple genetic clusters for an
3710 individual. The Northern population were less admixed. The cultivar populations were more similar
3711 to the Northern accessions (Figure 18) when $K \leq 3$. They begin to differentiate when $K = \leq 4$. Admixture
3712 was also observed with the Northern accessions showing most of the coefficient changes when
3713 $K = \leq 4$.

3714 In the LEA plot there was also observation of divergence in population E-12. This is observed
3715 by the diverging bars on population E-12, observed when $K \geq 3$ (Figure 18). The LEA agrees with the
3716 distinction of population 12 from the other groups observed in the other genetic structure analyses
3717 above.




3718 Figure 18. A LEA bar chart for every individual, separated into populations for K=2-6. The population names are
 3719 in alphabetic order of the most South latitude to the most North.

3720

3721 To observe this admixture even further, a map can be drawn with pie charts that reveals the
 3722 ancestry coefficients of different population based on their location (Figure 19). The pie charts on
 3723 the geographical map represents the genetic relatedness of each population to another based on an
 3724 LEA analysis. Just as the bar charts on Figure 14, it is revealed that admixture occurs within some of
 3725 the populations observed within this analysis. When K=3, admixture within the Southern wild
 3726 population can already be seen, with some of the more Southern population revealing LEA
 3727 association with the more Northern population highlighted in green for K=3. When K=4, the cultivars
 3728 was differentiated further. This is marked by the colour yellow on the pie chart. There seems to be
 3729 admixture in the supposedly cultivated population originating from France and in the Netherlands.
 3730 This was shown in K=4 whereby these populations are observed to have contained admixture from
 3731 every population but the population E-12 (Figure 19). Admixturing occurred in the cultivar

3732 populations “Suz” and “Tin”, suggesting genetic relatedness of these cultivars it’s wild ancestors
3733 (Figure 19).




3734 Figure 19. A geographical map of all the populations conveying ancestry differences based on LEA for K=2-6.

3735 4.4.3 Results: Population Divergence

3736 As observed in the LEA analysis, there was some potential divergence occurring within one
3737 or more population of our samples. To infer this further, we can look at measures such as population
3738 dissimilarity based on their SNPs. Another way of looking at diverging population is to observe their
3739 various allele frequencies. Wright's F_{st} values suggest differentiation between populations. F_{st} value
3740 of "0" infers no variance between the population whilst F_{st} value of "1" suggests complete variance
3741 between the population compared, illustrating potential differentiation and divergence (Wright.,
3742 1965; Weir., 2012; Bird K. et al., 2017).

3743 A matrix can be drawn from Wright's F_{st} values (Figure 20 on the next page). The matrix
3744 reveals a summary of Wright's F_{st} values against each of the population summarised within this
3745 study. As with the Nei's genetic distance matrix, the darker shade of orange reveals a higher Wright's
3746 F_{st} value. The matrix suggests higher Wright's F_{st} values between Southern and Northern Population
3747 in contrast with Northern to Northern population. This supports further the structuring observed
3748 within the genetic distance tree, the DAPC analysis and the LEA analysis. Variation in their Wright's
3749 F_{st} values suggest that population is more diverse and different than initially thought. In this
3750 analysis, it is further suggested that the Mediterranean population E-12 have a higher total genetic
3751 variance at a population level against other wild population. A high Wright's F_{st} value suggest a
3752 considerable degree of differentiation and divergence in population E-12 when compared to the rest
3753 of the population in this study.



3755 Figure 20. A matrix depicting Wright's F_{st} value. Higher value is highlighted in darker orange shades

3756

3757 4.4.3i Population Dissimilarity

3758 We observed dissimilarity of Sequence between individual population. It is with interest for
 3759 this study to observe the most divergence populations by observing the population which has the
 3760 most dissimilar SNPs with the other populations. As a starting point, variation and divergence can be
 3761 looked at for the individuals within the same population.

3762 The boxplot (Figure 21) reveals the overall sequence (Figure 21A) and SNP only (Figure 21B)

3763 dissimilarity of individuals within a population. The dissimilarity is low overall for all populations

3764 (>5% for overall sequence dissimilarity and >0.2 proportions for SNPs only). Although this was the

3765 case, there was some differences which can be observed. In the earlier more Southern populations,


3766 the SNP dissimilarity within populations were more diverse than that of most of the Northern

3767 populations, except for the French population “J-Mat”. Population 3 is showing the most

3768 dissimilarity within the Southern population. There is potential here that within population 3, either

3769 they are more diverging than other populations, or there was contamination within the sampling

3770 processes whereby other population may have been unintentionally selected as population 3.



3771 Figure 21. A boxplot to show Sequence (A) and SNP (B) dissimilarity between the populations in this study.

3772 For completion, a measure of nucleotide diversity (π), as was mentioned under section


3773 “4.4.2 *Genome-wide Diversity*” of this chapter can be compared and mean π for each population

3774 can be extracted from SambaR (Figure 22). The mean π was tested using a two sample t-test. The t-

3775 test revealed that there was a significant mean π difference between the wild population groups

3776 (Northern and Southern populations) ($P(T \leq t)$ two-tail = 0.019 (<0.05)). This illustrates divergence in
3777 the genetic diversity of Southern and Northern populations.

3778



3779 Figure 22. A bar chart illustrating mean pi for each population in our sample as outputted by SambaR.

3780

3781 4.4.3ii Minor Allele Frequency

3782 Another measure of population divergence can be done using the minor allele frequency
3783 (MAF). Minor Allele Frequency (MAF) is widely known as the frequency on which the second most
3784 common allele occurs in a sequence of a given population. They have been shown to play a role in
3785 population selection and divergence because MAF variants occurs once, and they drive a significant
3786 amount of selection (Hernandez et al., 2019). This would give an idea of how varied a genotype is for
3787 a given SNP. This can be used to differentiate between common and rare variants in the population
3788 (Linck & Battey., 2019). If the MAF is low, it may imply that the major allele for the SNP is conserved.
3789 A high presence of common alleles may also reflect signs of genetic bottlenecks (Marth et al., 2004).
3790 Alternatively, a high presence of rare alleles may suggest that a population is expanding (Marth et
3791 al., 2004).


3792 A matrix was drawn to illustrate MAF variance between the population samples within this
3793 study (Figure 23 on the next page). Higher variance in MAF variance is revealed in the more
3794 Southern population highlighted in shades of orange when MAF variance = >0.04. This reveals that
3795 major alleles maybe conserved more between the Northern population. This difference may suggest

3796 that certain genes are conserved in more Northern individuals. MAF variance are also highest in the
 3797 Mediterranean population “E-12”.

3798 For further comparisons, a mean MAF of each population (see appendix 29) were able to be
 3799 extracted from SambaR and a two-sample t-test were applied to the mean MAF dataset (Figure 24).
 3800 The t-test revealed that the difference in mean MAF values between Southern (A-G) and Northern
 3801 (H-N) populations were significant ($P(T \leq t) \text{ two-tail} = 0.009 (<0.05)$).

3802


3803



3804

3805 Figure 23. A matrix showing Minor Allele Frequency (MAF) between the different populations tested in this
 3806 thesis.

3807



3808 Figure 24. A Bar chart to represent the mean MAF values over each population.

3809

3810

3811 4.5 DISCUSSION AND CONCLUSION

3812 4.5.1 Genetic structure

3813 We hypothesized that within our population samples, there are population groupings based
3814 on their geographic origin. The Ward.D neighbour-joining tree revealed this, showing genetic
3815 structuring between northern and southern populations, and between wild and cultivars within our
3816 sample set. The PCoA based on Nei's genetic distance also supports this. This genetic structuring
3817 suggests possible genetic variation within Western European *Linum bienne* populations. The PcoA
3818 and CA analysis revealed grouping based on northern and southern regions, with some intermediary
3819 individuals in between the clusters. In addition, the pie-charts based on the DAPC agreed that for up
3820 to four clusters (K=4), there was structuring related to the geography of the *Linum* populations of
3821 this study. The sequence dissimilarity matrix revealed that the northern populations were
3822 differentiated from those of the southern populations. These results confirm structuring according
3823 to the geography of the wild *Linum*. Although this is the case, filtering for heterozygosity may impact
3824 measures such as the dissimilarity matrix and genetic distancing that maybe of true nature as
3825 opposed to paralogs. In this thesis, we were conservative towards avoidance of paralogs, based on
3826 the knowledge that our study models are highly-selfing and prone to gene duplication events such as
3827 polyploidy, which can complicate assembly of our sequences (Mastretta-Yanes et al., 2014). With a
3828 1% heterozygous filter we managed to remove the most heterozygous individuals whilst retaining
3829 heterozygosities that may suggest dissimilarities important for population diversity and divergence
3830 analyses. We also filtered at 5% heterozygosity where we begin to see populations dropping off for
3831 further analyses. Future studies may look at a range of heterozygous filter for comparison to this
3832 study.

3833 The geographic distribution of a plant population can lead to variation within species, so it is
3834 not surprising to find this in our wild *Linum* samples. In the model *Arabidopsis thaliana*, it has been
3835 illustrated that polymorphisms revealed differences within the population genetics of northern and
3836 southern populations (Fodorenko et al., 2001). In addition, studies of the model *Arabidopsis thaliana*
3837 have revealed patterns of genetic structuring that signals evolutionary processes such as migrations
3838 (Shirsekar et al., 2021). Additionally, Mediterranean populations of *Arabidopsis thaliana* revealed
3839 relationship that are closer to their relatives further South in Morocco and North Africa (Brennan et
3840 al., 2014). In that study, there was genetic structuring between northern and southern accessions.
3841 The more southern Mediterranean population were revealed to be related to populations further
3842 south of the range of the samples studied. There could be similar case made here when looking at
3843 the northern and southern clustering of wild *Linum* populations within our samples, with the
3844 Southern population suggesting genetic distinction from the Northern clusters. When looking at

3845 genetic structuring, these could be influenced by interaction of ecological and genetic processes
3846 such as local adaptation to seed dispersal. Ecological barriers to seed dispersal, due to the
3847 biogeography of Northern and Southern Europe may also play a part in the structuring seen in these
3848 Western European wild *Linum* populations. Seed dispersal maybe limited in Southern and Northern
3849 regions of Europe thus limiting gene flow between the two wild groups, causing this genetic
3850 structuring. This barrier to dispersal was observed in the Southeast Asian mangroves (Wee et al.,
3851 2020) and in peatmosses (*Sphagnum*) (Kyrkjeeide et al., 2016). Studies suggests that climate is the
3852 dominant determinant of plant range in Europe but in addition, species dispersal plays an important
3853 role in the genetic flow of a population (Normand et al., 2011).

3854 Another case made in the arctic-alpine plant species *Lloydia serotina*, was that reproductive
3855 biology can result in population structuring (Jones and Giddeon., 1999). Genetic differentiation that
3856 suggests variation in breeding system can also be observed in studies using three orchid species (Sun
3857 and Wong., 2001). In this study we observed signals of geneflow between Northern and Southern
3858 populations. We also saw that Southern population maybe more outcrossing in terms of breeding
3859 strategy. It maybe plausible that there is variation in breeding strategies between Northern and
3860 Southern clines of wild *L. bienne* observed in this study, this may be resulting from a seed dispersal
3861 barrier (ecological and geographical) ultimately resulting in genetic structuring. Studies on breeding
3862 strategies of wild *L. bienne* is scarce. We think that Southern population are more outcrossing based
3863 on higher heterozygosity. However, more samples needed to be studied in the future before this
3864 can be confirmed.

3865 Genetic structure analysis suggested that some individuals did not cluster with either
3866 northern or southern groups. This is potentially a signal for geneflow. If this signal is gene flow
3867 between Southern and Northern population, this would impede local adaptations within the
3868 Northern wild *Linum* populations. In *Arabidopsis*, it has been suggested that different populations
3869 under different environments showed this genetic structure (Hämälä et al., 2019). Another
3870 possibility is that historically, Northern populations diverged from these populations with some
3871 individuals not clustering under PCoA. However, the number of these individuals not clustering is
3872 limited to a few individuals. We saw no individuals forming a cline with the Mediterranean
3873 population E-12. This suggest that this population is divergent, and perhaps more possibility of local
3874 adaptation occurred in the Mediterranean populations.

3875 We saw in our neighbour joining tree, PcoA and DAPC analysis that the Northern and
3876 Southern wild population are more divergent from each other than the cultivars are to either
3877 population groups. The cultivars are more closely related to the Northern wild individuals. If these
3878 cultivars are genetically closer to the Northern groups, we expect them to have some phenotypic

3879 similarities observed in the other chapters such as their pollen viabilities and the relative gene
3880 expression to flowering initiation. We didn't find sensitivity to vernalization requirement in our
3881 cultivars as we did in the Northern wild populations. There wasn't any phenotypic case where it may
3882 suggest that the cultivar population are closely related to the wild Northern populations. However, it
3883 may be that the cultivars are genetically more related to the wild Northern populations than they
3884 are to the Southern populations due to cultivation from the Northern populations. Artificial selection
3885 in the cultivars may have implications on the variation on phenotypes between the two species.

3886 The cultivars "Suz" and "Tin" were found to be grouping with Northern wild populations as
3887 opposed to the cultivar group (Figure 12). We think that there's potential mislabelling, either in
3888 previous seed collections or in the laboratory, with regards to the cultivars "Suz" and "Tin" as seen in
3889 our Ward.D tree. These cultivars were grouped with the Northern population, however, are very
3890 unlikely to be doing so since there was no phenotypic case suggesting this. We take caution with any
3891 results regarding these cultivars.

3892

3893 4.5.2 Population Diversity and Divergence

3894 Genome-wide diversity measures looking at proportion of segregating sites suggests a lower
3895 level in most Northern populations and a higher population in the Southern population. A higher
3896 amount of segregating site can suggest a higher mutation rate for the Northern populations and
3897 expansion further North (Fu Y., 1995). Segregating sites can suggest evidence of positive selection in
3898 a population (Gilad Y. et al., 2002; Przeworski M., 2002; Booker T. et al., 2017). Suggestion of more
3899 occurrence in positive selection could be interpreted by a higher number of segregating sites within
3900 our Northern groups, potentially driven by the local adaptation to the environment (Figure 11).

3901 Positive selection is often associated with selective sweep, suggesting less genetic diversity in our
3902 Northern populations (Booker T. et al., 2017). Less genetic diversity could imply more inbreeding in
3903 these Northern population.

3904 Furthermore, studies using *Camellia sinensis* (Tea) and their wild relatives suggests genetic
3905 divergence in the wild relatives by looking at proportion of segregating sites and in addition, the
3906 proportion of heterozygous sites (Yang H. et al., 2016). We looked at the proportion of
3907 heterozygous sites for all sites in our study model. Most of these sites were filtered due to filtering
3908 of paralogs. However, for all sites, surviving heterozygous alleles were higher in the Southern
3909 populations, suggesting allelic richness (Figure 8C). This suggest that the Southern populations are
3910 genetically more diverse than the Northern population. Northern population may be more inclined

3911 to be conservative towards local adaptation to colder seasons. This was perhaps less desirable in the
3912 Southern populations due to warmer climates.

3913 The Mediterranean population E-12 was the most distinct among our samples. Based on
3914 Nei's genetic distance these Mediterranean individuals are clustering on their own as opposed to
3915 either Northern or Southern clines. It is thought that this Mediterranean population is genetically
3916 distinct from the rest of the Western European wild population. It is possible that the more
3917 genetically distinct population E-12 was closer to wild relatives in the eastern Europe region, but this
3918 was not tested in this study. We did not have materials for the wild *Linum* population originating in
3919 eastern European region such as Turkey and Italy, in which both wild and cultivated flax have been
3920 observed. There is even a possibility that population E-12 is more related to population as far as the
3921 eastern Mediterranean and the Middle East regions, because the cultivated flax (*L. usitatissimum*) is
3922 thought to be native to these regions before they were introduced to the more Northern climates
3923 (Sen, T. and Reddy, H.J., 2011). It would be of interest for future research to include more
3924 Mediterranean and Middle Eastern populations, to determine their genetic relationships with our
3925 samples. The distinction of population E-12 was also supported by the Tajima's D statistics, whereby
3926 population E-12 had a more negative Tajima's D than the rest of the populations sampled in this
3927 study. This may suggest that this population were diverged in the past, and this was a result of
3928 population expansion in wild *Linum*. In past studies, low Tajima's D in the fir species *Cunninghamia*
3929 *konishii* have been inferred as a post-glacial expansion with their populations (Hwang et al., 2003). A
3930 study in *L. flavum* suggested a post-glacial migration history in *Linum*, which may result in a
3931 population expansion during the post-glacial period (Plenk et al., 2017).

3932

3933 4.5.3 Breeding Strategies in Western European *L. bienne*

3934 Population diversity measures such as the measures of heterozygosity along with the multi
3935 locus heterozygosity also differed between the southern and northern populations. When filtering
3936 for excessive heterozygous alleles, remaining heterozygosity were observed mostly in Southern
3937 populations, with most of the Northern population inferring no heterozygosity (Figure 9). In addition
3938 to the structuring observed in the genetic distance tree and the PcoA, wild populations might also be
3939 differentiating in the way they breed. We observed a higher heterozygosity in Southern plants in all
3940 sites (Figure 8c and Figure 9). The more heterozygous populations suggest less inbreeding and more
3941 outcrossing than the less heterozygous northern populations. Sharry and Lord (1996) also explored
3942 heterozygosity, concluding that less heterozygous populations may have been more inbred than
3943 those of the more heterozygous populations (Sherry & Lord., 1996). We think that our Southern *L.*

3944 *bienne* populations are more outcrossing and that our Northern population are inbreeding, perhaps
3945 to conserve local adaptations or perhaps due to climatic barriers for seed dispersals. This
3946 differentiation would also be supported by the difference seen in Wright's F_{st} matrix. It is seen that
3947 F_{st} values between Northern and Southern populations are higher as highlighted in shades of
3948 orange. In the literature it is accepted that an F_{st} value greater than 0.15 between populations can
3949 be considered as significant in differentiating populations (Luo et al., 2019). This adds to the genetic
3950 structuring revealed by the genetic distance, PcoA and CA analysis that northern and southern
3951 populations are genetically different. In addition, there is potential that Southern population are
3952 more outcrossing in terms of their breeding strategy. This agrees with the findings on genetic
3953 structuring observed in section 4.41 "Results: Genetic Structure" and can so therefore observation in
3954 genetic diversity supports the finding that there is potential breeding strategy variation in Northern
3955 and Southern clines of *L. bienne*, causing genetic structuring between the two Western European
3956 wild *Linum* populations.

3957

3958 4.5.4 Genetic variation in Western European *L. bienne*

3959 Molecular diversity and association analyses in the past have revealed the potential of
3960 genetic variation in wild flax for the development of cultivated flax (Soto-Cerda et al., 2014). It is
3961 interesting that the cultivar populations within our sample sets seemed to be more genetically
3962 closely related to the Northern populations than to the Southern populations. The genetic distance
3963 tree and the ancestry coefficients on the LEA bar plot agreed that the cultivars in our samples were
3964 more closely related to the more northern wild accessions than so the southern accessions.
3965 Variation in genetic makeup within the *L. bienne* gene pool represents the potential of genetic
3966 variation for the improvement of cultivated flax in Northern Europe. Traits that are conserved in
3967 Northern populations in response to colder climate can be useful for flax cultivation in different
3968 growing seasons.

3969 To add to the potential genepool resources, we can infer some ecological interpretation for
3970 our wild populations. This includes selection and bottlenecks events that may have arisen in the
3971 past. Tajima's D suggest that there is less variation compared to what the population can sustain. In
3972 population genetics studies, negative Tajima's D have previously suggested population expansions
3973 and differentiation after a bottleneck (De Jong et al., 2011; Gunther et al., 2016). This negative
3974 Tajima's D was strongest in the more southern populations, particularly in population E-12. This may
3975 suggest that some southern populations, such as population E-12 have diverged through past
3976 expansions and potentially still expanding after a bottleneck. High and low divergent populations

3977 revealed by the F_{st} values could suggest population availability for breeding programmes
3978 (Baiakhmetov et al., 2021). The higher F_{st} values among the more southern populations and lower
3979 among the northern population suggested that there were potential differences in gene flow
3980 between populations. The minor allele frequency (MAF) measures between the northern and
3981 southern populations also revealed a significant divergence between the two groups. The lower
3982 observed MAF within the Northern populations might suggest that these populations are more
3983 conserved in terms of their genetic strategies, therefore may have resulted in the smaller sequence
3984 and nucleotide diversity whilst Southern population may have a larger diversity to provide a wider
3985 genepool for cultivar breeding programmes.

3986 In conclusion, the SNPs data revealed genetic structuring among the Western European wild
3987 relatives of flax as implied by the genetic distance tree. This was further supported by the PcoA,
3988 DAPC, and LEA analyses. There was also potential that this genetic differentiation could mean
3989 further differences in terms of local adaptations. Differences observed in heterozygosity and F_{st}
3990 values revealed that there maybe a difference in breeding strategies of these wild *Linum*
3991 populations. Population divergence measures also revealed that these northern and southern
3992 populations were divergent. It would be with further interest to observe SNPs diversity from a wider
3993 region such as Eastern Europe and the Eastern region of the Mediterranean. The only Mediterranean
3994 population observed in this case was population 12 and it was interesting to observe the difference
3995 in genetic structuring and genetic divergence here. However, we make these suggestions with
3996 caution as the number of individuals that were analysed was relatively small, especially in the
3997 southern region. There may also be potential human errors as was observed in some odd individuals,
3998 not grouping where they should be expected within their local range. It would be with greater
3999 interest to run this type of analysis with a wider sample range, more genotype data, and a more
4000 individuals in our sampled populations to strengthen these initial findings within our *Linum*
4001 populations.

4002

4003

4004 CHAPTER 5: VARIATION IN TRAITS BETWEEN VERNALIZED AND NON-
4005 VERNALIZED *LINUM*.

4006

4007 Environmental factors can affect more than one plant trait (Campetella et al., 2020). Plant
4008 traits can correlate differently between populations and geographic location even down to individual
4009 levels in the wild (Jiang et al., 2021). *Linum* is an oilseed and natural fiber crop, with a significant
4010 agricultural value. Oilseed and Fibre crops have different target traits in the interest of plant
4011 breeders. For oilseed types, oil content and number of the capsules and seed sizes are the most
4012 important traits for breeding (Çopur et al., 2006), whereas in fibre types, plant height and number of
4013 stems are the essential traits for fibre flax breeding (Xie D. et al., 2018). We already established that
4014 wild *L. bienne* could provide a wider genepool for breeding purposes. In this chapter we explored
4015 traits such as plant height, stem number and bud numbers due to vernalization as an environmental
4016 variable in wild our *L. bienne* samples. This can provide insights to how vernalization may affect traits
4017 in wild *Linum*.

4018 In chapter two, we saw how vernalization affected the number of days to flowering in both
4019 *Linum* species, suggesting variation in number of days to flowering between wild individuals from
4020 across Western Europe. In this chapter we explored how other traits could be affected by
4021 vernalization. In *Linum*, Genome-Wide Association (GWAS) have identified candidate genes fo traits
4022 such as plant height, number of branches and seed weight, revealing a biological basis for
4023 improvement amongst these traits for agricultural interests (Xie D. et al., 2018). In this chapter,
4024 phenotypic measures, such as plant height, plant stem numbers, and plant bud numbers will be
4025 quantified, analysed, visualized, and discussed further as an additional finding to support implication
4026 of wild *L. bienne* as a wider genepool. These traits are of importance, especially in terms of
4027 agriculture because the candidate genes for them have been identified which reveals potential for
4028 improvement in *L. usitatissimum*. We will test whether other traits were significantly affected due to
4029 vernalization treatments and investigate their relation to environmental variables such as latitude
4030 and local climate variables. We hypothesize that traits are significantly affected by vernalization
4031 treatments.

4032 A plant stem is one of the main structures of a vascular plant, the other being roots. They
4033 often support other major important floral organs such as the leaves, flowers, and fruits of the
4034 plants. They are also required for support, nutrient storage, and productions of new living tissues
4035 (Raven, 1982). Studies in the pepper species *Capsicum annum*, and maize identified changes in
4036 stem morphology due to varying light wavelength and soil temperature (Schuerger A. et al., 1997;

4037 Walker J., 1969). These early findings suggest that stem morphology is affected by light and
4038 temperatures as environmental variables. Due to wide latitudinal range in our *L. bienne* samples
4039 across Western Europe, there are possibility that stem formation can vary under different
4040 environment perhaps because of local adaptation. We hypothesized that the number of stems are
4041 varies under different vernalization treatments, and we further suggest that there is variation to
4042 correlation to the environment and to the number of days to flower when comparing vernalized and
4043 non-vernalized conditions. Reduction in stem numbers for vernalized individuals will suggest that
4044 vernalization acts as an abiotic stress and that control for stem numbers could act as a local
4045 adaptation for requirement to vernalize.

4046 The buds of a plant are widely known as the undeveloped embryonic shoot which is often
4047 found on the axil of a plant leaf (Walters, Keil and Walters, 1996) A *Linum* bud is a small lateral or
4048 terminal protuberance on the stem of a vascular plant that may develop into a flower, and
4049 subsequently form seeds (Trelease, 1931). This is therefore an important factor to the production of
4050 seeds and of interest to oilseed breeding. In *Arabis alpina*, it is suggested that exposure to colder
4051 environments initiates formation of flowering buds (Lazaro et al., 2018). This suggests potential
4052 adaptations and differentiation within this species due to vernalization mechanisms (Torång et al.,
4053 2015). In this chapter we observed whether development of buds is affected by vernalization for our
4054 *Linum* samples. We hypothesized that the number of buds able to form varied due to vernalization
4055 treatments. We further suggest that there is variation on bud formation under local climatic
4056 variables.

4057 In addition, we will observe seed areas of wild *L. bienne* seed samples from our vernalization
4058 experiment as well as seeds from the wild. Under abiotic stress, it is widely understood that seed sizes
4059 have important consequences for germination in plants. In *Anthoxanthum odoratum* it is observed
4060 that population with larger seeds had a higher probability of germinating (Roach D., 1987). In the
4061 winter annual plant *Dithyrea californica*, amount of precipitation (climatic variable) had a beneficial
4062 effect on plant fecundity and influenced seed-size survival selection (Larios E. et al., 2014). This
4063 suggests seed-size natural selection due to environmental factors in wild populations. Environmental
4064 factors during seed development have also been illustrated to influence seed germination in *Lotus*
4065 *tenuis* (Clua and Gimenez., 2003). On a species level, initial results using mountain alpiners indicates
4066 elevation-dependence seed production is specific to each species (Olejniczak P. et al., 2018). This
4067 reflects different resource allocation strategies for different species due to environmental variables
4068 introduced under different altitudes. More recently, research in naturally occurring sand rice
4069 (*Agriophyllum squarrosum*) suggests variation in seed sizes due to local environments. They suggest
4070 that large-seeded individuals were more competitive in semi-arid regions (Zhao P. et al., 2022). With

4071 all these in mind, it is evident that in the literature, variation in seed sizes occurs in many plant study
4072 species with local environmental variables, potentially playing a role in natural selection of seed sizes
4073 in plant populations.

4074 Seed size is a trait of interest in *Linum*. Variation in seed size due to vernalization treatment
4075 in the wild may indicate local adaptation to the environment in terms of seed sizes. In addition to this,
4076 seed sizes of *Linum* wild relatives can be useful in their cultivar relatives. This is because linseed is one
4077 of the biproduct of *Linum*. We will investigate if our wild *L. bienne* vary in seed sizes when under
4078 different vernalization treatments. Germination of plants are suggested to be related to their seed
4079 sizes, so seed size can also count for population fitness (Keddy & Constable., 1986). A study on onions
4080 (*Allium cepa* L.) also suggests that vernalization temperatures, duration and bulb size significantly
4081 influenced seed yield of all cultivars (Muthamia., 1994). This suggests that in many plant species
4082 requirements of vernalization will have a significant effect on seed health which suggests seed sizes
4083 and seed yield which is of interest in agriculture. We hypothesize that seed sizes for different *Linum*
4084 populations will have been affected by vernalization and this will have represented a correlation
4085 between seed size and environmental variables such as latitude of localities and climatic variables.

4086 With the above in mind, it is of both ecological and agricultural interest to investigate the
4087 consequences of vernalization in *Linum* traits. Findings here can be linked to local adaptation in the
4088 wild *L. bienne* and implications of potential loss of vernalization in cultivated *L. usitatissimum*. In this
4089 chapter we investigate whether vernalization treatments affects the fitness of wild *L. bienne*. For
4090 traits, we expect to find variation between vernalized and non-vernalized individuals. We expect seed
4091 sizes to also vary between Northern and Southern individuals, suggesting local adaptation in seed
4092 sizes. Implication of requirement of vernalization can then suggest that any adverse traits observed
4093 within the more Northern individuals due to vernalization stress are avoided by the potential loss of
4094 vernalization as was observed under chapter 2 of this thesis.

4095

4096

4097 5.2 METHODS

4098

4099 Regarding experimental setup, vernalization methods are the same as found in chapter 2 of
4100 this thesis (see section 2.2.1 “Samples and Experiment” for vernalization setup). For measures
4101 regarding plant height, plant stem number, and plant bud numbers, observations were made
4102 according to when an individual is observed as flowering for the first time. This is when the
4103 individual has had their first flower, which is fully opened, with petals revealing the flower’s sexual
4104 organs. This will coincide with the measure of the number of days a plant takes to flower and will
4105 reduce biases based on plant age. Measurement for flowering initiation and plant trait measures
4106 such as plant heights can be directly compared against each other. A measure of plant height is done
4107 using a 2-sided measuring tape, using Centimetres (Cm) as a unit of measurement. The height
4108 measurement was taken from where the plant has emerged from the soil onto the tallest upper
4109 stem of the plant, measuring the total observable height of plant material for the individual. Stems
4110 are stretched until a straight measure can be observed using the measuring tape. The measure of
4111 plant stems and buds were based on observations of the number of either plant stems or plant buds
4112 per individual at first flowering. Conditions for the vernalization and non-vernalization treatments
4113 can be found under section 2.2.1 “Samples and Experiment” of chapter 2 in this thesis.

4114 For measures regarding plant heights, stem numbers and bud numbers, data was only
4115 observed for the 2018 vernalization experiment. There was phenotypic measurement for both 2018
4116 and 2021 vernalization experiment underwent in chapter two of this thesis. However, the 2021
4117 vernalization experiment data for these were limited because of Covid-19 restrictions in place at the
4118 department of Biosciences during the time of the experiment. This caused phenotypic measures to
4119 be lost in most individuals during the 2021 vernalization experiment. There will also be a bias on
4120 earlier flowering plants, as plants with later flowering will have not had a chance to flower before
4121 the Bioscience departmental lockdown in March 2020 (see section 2.4 “Results: flowering time
4122 between experiments from different years” of this thesis). For this reason, the phenotypic measures
4123 for the 2020-21 vernalization experiment were discarded for future analysis. Environmental data
4124 such as the latitude of origin and climatic variables were used as was suggested in chapter 2 of this
4125 thesis.

4126 In addition, we observed seed sizes. For this observation we only used the wild *L. bienne*
4127 individuals as we were mainly interested in relationship between seed sizes and environmental
4128 variables. We observed seed sizes of vernalized and non-vernalized wild individuals along with wild
4129 SO individuals. We did this observation using non-fluorescence microscopy. A 10×10 objective was

4130 used to observe *Linum* seeds using a colour camera mounted on a Leica DMI-3000 Microscope. The
4131 Microscope was hardwired to a Windows computer running Leica's LAS X software, where
4132 measurements of seed length and width can take place. Measurement of the seed area is done by
4133 using ImageJ to measure pixels and calculate area. To do this, images of 5 seeds per-individual were
4134 captured using a microscope with a 10×10 objective. We used Leica's LAS X software to capture and
4135 save images before processing in ImageJ. In ImageJ, the images were individually measured. We
4136 calibrated ImageJ to each image by drawing a 1mm line and setting a scale to that line as 1mm. The
4137 image is then converted into 8-bit format and made binary (Process → Binary → Make Binary).
4138 Although ImageJ usually outputs binary images without holes, there are occasions where holes in
4139 the seed can be observed. Potential holes were able to manually be filled or automatically via the
4140 "fill holes" function (Process → Binary → Fill Holes). Images were then adjusted by threshold (Image
4141 → Adjust → Threshold) and analyzed by using the "Analyze particles" function of ImageJ. This
4142 function can analyze particles (pixels) and calculate area of a of each binary colour (black and white).
4143 The seeds were measured as the area in black. Values were inputted into Microsoft office and
4144 formatted for further analysis in R.

4145

4146 5.2.1 Data Analysis

4147 All phenotypic measures were statistically tested during this chapter and all additional data
4148 on this chapter were analysed using R v 4.1.2 (R Core Team, 2022) and Rstudio (Rstudio Team, 2020)
4149 as a graphical interface. For modelling and correlation purposes the R package "ggpubr"
4150 (Kassambara, 2022), "ggplot2" (Wickham and Sievert, n.d.), and "car" (Fox and Weisberg, 2011)
4151 were used to execute general-linear modelling (GLM). There are several assumptions made for this
4152 GLM; the data is normally distributed, and that values of trait measurements, and other variables
4153 are independent of each other. We also assumed that the variance in data residual is the same. For
4154 summary of statistics, model can be summarised in terms of adjusted r^2 , F-statistics, and p-values.
4155 The full R commands for the analysis during this chapter is available in appendix 30. Barcharts can be
4156 illustrated using the "barplot" function of R. A two-tailed t-test were then used to compare data
4157 under vernalized and non-vernalized treatments.

4158 For correlation analysis to environmental variables, data for latitude were collected for each
4159 population sampled from the wild at the place of collection by their respective collectors, as per
4160 chapter two of this thesis. Principle component values variables were able to be gathered for
4161 different climatic variables as was described under chapter 2 of this thesis. Scatterplot for
4162 correlation between traits and environments were illustrated using "ggscatter" function of R with a

4163 GLM modelling with a Pearson's correlation method. The same is applied for analyses of seed sizes
4164 against environmental variables.

4165

4166


4167 5.3 RESULTS

4168 5.3.1 Plant Traits vs. Days to Flowering

4169 In chapter two of this thesis, we looked at the relationship between the number of days to
4170 flowering and latitude for our 2018 vernalization experiments. The results demonstrates a positive
4171 relationship between the number of days a plant takes to flower and the latitude of locality to each
4172 plant. During this experiment, we also examined other plant traits such as plant height, and the
4173 number of stems. We ask whether the number of days to flowering will have other effects in other
4174 traits. This is interesting as in chapter one we discussed that different plant traits in *Linum* may
4175 correlate with each other. We measured three different traits in the 2018 vernalization experiments
4176 (plant height, stem numbers, bud numbers). These measures were plotted against the number of
4177 days the plant takes to flower and faceted based on treatments. The measurements were made in
4178 relation to the first flowering. As individual plants were first flowering, the plant height, number of
4179 stems and number of buds were also observed and recorded. The raw count data for this can be
4180 found under appendix 31.

4181 Correlation reveals the relationship between the flowering time initiation (number of days
4182 to flower (labelled on y-axis as “Days_to_fl”) and overall plant height (labelled on the x-axis as
4183 “Height_(Cm)”) for no-vernalization and vernalization treatments (Figure 1). A Pearson’s correlation
4184 reveals that the number of days to flowering is negatively correlated with the plant height at first
4185 flowering, under the no-vernalization treatments ($R=-0.28$, $p<0.001$). This contrasted with what is
4186 observed in the vernalization treatment, whereby, there were no trends observed ($R=-0.022$, $p=0.79$)
4187 (Figure 1). This illustrates that, with no-vernalization the number of days to flowering were
4188 negatively affecting plant height (i.e., the longer a plant takes to flower the smaller (in height) they
4189 tend to be at first flowering). This effect was reduced when the plants were vernalized.


4190



4191 Figure 1. Two scatterplots to illustrate the relationship between the number of days a plant takes to flower
 4192 (Days_to_fl) and he plant height (Height_(Cm)). The plot revealed a contrast between no-vernialization and
 4193 vernalization treatments, with a significant negative correlation under the no-vernialization treatment ($R=-$
 4194 $0.28, p=<0.001$).

4195

4196 Furthermore, a comparison between the plant height and the different treatments was
 4197 illustrated using a two-sample t-test on Figure 2 (assuming unequal variance). The t-test suggests
 4198 that there was a significant difference on the plant height between the two treatments ($P(T<=t)$ two-
 4199 tail= <0.001).



4200 Figure 2. A bar chart to illustrate the significant difference between no-vernialization and vernalization
 4201 treatments on mean plant height (Height (Cm)) ($P(T<=t)$ two-tail= <0.001).

4202 Another trait we measured during the study is the number of stems at first flowering. The
 4203 result suggests that when no-vernalization and vernalization treatments were applied to the *Linum*
 4204 samples, a positive correlation can be observed between the number of days a plant takes to flower
 4205 and the number of observed stems on first flowering (Figure 3). However, the strength of the
 4206 relationship seems to have been reduced when plants were vernalized. R values for the vernalized
 4207 treatments were reduced from $R=0.69$ to $R=0.18$.
 4208





Figure 3. Scatterplots to illustrate the relationship between the number of days a plant takes to flower (Days_to_fl) and the number of observed stems at first flowering (Stem_no) for no-vernalization and vernalization treatments. For both treatments, the correlation was significantly positive (no-vernalization $R=0.69$, $p<0.001$; vernalization $R=0.18$, $p=0.011$).


4209
 4210 To observe whether this change of stem number was significant, a two-sample t-test
 4211 (assuming unequal variance) can be implied for the two sets of data (Figure 4). The t-test revealed
 4212 that the number of stems observed after first flowering was significantly lower when individuals
 4213 were vernalized ($P(T<=t)$ two-tail= <0.001). These findings suggest that the number of stems
 4214 observed when vernalized is significantly reduced, and the positive correlation between the number
 4215 of days to flowering and the number of stems observed was also reduced. However, the correlation
 4216 was still significant between the two traits, even when vernalized ($p<0.05$ for both treatments).
 4217



4218 Figure 4. A bar chart to illustrate the mean stem number differences between the two treatments, showing a
 4219 significant reduction in stem number when vernalization occurred ($P(T<=t)$ two-tail= <0.001).

4220

4221 Another plant traits which were observed during the vernalization study is the number of
 4222 buds. The relationship between the number of days to flowering and the number of observed buds
 4223 on the first flowering suggests that there was a relationship when plants were introduced to
 4224 vernalization treatments (Figure 5). Interestingly, this reduction corresponds with the of number of
 4225 days to flower due to vernalization where it was also reduced. Perhaps vernalization is also affecting
 4226 the ability of buds to be formed. The correlation between the number of days a plant takes to flower
 4227 and the observed number of buds on first flowering suggests that when vernalized they're negatively
 4228 correlating with each other ($R=-0.25$, $p<0.001$). The result in Figure 5 demonstrates that under
 4229 vernalization, plants that flowers longer tend to have less buds. When non-vernalized this trend was
 4230 not observed.




4231 Figure 5. Scatterplots with a linear model to illustrate the relationship between the number of days a plant
 4232 takes to flower (Days_to_fl) to the number of observed buds on first flowering (Bud_no). The correlation
 4233 suggests that there was no significant relationship under the no-vernialization treatments ($R= 0.046, p=0.57$).
 4234 When vernialized, there was a negative correlation observed ($R=-0.25, p<0.001$).

4235

4236 When samples were introduced to vernialization, it seems that the number of days a plant
 4237 takes to flower is less sporadic. However, when looking at the difference between the two
 4238 treatments and the number of buds observed on first flowering, it is suggested that they are
 4239 significantly different under a two-sample t-test ($P(T<=t) two-tail=0.021$) (Figure 6).

Bud Number vs Treatments




4240 Figure 6. A barchart to illustrate the difference in mean bud number under the two vernialization treatments,
 4241 revealing a significant difference between the two treatments ($P(T<=t) two-tail=0.021$).

4242 **5.3.2 Plant traits vs. Latitude**

4243 To look at whether any of the plant traits under the vernalization treatments were correlated with
4244 the local environmental variables such as latitude, we modelled the relationship between the three
4245 traits (plant height, number of stems, and the number of buds) against latitude and climatic
4246 variables. For this purpose, only the wild individuals from either the no-vernalization or vernalization
4247 treatments were examined. This is because environmental data for the cultivars were not reliable for
4248 our dataset, as was in chapter 2 of this thesis.


4249 A linear model fitted on a scatterplot suggests there was correlation between the height of
4250 the plants (Cm) and latitude ($^{\circ}$ N) (Figure 7). The scatterplot was split by either non-vernalization or
4251 vernalization treatments. They revealed that with both treatments, there were no significant
4252 correlation between plant height and the latitude of locality of each of the sampled plants (for no-
4253 vernalization $R=0.1$, $p=0.31$, for vernalization $R=0.13$, $p=0.11$). This result suggests that in terms of
4254 plant height, neither no-vernalization treatments and/or vernalized treatments is correlated with
4255 the latitude of localities of each respected sample.



4256 Figure 7. Scatterplots with a linear model to illustrate the relationship between the height of individual plants
4257 and their latitude of locality, showing no significant correlation between plant height and latitude (for no-
4258 vernalization $R=0.1$, $p=0.31$, for vernalization $R=0.13$, $p=0.11$).

4259 The number of stems were also observed in relation with the latitude of the individuals in
4260 the two separate experiments. The stem numbers were observed to be less strongly correlated with
4261 the latitude of individuals when *Linum* samples were vernalised.

4262 The result suggests relationship between the number of stems observed and the latitude
4263 locality for individuals under both no-vernalization and vernalization treatments (Figure 8). Under
4264 both treatments the relationship between stem number and the latitude is positive, with
4265 significance under no-vernalization treatment ($R=0.4, p<0.001$). However, the significance of this
4266 relationship was not seen under the vernalization experiment ($R=0.15, p=0.06$). This suggests that
4267 when *Linum* samples are treated with vernalization, the stem number observed in some individuals
4268 is reduced to the point that significant correlation with larger stem number observed in more
4269 Northern individuals were marginally observable under vernalization.




4270 Figure 8. Scatterplots with linear model to illustrate the number of stems observed over the two treatments.
4271 The model reveals that when under no-vernalization treatments, the number of stems observed, is positively
4272 correlated with latitude ($R=0.4, p<0.001$). This significance was not seen in the vernalization treatments
4273 ($R=0.15, p=0.06$).

4274

4275 Another trait which was observed during this study was the number of buds for every
4276 individual. This was also observed based on the treatments of the individuals, to see whether there
4277 was a trend between the number of buds observed and the latitude of different individuals.

4278 Linear modelling on a scatterplot illustrates that the number of buds observed was
4279 significantly correlated with the latitude of locality under the no-vernalization treatments ($R=0.23,$
4280 $p=0.017$) (Figure 9). However, this relationship was contradicted when *Linum* samples were

4281 vernalized. There was a negative correlation between the number of observed buds and the latitude
4282 of individuals. Although this was the case, the negative correlation observed was not significant ($R=-$
4283 0.086 , $p=0.29$). This result suggests that correlation between the bud number and latitude differs in
4284 the two treatments. There is a positive trend, that the more northern an individual is localized, the
4285 more buds were observed when treated with no vernalization. However, when vernalized, this trend
4286 was not observed,



4287 Figure 9. Scatterplots with a linear model to reveal the relationship between the number of buds observed and
4288 latitude between the individuals under no-vernalization and vernalization treatments. The Figure revealed that
4289 there is a positive correlation between the number of bud and latitude on the no-vernalization treatments
4290 ($R=0.23$, $p=0.017$). The trend became negative when vernalized, although not significantly correlated ($R=-$
4291 0.086 , $p=0.29$).


4292

4293

4294 5.3.3 Plant traits vs Climate Variables (pc1)

4295 To relate the latitude to environmental variables, it is viable to relate it to climatic variables
4296 of the local climates to latitude. This is to see whether local climatic variable may affect plant traits
4297 which were measured during this study. This can then be related to whether any vernalization
4298 treatments may have affected traits that are measured in this study.

4299 We saw that plant height is significantly correlated to climate variables, only in the no-
4300 vernalization treatments ($R=-0.17, p=0.006$) (Figure 10). This correlation was reduced when plants
4301 were vernalized, and the correlation became statistically insignificant ($R=-0.15, p=0.062$). This
4302 reveals that vernalization treatments have effects on the height of the overall plants in that it is
4303 reducing correlation with local climate variables seen under no-vernalization treatment. This is
4304 suggesting that plant height is affected by vernalization.




4305 Figure 10. Scatterplots with a linear model fitted to illustrate relationships between plant height (Cm) against
4306 climate variable (pc1) for different vernalization treatments. There were negative correlations between the no-
4307 vernalization and vernalization treatments with the correlation on the no-vernalization treatments showing a
4308 significant correlation ($R=-0.17, p=0.006$).

4309

4310 Another measured trait was the stem number of individuals. This was also measured and
4311 modelled against climate variables (pc1). The model suggests that there was a negative correlation
4312 between stem numbers and climatic variables under the no-vernalization treatments. This
4313 correlation was statistically significant ($R=-0.4, p<0.001$). When compared against the vernalization
4314 treatments, the significance seen under the no-vernalization treatments was observed less, although
4315 still significant. Under vernalization, it seems that stem number of individuals were drastically

4316 reduced for individuals experiencing more negative pc1 climatic variable values. The correlation
4317 between the stem number and the climatic variable are still found to be statistically significant under
4318 $\alpha=0.05$ ($R=-0.18$, $p=0.028$).




4319

4320 Figure 11. Scatterplot with a fitted linear model to suggest the correlation between stem number and the
4321 climatic variables of their local environments (in terms of pc1). The correlation reveals negative correlation
4322 between the no-vernialization and vernialization treatments, with a bigger significance under the no-
4323 vernialization treatments ($R=-0.4$, $p<0.001$).

4324

4325 The number of buds observed was also seen to be negatively correlated with climatic
4326 variables (pc1) under the no-vernialization treatment (Figure 12). This correlation was found to be
4327 statistically significant when modelled ($R=-0.22$, $p=0.028$). In contrast with what was observed with
4328 the stem number though, the correlation between bud numbers and pc1 seems to have changed
4329 directions under the vernialization treatments. Albeit, not statistically significant ($R=0.12$, $p=0.15$).
4330 This difference suggests that vernialization affects the number of buds observed during this study to
4331 a point that it no longer correlates with local environments as observed under the no-vernialization
4332 treatments. This reveals when all plants were vernialized, the individuals with the most numbers of
4333 buds observed under the no-vernialization treatments, reduced their number of buds drastically
4334 through the observed effects of vernialization.



4335 Figure 12. Scatterplots with fitted linear models which suggests negative correlation between observed bud
 4336 numbers and climatic variables (pc1) under the no-vernalization treatments ($R=-0.22$, $p=0.028$). The
 4337 correlation changed directions when vernalization was introduced. However, this correlation was not
 4338 statistically significant ($R=0.12$, $p=0.15$).

4339


4340 5.3.4 Seed Area vs Latitude

4341 As a part of this chapter, we examined seed size variation in wild *Linum* based on different
 4342 latitudes. It is of interest to see whether seed sizes of the different wild *Linum* populations suggests
 4343 significant difference to each other. We investigated whether seed size (seed area) forms a
 4344 correlation to environmental variables. We investigate the question of; “to what extent can
 4345 localization to a local environment in wild *Linum* be of an affect to their seed size?”

4346 To explore this, a preliminary seed size measurements were able to be carried out, using data from
 4347 non-vernalization, vernalization and as an addition, wild individuals. Microscopic images of 5 seeds
 4348 were taken. The images were taken with the same light levels, making sure all 5 seeds were visible
 4349 on the image. After the image were captured, it was saved into a drive and were further processed
 4350 using ImageJ (Schneider C. et al., 2012). Seed areas were able to be highlighted and calculated in
 4351 ImageJ. A macro was written to semi-automate the process. The macro can be found under
 4352 appendix 32. In this case, vernalized individuals were labelled as “Vern18”, and non-vernalized
 4353 individuals were labelled as “nonvern18”. Wild 50 individuals were also able to be counted observed,
 4354 labelled as “Wild”.

4355

4356 When looking at the correlation between seed area and latitude, it is found that in all
 4357 treatments, seed areas are significantly correlated for all treatments (nonvern18 $R=0.51$, $p<0.001$;
 4358 vern18 $R=0.44$, $p<0.001$; Wild $R=0.61$, $p<0.001$) (Figure 13). The wild seeds were also correlating
 4359 stronger than the vernalized seeds. This suggests in all cases seed area is positively correlated to
 4360 latitude and that initial vernalization treatments will result in no significant change to this
 4361 correlation.




4362 Figure 13. Scatterplots with a linear model to show the seed area (Cm) in relation to latitude for our individual
 4363 *Linum* samples. The correlation illustrates that seed area is significantly correlated with latitude of locality for
 4364 all our treatments, the correlation was considerable higher in the wild (nonvern18 $R=0.51$, $p<0.001$; vern18
 4365 $R=0.44$, $p<0.001$; Wild $R=0.61$, $p<0.001$).

4366

4367 5.3.5 Seed Area vs Climate Variable (pc1)

4368 For completion, it is with interest to further see whether seed size in terms of total seed
4369 area observed for different individual sampled during this study. The scatterplots with linear
4370 modelling suggest negative correlations between seed area and climatic variables (Figure 14).
4371 Negative pc1 values represent loading values for colder climates (see chapter 2, under section 2.2.1
4372 Samples and Experiment”). This data suggests that in all treatments, the seed sizes seem to be larger
4373 in area in individuals found in a colder climate and that vernalization have little effects to change this
4374 significance. In the wild individuals, the correlation seems to be stronger.



4375 Figure 14. Scatterplot with a linear model to illustrate the correlation between the seed area (Cm) and the
4376 climatic variables (pc1). The correlation suggests that there is a negative correlation between seed area and
4377 climatic variables in all treatments (nonvern18 $R=-0.47, p=0.0025$; vern18).

4378

4379

4380 **5.4 DISCUSSIONS AND CONCLUSIONS**

4381

4382 Additional results to investigate whether traits maybe associated with the number of days a
4383 plant may take to flower suggests that there is a relationship between the number of days individual
4384 plants takes to flower and plant height and stem numbers. It seems that, under no-vernalization,
4385 when individuals flowered earlier, they tend to be taller at first flowering. In the literature, there are
4386 many cases where number of days a plant takes to flower is positively correlated with plant height
4387 (Singh N. et al., 1995; Gupta S. et al., 2021). In *L. bienne* we expected that this would be the case
4388 when they are under no vernalization stress. We hypothesized that under vernalization this
4389 correlation between number of days to flowering and plant height would be reduced. We suggest
4390 that vernalization acts as an abiotic stress in wild *L. bienne*.

4391

4392 **5.4.1 Number of Days to Flower and Plant Heights**

4393 When vernalized we observed that, correlation between number of days to flower and plant
4394 height was not significant, as the number of days for some individuals to flower were reduced
4395 greatly. This illustrates that under overall vernalization treatments, correlation between the number
4396 of days to flower and plant height at first flowering were not observed. When comparing average
4397 plant heights, there is a significant difference between vernalized and non-vernalized individuals.
4398 This suggest that under vernalization, both number of days (as found in chapter two) and plant
4399 heights were significantly reduced. This illustrates vernalization as an abiotic stress in *L. bienne*
4400 samples found across Western Europe. In literature, there are many examples where vernalization
4401 reduces plant heights (Dole & Wilkins, 1994; Clough et al., 2001; Wu et al., 2016). In other
4402 suggestions, some of the shorter plants during first flowering was able to sense vernalization and
4403 flower much quicker when compared to under no-vernalization treatments. A study using a variety
4404 of grass species found that flowering phenology is associated with the overall plant height under
4405 different environments (Soto-Cerda et al., 2014). This suggests that plants which can flower quicker
4406 may have different overall height to plants that are slower to flower. In our *Linum bienne* samples, it
4407 is illustrated that reduction of the number of days to flower is also reflected by the reduction in
4408 plant height observed, which was significantly different between vernalized and non-vernalized
4409 individuals. We accepted that under vernalization, both number of days to flowering and plant
4410 height are significantly reduced in *L. bienne* found across Western Europe.

4411

4412 5.4.2 Number of Days to Flower and Stem Numbers

4413 There was also a positive correlation between the number of days individuals takes to flower
4414 and stem number under the no-vernalization treatments. It seems that the longer an individual
4415 takes to flower the more stem it can develop before first flowering. This correlation was reduced
4416 when vernalization was implied to the wild individuals, with individuals under vernalization
4417 treatments significantly forming a smaller number of stems. However, the correlation between the
4418 number of days to flower and the number of stems was still significant. This shows that stem
4419 number is highly correlated to the number of days individuals can flower initially even when wild
4420 individuals were exposed to vernalization. When testing against different treatments, individuals
4421 exposed to the vernalization experiments have significantly a smaller number of stems. This suggests
4422 an effect of vernalization in the number of stems wild *Linum* individuals can form. There is
4423 suggestion in the literature that different plant populations may have different number of stems
4424 according to their local environment (Sultan, 2000; Szakiel et al., 2010; ATEs., 2011). Vernalization
4425 suggests stark differences in stem number when compared to non-vernalization treatments, which
4426 suggests that in *L. bienne*, this is also true. In other cases, vernalization affects the morphology of
4427 stem formation in *Phleum pratense* . (Seppänen et al., 2010) This illustrates the potential of
4428 vernalization as an abiotic stress to affect developments of stems in wild *L. bienne* found across
4429 Western Europe.

4430

4431 5.4.3 Number of Days to Flower and Bud Numbers

4432 In terms of the number of buds forming, our results suggest that under the no-vernalization
4433 treatment this measure was not significantly correlated to the number of days an individual is able
4434 to form their first flower. When exposed to vernalization however, the number of buds observed
4435 seems to negatively correlate with the number of first flowering as they were significantly reduced
4436 due to vernalization as an abiotic stress. This suggests that under vernalization requirements, there
4437 may be groupings of plants which can form a higher number of buds with a shortened number of
4438 days to first flowering. In all three cases of the trait measures, including number of buds, variation in
4439 correlation between the number of days to flower and traits measured in this chapter occurred . In
4440 all cases though, vernalization reduced the number of observed bud numbers significantly. This
4441 suggests that vernalization is an abiotic stress that wild population are potentially locally adapting
4442 into.

4443

4444 5.4.4 Effects of Vernalization in Plant Traits

4445 With the above additional results for the plant traits (plant height, stem number, bud
4446 number) considered, it is suggested that the effect of vernalization leads to significantly lower values
4447 in all three trait measures. This suggests that under vernalization, wild *Linum* samples in this study
4448 experienced significant reductions in desired plants traits such as height and plant stem number. In
4449 the model *Arabidopsis thaliana*, it has been observed that traits such as leaf sizes were reduced
4450 under vernalization (Hopkins., 2008). The study also found a latitudinal cline to this response in
4451 vernalization. This suggests that morphology of plant traits may be affected by vernalization and that
4452 environmental variables may also play a part. In the study for this chapter, it is found in wild *Linum*
4453 *bienne* samples across Western Europe that plant height, stem number, and bud number is negative
4454 affected by vernalization. Findings in these traits under vernalization, adds to the suggestion that
4455 vernalization is an abiotic stress and that variation in sensitivity observed can be seen as potential
4456 local adaptation mechanisms between wild *L. bienne* populations found across Western Europe. To
4457 thrive in a colder niche there could be a case that some wild *L. bienne* populations are more locally
4458 adapted to colder climates through requirement of vernalization. In addition, manipulation to
4459 flowering under different seasons/environments can also be studied further as suggested under
4460 other studies observing vernalization in *Calandrinia* (Cave & Johnston, 2010). Reduction in desirable
4461 traits also suggests that the loss of vernalization under *L. usitatissimum*, observed in chapter two of
4462 this thesis could be a benefit in terms of agriculture. Loss of vernalization could mean that negative
4463 effects on traits such as plant height, stem numbers and bud numbers observed under vernalization
4464 in the wild *L. bienne* is selectively avoided in Northern European Winter and Spring *L. usitatissimum*
4465 types. Although this was the case, we assumed that the data is normally distributed, and that values
4466 of trait measurements, and other variables are independent of each other. Traits observed may be
4467 dependent of each other and that the same genes may influence multiple traits observed in this
4468 chapter. These analyses do not consider these possibilities.

4469 The response to plant height in both vernalization treatments, suggests there was no
4470 correlation with the latitude of locality for these plants. However, when looking at local climatic
4471 variables, there was negative correlation between the local climate variable against plant height
4472 when no-vernalization occurred, with individuals from warmer climate (bigger pc1 values) being
4473 taller. This correlation became less significant under vernalization treatment. This suggests that
4474 perhaps climatic variables are having stronger effects to plant height, in our case, individuals under
4475 no-vernalization seems to be taller in colder climate (smaller pc1 values). Vernalization also plays a
4476 part in this, when plants were exposed to vernalization, it seems that this correlation became less
4477 significant. It would be viable to suggest that the process of vernalization induces stress to wild *L.*

4478 *bienne* individuals. To undergo vernalization and flower at the correct time, individual requiring
4479 vernalization may have less favourable traits. Loss of vernalization requirement in the cultivated *L.*
4480 *usitatissimum* in Northern Europe (as discussed in chapter two), could mean that this process is
4481 bypassed in some cultivar types and stress induced by cold for plants to require vernalization is not a
4482 factor. This may form as one of the factors enabling the cultivars to have form better in traits of
4483 agricultural interests for their biproducts. We didn't compare traits between cultivars and wild
4484 species when vernalized as cultivated samples were not included in this chapter for analysis. This will
4485 be of interest for future study.

4486

4487 5.4.5 Effects of Climate in Plant traits for wild *L. bienne*

4488 When looking at other traits against latitude, such as the stem number, there was also
4489 suggestions that the number of stems observed is correlated with the latitude of locality for the
4490 individuals for the no-vernalization treatment. In this case the correlation seems to be positive, with
4491 more Northern plants revealing a greater number of stems. This was also supported when looking at
4492 the climatic variable. A negative correlation was observed which suggests that individual from
4493 warmer climates (higher pc1 values) having a smaller number of stems. When looking at individuals
4494 experiencing vernalization, the stem number able to be formed was not significantly correlated to
4495 either latitude of locality or local climatic variables. This suggests varied morphology in response to
4496 vernalization for our *Linum* collection. The bud number observations suggest there was a positive
4497 correlation between the observed bud number and latitude of localities under no-vernalization
4498 treatments, this suggests more Northern individuals were able to form more buds under no-
4499 vernalization. When vernalized however, the correlation became negative, with more Northern
4500 individuals suggesting for be forming less buds. However, this change of direction in correlation, was
4501 not observed to be significant. When looking at bud number and pc1, it is supporting the traits seen
4502 with latitude, with the two treatments showing contrasting correlation. In this case negative
4503 correlation between bud number and climatic variable (pc1) suggests that individuals found in
4504 warmer climate seems to form less buds. The correlation changed direction under vernalization
4505 treatments, however, as seen with the latitudinal correlation, this change in correlation direction
4506 was not statistically significant.

4507 In all three cases of the measured traits, there is suggestion of differentiation in strategy for
4508 different individuals and when no-vernalization and vernalization treatments were implied. This
4509 suggests there may be genetic interest for looking at these differences in morphology, as was found
4510 under chapter two of this thesis when looking at relative gene expression of flowering time genes

4511 and the number of days an individual takes to flower. In a study with the majorly cultivated durum
4512 wheat, a study has found different phenology and trait strategies in different landraces. This was
4513 also linked to differences in genetic variation responses for the vernalization gene (*VRN1*) and
4514 photoperiod sensitivity gene (*PPD1*) (Royo et al., 2020). There is suggestion in the literatures which
4515 highlight different morphology and phenotype of species of plants due to vernalization and suggests
4516 that this is also linked to genetic variation. It is likely that differences seen in plant height, stem
4517 number, and bud number in wild *Linum* is a result of genetic variation due to vernalization
4518 responses. It will be with great interest in the future to look at specific genes which may play a part
4519 in plant growth, specifically for stems and buds' formation to see whether variation between
4520 individuals still correlate and to see whether variation seen between no-vernalization and
4521 vernalization treatments can be linked genetic factors.

4522

4523 5.4.6 Effects of Climates in Seed Areas for wild *L. bienne*

4524 We also observed seed sizes in terms of seed areas, against latitude of locality for every
4525 individual as well as local climatic variable for them. It is suggested that there was significant
4526 variation between seed area, and these were all correlated with latitude and local climatic variables
4527 (pc1). The correlation was still retained under no-vernalization and vernalization treatments. Positive
4528 correlation with latitude suggests that in all treatment cases, the more Northern individual's seeds
4529 were bigger than those of their Southern counterparts. This also translates to the correlation seen in
4530 climatic variables, with negative correlations, suggesting that individuals from colder regions having
4531 the bigger seeds. Bigger seeds may imply better fitness in plant reproduction, smaller seeds are
4532 however, easier to produce, suggesting this trade-off between seed size and the number of seeds a
4533 plant can make for a given amount of energy (MOLES et al., 2004). In a study with wild barley
4534 (*Hordeum vulgare ssp.*), there is a suggestion that variation in seed sizes contributes to plant growth
4535 and reproduction, with larger seeds often found to have better fitness (Giles., 1990). A study on
4536 different *Glycine* species found that seed mass tends to be larger at Northern latitudes. They also
4537 found that the major contributing climatic variable to this are temperature and daylight availability
4538 (Murray et al., 2004). These agree with our finding that *Linum* seeds also found to be larger in
4539 Northern populations. When looking at the wild seeds in our observations, the correlation between
4540 seed size and latitude and climate variables were found to be stronger. In sand rice (*Agriophyllum*
4541 *squarrosum*), there was also geographic variation of seed size due in wild populations. In *Linum*, the
4542 seed sizes are strongly correlated to local variation in climatic variables. It would be of interest to
4543 genetically evaluate this to see whether the strong correlation found in this thesis had genetic
4544 implication for genetic basis which may reveal evidence of genetic variance. In the past, studies

4545 looking at genetic variance for variation in seed sizes in different plant species have found no
4546 correlation between genetic variation and variation in seed sizes, suggesting that seed size could not
4547 have evolved due to natural selection (Wolfe., 1995) (Schwaegerle & Levin., 1990). It would be of
4548 good interest to genetically evaluate the seed size variation under different environmental variables
4549 as illustrated in this thesis.

4550

4551

4552 GENERAL DISCUSSION

4553

4554 Experiments in this thesis were set out to investigate adaptive strategies in *Linum*, using two
4555 species (*L. usitatissimum* and *L. bienne*). We were interested in variation in traits and genetic
4556 materials of both wild and cultivated species to observe signals of local adaptation. We were
4557 particularly interested in flowering mechanisms. We found that as a temperate plant, requirement
4558 to vernalize varies in wild *L. bienne* found across Western Europe. However, we saw no significant
4559 changes to the flowering initiation when we vernalized the cultivars *L. usitatissimum*. Examples in
4560 the literature suggests that there is variation in vernalization sensitivity between different *L.*
4561 *usitatissimum* varieties grown under different seasons (Darapuneni M. et al., 2014) (see section 2.5
4562 of chapter two). We suggest that there is no sensitivity to vernalization in winter and spring types *L.*
4563 *usitatissimum* found in the Northern part of Western Europe. Comparison against lines from
4564 Southeastern European regions such as Turkey, where *L. usitatissimum* is cultivated in different
4565 seasons, will be of interest for observation of vernalization sensitivity variation in *L. usitatissimum*
4566 types found across two different growing climates across Europe.

4567 In *L. bienne*, we observed a positive trend on the number of days individuals took to flower
4568 and their latitude of origin. The finding suggests that wild individuals further North are more
4569 sensitive towards the requirement to vernalize, with substantial reduction in the number of days to
4570 flower when vernalized. This was further reinforced by genetic expression studies of 3 flowering
4571 time genes (*FT*, *CO*, and *GI*). We found no significance between the expression response of each of
4572 the tested genes when looking at vernalized and non-vernalized treatments. However, we observed
4573 the relative expression of *FT* to be positively correlated with latitude of origin in wild *L. bienne*
4574 individuals. With this observation, we accepted the hypothesis that variation in flowering initiation
4575 occurs in wild *Linum* across Western Europe, and this is reflected in the relative gene expression of
4576 the floral integrator *FT* against latitude and climate of origin of samples. This trend was not observed
4577 in any of the other flowering time genes tested.

4578 We also acknowledge that the sample size for the relative gene expression study is relatively
4579 small. When we removed the Southern-most individual, we no longer observed a positive
4580 correlation between *FT* expression and latitude. Interestingly, flowering initiation were still positively
4581 correlated with latitude, suggesting that perhaps expression of other floral integrators and/or
4582 repressors could be affecting the flowering initiation. Furthermore, we also saw a relationship
4583 between the number of days individuals took to flower and the local climate. We saw that
4584 expression of *FT* were correlated with climatic variables. This introduces the possibility that, wild

4585 population are locally adapting to the environmental variables such as precipitation, solar radiation,
4586 and temperatures. Wang J. et al., 2018 illustrates that for perennial plants, the locus *FT* forms part of
4587 the genomic region responsible for local adaptation (Wang J. et al., 2018). We make a case that
4588 evidence of local adaptation can be observed in wild *L. bienne* across Western Europe, based on the
4589 relationship seen between *FT* expression and latitude and climatic variables. Although this is the
4590 case, there is a possibility that variation in requirement to vernalize can arise from mechanisms such
4591 as genetic drift. We didn't have precise location data for our cultivars. It would be of further interest
4592 to compare the relationship of *FT* expression with latitude/climatic variables for the different *Linum*
4593 species. We expect that, due to domestication and the loss of requirement to vernalize observed in
4594 this study, our *L. usitatissimum* variants would not form a relationship with latitude/climatic
4595 variables. This would suggest a contrast between the wild *L. bienne* and the cultivated *L.*
4596 *usitatissimum*. Whilst the requirement to vernalize may be artificially selected in the cultivars, the
4597 wild suggests that flowering initiation and requirement to vernalize may be driven by local
4598 adaptation under various climate. Although this was our finding, we suggest taking the results with
4599 precautions. Along with a relatively small sample size, relative gene expressions were comparative
4600 only against one HKG. We propose a future study with bigger sample sizes from a wide range of
4601 latitude, specifically more Southern populations and their floral gene relative expression compared
4602 against multiple HKG's. This is to robustly reflect the correlation with *FT* expression and latitude. In
4603 addition to this, it would be ideal to observe expression of genes directly in the vernalization
4604 pathway such as *FRI*, *VRN1* and *FLC*. We expect that the relative expression of these genes is related
4605 to the latitude and local climate in which the wild population are found since they affect *FT*
4606 expression.

4607 We saw in chapter two that the number of days wild *L. bienne* takes to flower is related to
4608 the latitude. This implies that temperature, as a variable, may control for the number of days to
4609 flower. To investigate whether temperature had further effects on flowering mechanisms, we
4610 continued to observe responses in floral organs. We explored pollen viability in response to
4611 reduction (-5°C from initial temperature) and addition (+5°C from initial temperature). We see a
4612 response in reduction of pollen count for both reduction and addition of temperature. When looking
4613 at pollen tube count, this varied within all our temperature treated samples. However, when scaling
4614 this with the pollen (proportion of pollen tube forming to pollen count) only the addition in
4615 temperature shows significant reduction in the proportion of pollen tube forming. This suggests that
4616 *Linum* pollen tubes are potentially sensitive to increases in temperature. To reinforce this, we found
4617 the proportion of pollen tube able to reach the ovary to be significantly reduced under the +5°C
4618 temperature treatment. This was observed in both species. This may suggest that pollen ability to

4619 germinate is limited to temperature or the female floral organs are not able to sustain pollen tubes
4620 under heat stress. This suggests the interest in looking at ability of female counterparts to sustain
4621 pollen tube growth under different temperature for future studies in this area.

4622 We concluded this chapter by observing whether latitude/climate variables are correlated
4623 with the amount of pollen and pollen tubes observed in *L. bienne* when treated to the different
4624 temperature treatments. We found no correlation between the number of pollen and
4625 latitude/climate. However, the number of pollen tube and its proportion to the number of pollen
4626 observed a correlation with latitude and climate in the 20°C (-5°C) treatment. Furthermore,
4627 correlation with latitude is found to be positive, illustrating that more Northern individual can form
4628 more pollen tubes. This may suggest potential adaptation to the colder temperature for the more
4629 Northern individual in relation to its ability to form pollen tubes and thus increasing its chances of
4630 germinating. We suggest that this also formed because of local adaptation found in the Northern
4631 population of Western European *L. bienne*. From this chapter we reinforced the suggestion that local
4632 adaptation between the North and South wild *L. bienne* in Western Europe occurs, as was illustrated
4633 in chapter two. For future research, it would be of interest to look at this from the female
4634 counterpart's perspective. We suspect that ovary would be sensitive to heat-stress and stigma will
4635 have varying ability to sustain pollen tube growth. In addition to this, it would also be beneficial to
4636 conduct a GWAS study to identify the genes which may play a role in cold resistance in the wild
4637 which maybe linked to the higher amount of pollen tube able to be observed under wild individuals.

4638 In chapter four we analysed 100 Western European *Linum* individuals and their population
4639 genetics. One of the main results from this chapter was the observation of Northern and Southern
4640 groups in the wild *L. bienne*. There seems to be structuring between the Western European wild *L.*
4641 *biene* population as was inferred in section 4.4.1 "Results: Genetic Structure". We observed initial
4642 signal of potential variation in breeding strategies of *L. bienne* found across Western Europe in our
4643 population diversity analysis. There is potential that the Southern group of wild *L. bienne* is more
4644 outcrossing in terms of their breeding strategy and thus causing this genetic structuring we see
4645 when illustrating genetic distance tree. In the literatures, it is found that various mode of seed
4646 dispersal often affects a plant's breeding strategy. They found that animal-dispersed plant species
4647 exhibited higher levels of genetic diversity and lack of inbreeding (Nazareno et al., 2021). This further
4648 suggests that there is potentially more variation within Southern *L. bienne* individuals in Western
4649 Europe. All measures of population diversity suggests that heterozygosity is higher in the Southern
4650 cline, with relatively little amount of heterozygosity observed in the Northern group. Measures of
4651 population divergence such as Tajima's D also suggest that Southern population is more recently
4652 expanded and are expanding more than the Northern group. These findings suggests that there is

4653 potential variation in breeding strategies and dispersal of North and South groups of *L. bienne* found
4654 in Western Europe. The implication of variation in breeding strategies could result in the variation
4655 we saw in chapter two whereby Northern individuals tends to be more sensitive to vernalization as
4656 was expressed by their variation in relative ge expression under different latitudes. Furthermore, we
4657 saw that ability of pollen tube forming and germination processes is correlated with individual's
4658 latitude when treated to "cold" treatments (see section 3.4.8 "Results: Pollen and Latitude of
4659 Origin"). Genetic structuring and diversity analysis in chapter four supports the findings that
4660 Northern and Southern clines of *L. bienne* found in Western Europe is potentially locally adapted to
4661 their environment.

4662 In addition to the observation between Northern and Southern clusters, we observed a
4663 Mediterranean population which is significantly diverged from other populations. This could be a
4664 result of increased rate of outcrossing in nature we saw in the Southern clines when looking at
4665 population diversity measures. It would be of interest to include population from the Eastern
4666 European region and see if this cline between North and South population could also be observed
4667 between Western and Eastern European wild *L. bienne*, with Eastern population potentially more
4668 outcrossing in breeding strategy. In future studies investigating Western and Eastern *L. bienne*
4669 population, wild *L. bienne* seeds needs to be collected from Eastern European regions such as Turkey
4670 and the Eastern Mediterranean. There is no availability of SO seeds from these wild population to
4671 date.

4672 We also acknowledge that we may have lost some valuable data by mapping to the *L.*
4673 *usitatissimum* whole genome. This was inferred from the higher number of SNPs gained from *de*
4674 *novo* mapping (see section 4.5.4 "*De_novo* mapping"). In addition, it is worth exploring other NGS
4675 techniques to imply further structuring based on a larger coverage of the genome. This is because of
4676 disadvantages to using ddRADSeq to sequence our short reads. Our ddRADSeq only studies 250-
4677 500bp of sequences per individual cut by the digestion process of ddRADSeq and require high quality
4678 DNA which may impact the number of individuals able to be analysed and outputted from the
4679 ddRADSeq procedures. We also looked at two species with potential divergence in our wild
4680 populations. Other recent short-reads NGS techniques that can be looked at for observation of
4681 multiple species is Sequence capture. Sequence capture holds more promise for obtaining data sets
4682 that are comparable across species and for calibrating parameter estimates for demographic or
4683 phylogenetic studies (Harvey et al., 2016). Sequence capture also doesn't require a high quality of
4684 DNA as opposed to RAD Sequencing processes. There is potential that individuals with a relatively
4685 lower quality of DNA could be analysed further with Sequence capture (Harvey et al., 2016). In
4686 addition, studies using Sequence capture can capture larger sizes of sequences (300-1200bp) which

4687 covers 972 genes (Sanderson et al., 2020). Future studies can look at the potential of using Sequence
4688 capture for population genetic analysis in *L. bienne* across Western Europe.

4689 When looking at traits in chapter five, we found evidence that plant height, stem number,
4690 and bud number is significantly reduced by the introduction of vernalization. This suggest that
4691 vernalization is an abiotic stress variable in wild *L. bienne* found across Western Europe. When under
4692 no-vernalization, we saw positive correlation between stem and buds formed at first flowering and
4693 latitude. This illustrates that under no vernalization stress, more Northern individuals can form more
4694 stems and buds. However, under vernalization there was no observed correlation between traits and
4695 latitude/climate. This suggest that, in the wild, vernalization acts as abiotic stress and as was found
4696 in chapter two, vernalization responses varies across the wild *L. bienne*. In terms of local adaptation
4697 in wild *L. bienne*, the ability Northern individuals to form more stems and buds, under no-
4698 vernalization can suggest that seasonal queues, especially regarding vernalization, may occur in the
4699 Northern cline. This ability to form more stems and buds under no-vernalization was seen less in the
4700 Southern cline, suggesting the potential for local adaptation regarding vernalization requirement in
4701 the Northern cline, which are expected to be more sensitive to vernalization due to colder climates.
4702 This was also supported by higher expression of the floral integrator *FT* as was suggested in chapter
4703 two. In future studies, it is of interest to look at expression of genes directly affecting number of
4704 stems and buds to reinforce the variation seen within this thesis with genetic expression studies.

4705 We further support the potential local adaptation between Northern and Southern clines of
4706 *L. bienne* population using seed sizes. Under three growth conditions, seeds areas are positively
4707 correlated with latitude and negatively correlated with climatic variables, mostly attributed to colder
4708 temperatures. This suggest that Northern individuals seems to have a bigger seed area to the
4709 Southern individuals, which can infer that trade-off between seed size and potential to produce
4710 more seeds between the two clines. In the wild S0 seeds, this correlation between seed area and
4711 latitude is more strongly observed, which may suggest that seed area is a variable for measuring
4712 seed size variation in wild *L. bienne* for future studies regarding local adaptation and breeding
4713 strategies of wild *L. bienne*.

4714 With the above five chapters concluded, we illustrated potential signal of local adaptation in
4715 wild *L. bienne* to Northern climates, requiring vernalization. We observed no sensitivity of *L.*
4716 *usitatissimum* in our sample set to vernalization. We expect there are variation between various *L.*
4717 *usitatissimum* variety regarding vernalization sensitivity, especially for non-winter types. As was seen
4718 in *L. bienne*, this vernalization requirement can suggest reduction in traits that are beneficial to
4719 agriculture, and so loss of vernalization in winter types maybe beneficial for these traits of interest.
4720 We imply that the loss of vernalization is agriculturally beneficial, however, this may limit growth

4721 time and cycle of plants under different seasons, thus resulting in seasonal types of *L. usitatissimum*
4722 varieties. Wild Northern population could be studied to understand mechanisms of vernalization in
4723 *Linum*, this may add to breeding purposes, providing *Linum* cultivars that are perhaps more resistant
4724 to colder climates and can be grown in different seasons.

4725 In terms of wild *L. bienne* population found across Western Europe, population structure
4726 suggests that gene flow between the Northern and Southern group in our samples are limited due to
4727 the grouping suggested by the Ward's D tree based on Nei's genetic. Genetically, we saw little
4728 evidence of geneflow between the wild population which is a signal of locally adapting populations
4729 (Boshier et al., 2015). Phenotypically, we saw variation for flowering traits for Northern and Southern
4730 group of population within our sample set as a signal of this local adaptation.

4731 Although this was the case, in our PCoA we saw a few outliers to these groupings suggesting
4732 a cline between the Southern and Northern populations. This suggests a limited geneflow between
4733 the two groups, albeit we think that this geneflow is limited to one or two individuals in our sample
4734 set. We suggest future studies to look at more wild population originating in Northern Spain, as this
4735 is where we saw potential outliers which may suggest gene flow between the Southern and
4736 Northern groups of wild *Linum* observed in this study. Further divergence can be seen in our
4737 Mediterranean population. Population divergence analysis suggests that the population E-12 in our
4738 sample set is distinct to both Northern and Southern group with a higher number of private alleles
4739 and heterozygosities. This suggests that perhaps this population is part of a diverging group of wild
4740 *L. bienne*. Higher allelic richness can also suggest a different approach to breeding strategies than
4741 seen in the rest of the Western European wild population which may allude to differentiation in
4742 mechanisms such as seed dispersals.

4743 Less heterozygous alleles in the Northern group of population also infer less allelic richness
4744 and potentially more inbreeding, suggesting that these Northern group have a more conserved
4745 genetic makeup, potentially counting for local adaptation to a colder climate as we initially observed
4746 under flowering mechanisms such as vernalization. However, it is interesting to find that there are
4747 intermediate individuals forming a cluster between the Southern and Northern population in our
4748 PCoA analysis. This suggests that perhaps several populations based in Northern Spain acts as a
4749 geneflow mediator, connecting the two distinct Western European wild flax population. This could
4750 suggest that, although the Northern group of wild *L. bienne* population are more conserved and
4751 locally adapted to Northern climates, they have diverged from a Southern population. This suggests
4752 implications for *Linum* to diverge into colder climates and are locally adapting to these climates. This
4753 implies the importance of colder climates for natural selection of wild plants and their
4754 diversification. This could impact availability of a wider genepool for study on mechanisms to

4755 counter cold climates in plants, vernalization to name one. These wider genepool are useful for
4756 future breeding purposes agriculturally. In a warming climate, these wider genepool may be
4757 threatened as they are found to be more conservative and inbred than their Southern relatives.

4758 Other studies suggests that the Mediterranean is a “hotspot” for recent plant diversification
4759 (Buirra et al., 2020). We think that wild *L. bienne* population originating further in the Mediterranean
4760 as opposed to Western Europe may be more diverse and distinct than previously thought for *L.*
4761 *bienne* across Western European region. Diverging Mediterranean populations may also adapt at a
4762 different rate to their environment and are more outcrossing (as was observed in population E-12).
4763 We think that Mediterranean wild *L. bienne* population may provide a wider genepool for genetic
4764 study of *Linum* and have the potential to provide diversification to breeding strategies in *Linum*. To
4765 develop this further, we suggest collection of further wild *L. bienne* from across the Mediterranean
4766 region to confirm this genetic diversity from the Western European relatives, seen in this study.

4767 We take these implications within this thesis with precautions as the number of sample sets
4768 are relatively small and minimum representative for population genetics purposes was only seen to
4769 three individuals. For future relative gene expression studies, a confirmation using multiple HKG’s
4770 needs to be conducted as future research towards the results found within this thesis. However,
4771 signals of local adaptations can still be observed even with these limitations in the wild *L. bienne*
4772 found across Western Europe. This would imply potential traits of interest in relation to local
4773 adaptation which maybe playing a role towards resistance to abiotic factors such as vernalization.
4774 There are also implications for further studies observing cold-resistance genes and pollen viability in
4775 wild *L. bienne* to observe mechanisms of cold resistance in future *Linum* GWAS studies that maybe
4776 beneficial for implication to *Linum* agriculture.

4777

4778

4779 APPENDICES

4780

4781 **Appendix 1: A list of *L. bienne* and *L. usitatissimum* samples collected for this thesis**4782 ***L. bienne***

| Pop | Lat | Long | collected | collector | details |
|-----|----------|----------|------------|--------------------|---|
| 1 | 36.80044 | -5.39258 | 04/06/2016 | RPB | Llanos del Rabel trail |
| 2 | 36.08092 | -5.62553 | 05/06/2016 | RPB | Virgen de la Luz Sanctuary Facinas |
| 3 | 36.03633 | -5.55589 | 05/06/2016 | RPB | Guadalmesi |
| 4 | 36.15083 | -5.70494 | 05/06/2016 | RPB | El Nene, facinas |
| 5 | 37.25853 | -6.09722 | 07/06/2016 | RPB | Puebla del Río-Aznalcazar, Sevilla |
| 8 | 37.70194 | -5.83206 | 08/06/2016 | RPB | Road 432, Km 12 Road to El Pedroso |
| 6 | 37.93551 | -5.71117 | 08/06/2016 | RPB | Constantina-Cazalla de la Sierra, Sevilla (trail) |
| 7 | 38.25336 | -4.31739 | 08/06/2016 | RPB | Cardeña-Villa del Río, Cortijo Tejoneras, Córdoba |
| 9 | 37.8085 | -6.42881 | 09/06/2016 | RPB | N-433 before exit to Zufre-La Granada de Riotinto, Huelva |
| 10 | 37.88211 | -6.61781 | 09/06/2016 | RPB | Linares de la Sierra, Huelva |
| 11 | 38.33105 | -3.58086 | 14/06/2016 | RPB | La Aliseda, Finca La Inmediata (Km 3), Jaen |
| 12 | 42.31008 | 3.151889 | 17/06/2016 | RPB | Palau-Savereda |
| 13 | 43.02902 | -3.24035 | 19/06/2016 | RPB | Quincoces de Yuso-Relloso, Burgos |
| 14 | 42.79404 | -3.42474 | 20/06/2016 | RPB | Tartales de Cilla |
| 15 | 43.46278 | -3.65331 | 20/06/2016 | RPB | Cantabria-Carriazo.Galizano |
| 16 | 42.17089 | -8.68382 | 21/06/2016 | RPB | Universidad de Vigo |
| 17 | 50.69147 | -1.0954 | 24/09/2016 | RPB | Bembridge, 1st stop, Isle of Wight |
| 18 | 50.68183 | -1.07492 | 24/09/2016 | RPB | Bembridge, 2nd stop, Isle of Wight |
| 19 | 42.17089 | -8.68382 | | RPB | Universidad de Vigo |
| Tor | 36.73956 | -3.92635 | 11/07/2013 | ACB | Torroxo Costa, Malaga |
| Lla | 43.40738 | -4.68753 | 21/07/2014 | ACB | Llanes, Asturias |
| Mat | 48.35697 | 4.458631 | 29/06/2016 | ACB | Mathaux, Aube |
| Vil | 45.09393 | -1.05034 | 02/07/2016 | ACB | Villeneuve, Charente Maritime |
| Bro | 47.16704 | -0.20705 | 04/07/2016 | ACB | Brossay, Maine et Loire |
| Roc | 47.38702 | -0.52574 | 04/07/2016 | ACB | Domaine de Rochambeau, Maine et Loire |
| Saf | 47.4961 | -1.59254 | 05/07/2016 | ACB | Saffre, Loire Atlantique |
| Tal | 47.6997 | -3.45465 | 06/07/2016 | ACB | Pointe du Talude, Morbihan |
| Fro | 53.30782 | -2.71877 | 02/09/2016 | ACB | Frodsham, Cheshire |
| Tym | 53.30307 | -3.55328 | 02/09/2016 | ACB | Tyr_Mawr_Holiday_Park, Denbighshire |
| Sut | 53.35291 | -0.95927 | 09/09/2016 | ACB | Sutton_Cum_Lound, Nottinghamshire |
| Man | 53.13731 | -1.14367 | 10/09/2016 | ACB | Mansfield, Nottinghamshire |
| Dor | 50.6 | -2.01 | | Emorsgate Seeds | Dorset |

4783

4784

4785

4786 *L. usitatissimum*

| Shortname | Name | Type | Subsp | Country | Source | Code | centreleft | centreright |
|-----------|----------------|----------------|---------------|-------------|----------|--------|------------|-------------|
| Pri | Primus | oilseed | mediterraneum | Italy | IPK | 247707 | 41.87194 | 12.56738 |
| Rab | Raba0189 | oilseed | mediterraneum | Morocco | IPK | 247713 | 31.7917 | -7.09262 |
| Gis | Gisa | oilseed | caesium | Italy | IPK | 260080 | 41.87194 | 12.56738 |
| Tin | TineTammesLila | oilseed | caesium | Netherlands | IPK | 236553 | 52.13263 | 5.291266 |
| Mon | Monarch | fibre | elongatum | UK | IPK | 255846 | 55.37805 | -3.43597 |
| Ome | Omegalin | oilseed spring | | France | TDL | | 46.22764 | 2.213749 |
| Lir | LiralCrown | oilseed | caesium | UK | IPK | 231715 | 55.37805 | -3.43597 |
| Ble | Blenda04C | fibre | elongatum | Netherlands | IPK | 225632 | 52.13263 | 5.291266 |
| Ara | Aramis | fibre spring | | France | TDL | | 46.22764 | 2.213749 |
| Ari | Ariane | fibre | elongatum | France | IPK | 254727 | 46.22764 | 2.213749 |
| Bol | Bolchoi | fibre spring | | France | TDL | | 46.22764 | 2.213749 |
| Boo | Boothby Grafoe | oilseed | | UK | own | | 55.37805 | -3.43597 |
| Ede | Eden | fibre spring | | France | TDL | | 46.22764 | 2.213749 |
| Mar | Marmalade | oilseed | | Canada | Flaxland | | 56.13037 | -106.347 |
| Olg | Olga | fibre winter | | France | TDL | | 46.22764 | 2.213749 |
| Suz | Suzanne | fibre | | Netherlands | Flaxland | | 52.13263 | 5.291266 |
| Vol | Volga | oilseed winter | | France | TDL | | 46.22764 | 2.213749 |

4787

4788

4789 **Appendix 2: Sample lists for vernalization study**

| Individual | Population | Species |
|------------|------------|---------|
| 2_29 | 2 | Bienne |
| 2_3 | 2 | Bienne |
| 3_13 | 3 | Bienne |
| 3_15 | 3 | Bienne |
| 4_23 | 4 | Bienne |
| 4_27 | 4 | Bienne |
| 5_4 | 5 | Bienne |
| 6_1 | 6 | Bienne |
| 6_26 | 6 | Bienne |
| 6_29 | 6 | Bienne |
| 7_17 | 7 | Bienne |
| 9_27 | 9 | Bienne |
| 9_23 | 9 | Bienne |
| 9_24 | 9 | Bienne |

| | | |
|----------|------|--------|
| 10_26 | 10 | Bienne |
| 10_30 | 10 | Bienne |
| 11_23 | 11 | Bienne |
| 13_12 | 13 | Bienne |
| 14_21 | 14 | Bienne |
| 14_6 | 14 | Bienne |
| 15_27 | 15 | Bienne |
| 15_28 | 15 | Bienne |
| 15_29 | 15 | Bienne |
| 15_32 | 15 | Bienne |
| 19_26 | 19 | Bienne |
| 19_30 | 19 | Bienne |
| Dor_B | Dor | Bienne |
| low1_10 | low1 | Bienne |
| IOW1_11 | low1 | Bienne |
| low1_17 | low1 | Bienne |
| low2_2 | low2 | Bienne |
| low2_25 | low2 | Bienne |
| low2_26 | low2 | Bienne |
| low2_30 | low2 | Bienne |
| Lla_17 | Lla | Bienne |
| Lla_22 | Lla | Bienne |
| Lla_25 | Lla | Bienne |
| Lla_20 | Lla | Bienne |
| Lla_43_A | Lla | Bienne |
| Lla_B | Lla | Bienne |
| Lla_33 | Lla | Bienne |
| Man_5 | Man | Bienne |
| Man_6 | Man | Bienne |
| Man_4 | Man | Bienne |
| Man_8 | Man | Bienne |
| Mat_17 | Mat | Bienne |
| Mat_2 | Mat | Bienne |
| Mat_23 | Mat | Bienne |
| Mat_24 | Mat | Bienne |
| Mat_14 | Mat | Bienne |
| Roc_12 | Roc | Bienne |
| Saf_10 | Saf | Bienne |
| Saf_19 | Saf | Bienne |
| Saf_9 | Saf | Bienne |
| Saf_16 | Saf | Bienne |
| Tal_10 | Tal | Bienne |
| Tal_25 | Tal | Bienne |
| Tal_28 | Tal | Bienne |
| Tal_4 | Tal | Bienne |

| | | |
|--------|-----|---------------|
| Tor_4 | Tor | Bienne |
| Tym_26 | Tym | Bienne |
| Tym_5 | Tym | Bienne |
| Vil_21 | Vil | Bienne |
| Vil_25 | Vil | Bienne |
| Vil_36 | Vil | Bienne |
| Vil_27 | Vil | Bienne |
| Ara | Ara | Usitatissimum |
| Bey | Bey | Usitatissimum |
| Ble | Ble | Usitatissimum |
| Ede | Ede | Usitatissimum |
| Gis | Gis | Usitatissimum |
| Lir | Lir | Usitatissimum |
| Mon | Mon | Usitatissimum |
| Olg | Olg | Usitatissimum |
| Ome | Ome | Usitatissimum |
| Rab | Rab | Usitatissimum |
| Tin | Tin | Usitatissimum |
| Suz | Suz | Usitatissimum |

4790

4791

4792

4793

4794

4795 **Appendix 3: Principal component for each population and its climatic variable loading values**

4796

| Principal Components | Population | PC1 | PC2 | PC3 |
|----------------------|------------|--------|--------|--------|
| | 3 | 5.367 | 3.009 | 0.541 |
| | 2 | 6.383 | 3.188 | -0.130 |
| | 4 | 6.533 | 2.432 | -0.611 |
| | 1 | 1.846 | -1.332 | -0.190 |
| | 5 | 6.354 | -1.009 | 0.370 |
| | 8 | 5.655 | -1.622 | 0.537 |
| | 9 | 3.087 | -2.111 | 0.259 |
| | 10 | 2.429 | -2.026 | 0.143 |
| | 6 | 2.619 | -2.285 | 0.324 |
| | 7 | 1.175 | -3.868 | 0.203 |
| | 11 | 1.613 | -4.139 | 0.844 |
| | 19 | -1.551 | 2.828 | -5.173 |
| | 12 | 0.120 | 1.792 | 3.345 |
| | 14 | -3.398 | -3.411 | -1.836 |
| | 13 | -3.830 | -2.266 | -2.315 |
| | 15 | -0.135 | 1.706 | -2.983 |
| | IOW2 | -5.447 | 2.161 | 1.706 |
| | IOW1 | -5.463 | 2.189 | 1.743 |
| | Tor | 6.561 | 1.674 | 2.282 |

Loadings

| | | | |
|----------|--------|--------|--------|
| CGa1 | 5.845 | 2.724 | 2.514 |
| Lla | -0.093 | 0.232 | -1.601 |
| Vil | -1.449 | 0.359 | -0.876 |
| Bro | -3.291 | -0.895 | 0.205 |
| Roc | -3.057 | -0.802 | 0.147 |
| Saf | -2.820 | -0.663 | -0.468 |
| Tal | -3.097 | 3.614 | 1.631 |
| Mat | -4.574 | -2.090 | -0.497 |
| BH | -5.397 | 3.754 | 2.368 |
| Lil | -5.769 | 1.930 | 0.624 |
| CR | -5.348 | 0.097 | -0.455 |
| Tym | -6.617 | 1.478 | 1.481 |
| Sut | -6.676 | -0.878 | 1.388 |
| Man | -7.323 | -0.454 | 0.932 |
| LJLb1 | 0.054 | -4.368 | 1.209 |
| Dor | -6.279 | 2.961 | 2.316 |
| prec_DJF | 0.174 | 0.529 | -0.681 |
| prec_JJA | -0.869 | 0.209 | -0.273 |
| prec_MAM | -0.094 | 0.445 | -0.849 |
| prec_SON | -0.233 | 0.615 | -0.694 |
| srad_DJF | 0.930 | -0.227 | -0.008 |
| srad_JJA | 0.894 | -0.269 | 0.051 |
| srad_MAM | 0.930 | -0.224 | 0.020 |
| srad_SON | 0.940 | -0.200 | -0.032 |
| tavg_DJF | 0.922 | 0.352 | -0.029 |
| tavg_JJA | 0.923 | -0.298 | 0.158 |
| tavg_MAM | 0.991 | 0.035 | 0.024 |
| tavg_SON | 0.992 | 0.088 | 0.046 |
| tmax_DJF | 0.964 | 0.125 | -0.042 |
| tmax_JJA | 0.807 | -0.523 | 0.128 |
| tmax_MAM | 0.947 | -0.253 | 0.003 |
| tmax_SON | 0.974 | -0.167 | 0.025 |
| tmin_DJF | 0.783 | 0.591 | -0.012 |
| tmin_JJA | 0.959 | 0.111 | 0.183 |
| tmin_MAM | 0.913 | 0.387 | 0.042 |
| tmin_SON | 0.911 | 0.384 | 0.068 |
| vapr_DJF | 0.827 | 0.501 | -0.030 |
| vapr_JJA | 0.638 | 0.589 | 0.049 |
| vapr_MAM | 0.810 | 0.523 | -0.018 |
| vapr_SON | 0.805 | 0.532 | -0.002 |
| wind_DJF | -0.545 | 0.684 | 0.434 |
| wind_JJA | -0.377 | 0.625 | 0.544 |
| wind_MAM | -0.453 | 0.690 | 0.489 |
| wind_SON | -0.517 | 0.653 | 0.504 |

4797

4798

4799 **Appendix 4: List of Primers and its sequences for flowering time experiments in Linum**

4800
4801
4802
4803
4804
4805
4806
4807

| Primers | Sequence | Conc. |
|-------------|------------------------------|----------------------|
| LuGAPDH_for | AGGTTCTCCCGCTCTCAAT | 25nm ⁴⁸⁰⁸ |
| LuGAPDH_rev | CCTCCTTGATAGCAGCCTTG | 25nm ⁴⁸⁰⁹ |
| LuUBI2_for | CCAAGATCCAGGACAAGGAA | 25nm |
| LuUBI2_rev | GAACCAGGTGGAGAGTCGAT | 25nm ⁴⁸¹⁰ |
| LuCO1-pr | AGGCTCCGGTCATGATGAATGACCACTG | 25nm |
| LuCO2-pr | TGCTCCCGTCATGAATATGAATGACCAC | 25nm ⁴⁸¹¹ |
| LuCO-rev | AGATACGCTGTGGCTCAAG | 25nm ⁴⁸¹² |
| LuGI1.1-pr | CTCTACTCTCCGCATCCTGTCA | 25nm |
| LuGI1.2-pr | TGATGGAGTTGAAGTACAGCATGAACC | 25nm ⁴⁸¹³ |
| LuGI2-pr | CACTACGCCAAGTTGATTGCATCG | 25nm ⁴⁸¹⁴ |
| LuGI-rev | GTATGTACAAGTTCATGACA | 25nm |
| LuFT1-pr | AACTCTACAAGTTAGGTTCCCCCGTTG | 25nm |
| LuFT2-pr | AACTCTACAAGTTAGGTTCCGCCTGTTG | 25nm |
| LuFT-rev | GTCTCTCGTTGGCAGTAAA | 25nm |

Appendix 5:
Primer efficiency
comparisons:
(We used #3)

#1

| Primer Efficiency | | | | | | |
|-------------------|---------|----------------|---------|----------------|-----------|----------|
| Sample | Primer | R ² | Slope | Efficiency (%) | Converted | Value |
| | LuGAPDH | 0.97378 | 3.86418 | 81.4625 | 1.81463 | 1.814625 |
| | LuUBI2 | 0.93874 | 2.83438 | 125.326 | 2.25326 | 2.253256 |
| | LuGI1.1 | 0.99999 | 0.60742 | 4329.12 | 44.2912 | 44.29123 |
| | LuCO1 | 0.97491 | 3.12638 | 108.862 | 2.08862 | 2.088616 |
| | LuCO2 | 0.9966 | 2.68462 | 135.772 | 2.35772 | 2.357722 |
| | LuFT1 | 0.93924 | 2.75478 | 130.678 | 2.30678 | 2.306777 |
| | LuFT2 | 0.9502 | 2.94433 | 118.593 | 2.18593 | 2.185926 |

#2

| Primer Efficiency (factor 10) | | | | | | |
|--------------------------------------|----------------|----------------|----------------|-----------------------|-------------------|-----------------|
| <i>Sample</i> | <i>Primer</i> | <i>R^2</i> | <i>Slope</i> | <i>Efficiency (%)</i> | <i>Conversion</i> | <i>Value</i> |
| | | | - | | | |
| | <i>LuGAPDH</i> | <i>0.97378</i> | <i>3.86418</i> | <i>88.6802</i> | <i>1.8868</i> | <i>1.886802</i> |
| | | | - | | | |
| | <i>LuUBI2</i> | <i>0.93874</i> | <i>2.83438</i> | <i>160.188</i> | <i>2.60188</i> | <i>2.601882</i> |
| | | | - | | | |
| | <i>LuGI1.1</i> | <i>0.99999</i> | <i>0.60742</i> | <i>4834.91</i> | <i>49.3491</i> | <i>49.34912</i> |
| | | | - | | | |
| | <i>LuCO1</i> | <i>0.97491</i> | <i>3.12638</i> | <i>-99.8688</i> | <i>0.00131</i> | <i>0.001312</i> |
| | | | - | | | |
| | <i>LuCO2</i> | <i>0.9966</i> | <i>2.68462</i> | <i>97.5082</i> | <i>1.97508</i> | <i>1.975082</i> |
| | | | - | | | |
| | <i>LuFT1</i> | <i>0.93924</i> | <i>2.75478</i> | <i>207.128</i> | <i>3.07128</i> | <i>3.071284</i> |
| | | | - | | | |
| | <i>LuFT2</i> | <i>0.9502</i> | <i>2.94433</i> | <i>214.621</i> | <i>3.14621</i> | <i>3.146207</i> |

#3 –

| Primer Efficiency | | | | | | |
|--------------------------|----------------|----------------|----------------|-----------------------|------------------|-----------------|
| <i>Sample</i> | <i>Primer</i> | <i>R^2</i> | <i>Slope</i> | <i>Efficiency (%)</i> | <i>Converted</i> | <i>Value</i> |
| | | | - | | | |
| | <i>LuGAPDH</i> | <i>0.99505</i> | <i>3.67007</i> | <i>87.2728</i> | <i>1.87273</i> | <i>1.872728</i> |
| | | | - | | | |
| | <i>LuUBI2</i> | <i>0.90322</i> | <i>5.54602</i> | <i>51.464</i> | <i>1.51464</i> | <i>1.51464</i> |
| | | | - | | | |
| | <i>LuGI1.1</i> | <i>0.33336</i> | <i>0.55935</i> | <i>6034.64</i> | <i>61.3464</i> | <i>61.34644</i> |
| | | | - | | | |
| | <i>LuCO1</i> | <i>0.55246</i> | <i>3.56356</i> | <i>90.8175</i> | <i>1.90818</i> | <i>1.908175</i> |
| | | | - | | | |
| | <i>LuCO2</i> | <i>0.96165</i> | <i>3.61371</i> | <i>89.1143</i> | <i>1.89114</i> | <i>1.891143</i> |
| | | | - | | | |
| | <i>LuFT1</i> | <i>0.79856</i> | <i>3.08473</i> | <i>110.949</i> | <i>2.10949</i> | <i>2.109493</i> |
| | | | - | | | |
| | <i>LuFT2</i> | <i>0.98005</i> | <i>2.38037</i> | <i>163.09</i> | <i>2.6309</i> | <i>2.630896</i> |

4815

4816 **Appendix 6: Relative Gene Expression R commands:**

4817 `library(ggpubr)`

4818 `library("gridExtra")`

4819 `library("ggplot2")`

4820 `library("cowplot")`

4821 `library("dplyr")`

```

4822 library("ggpubr")
4823 library("viridis")
4824 library(readr)
4825 combined_vern_same_ind_clean1 <- read_csv("combined_vern_same_ind_clean1.csv",
4826           col_types = cols(Days_to_flower = col_number(),
4827           Experiment = col_character()))
4828 View(combined_vern_same_ind_clean1)
4829
4830 ggplot(combined_vern_same_ind_clean1, aes(x=Experiment, y=Days_to_flower, fill=Experiment)) +
4831   geom_boxplot() + labs(title = "Difference in Days to Flower Between Vernalization Years", x = "Vernalization
4832   Year", y = "Number of Days to Flower")
4833
4834 #multi-gene:
4835 library(ggplot2)
4836 ggplot(rge_input_luco1, aes(x=group, y=RGE, fill=group)) + geom_boxplot() + labs(title = "Relative Gene
4837   Expression - LuCO1", x = "Sample", y = "Relative Gene Expression")+ stat_compare_means(method = "anova")
4838 ggplot(rge_input_luco2, aes(x=group, y=RGE, fill=group)) + geom_boxplot() + labs(title = "Relative Gene
4839   Expression - LuCO2", x = "Sample", y = "Relative Gene Expression")
4840 ggplot(rge_input_luft1, aes(x=group, y=RGE, fill=group)) + geom_boxplot() + labs(title = "Relative Gene
4841   Expression - LuFT1", x = "Sample", y = "Relative Gene Expression")
4842 ggplot(rge_input_luft2, aes(x=group, y=RGE, fill=group)) + geom_boxplot() + labs(title = "Relative Gene
4843   Expression - LuFT2", x = "Sample", y = "Relative Gene Expression")
4844 ggplot(rge_input_lugi11, aes(x=group, y=RGE, fill=group)) + geom_boxplot() + labs(title = "Relative Gene
4845   Expression - LuGI1.1", x = "Sample", y = "Relative Gene Expression")
4846
4847
4848 #delta-delta:
4849 setwd("G:/Linum Project/rdir")
4850 #luco1:
4851 library(readr)

```

```

4852 rge_input_luco1_deltadelta <- read_csv("rge_input_luco1_deltadelta_withoutoutlyer.csv",
4853           col_types = cols(RGE = col_number()))
4854 View(rge_input_luco1_deltadelta)
4855 library(ggplot2)
4856 ggplot(rge_input_luco1_deltadelta, aes(x=group, y=RGE, fill=group)) + geom_boxplot() + labs(title = "Relative
4857 Gene Expression - LuCO1", x = "Sample", y = "Relative Gene Expression")+ stat_compare_means(label =
4858 "p.signif", method = "t.test",ref.group = ".all.")
4859 #luco2:
4860 library(readr)
4861 rge_input_luco2_deltadelta <- read_csv("rge_input_luco2_deltadelta_withoutoutlyer.csv",
4862           col_types = cols(RGE = col_number()))
4863 View(rge_input_luco2_deltadelta)
4864 library(ggplot2)
4865 ggplot(rge_input_luco2_deltadelta, aes(x=group, y=RGE, fill=group)) + geom_boxplot() + labs(title = "Relative
4866 Gene Expression - LuCO2", x = "Sample", y = "Relative Gene Expression")+stat_compare_means(label =
4867 "p.signif", method = "t.test",ref.group = ".all.")
4868 #lugi1.1:
4869 library(readr)
4870 rge_input_lugi1_1_deltadelta <- read_csv("rge_input_lugi1_1_deltadelta_withoutoutlyer.csv",
4871           col_types = cols(RGE = col_number()))
4872 View(rge_input_lugi1_1_deltadelta)
4873 library(ggplot2)
4874 ggplot(rge_input_lugi1_1_deltadelta, aes(x=group, y=RGE, fill=group)) + geom_boxplot() + labs(title = "Relative
4875 Gene Expression - LuGI1.1", x = "Sample", y = "Relative Gene Expression")+stat_compare_means(label =
4876 "p.signif", method = "t.test",ref.group = ".all.")
4877 #Luft1:
4878 #without Saf_10:
4879 library(readr)
4880 rge_input_luft1_deltadelta_withoutoutlyer <- read_csv("rge_input_luft1_deltadelta_withoutoutlyer.csv",
4881           col_types = cols(RGE = col_number()))

```

```

4882 View(rge_input_luft1_deltadelta_withoutoutlyer)
4883
4884 library(ggplot2)
4885 ggplot(rge_input_luft1_deltadelta_withoutoutlyer, aes(x=group, y=RGE, fill=group)) + geom_boxplot() +
4886 labs(title = "Relative Gene Expression - Luft1", x = "Sample", y = "Relative Gene
4887 Expression")+stat_compare_means(label = "p.signif", method = "t.test",ref.group = ".all.")
4888
4889 #luft2:
4890 library(readr)
4891 rge_input_luft2_deltadelta <- read_csv("rge_input_luft2_deltadelta_withoutoutlier.csv",
4892           col_types = cols(RGE = col_number()))
4893 View(rge_input_luft2_deltadelta)
4894 #without saf_10:
4895 library(readr)
4896 rge_input_luft2_deltadelta_withoutoutlier <- read_csv("rge_input_luft2_deltadelta_withoutoutlier.csv",
4897           col_types = cols(RGE = col_number()))
4898 View(rge_input_luft2_deltadelta_withoutoutlier)
4899 library(ggplot2)
4900 ggplot(rge_input_luft2_deltadelta_withoutoutlier, aes(x=group, y=RGE, fill=group)) + geom_boxplot() +
4901 labs(title = "Relative Gene Expression - Luft2", x = "Sample", y = "Relative Gene
4902 Expression")+stat_compare_means(label = "p.signif", method = "t.test",ref.group = ".all.")
4903
4904
4905 #luco1:
4906 rge_input_luco1$group <- factor(rge_input_luco1$group, levels = c("control_cul", "vern_cul", "control_wil",
4907 "vern_wil"))
4908 meanRGEluco1 <- tapply(rge_input_luco1$RGE, rge_input_luco1$group, mean)
4909 sdevRGEluco1 <- tapply(rge_input_luco1$RGE, rge_input_luco1$group, sd)
4910 numberRGEluco1 <- tapply(rge_input_luco1$RGE, rge_input_luco1$group, length)
4911 data.frame(mean=meanRGEluco1, std.dev=sdevRGEluco1, n=numberRGEluco1)

```

```

4912
4913 #anova:
4914 rge_luco1_anova <- lm(RGE ~ group, data = rge_input_luco1)
4915 anova(rge_luco1_anova)
4916
4917 #luco1 cultivar comparison:
4918 library(readr)
4919 rge_input_luco1_cul <- read_csv("rge_input_luco1_cul.csv",
4920                               + col_types = cols(RGE = col_number()))
4921
4922 #t-test:
4923 t.test(RGE~group, data = rge_input_luco1_cul)
4924
4925 #luco1 wild comparison:
4926 library(readr)
4927 rge_input_luco1_wil <- read_csv("rge_input_luco1_wil.csv",
4928                               col_types = cols(RGE = col_number()))
4929 View(rge_input_luco1_wil)
4930
4931 #t-test:
4932 t.test(RGE~group, data = rge_input_luco1_wil)
4933
4934
4935 #luco2:
4936 rge_input_luco2$group <- factor(rge_input_luco2$group, levels = c("control_cul", "vern_cul", "control_wil",
4937 "vern_wil"))
4938 meanRGEluco2 <- tapply(rge_input_luco2$RGE, rge_input_luco2$group, mean)
4939 sdevRGEluco2 <- tapply(rge_input_luco2$RGE, rge_input_luco2$group, sd)
4940 numberRGEluco2 <- tapply(rge_input_luco2$RGE, rge_input_luco2$group, length)

```

```

4941 data.frame(mean=meanRGEluco2, std.dev=sdevRGEluco2, n=numberRGEluco2)
4942
4943 #anova:
4944 rge_luco2_anova <- lm(RGE ~ group, data = rge_input_luco2)
4945 anova(rge_luco2_anova)
4946
4947 #luco2 cultivar comparison:
4948 library(readr)
4949 rge_input_luco2_cul <- read_csv("rge_input_luco2_cul.csv",
4950                               col_types = cols(RGE = col_number()))
4951 View(rge_input_luco2_cul)
4952
4953 #t-test:
4954 rge_input_luco2_cul$group <- factor(rge_input_luco2_cul$group, levels = c("control_cul", "vern_cul"))
4955 rge_input_luco2_cul_ttest <- lm(RGE ~ group, data = rge_input_luco2_cul)
4956 t.test(RGE~group, data = rge_input_luco2_cul)
4957
4958 #luco2 wild comparison:
4959 library(readr)
4960 rge_input_luco2_wil <- read_csv("rge_input_luco2_wil.csv",
4961                               col_types = cols(`1.053672824` = col_number()))
4962 View(rge_input_luco2_wil)
4963
4964 t.test(RGE~group, data = rge_input_luco2_wil)
4965
4966 #luft2:
4967 rge_input_luft2$group <- factor(rge_input_luft2$group, levels = c("control_cul", "vern_cul", "control_wil",
4968 "vern_wil"))
4969 meanRGEluft2 <- tapply(rge_input_luft2$RGE, rge_input_luft2$group, mean)

```



```

4970 sdevRGEluft2 <- tapply(rge_input_luft2$RGE, rge_input_luft2$group, sd)
4971 numberRGEluft2 <- tapply(rge_input_luft2$RGE, rge_input_luft2$group, length)
4972 data.frame(mean=meanRGEluft2, std.dev=sdevRGEluft2, n=numberRGEluft2)
4973
4974 #anova
4975 rge_luft2_anova <- lm(RGE ~ group, data = rge_input_luft2)
4976 anova(rge_luft2_anova)
4977
4978 #luft2 cultivar comparison:
4979 library(readr)
4980 rge_input_luft2_cul <- read_csv("rge_input_luft2_cul.csv",
4981                               col_types = cols(RGE = col_number()))
4982 View(rge_input_luft2_cul)
4983
4984 #t-test:
4985 rge_input_luft2_cul$group <- factor(rge_input_luft2_cul$group, levels = c("control_cul", "vern_cul"))
4986 rge_input_luft2_cul_ttest <- lm(RGE ~ group, data = rge_input_luft2_cul)
4987 t.test(RGE~group, data = rge_input_luft2_cul)
4988
4989 #luft2 wild comparison:
4990 library(readr)
4991 rge_input_luft2_wil <- read_csv("rge_input_luft2_wil.csv",
4992                               + col_types = cols(Ind = col_character(),
4993                               + RGE = col_number(), group = col_character()))
4994 View(rge_input_luft2_wil)
4995
4996
4997 #luft1:

```

```

4998 rge_input_luft1$group <- factor(rge_input_luft1$group, levels = c("control_cul", "vern_cul", "control_wil",
4999 "vern_wil"))
5000 meanRGEluft1 <- tapply(rge_input_luft1$RGE, rge_input_luft1$group, mean)
5001 sdevRGEluft1 <- tapply(rge_input_luft1$RGE, rge_input_luft1$group, sd)
5002 numberRGEluft1 <- tapply(rge_input_luft1$RGE, rge_input_luft1$group, length)
5003 data.frame(mean=meanRGEluft1, std.dev=sdevRGEluft1, n=numberRGEluft1)
5004
5005 #anova
5006 rge_luft1_anova <- lm(RGE ~ group, data = rge_input_luft1)
5007 anova(rge_luft1_anova)
5008
5009
5010 #t-test:
5011 rge_input_luft2_wil$group <- factor(rge_input_luft2_wil$group, levels = c("control_cul", "vern_cul"))
5012 rge_input_luft2_wil_ttest <- lm(RGE ~ group, data = rge_input_luft2_wil)
5013 t.test(RGE~group, data = rge_input_luft2_wil)
5014
5015 #luft1 cultivar comparison:
5016 library(readr)
5017 rge_input_luft1_cul <- read_csv("rge_input_luft1_cul.csv",
5018                               col_types = cols(...1 = col_character(),
5019                                                  RGE = col_number(), group = col_character()))
5020 View(rge_input_luft1_cul)
5021
5022 #t-test:
5023 t.test(RGE~group, data = rge_input_luft1_cul)
5024
5025 #luft1 wild comparison:
5026 library(readr)

```

```

5027 rge_input_luft1_wil <- read_csv("rge_input_luft1_wil.csv",
5028                               col_types = cols(RGE = col_number()))
5029 View(rge_input_luft1_wil)
5030
5031 #t-test:
5032 t.test(RGE~group, data = rge_input_luft1_wil)
5033
5034 #lugi11:
5035 rge_input_lugi11$group <- factor(rge_input_lugi11$group, levels = c("control_cul", "vern_cul", "control_wil",
5036 "vern_wil"))
5037 meanRGElugi11 <- tapply(rge_input_lugi11$RGE, rge_input_lugi11$group, mean)
5038 sdevRGElugi11 <- tapply(rge_input_lugi11$RGE, rge_input_lugi11$group, sd)
5039 numberRGElugi11 <- tapply(rge_input_lugi11$RGE, rge_input_lugi11$group, length)
5040 data.frame(mean=meanRGElugi11, std.dev=sdevRGElugi11, n=numberRGElugi11)
5041
5042 #anova
5043 rge_lugi11_anova <- lm(RGE ~ group, data = rge_input_lugi11)
5044 anova(rge_lugi11_anova)
5045
5046 #lugi11 cultivar comparison:
5047 library(readr)
5048 rge_input_lugi11_cul <- read_csv("rge_input_lugi11_cul.csv",
5049                               col_types = cols(RGE = col_number()))
5050 View(rge_input_lugi11_cul)
5051
5052 #t-test:
5053 t.test(RGE~group, data = rge_input_lugi11_cul)
5054
5055 #lugi11 wild comparison:

```

```

5056 library(readr)
5057 rge_input_lugi11_wil <- read_csv("rge_input_lugi11_wil.csv",
5058                               col_types = cols(RGE = col_number()))
5059 View(rge_input_lugi11_wil)
5060
5061 #t-test:
5062 t.test(RGE~group, data = rge_input_lugi11_wil)
5063
5064 Appendix 7: Relative Gene Expression GLM R Commands:
5065 library(readr)
5066 combined_vern_same_ind <- read_csv("combined_vern_same_ind.csv",
5067                                   col_types = cols(Lat = col_number(),
5068                                                   Alt = col_number(), Height = col_number(),
5069                                                   Stem_no = col_number(), Bud_no = col_number(),
5070                                                   Days_to_fl = col_number(), pc_1 = col_number(),
5071                                                   pc_2 = col_number(), pc_3 = col_number()))
5072 View(combined_vern_same_ind)
5073
5074 #plot:
5075 x <- combined_vern_same_ind$Height
5076 y <- combined_vern_same_ind$Days_to_fl
5077 plot(x, y, main = "Height against Days to Flowering",
5078      xlab = "Plant Height (Cm)", ylab = "Days to Flowering",
5079      pch = 19, frame = FALSE)
5080
5081 #scatterplot
5082 install.packages("car")
5083 library("car")
5084 #height:

```

```

5085 View(combined_vern_same_ind)
5086 scatterplot(Days_to_fl ~ Height | Experiment, data = combined_vern_same_ind,
5087             smooth = TRUE, regLine = TRUE, grid = FALSE, frame = FALSE, xlab = "Plant Height (Cm)", ylab = "Days to
5088 Flowering")
5089 abline(lm(Height ~ Days_to_fl data = combined_vern_same_ind),col="red")
5090
5091 ggscatter(combined_vern_same_ind, x = "Height", y = "Days_to_fl", size = 2.0,
5092           rug = TRUE,                # Add marginal rug
5093           color = "Experiment", palette = "jco", facet.by = "Experiment",add = "reg.line", conf.int = TRUE) +
5094           stat_cor(aes(color = Experiment), method = "pearson")
5095
5096 #stem no:
5097 scatterplot(Days_to_fl ~ Stem_no | Experiment, data = combined_vern_same_ind,
5098             smooth = TRUE, regLine = TRUE, grid = FALSE, frame = FALSE, xlab = "Stem number", ylab = "Days to
5099 Flowering")
5100 abline(lm(Stem_no ~ Days_to_fl data = combined_vern_same_ind),col="red")
5101
5102
5103 ggscatter(combined_vern_same_ind, x = "Stem_no", y = "Days_to_fl", size = 2.0,
5104           rug = TRUE,                # Add marginal rug
5105           color = "Experiment", palette = "jco", facet.by = "Experiment",add = "reg.line", conf.int = TRUE) +
5106           stat_cor(aes(color = Experiment), method = "pearson")
5107
5108 #bud no:
5109 scatterplot(Days_to_fl ~ Bud_no | Experiment, data = combined_vern_same_ind,
5110             smooth = TRUE, regLine = TRUE, grid = FALSE, frame = FALSE, xlab = "Bud number", ylab = "Days to
5111 Flowering")
5112 abline(lm(Stem_no ~ Days_to_fl data = combined_vern_same_ind),col="red")
5113

```

```

5114
5115 ggscatter(combined_vern_same_ind, x = "Bud_no", y = "Days_to_fl", size = 2.0,
5116         rug = TRUE,                # Add marginal rug
5117         color = "Experiment", palette = "jco", facet.by = "Experiment", add = "reg.line", conf.int = TRUE) +
5118         stat_cor(aes(color = Experiment), method = "pearson")
5119
5120 #pc1:
5121 scatterplot(pc_1 ~ Days_to_fl | Experiment, data = combined_vern_same_ind,
5122            smooth = FALSE, grid = FALSE, frame = FALSE, xlab = "Days to First Flower", ylab = "Climate (PC1)")
5123 abline(lm(pc_1 ~ Days_to_fl data = combined_vern_same_ind), col="red")
5124
5125 plot(pc_1 ~ Days_to_fl, data = combined_vern_same_ind)
5126 abline(lm(pc_1 ~ Days_to_fl, data = combined_vern_same_ind), col="red")
5127 summary(abline)
5128
5129 #Alt:
5130 scatterplot(Days_to_fl ~ Alt | Experiment, data = combined_vern_same_ind,
5131            smooth = FALSE, grid = FALSE, frame = FALSE, xlab = "Days to First Flower", ylab = "Altitude (M)")
5132 abline(lm(Alt ~ Days_to_fl data = combined_vern_same_ind), col="red")
5133
5134 plot(Alt ~ Days_to_fl, data = combined_vern_same_ind)
5135 abline(lm(Alt ~ Days_to_fl, data = combined_vern_same_ind), col="red")
5136 summary(abline)
5137
5138
5139 #Model summary
5140 summary1 <- summary(glm(Stem_no ~ Days_to_fl, data = combined_vern_same_ind))
5141 adjRsqr <- summary1$adj.r.squared
5142 fStat <- summary1$statistic

```

```

5143  pValue <- pf(fStat[summary1])
5144  summary(summary1)
5145  #PC1 - Days to fl LM:
5146  pc1daysmod<-cbind(combined_vern_same_ind$Days_to_fl, combined_vern_same_ind$pc_1)
5147  pc1daysmod2<-pc1daysmodlm<-lm(pc1daysmod~Experiment,data=combined_vern_same_ind)
5148  summary(pc1daysmodlm)
5149
5150
5151  #Alt - Days to fl GLM:
5152  altdaysmod<-cbind(combined_vern_same_ind$Days_to_fl, combined_vern_same_ind$Alt)
5153  altdaysmod2<-altdaysmodlm<-lm(altdaysmod~Experiment,data=combined_vern_same_ind)
5154  summary(altdaysmodlm)
5155
5156  #Lat - Days to fl GLM:
5157  latdaysmod<-cbind(combined_vern_same_ind$Days_to_fl, combined_vern_same_ind$Lat)
5158  latdaysmod2<-latdaysmodlm<-lm(altdaysmod~Experiment,data=combined_vern_same_ind)
5159  summary(latdaysmodlm)
5160
5161  #RGE - Days to fl GLM:
5162  library(readr)
5163  combined_loci_wild_deltadelta_ft2clear <- read_csv("combined_loci_wild_deltadelta_ft2clear.csv",
5164          col_types = cols(`RGE_c` = col_number(),
5165          `RGE_v` = col_number(), `RGE_Difference` = col_number(),
5166          Days_to_fl = col_number(), Lat = col_number(),
5167          Alt = col_number(), pc = col_number(),
5168          pc2 = col_number(), pc3 = col_number()))
5169  View(combined_loci_wild_deltadelta_ft2clear)
5170
5171  #Cultivars:

```

```

5172 library(readr)
5173 combined_loci_cult_deltadelta_fl <- read_csv("combined_loci_cult_deltadelta_fl.csv",
5174           col_types = cols(`RGE_c` = col_number(),
5175           `RGE_v` = col_number(), `RGE_Difference` = col_number(),
5176           Lat = col_number(), Days_to_fl = col_number(),
5177           `Height_(Cm)` = col_number()))
5178 View(combined_loci_cult_deltadelta_fl)
5179
5180
5181 #Gene expression influence in flowering time:
5182 #Wild
5183 #subset data to loci first:
5184 Luco1_data <- subset(combined_loci_wild_deltadelta_ft2clear, Loci == "LuCO1")
5185 Luco2_data <- subset(combined_loci_wild_deltadelta_ft2clear, Loci == "LuCO2")
5186 Luft1_data <- subset(combined_loci_wild_deltadelta_ft2clear, Loci == "LuFT1")
5187 Luft2_data <- subset(combined_loci_wild_deltadelta_ft2clear, Loci == "LuFT2")
5188 Lugi11_data <- subset(combined_loci_wild_deltadelta_ft2clear, Loci == "LuGI1.1")
5189
5190 #GLM:
5191 modluco1wild<-glm(Luco1_data$Days_to_fl ~ Luco1_data$"RGE_Difference")
5192 modluco2wild<-glm(Luco2_data$Days_to_fl ~ Luco2_data$"RGE_Difference")
5193 modluft1wild<-glm(Luft1_data$Days_to_fl ~ Luft1_data$"RGE_Difference")
5194 modluft2wild<-glm(Luft2_data$Days_to_fl ~ Luft2_data$"RGE_Difference")
5195 modlugi11wild<-glm(Lugi11_data$Days_to_fl ~ Lugi11_data$"RGE_Difference")
5196
5197 summary(modluco1wild)
5198 summary(modluco2wild)
5199 summary(modluft1wild)
5200 summary(modluft2wild)

```



```

5201 summary(modlugi11wild)
5202
5203 #Cultivars
5204 Luco1_cult_data <- subset(combined_loci_cult_deltadelta_fl, Loci == "LuCO1")
5205 Luco2_cult_data <- subset(combined_loci_cult_deltadelta_fl, Loci == "LuCO2")
5206 Luft1_cult_data <- subset(combined_loci_cult_deltadelta_fl, Loci == "LuFT1")
5207 Luft2_cult_data <- subset(combined_loci_cult_deltadelta_fl, Loci == "LuFT2")
5208 Lugi11_cult_data <- subset(combined_loci_cult_deltadelta_fl, Loci == "LuGI1.1")
5209
5210 #GLM:
5211 modluco1cul<-glm(Luco1_cult_data$Days_to_fl ~ Luco1_cult_data$"RGE_Difference")
5212 modluco2cul<-glm(Luco2_cult_data$Days_to_fl ~ Luco2_cult_data$"RGE_Difference")
5213 modluft1cul<-glm(Luft1_cult_data$Days_to_fl ~ Luft1_cult_data$"RGE_Difference")
5214 modluft2cul<-glm(Luft2_cult_data$Days_to_fl ~ Luft2_cult_data$"RGE_Difference")
5215 modlugi11cul<-glm(Lugi11_cult_data$Days_to_fl ~ Lugi11_cult_data$"RGE_Difference")
5216
5217 summary(modluco1cul)
5218 summary(modluco2cul)
5219 summary(modluft1cul)
5220 summary(modluft2cul)
5221 summary(modlugi11cul)
5222
5223 #charts:
5224 View(Luco1_data)
5225
5226 write.csv(Luco1_data,"G:/Linum Project/rdir/Luco1_data_only.csv", row.names = FALSE)
5227
5228 #
5229 #Delta-delta data:

```

```

5230 #Load packages
5231 library(ggplot2)
5232 library(ggpubr)
5233 library(viridis)
5234 library(readr)
5235 #Cultivars
5236 setwd("G:/Linum Project/rdir")
5237
5238
5239 ggscatter(combined_loci_cult_deltadelta_fl, x = "Days_to_fl", y = "RGE_Difference", size = 1.0,
5240           rug = TRUE,                # Add marginal rug
5241           color = "Loci", palette = "uchicago", facet.by = "Loci", add = "reg.line", conf.int = TRUE) +
5242           stat_cor(aes(color = Loci), label.sep = ";", label.x.npc = "left", label.y.npc=0.95, method =
5243 "pearson")+geom_point(aes(color = Loci))+stat_smooth(method="glm",se=FALSE)+
5244 theme_light()+ggtitle("Cultivar RGE-Flowering")
5245
5246 #Lat:
5247 ggscatter(combined_loci_cult_deltadelta_fl, x = "Lat", y = "RGE_Difference", size = 1.0,
5248           rug = TRUE,                # Add marginal rug
5249           color = "Loci", palette = "uchicago", facet.by = "Loci", add = "reg.line", conf.int = TRUE) +
5250           stat_cor(aes(color = Loci), label.sep = ";", label.x.npc = "left", label.y.npc=0.95, method =
5251 "pearson")+geom_point(aes(color = Loci))+stat_smooth(method="glm",se=FALSE)+
5252 theme_light()+ggtitle("Cultivar RGE-Latitude")
5253
5254 #GLM:
5255 modluco1cullat<-glm(Luco1_cult_data$Lat ~ Luco1_cult_data$"RGE_Difference")
5256 modluco2cullat<-glm(Luco2_cult_data$Lat ~ Luco2_cult_data$"RGE_Difference")
5257 modluft1cullat<-glm(Luft1_cult_data$Lat ~ Luft1_cult_data$"RGE_Difference")
5258 modluft2cullat<-glm(Luft2_cult_data$Lat ~ Luft2_cult_data$"RGE_Difference")
5259 modlugi11cullat<-glm(Lugi11_cult_data$Lat ~ Lugi11_cult_data$"RGE_Difference")

```

```

5260
5261 summary(modluco1cullat)
5262 summary(modluco2cullat)
5263 summary(modluft1cullat)
5264 summary(modluft2cullat)
5265 summary(modlugi11cullat)
5266
5267 #Wild:
5268 ggscatter(combined_loci_wild_deltadelta_ft2clear, x = "Days_to_fl", y = "RGE_Difference", size = 1.0,
5269           rug = TRUE,                # Add marginal rug
5270           color = "Loci", palette = "uchicago", facet.by = "Loci", add = "reg.line", conf.int = TRUE) +
5271           stat_cor(aes(color = Loci), label.sep = ";", label.x.npc = "left", label.y.npc=0.32, method =
5272 "pearson")+geom_point(aes(color = Loci))+stat_smooth(method="glm",se=FALSE)+
5273 theme_light()+ggtitle("Wild RGE-Flowering")
5274
5275 #Lat
5276 ggscatter(combined_loci_wild_deltadelta_ft2clear, x = "Lat", y = "RGE_Difference", size = 1.0,
5277           rug = TRUE,                # Add marginal rug
5278           color = "Loci", palette = "uchicago", facet.by = "Loci", add = "reg.line", conf.int = TRUE) +
5279           stat_cor(aes(color = Loci), label.sep = ";", label.x.npc = "left", label.y.npc="centre", method =
5280 "pearson")+geom_point(aes(color = Loci))+stat_smooth(method="glm",se=FALSE)+
5281 theme_light()+ggtitle("Wild RGE-Latitude")
5282
5283 #GLM:
5284 modluco1willat<-glm(Luco1_data$Lat ~ Luco1_data$"RGE_Difference")
5285 modluco2willat<-glm(Luco2_data$Lat ~ Luco2_data$"RGE_Difference")
5286 modluft1willat<-glm(Luft1_data$Lat ~ Luft1_data$"RGE_Difference")
5287 modluft2willat<-glm(Luft2_data$Lat ~ Luft2_data$"RGE_Difference")
5288 modlugi11willat<-glm(Lugi11_data$Lat ~ Lugi11_data$"RGE_Difference")
5289

```

```

5290 summary(modluco1willat)
5291 summary(modluco2willat)
5292 summary(modluft1willat)
5293 summary(modluft2willat)
5294 summary(modlugi11willat)
5295
5296
5297 #PC1:
5298 ggscatter(combined_loci_wild_deltadelta_ft2clear, x = "pc", y = "RGE_Difference", size = 1.0,
5299           rug = TRUE,                # Add marginal rug
5300           color = "Loci", palette = "uchicago", facet.by = "Loci", add = "reg.line", conf.int = TRUE) +
5301           stat_cor(aes(color = Loci), label.sep = ";", label.x.npc = "left", label.y.npc=0.38, method =
5302 "pearson")+geom_point(aes(color = Loci))+stat_smooth(method="glm",se=FALSE)+
5303 theme_light()+ggtitle("Wild RGE-PC1")
5304
5305 #GLM:
5306 modluco1wilpc<-glm(Luco1_data$pc ~ Luco1_data$"RGE_Difference")
5307 modluco2wilpc<-glm(Luco2_data$pc ~ Luco2_data$"RGE_Difference")
5308 modluft1wilpc<-glm(Luft1_data$pc ~ Luft1_data$"RGE_Difference")
5309 modluft2wilpc<-glm(Luft2_data$pc ~ Luft2_data$"RGE_Difference")
5310 modlugi11wilpc<-glm(Lugi11_data$pc ~ Lugi11_data$"RGE_Difference")
5311
5312 summary(modluco1wilpc)
5313 summary(modluco2wilpc)
5314 summary(modluft1wilpc)
5315 summary(modluft2wilpc)
5316 summary(modlugi11wilpc)
5317
5318 #scatterplot

```

```



5319 install.packages("tidyverse")
5320 library(tidyverse)
5321 library(ggplot2)
5322
5323 ggplot(combined_loci_cult_deltadelta_fl, aes(x=RGE_Difference, y=Days_to_fl, shape=Loci, color=Loci)) +
5324   geom_point(size=2.5, shape=18) + ggtitle("Cultivar Flowering initiation")+labs(y= "Days to Flowering", x =
5325 "Relative Gene Expression Difference")
5326
5327 #wild
5328 ggplot(combined_loci_wild_deltadelta_ft2clear, aes(x=RGE_Difference, y=Days_to_fl, shape=Loci, color=Loci))
5329 +
5330   geom_point(size=2.5, shape=18) + ggtitle("Wild Flowering initiation")+labs(y= "Days to Flowering", x =
5331 "Relative Gene Expression Difference")
5332
5333 #
5334 s3d <- scatterplot3d(combined_vern_same_ind, pch = 16, color=colors)
5335 legend(s3d$xyz.convert(7.5, 3, 4.5), legend = levels(combined_vern_same_ind$Experiment),
5336       col = c("#999999", "#E69F00", "#56B4E9"), pch = 16)
5337

```

5338 **Appendix 8: Relative gene expression difference based on using two HKGs**




5339 Relative gene expression comparison for the different genes tested in relation to each
5340 species and treatment:

5341



5349

5350



5359

5360

5361

5362 **Appendix 9: Scatterplot for gene expression and latitude correlation under two**

5363 **Housekeeping genes**


5364 Scatterplot showing relationships between latitude and relative gene expression (RGE) differences

5365 for five flowering time genes. A = wild types, B = Cultivars. RGE differences were calculated using 2


5366 HKGs:

5367

5368 **A**



B



5378

5379

5380

5381

5382

| Gene | Group test | Cultivar t-test | Wild t-test |
|---------|------------|-----------------|-------------|
| LuCO1 | 0.4672 | 0.9027 | 0.7722 |
| LuCO2 | 0.194 | 0.09186 | 0.5341 |
| LuGI1.1 | 0.9434 | 0.9163 | 0.6533 |
| LuFT1 | 0.2373 | 0.08303 | 0.4711 |
| LuFT2 | 0.354 | 0.2029 | 0.6945 |

5383

5384

5385

5386

5387

5388

5389

5390 **Appendix 10: List of *Linum* individuals for pollen viability germination experiments in chapter 3 of**
 5391 **the thesis with number of individuals in each treatments for each individuals.**

5392

| Individual | Population | Species | Numbers in 20C | Numbers in 25C | Numbers in 30C |
|------------|------------|---------------|----------------|----------------|----------------|
| Ara | Ara | Usitatissimum | 0 | 0 | 13 |
| Bey | Bey | Usitatissimum | 0 | 3 | 0 |
| Ble | Ble | Usitatissimum | 4 | 0 | 0 |
| Bol | Bol | Usitatissimum | 0 | 0 | 0 |
| Boo | Boo | Usitatissimum | 0 | 22 | 0 |
| Ede | Ede | Usitatissimum | 0 | 4 | 0 |
| Kar | Kar | Usitatissimum | 3 | 0 | 10 |
| Olg | Olg | Usitatissimum | 0 | 16 | 3 |
| Ome | Ome | Usitatissimum | 30 | 11 | 0 |
| Rab | Rab | Usitatissimum | 0 | 7 | 0 |
| Sar | Sar | Usitatissimum | 2 | 6 | 0 |
| Suz | Suz | Usitatissimum | 0 | 3 | 0 |
| Tin | Tin | Usitatissimum | 6 | 0 | 7 |
| Vol | Vol | Usitatissimum | 0 | 0 | 8 |
| 3_12 | 3 | bienne | 5 | 2 | 0 |
| 6_17 | 6 | bienne | 7 | 0 | 6 |
| 10_24 | 10 | bienne | 1 | 12 | 0 |
| 15_31 | 15 | bienne | 4 | 0 | 0 |
| 15_32 | 15 | bienne | 1 | 0 | 5 |
| 19_22 | 19 | bienne | 3 | 0 | 2 |
| 19_30 | 19 | bienne | 1 | 0 | 0 |
| 19_28 | 19 | bienne | 3 | 0 | 2 |
| Man_6 | Man | bienne | 7 | 0 | 0 |
| Mat_2 | Mat | bienne | 12 | 2 | 0 |
| Saf_19 | Saf | bienne | 4 | 0 | 0 |
| Sut_41 | Sut | bienne | 2 | 6 | 0 |
| Tym_21 | Tym | bienne | 1 | 0 | 0 |
| 6_30 | 6 | bienne | 0 | 1 | 0 |
| 13_6 | 13 | bienne | 0 | 3 | 0 |
| 15_17 | 15 | bienne | 0 | 3 | 0 |
| 19_23 | 19 | bienne | 0 | 7 | 0 |
| 19_32 | 19 | bienne | 0 | 1 | 0 |
| Ezc_7 | Ezc | bienne | 0 | 5 | 0 |
| Lla_33 | Lla | bienne | 0 | 1 | 0 |
| Lla_13 | Lla | bienne | 0 | 1 | 0 |
| Lla_31 | Lla | bienne | 0 | 5 | 0 |
| Lla_18 | Lla | bienne | 0 | 4 | 0 |
| Lla_22 | Lla | bienne | 0 | 1 | 0 |
| 8_16 | 8 | bienne | 0 | 0 | 6 |
| 9_27 | 9 | bienne | 0 | 0 | 1 |


| | | | | | |
|---------|------|--------|---|---|---|
| 10_30 | 10 | bienne | 0 | 0 | 1 |
| 13_12 | 13 | bienne | 0 | 0 | 2 |
| Dor_B | Dor | bienne | 0 | 0 | 8 |
| Ezc_9 | Ezc | bienne | 0 | 0 | 3 |
| low2_26 | low2 | bienne | 0 | 0 | 2 |
| low2_25 | low2 | bienne | 0 | 0 | 1 |
| Lla_20 | Lla | bienne | 0 | 0 | 1 |
| Lla_29 | Lla | bienne | 0 | 0 | 2 |
| Man_8 | Man | bienne | 0 | 0 | 6 |
| Saf_9 | Saf | bienne | 0 | 0 | 3 |
| Vil_36 | Vil | bienne | 0 | 0 | 2 |

5393

5394

5395 **Appendix 11: Fluorescent microscopy observation of a *Linum* flower conserved at 4°C in 70%EtOH**
5396 **solution, showing pollen and tube observations after 72 hours preservation.**

5397



5398

5399

5400

5401

5402


5403

5404


5405 **Appendix 12: 10×20 magnification of fluorescence observed after flower preservation using both**
5406 **FAA (A) and 70% EtOH (B) solutions. The flowers were from the same individual observed in the**
5407 **glasshouse conditions.**

5408

5409 A



B



5410

5411

5412

5413

5414


5415

5416

5417

5418 **Appendix 13: Pollen Tube Observation Using 0.5% (w/v) Aniline blue in Potassium Phosphate**

5419 **(KH₂PO₄)**



5420

5421

5422 **Appendix 14: Pollen Modelling Commands**

```
5423 #test for normality (pollen tube)
5424 plot(density(Pollen$Percent))
5425 hist(Pollen$Percent, main="Histogram for Percentage of Pollen Tube %", xlab="% of Pollen Tube Counts")
5426 densityPollen<-density(Pollen$Percent)
5427 lines(densityPollen$x,densityPollen$y*5000)
5428 shapiro.test(Pollen$Percent)
5429 #data is not normally distributed
5430 #adjust for non-normally distributed data (tukey's)
5431
5432 #Data for 20C - Cold (pollen tube)
5433 setwd("G:/Linum Project/Pollen germination")
5434 library(readxl)
5435 Pollencold <- as.data.frame(read_excel("Pollen_and_tube_count2.xlsx", sheet = "20c", na = "NA"))
5436 str(Pollencold)
5437 View(Pollencold)
5438 Pollencold$Treatment<-as.factor(Pollencold$Treatment)
5439
5440 #test for normality - Cold
5441 plot(density(Pollencold$Percent))
5442 hist(Pollencold$Percent, main="Histogram for Percentage of Pollen Tube %", xlab="% of Pollen Tube Counts")
5443 densityPollencold<-density(Pollencold$Percent)
5444 lines(densityPollencold$x,densityPollencold$y*5000)
5445 shapiro.test(Pollencold$Percent)
5446 #data is not normally distributed
5447 #adjust for non-normally distributed data (tukey's)
5448
5449 #Pollen number:
5450 plot(density(Pollen$Count))
```

```

5451 hist(Pollen$Count, main = "Pollen Count Distribution", xlab = "Pollen Count")
5452 densityCount<-density(Pollen$Count)
5453 lines(densityCount$x,densityCount$y*5000)
5454 shapiro.test(Pollen$Count)
5455
5456 #test pollen count versus treatment, quasibinomial model - TOTAL
5457 count<-Pollen$Count
5458 modcount<-lm(Pollen$Count~Treatment, data=Pollen)
5459 summary(modcount)
5460 tukcount<-glht(modcount,mcp(Treatment="Tukey"))
5461 summary(tukcount)
5462 Props<-Pollen$Tube/Pollen$Count
5463 Pollen<-data.frame(Pollen,count)
5464 str(Pollen)
5465 barcentres<-
5466 barplot(tapply(Pollen$Count,Pollen$Treatment,mean),ylim=c(0,max(tapply(Pollen$Count,Pollen$Treatment,m
5467 ean)+5)), ylab = "Pollen Count", xlab = "Treatment", main = "Number of Pollen vs Treatment")
5468 means<-tapply(Pollen$Count,Pollen$Treatment,mean)
5469 ses<-tapply(Pollen$Count,Pollen$Treatment,sd)/sqrt(tapply(Pollen$Count,Pollen$Treatment,length))
5470 arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
5471
5472 #test pollen count versus treatment, Poisson correction - TOTAL
5473 count<-Pollen$Count
5474 modcountpoisson<-glm(Pollen$Count~Treatment, family=poisson(), data=Pollen,)
5475 summary(modcountpoisson)
5476 tukcount<-glht(modcountpoisson,mcp(Treatment="Tukey"))
5477 summary(tukcount)
5478 Props<-Pollen$Tube/Pollen$Count
5479 Pollen<-data.frame(Pollen,count)

```

```

5480 str(Pollen)
5481 barcentres<-
5482 barplot(tapply(Pollen$Count,Pollen$Treatment,mean),ylim=c(0,max(tapply(Pollen$Count,Pollen$Treatment,m
5483 ean)+5)), ylab = "Pollen Count", xlab = "Treatment", main = "Number of Pollen vs Treatment")
5484 means<-tapply(Pollen$Count,Pollen$Treatment,mean)
5485 ses<-tapply(Pollen$Count,Pollen$Treatment,sd)/sqrt(tapply(Pollen$Count,Pollen$Treatment,length))
5486 arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
5487
5488 #test pollen tube count versus treatment, Poisson correction - TOTAL
5489 tubecount<-Pollen$Tube
5490 modtubepoisson<-glm(Pollen$Tube~Treatment, family=poisson(), data=Pollen,)
5491 summary(modtubepoisson)
5492 tuktube<-glht(modtubepoisson,mcp(Treatment="Tukey"))
5493 summary(tuktube)
5494 Props<-Pollen$Tube/Pollen$Count
5495 Pollentube<-data.frame(Pollen,tubecount)
5496 str(Pollentube)
5497 barcentres<-
5498 barplot(tapply(Pollen$Tube,Pollen$Treatment,mean),ylim=c(0,max(tapply(Pollen$Tube,Pollen$Treatment,mea
5499 n)+2)), ylab = "Pollen Tube Count", xlab = "Treatment", main = "Number of Pollen Tubes vs Treatment")
5500 means<-tapply(Pollen$Tube,Pollen$Treatment,mean)
5501 ses<-tapply(Pollen$Tube,Pollen$Treatment,sd)/sqrt(tapply(Pollen$Tube,Pollen$Treatment,length))
5502 arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
5503
5504 #test reached versus treatment, binomial model
5505 mod1<-glm(Reached~Treatment,family=binomial(link='logit'),data=Pollen)
5506 summary(mod1)
5507 library(multcomp)
5508 tuk1<-glht(mod1,mcp(Treatment="Tukey"))
5509 summary(tuk1)

```

```

5510  barplot(tapply(Pollen$Reached,Pollen$Treatment,mean))
5511  barcentres<-
5512  barplot(tapply(Pollen$Reached,Pollen$Treatment,mean),ylim=c(0,max(tapply(Pollen$Reached,Pollen$Treatm
5513  ent,mean)+0.1)), ylab = "Proportion of Pollen Reaching Ovary", xlab = "Treatment", main = "Proportion of
5514  Pollen Reaching Ovary vs Treatment")
5515  means<-tapply(Pollen$Reached,Pollen$Treatment,mean)
5516  ses<-tapply(Pollen$Reached,Pollen$Treatment,sd)/sqrt(tapply(Pollen$Reached,Pollen$Treatment,length))
5517  arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
5518
5519  #test prop tube versus treatment, quasibinomial model
5520  Prop<-cbind(Pollen$Tube, Pollen$Count)
5521  mod2<-glm(Prop~Treatment, family=quasibinomial(link = 'logit'), data=Pollen)
5522  summary(mod2)
5523  tuk2<-glht(mod2,mcp(Treatment="Tukey"))
5524  summary(tuk2)
5525  Props<-Pollen$Tube/Pollen$Count
5526  Pollen<-data.frame(Pollen,Props)
5527  str(Pollen)
5528  barcentres<-
5529  barplot(tapply(Pollen$Props,Pollen$Treatment,mean),ylim=c(0,max(tapply(Pollen$Props,Pollen$Treatment,m
5530  ean)+0.1)), ylab = "Proportion of Pollen Tube Number", xlab = "Treatment", main = "Proportion of Pollen Tube
5531  vs Treatment")
5532  means<-tapply(Pollen$Props,Pollen$Treatment,mean)
5533  ses<-tapply(Pollen$Props,Pollen$Treatment,sd)/sqrt(tapply(Pollen$Props,Pollen$Treatment,length))
5534  arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
5535
5536  #test proptubes versus treatment x latitude
5537  modlat<-glm(Prop~Treatment*Lat, family=quasibinomial(link = 'logit'), data=Pollen)
5538  summary(modlat)
5539  anova(modlat,test = "F")

```

```

5540 interactlat<-interaction(Pollen$Treatment,Pollen$Lat)
5541 Pollen<-data.frame(Pollen,interactlat)
5542 View(Pollen)
5543 modlatx<-glm(Prop~1+interactlat, family=quasibinomial(link = 'logit'), data=Pollen)
5544 summary(modlatx)
5545 tuklatx<-glht(modlatx,mcp(interactlat="Tukey"))
5546 summary(tuklatx)
5547 Props<-Pollen$Tube/Pollen$Count
5548 Pollen<-data.frame(Pollen,Props)
5549 barcentres<-
5550 barplot(tapply(Pollen$Props,Pollen$interactlat,mean),ylim=c(0,max(tapply(Pollen$Props,Pollen$interactlat,mean
5551 )+1.0)), ylab = "Proportion of Pollen Tube Number/Pollen Count", xlab = "Treatment and Latitude", main =
5552 "Proportion of Pollen Tube per Latitude and Treatment")
5553
5554 #test prop tubes/count versus treatment x species
5555 mod3<-glm(Props~Treatment*Species, family=quasibinomial(link = 'logit'), data=Pollen)
5556 summary(mod3)
5557 anova(mod3,test="F")
5558 interact<-interaction(Pollen$Treatment,Pollen$Species)
5559 Pollen<-data.frame(Pollen,interact)
5560 mod3x<-glm(Props~1+interact, family=quasibinomial(link = 'logit'), data=Pollen)
5561 tuk3x<-glht(mod3x,mcp(interact="Tukey"))
5562 summary(tuk3x)
5563 Props<-Pollen$Tube/Pollen$Count
5564 Pollen<-data.frame(Pollen,Props)
5565 str(Pollen)
5566 barcentres<-
5567 barplot(tapply(Pollen$Props,Pollen$interact,mean),ylim=c(0,max(tapply(Pollen$Props,Pollen$interact,mean)+
5568 0.1)), ylab = "Proportion of Pollen Tube Number/Pollen Count", xlab = "Treatment and Species", main =
5569 "Proportion of Pollen Tube per species and treatment")

```

```

5570 means<-tapply(Pollen$Props,Pollen$interact,mean)
5571 ses<-tapply(Pollen$Props,Pollen$interact,sd)/sqrt(tapply(Pollen$Props,Pollen$interact,length))
5572 arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
5573
5574
5575
5576 #Analyse species separately for comparison:
5577 #bienne:
5578 library(readr)
5579 bienne_all <- read_csv("bienne_all.csv",
5580                       col_types = cols(Count = col_number(),
5581                                         Tube = col_number(), Percent = col_number(),
5582                                         Reached = col_number(), Lat = col_number(),
5583                                         pc1 = col_number()))
5584 View(bienne_all)
5585 bienne_all$Treatment<-as.factor(bienne_all$Treatment)
5586
5587 Propsbienne<-bienne_all$Tube/bienne_all$Count
5588 Propbienne<-cbind(bienne_all$Tube, bienne_all$Count)
5589 modpropbienne<-glm(Propbienne~Treatment, family=quasibinomial(link = 'logit'), data=bienne_all)
5590 summary(modpropbienne)
5591 tukpropbienne<-glht(modpropbienne,mcp(Treatment="Tukey"))
5592 summary(tukpropbienne)
5593 Propsb<-bienne_all$Tube/bienne_all$Count
5594 bienne_all<-data.frame(bienne_all,Propsb)
5595 str(bienne_all)
5596 barcentres<-
5597 barplot(tapply(bienne_all$Propsb,bienne_all$Treatment,mean),ylim=c(0,max(tapply(bienne_all$Propsb,bienn

```



```

5598 e_all$Treatment,mean)+0.1)), ylab = "Proportion of Pollen Tube Number", xlab = "Treatment", main =
5599 "Proportion of Pollen Tube vs Treatment for L.bienne")
5600 means<-tapply(bienne_all$Propsb,bienne_all$Treatment,mean)
5601 ses<-
5602 tapply(bienne_all$Propsb,bienne_all$Treatment,sd)/sqrt(tapply(bienne_all$Propsb,bienne_all$Treatment,len
5603 gth))
5604 arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
5605
5606 #usitatissimum:
5607 library(readr)
5608 usitatissimum_all <- read_csv("usitatissimum_all.csv",
5609                               col_types = cols(Count = col_number(),
5610                                                  Tube = col_number(), Percent = col_number(),
5611                                                  Reached = col_number(), Lat = col_number(),
5612                                                  pc1 = col_number()))
5613 View(usitatissimum_all)
5614 usitatissimum_all$Treatment<-as.factor(usitatissimum_all$Treatment)
5615
5616 Propscul<-usitatissimum_all$Tube/usitatissimum_all$Count
5617 Propcul<-cbind(usitatissimum_all$Tube, usitatissimum_all$Count)
5618 modpropcul<-glm(Propcul~Treatment, family=quasibinomial(link = 'logit'), data=usitatissimum_all)
5619 summary(modpropcul)
5620 tukpropcul<-glht(modpropcul,mcp(Treatment="Tukey"))
5621 summary(tukpropcul)
5622 Propsc<-usitatissimum_all$Tube/usitatissimum_all$Count
5623 usitatissimum_all<-data.frame(usitatissimum_all,Propsc)
5624 str(usitatissimum_all)
5625 barcentres<-
5626 barplot(tapply(usitatissimum_all$Propsc,usitatissimum_all$Treatment,mean),ylim=c(0,max(tapply(usitatissim

```

```

5627 um_all$Propsc,usitatissimum_all$Treatment,mean)+0.1)), ylab = "Proportion of Pollen Tube Number", xlab =
5628 "Treatment", main = "Proportion of Pollen Tube vs Treatment for L.usitatissimum")
5629 means<-tapply(usitatissimum_all$Propsc,usitatissimum_all$Treatment,mean)
5630 ses<-
5631 tapply(usitatissimum_all$Propsc,usitatissimum_all$Treatment,sd)/sqrt(tapply(usitatissimum_all$Propsc,usitati
5632 ssimum_all$Treatment,length))
5633 arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
5634
5635
5636 ###test reached versus treatment x species with mixed model, ind as random effect
5637 library(lmerTest)
5638 library("MuMIn")
5639 mod4<-glmer(Reached~Treatment*Species+(1|Ind), family=binomial(link='logit'), data=Pollen)
5640 summary(mod4)
5641 anova(mod4,test="F")
5642 r.squaredGLMM(mod4)
5643 mod4x<-glm(Reached~Treatment*Species, family=binomial(link='logit'), data=Pollen)
5644 anova(mod4, mod4x, test="LRT") #likelihood ratio test
5645 library(multcomp)
5646 tuk4<-glht(mod4,mcp(Treatment="Tukey"))
5647 summary(tuk4)
5648 interact<-interaction(Pollen$Treatment,Pollen$Species)
5649 Pollen<-data.frame(Pollen,interact)
5650 mod4y<-glmer(Reached~-1+interact+(1|Ind), family=binomial(link='logit'), data=Pollen)
5651 tuk4y<-glht(mod4y,mcp(interact="Tukey"))
5652 summary(tuk4y)
5653 barcentres<-
5654 barplot(tapply(Pollen$Reached,Pollen$interact,mean),ylim=c(0,max(tapply(Pollen$Reached,Pollen$interact,m
5655 ean)+0.1)),ylab = "Proportion of Pollen Tube Reached", xlab = "Treatment and Species", main = "Proportion of
5656 Pollen Tube Reaching Ovaries For Each Treatment and Species")

```

```

5657 means<-tapply(Pollen$Reached,Pollen$interact,mean)
5658 ses<-tapply(Pollen$Reached,Pollen$interact,sd)/sqrt(tapply(Pollen$Reached,Pollen$interact,length))
5659 arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
5660 write.table(tuk4y)
5661
5662 #Pollen Reached for bienne:
5663 library(readr)
5664 bienne_all <- read_csv("bienne_all.csv",
5665                       col_types = cols(Count = col_number(),
5666                                         Tube = col_number(), Percent = col_number(),
5667                                         Reached = col_number(), Lat = col_number(),
5668                                         pc1 = col_number()))
5669 View(bienne_all)
5670 bienne_all$Treatment<-as.factor(bienne_all$Treatment)
5671
5672 modbreach<-glm(Reached~Treatment,family=binomial(link='logit'),data=bienne_all)
5673 summary(modbreach)
5674 library(multcomp)
5675 tukbreach<-glht(modbreach,mcp(Treatment="Tukey"))
5676 summary(tukbreach)
5677 barplot(tapply(bienne_all$Reached,bienne_all$Treatment,mean))
5678 barcentres<-
5679 barplot(tapply(bienne_all$Reached,bienne_all$Treatment,mean),ylim=c(0,max(tapply(bienne_all$Reached,bie
5680 nne_all$Treatment,mean)+0.1)), ylab = "Proportion of Pollen Reaching Ovary", xlab = "Treatment", main =
5681 "Proportion of Pollen Reaching Ovary vs Treatment for L.bienne")
5682 means<-tapply(bienne_all$Reached,bienne_all$Treatment,mean)
5683 ses<-
5684 tapply(bienne_all$Reached,bienne_all$Treatment,sd)/sqrt(tapply(bienne_all$Reached,bienne_all$Treatment,l
5685 ength))
5686 arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)

```

```

5687
5688 #Pollen Reached for usitatissimum:
5689 library(readr)
5690 usitatissimum_all <- read_csv("usitatissimum_all.csv",
5691                               col_types = cols(Count = col_number(),
5692                                                Tube = col_number(), Percent = col_number(),
5693                                                Reached = col_number(), Lat = col_number(),
5694                                                pc1 = col_number()))
5695 View(usitatissimum_all)
5696 usitatissimum_all$Treatment<-as.factor(usitatissimum_all$Treatment)
5697
5698 modcreach<-glm(Reached~Treatment,family=binomial(link='logit'),data=bienne_all)
5699 summary(modcreach)
5700 library(multcomp)
5701 tukcreach<-glht(modcreach,mcp(Treatment="Tukey"))
5702 summary(tukcreach)
5703 barplot(tapply(usitatissimum_all$Reached,usitatissimum_all$Treatment,mean))
5704 barcentres<-
5705 barplot(tapply(usitatissimum_all$Reached,usitatissimum_all$Treatment,mean),ylim=c(0,max(tapply(usitatissi
5706 mum_all$Reached,usitatissimum_all$Treatment,mean)+0.1)), ylab = "Proportion of Pollen Reaching Ovary",
5707 xlab = "Treatment", main = "Proportion of Pollen Reaching Ovary vs Treatment for L.usitatissimum")
5708 means<-tapply(usitatissimum_all$Reached,usitatissimum_all$Treatment,mean)
5709 ses<-
5710 tapply(usitatissimum_all$Reached,usitatissimum_all$Treatment,sd)/sqrt(tapply(usitatissimum_all$Reached,us
5711 itatissimum_all$Treatment,length))
5712 arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
5713
5714
5715 #test proportion tubes versus treatment x species with mixed model, ind as random effect
5716 Prop<-cbind(Pollen$Tube, Pollen$Count)

```

```

5717 mod5<-glmer(Prop~Treatment*Species+(1|Ind), family=quasibinomial(link='logit'), data=Pollen)
5718 summary(mod5)
5719 tuk5<-glht(mod5,mcp(Treatment="Tukey"))
5720 summary(tuk5)
5721 interact<-interaction(Pollen$Treatment,Pollen$Species)
5722 mod5y<-glmer(Prop~-1+interact+(1|Ind), family=binomial(link='logit'), data=Pollen)
5723 summary(mod5y)
5724 tuk5y<-glht(mod5y,mcp(interact="Tukey"))
5725 summary(tuk5y)
5726 Props<-Pollen$Tube/Pollen$Count
5727 Pollen<-data.frame(Pollen,Props)
5728 str(Pollen)
5729 barcentres<-
5730 barplot(tapply(Pollen$Props,Pollen$interact,mean),ylim=c(0,max(tapply(Pollen$Props,Pollen$interact,mean)+
5731 0.1)))
5732 means<-tapply(Pollen$Props,Pollen$interact,mean)
5733 ses<-tapply(Pollen$Props,Pollen$interact,sd)/sqrt(tapply(Pollen$Props,Pollen$interact,length))
5734 arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
5735
5736 Appendix 15 : Adapter sequences used for the ddRADSeq process
5737 Adapter_1 = AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCT --
5738 Adapter_2 = CAAGCAGAAGACGGCATAACGAGATCGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCT
5739

```

5740 **Appendix 16 : P1 and P2 Barcode sequences for each samples**

| 5741 | P1 | P2 | Sample |
|------|--------|--------|------------|
| 5742 | gcatga | tcatgc | Sample_18 |
| 5743 | gcatga | cggatc | Sample_19 |
| 5744 | gcatga | taagac | Sample_20 |
| 5745 | gcatga | actccg | Sample_49 |
| 5746 | gcatga | aacgtg | Sample_50 |
| 5747 | gcatga | ctgtat | Sample_69 |
| 5748 | gcatga | gtaaca | Sample_70 |
| 5749 | gcatga | gagcgt | Sample_71 |
| 5750 | gcatga | tgccaa | Sample_106 |
| 5751 | gcatga | gttact | Sample_83 |
| 5752 | gcatga | cgcttg | Sample_21 |
| 5753 | gcatga | tagcat | Sample_22 |
| 5754 | gatccg | tcatgc | Sample_23 |
| 5755 | gatccg | cggatc | Sample_74 |
| 5756 | gatccg | taagac | Sample_84 |
| 5757 | gatccg | actccg | Sample_85 |
| 5758 | gatccg | aacgtg | Sample_92 |
| 5759 | gatccg | ctgtat | Sample_93 |
| 5760 | gatccg | gtaaca | Sample_107 |
| 5761 | gatccg | gagcgt | Sample_110 |
| 5762 | gatccg | tgccaa | Sample_111 |
| 5763 | gatccg | tagcat | Sample_87 |
| 5764 | gatccg | cgcttg | Sample_86 |
| 5765 | gtctta | tcatgc | Sample_88 |
| 5766 | gtctta | cggatc | Sample_89 |

5767 gtctta taagac Sample_90
5768 gtctta actccg Sample_91
5769 gtctta aacgtg Sample_104
5770 gtctta ctgtat Sample_105
5771 gtctta gagcgt Sample_108
5772 gtctta tgccaa Sample_109
5773 gtctta cgcttg Sample_54
5774 gtctta tagcat Sample_55
5775 cggagt tcatgc Sample_56
5776 cggagt cggatc Sample_57
5777 cggagt taagac Sample_40
5778 cggagt actccg Sample_45
5779 cggagt aacgtg Sample_46
5780 cggagt ctgtat Sample_47
5781 cggagt gtaaca Sample_48
5782 cggagt gagcgt Sample_52
5783 cggagt tgccaa Sample_61
5784 cggagt gttact Sample_77
5785 cggagt cgcttg Sample_81
5786 cggagt tagcat Sample_62
5787 cacgtt tcatgc Sample_63
5788 cacgtt cggatc Sample_64
5789 cacgtt taagac Sample_65
5790 cacgtt aacgtg Sample_67
5791 cacgtt ctgtat Sample_51
5792 cacgtt gtaaca Sample_80
5793 cacgtt gagcgt Sample_94

| | | | |
|------|--------|--------|------------|
| 5794 | cacgtt | tgccaa | Sample_95 |
| 5795 | cacgtt | gttact | Sample_96 |
| 5796 | cacgtt | cgcttg | Sample_103 |
| 5797 | cacgtt | tagcat | Sample_112 |
| 5798 | atacag | tcatgc | Sample_113 |
| 5799 | atacag | cggatc | Sample_42 |
| 5800 | atacag | taagac | Sample_25 |
| 5801 | atacag | actccg | Sample_26 |
| 5802 | atacag | aacgtg | Sample_27 |
| 5803 | atacag | ctgtat | Sample_28 |
| 5804 | atacag | gtaaca | Sample_29 |
| 5805 | atacag | gagcgt | Sample_30 |
| 5806 | atacag | tgccaa | Sample_31 |
| 5807 | atacag | gttact | Sample_75 |
| 5808 | atacag | cgcttg | Sample_100 |
| 5809 | atacag | tagcat | Sample_101 |
| 5810 | tgttac | cggatc | Sample_114 |
| 5811 | tgttac | taagac | Sample_115 |
| 5812 | tgttac | actccg | Sample_24 |
| 5813 | tgttac | aacgtg | Sample_43 |
| 5814 | tgttac | ctgtat | Sample_82 |
| 5815 | tgttac | gtaaca | Sample_58 |
| 5816 | tgttac | gagcgt | Sample_60 |
| 5817 | tgttac | tgccaa | Sample_44 |
| 5818 | tgttac | gttact | Sample_12 |
| 5819 | tgttac | cgcttg | Sample_13 |
| 5820 | tgttac | tagcat | Sample_14 |

5821 acgctc tcatgc Sample_15
5822 acgctc cggatc Sample_16
5823 acgctc taagac Sample_17
5824 acgctc actccg Sample_73
5825 acgctc aacgtg Sample_41
5826 acgctc ctgtat Sample_33
5827 acgctc gtaaca Sample_34
5828 acgctc gagcgt Sample_35
5829 acgctc tgccaa Sample_36
5830 acgctc gttact Sample_37
5831 acgctc cgcttg Sample_38
5832 acgctc tagcat Sample_39
5833 ttggca cggatc Sample_79
5834 ttggca aacgtg Sample_99
5835 ttggca ctgtat Sample_59
5836 ttggca gtaaca Sample_6
5837 ttggca gagcgt Sample_7
5838 ttggca tgccaa Sample_8
5839 ttggca gttact Sample_9
5840 ttggca cgcttg Sample_10
5841 ttggca tagcat Sample_11
5842 agtaac tcatgc Sample_116
5843 agtaac cggatc Sample_117
5844 agtaac taagac Sample_118
5845 agtaac actccg Sample_119
5846 agtaac aacgtg Sample_120
5847 agtaac ctgtat Sample_1

5848 agtaac gtaaca Sample_2
5849 agtaac gagcgt Sample_3
5850 agtaac tgccaa Sample_4
5851 agtaac gttact Sample_5
5852 agtaac cgcttg Sample_32
5853 agtaac tagcat Sample_76
5854 caagcg tcatgc Sample_121
5855 caagcg cggatc Sample_122
5856 caagcg taagac Sample_123
5857 caagcg actccg Sample_124
5858 caagcg aacgtg Sample_125
5859 caagcg ctgtat Sample_126
5860 caagcg gtaaca Sample_150
5861 caagcg gagcgt Sample_151
5862 caagcg tgccaa Sample_152

5863

5864 **Appendix 17: Process radtags command**

5865 process_radtags -P -1 /nobackup/mnkz72/raw/merged_rad_raw_R1.fastq.gz -2
5866 /nobackup/mnkz72/raw/merged_rad_raw_R2.fastq.gz -b /nobackup/mnkz72/barcodes/index_p1p2new.txt -o
5867 /nobackup/mnkz72/raw/demultiplexed -r -c -q --barcode_dist_2 2 --disable_rad_check --inline_inline --renz_1
5868 pstl --renz_2 msel --adapter_1 AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTCTTCCGATCT -
5869 -adapter_2 CAAGCAGAAGACGGCATACGAGATCGGTCTCGGCATTCCTGCTGAACCGCTCTTCCGATCT

5870

5871 **Appendix 18: BWA-MEM Command**

5872 bwa mem -t16 /nobackup/mnkz72/1st_genome_asm/cultivar/GCA_000224295.2_ASM22429v2_genomic.fna
5873 /nobackup/mnkz72/raw/demultiplexed_run2/clean/Sample_1.1.fq.gz
5874 /nobackup/mnkz72/raw/demultiplexed_run2/clean/Sample_1.1.fq.gz | samtools view -b -h | samtools sort -
5875 @12 -o /nobackup/mnkz72/raw/aligned_run2/Sample_1.bam

5876

5877 **Appendix 19: Population list for popmap file**

| | | | |
|------|------------|-----|-----|
| 5878 | Sample_1 | Vil | FRA |
| 5879 | Sample_10 | Tym | UKN |
| 5880 | Sample_100 | Mat | FRA |
| 5881 | Sample_101 | Mat | FRA |
| 5882 | Sample_103 | Lla | SNO |
| 5883 | Sample_104 | 15 | SNO |
| 5884 | Sample_105 | 15 | SNO |
| 5885 | Sample_106 | 15 | SNO |
| 5886 | Sample_107 | 12 | MED |
| 5887 | Sample_108 | 15 | SNO |
| 5888 | Sample_109 | 15 | SNO |
| 5889 | Sample_11 | Tym | UKN |
| 5890 | Sample_110 | 12 | MED |
| 5891 | Sample_111 | 12 | MED |
| 5892 | Sample_112 | Lla | SNO |
| 5893 | Sample_113 | Lla | SNO |
| 5894 | Sample_114 | Mat | FRA |
| 5895 | Sample_115 | Mat | FRA |
| 5896 | Sample_116 | Tym | UKN |
| 5897 | Sample_117 | Tym | UKN |
| 5898 | Sample_118 | Tym | UKN |
| 5899 | Sample_119 | Tym | UKN |
| 5900 | Sample_12 | Sut | UKN |
| 5901 | Sample_120 | Tym | UKN |
| 5902 | Sample_121 | Vil | FRA |
| 5903 | Sample_122 | Vil | FRA |

| | | | |
|------|------------|-----|-----|
| 5904 | Sample_123 | Vil | FRA |
| 5905 | Sample_124 | Vil | FRA |
| 5906 | Sample_125 | Vil | FRA |
| 5907 | Sample_126 | Vil | FRA |
| 5908 | Sample_13 | Sut | UKN |
| 5909 | Sample_14 | Sut | UKN |
| 5910 | Sample_15 | Sut | UKN |
| 5911 | Sample_16 | Sut | UKN |
| 5912 | Sample_17 | Sut | UKN |
| 5913 | Sample_18 | 1 | SSA |
| 5914 | Sample_19 | 1 | SSA |
| 5915 | Sample_2 | Vil | FRA |
| 5916 | Sample_20 | 1 | SSA |
| 5917 | Sample_21 | 6 | SSB |
| 5918 | Sample_22 | Ome | CUL |
| 5919 | Sample_23 | 6 | SSB |
| 5920 | Sample_24 | Mon | CUL |
| 5921 | Sample_25 | Mat | FRA |
| 5922 | Sample_26 | Mat | FRA |
| 5923 | Sample_27 | Mat | FRA |
| 5924 | Sample_28 | Mat | FRA |
| 5925 | Sample_29 | Mat | FRA |
| 5926 | Sample_3 | Vil | FRA |
| 5927 | Sample_30 | Mat | FRA |
| 5928 | Sample_31 | Mat | FRA |
| 5929 | Sample_32 | Vil | FRA |
| 5930 | Sample_33 | Tal | FRA |

| | | | |
|------|-----------|-----|-----|
| 5931 | Sample_34 | Tal | FRA |
| 5932 | Sample_35 | Tal | FRA |
| 5933 | Sample_36 | Tal | FRA |
| 5934 | Sample_37 | Tal | FRA |
| 5935 | Sample_38 | Tal | FRA |
| 5936 | Sample_39 | Tal | FRA |
| 5937 | Sample_40 | Bol | CUL |
| 5938 | Sample_41 | Suz | CUL |
| 5939 | Sample_42 | Mar | CUL |
| 5940 | Sample_43 | Olg | CUL |
| 5941 | Sample_44 | Rab | CUL |
| 5942 | Sample_45 | Dor | UKS |
| 5943 | Sample_46 | Dor | UKS |
| 5944 | Sample_47 | Dor | UKS |
| 5945 | Sample_48 | Dor | UKS |
| 5946 | Sample_49 | 3 | SSA |
| 5947 | Sample_5 | Vil | FRA |
| 5948 | Sample_50 | 3 | SSA |
| 5949 | Sample_51 | Lir | CUL |
| 5950 | Sample_52 | Gis | CUL |
| 5951 | Sample_54 | 19 | SNO |
| 5952 | Sample_55 | 19 | SNO |
| 5953 | Sample_56 | 19 | SNO |
| 5954 | Sample_57 | 19 | SNO |
| 5955 | Sample_59 | Tin | CUL |
| 5956 | Sample_6 | Tym | UKN |
| 5957 | Sample_60 | Pri | CUL |

| | | | |
|------|-----------|------|-----|
| 5958 | Sample_61 | low2 | UKS |
| 5959 | Sample_62 | low2 | UKS |
| 5960 | Sample_63 | low2 | UKS |
| 5961 | Sample_64 | low2 | UKS |
| 5962 | Sample_65 | low2 | UKS |
| 5963 | Sample_67 | low2 | UKS |
| 5964 | Sample_7 | Tym | UKN |
| 5965 | Sample_70 | 3 | SSA |
| 5966 | Sample_71 | 3 | SSA |
| 5967 | Sample_73 | Sut | UKN |
| 5968 | Sample_74 | 10 | SSB |
| 5969 | Sample_75 | Mat | FRA |
| 5970 | Sample_76 | Vil | FRA |
| 5971 | Sample_77 | low2 | UKS |
| 5972 | Sample_79 | Tal | FRA |
| 5973 | Sample_8 | Tym | UKN |
| 5974 | Sample_80 | Lla | SNO |
| 5975 | Sample_81 | low2 | UKS |
| 5976 | Sample_82 | Olg | CUL |
| 5977 | Sample_83 | 3 | SSA |
| 5978 | Sample_84 | 10 | SSB |
| 5979 | Sample_85 | 10 | SSB |
| 5980 | Sample_86 | 13 | SNO |
| 5981 | Sample_87 | 13 | SNO |
| 5982 | Sample_88 | 13 | SNO |
| 5983 | Sample_89 | 13 | SNO |
| 5984 | Sample_9 | Tym | UKN |

5985 Sample_90 13 SNO
5986 Sample_91 13 SNO
5987 Sample_92 10 SSB
5988 Sample_93 3 SSB
5989 Sample_94 Lla SNO
5990 Sample_95 Lla SNO
5991 Sample_96 Lla SNO
5992 Sample_99 Tal FRA
5993 Sample_97 Tal FRA
5994 Sample_98 Tal FRA

5995

5996 **Appendix 20: ref_map.pl pipeline command**

5997 ref_map.pl -T 16 -r 0.8 -X populations:--fstats --structure --plink --vcf --samples
5998 /nobackup/mnkz72/raw/aligned_samples --popmap
5999 /nobackup/mnkz72/popmap/popmap/popmap_new_0322.txt -o /nobackup/mnkz72/new_refmap/run_2

6000

6001 **Appendix 21: PLINK command**

6002 module load bioinformatics
6003 module load plink
6004 #Change path to workind directory:
6005 cd /nobackup/mnkz72/new_refmap/rdir/run3
6006 #Allow for extra chromosomes
6007 plink --file populations.plink --out run3_pops --recodeA --allow-extra-chr --noweb
6008 #Make BED:
6009 plink --file populations.plink --out run3_pops --make-bed --allow-extra-chr --noweb
6010

6011 **Appendix 22: Geographical information of the different population**

| sample | longitude | latitude |
|---------|-----------|----------|
| 1_1 | -5.39258 | 36.80044 |
| 1_26 | -5.39258 | 36.80044 |
| 1_6 | -5.39258 | 36.80044 |
| 10_15 | -6.61781 | 37.88211 |
| 10_26 | -6.61781 | 37.88211 |
| 10_27 | -6.61781 | 37.88211 |
| 10_30 | -6.61781 | 37.88211 |
| 12_10 | 3.151889 | 42.31008 |
| 12_20 | 3.151889 | 42.31008 |
| 12_8 | 3.151889 | 42.31008 |
| 13_1 | -3.24035 | 43.02902 |
| 13_12 | -3.24035 | 43.02902 |
| 13_19 | -3.24035 | 43.02902 |
| 13_22 | -3.24035 | 43.02902 |
| 13_23 | -3.24035 | 43.02902 |
| 3_11 | -5.55589 | 36.03633 |
| 3_13 | -5.55589 | 36.03633 |
| 3_14 | -5.55589 | 36.03633 |
| 3_15 | -5.55589 | 36.03633 |
| Gis | 12.56738 | 41.87194 |
| low2_10 | -1.07492 | 50.68183 |
| low2_13 | -1.07492 | 50.68183 |
| low2_14 | -1.07492 | 50.68183 |
| low2_2 | -1.07492 | 50.68183 |
| low2_25 | -1.07492 | 50.68183 |
| low2_26 | -1.07492 | 50.68183 |
| low2_7 | -1.07492 | 50.68183 |
| low2_9 | -1.07492 | 50.68183 |
| Lir | -3.43597 | 55.37805 |
| Lla_17 | -4.68753 | 43.40738 |
| Lla_20 | -4.68753 | 43.40738 |
| Lla_23 | -4.68753 | 43.40738 |
| Lla_33 | -4.68753 | 43.40738 |
| Lla_A | -4.68753 | 43.40738 |
| Lla_B | -4.68753 | 43.40738 |
| Lla_C | -4.68753 | 43.40738 |
| Suz | 5.291266 | 52.13263 |
| Mat_12 | 4.458631 | 48.35697 |
| Mat_13 | 4.458631 | 48.35697 |
| Mat_2 | 4.458631 | 48.35697 |
| Mat_20 | 4.458631 | 48.35697 |
| Mat_22 | 4.458631 | 48.35697 |
| Mat_24 | 4.458631 | 48.35697 |

| | | |
|---------|----------|----------|
| Mat_25 | 4.458631 | 48.35697 |
| Mat_4 | 4.458631 | 48.35697 |
| Mat_7 | 4.458631 | 48.35697 |
| Mon | -3.43597 | 55.37805 |
| Olg | 2.213749 | 46.22764 |
| Olg_1 | 2.213749 | 46.22764 |
| Ome | 2.213749 | 46.22764 |
| Pri | 12.56738 | 41.87194 |
| Rab | -7.09262 | 31.7917 |
| Sut_29 | -0.95927 | 53.35291 |
| Sut_33 | -0.95927 | 53.35291 |
| Sut_34 | -0.95927 | 53.35291 |
| Sut_4 | -0.95927 | 53.35291 |
| Sut_41 | -0.95927 | 53.35291 |
| Tal_10 | -3.45465 | 47.6997 |
| Tal_11 | -3.45465 | 47.6997 |
| Tal_14 | -3.45465 | 47.6997 |
| Tal_28 | -3.45465 | 47.6997 |
| Tal_3 | -3.45465 | 47.6997 |
| Tal_33 | -3.45465 | 47.6997 |
| Tal_34 | -3.45465 | 47.6997 |
| Tal_38 | -3.45465 | 47.6997 |
| Tal_40 | -3.45465 | 47.6997 |
| Vil_1 | -1.05034 | 45.09393 |
| Vil_15 | -1.05034 | 45.09393 |
| Vil_2 | -1.05034 | 45.09393 |
| Vil_21 | -1.05034 | 45.09393 |
| Vil_25p | -1.05034 | 45.09393 |
| Vil_27 | -1.05034 | 45.09393 |
| Vil_32 | -1.05034 | 45.09393 |
| Vil_35 | -1.05034 | 45.09393 |
| Vil_36g | -1.05034 | 45.09393 |
| Vil_36p | -1.05034 | 45.09393 |
| Vil_10 | -1.05034 | 45.09393 |
| Tym_16 | -3.55328 | 53.30307 |
| Tym_19 | -3.55328 | 53.30307 |
| Mat_17 | 4.458631 | 48.35697 |
| Tym_3 | -3.55328 | 53.30307 |
| Tym_26g | -3.55328 | 53.30307 |
| Tym_26p | -3.55328 | 53.30307 |
| Tym_41 | -3.55328 | 53.30307 |
| Sut_17 | -0.95927 | 53.35291 |
| Sut_36 | -0.95927 | 53.35291 |
| 6_29 | -5.71117 | 37.93551 |
| 6_1 | -5.71117 | 37.93551 |
| Mat_11 | 4.458631 | 48.35697 |

| | | |
|--------|----------|----------|
| Dor_D | -2.01 | 50.6 |
| Dor_C | -2.01 | 50.6 |
| Dor_B | -2.01 | 50.6 |
| Dor_A | -2.01 | 50.6 |
| Tin | 5.291266 | 52.13263 |
| Tym_40 | -3.55328 | 53.30307 |
| Tym_5 | -3.55328 | 53.30307 |
| 3_27 | -5.55589 | 36.03633 |
| Tym_20 | -3.55328 | 53.30307 |
| 13_8 | -3.24035 | 43.02902 |
| Tym_30 | -3.55328 | 53.30307 |

6012

6013

6014 **Appendix 23: Command line to filter for heterozygosity**

6015 #Prepare and format file in bcftools

6016 #First, convert multiallelic sites to biallelic using bcftools:

6017 bcftools norm -m - "/nobackup/mnkz72/new_refmap/bcf_out/populations.snps.vcf" >

6018 allsamples_biallelic_convert.vcf

6019 #Filter the alternative alleles under certain value

6020 bcftools view -e "FORMAT/AD[:1]<2 && INFO/AD[1]<5"

6021 "/nobackup/mnkz72/new_refmap/bcf_out/allele_filter/allsamples_biallelic_convert.vcf" > biallelic-filtered.vcf

6022 #Compress vcf output with bgzip (Samtools):

6023 bgzip -@8 "/nobackup/mnkz72/new_refmap/bcf_out/allele_filter/allsamples_biallelic_convert.vcf"

6024 bgzip -@8 "/nobackup/mnkz72/new_refmap/bcf_out/populations.snps.vcf"

6025 #Produce vcf file statistics for files:

6026 bcftools stats "/nobackup/mnkz72/new_refmap/bcf_out/populations.snps.vcf.gz" > run3_stats.txt

6027 bcftools stats "/nobackup/mnkz72/new_refmap/bcf_out/allele_filter/allsamples_biallelic_convert.vcf.gz" >

6028 biallelic_convert_stats.txt

6029 #het-filtered vcf:

6030 bcftools stats "/nobackup/mnkz72/new_refmap/bcf_out/allpopulations.hetfiltered.vcf.gz" >

6031 hetfiltered_stats.txt

6032 bcftools stats "/nobackup/mnkz72/new_refmap/bcf_out/allsamples.hetfiltered.vcf.gz" > het_bigfile_stats.txt

6033 #Extract Genotypes per sample:

```

6034 bcftools query -f '%CHROM %POS[\t%GT]\n' "/nobackup/mnkz72/new_refmap/bcf_out/allele_filter/biallelic-
6035 filtered.vcf" > genotypesbysample_run3.txt
6036 head -n 1000 "/nobackup/mnkz72/new_refmap/bcf_out/allele_filter/genotypesbysample_run3.txt" | cut -c 1-
6037 1000 "/nobackup/mnkz72/new_refmap/bcf_out/allele_filter/genotypesbysample_run3.txt" >
6038 genotypesbysample_cut_1000.txt
6039 bcftools query -f '%CHROM %POS[\t%GT]\n'
6040 "/nobackup/mnkz72/new_refmap/vcftools/allele_filter/biallelic_filtered_populations_snps.vcf" >
6041 genotypesbysample_biallelic.txt
6042 head -n 1000 "/nobackup/mnkz72/new_refmap/bcf_out/allele_filter/genotypesbysample_run3.txt" | cut -c 1-
6043 1000 "/nobackup/mnkz72/new_refmap/bcf_out/allele_filter/genotypesbysample_run3.txt" >
6044 genotypesbysample_run3_cut_1000.txt
6045 #score hets as 1, homs as 0
6046 sumx<-rep(NA,nrow(genosx))
6047 sumy<-NULL
6048 for (i in c(1:ncol(genosx))){
6049   for (j in c(1:nrow(genosx))){
6050     if(genosx[j,i]=="1/0" | genosx[j,i]=="0/1"){
6051       sumx[j]<-1
6052     } else if (genosx[j,i]=="0/0" | genosx[j,i]=="1/1"){
6053       sumx[j]<-0
6054     }
6055   }
6056   sumy<-cbind(sumy,sumx)
6057   sumx<-rep(NA,nrow(genosx))
6058 }
6059 sumy
6060
6061 plot(c(1:nrow(sumy)),sumy[,3])
6062
6063 #proportion of hets per ind moving average

```

```

6064 sumz<-NULL
6065 mod<-50 # number of SNPs for moving average
6066 means<-NULL
6067 sumzz<-NULL
6068 inds<-NULL
6069 for (k in c(1:ncol(sumy))){
6070   for (l in c(1:nrow(sumy))){
6071     if (l %% mod == 0){
6072       meanx<-sum(sumy[c((l-mod):l),k],na.rm=T)/mod #mean over window size including missing data
6073       means<-c(means,meanx)
6074     }
6075   }
6076   sumz<-cbind(sumz,means)
6077   ind<-rep(k,l %% mod)
6078   inds<-c(inds,ind)
6079   sumzz<-c(sumzz,means)
6080   means<-NULL
6081 }
6082 sumz
6083 sumzzz<-data.frame(inds,sumzz)
6084 head(sumzzz)
6085
6086 boxplot(sumzzz$sumzz~sumzzz$inds,xlab="sample",ylab="prop het")
6087 #view individuals of interest from boxplot to find regions of high het (replace number in sumz[,])
6088 plot(c(1:nrow(sumz)),sumz[,15],type="l",xlab="SNP_position",ylab="prop het")
6089 #view genos in region of interest (SNP_position*mod)
6090 sumy[(13*mod):(14*mod),15]
6091
6092 #plot all inds

```

```

6093 plot(c(1:nrow(sumz)),rep(max(sumz,na.rm=T),nrow(sumz)),ylim=c(-
6094 0.05,max(sumz,na.rm=T)),type="l",col="white",xlab="SNP_position",ylab="prop het")
6095 for (m in c(1:ncol(sumz))) {
6096   lines(c(1:nrow(sumz)),sumz[,m],type="l")
6097 }
6098
6099 #proportion of hets per location moving average
6100 pop<-c(1:19)# list of inds to include (same population), numbers are order in the sample list
6101 lenp<-length(pop)
6102 sump<-sumy[,pop]
6103 meanp<-NULL
6104 for (m in c(1:nrow(sump))) {
6105   meany<-sum(sump[m,],na.rm=T)/lenp #mean over population including missing data
6106   meanp<-c(meanp,meany)
6107 }
6108 meanp
6109
6110 plot(meanp,type="l",xlab="SNP_position",ylab="prop het")
6111
6112 #find het loci
6113 hetlocs<-which(meanp>quantile(meanp,0.99))
6114 hetvals<-meanp[meanp>quantile(meanp,0.99)]
6115 hetgenos<-data.frame(hetlocs,hetvals,sump[hetlocs,])
6116 hetgenos
6117
6118 #test with two pops
6119 #proportion of hets per location moving average
6120 pop<-list(c(1:8),c(9:19))# list of inds to include (same population), numbers are order in the sample list
6121 meanp<-NULL

```

```

6122  meanps<-NULL
6123  for (n in 1:length(pop)){
6124    lenp<-length(pop[[n]])
6125    sump<-sumy[,pop[[n]]]
6126    for (m in c(1:nrow(sump))){
6127      meany<-sum(sump[m,],na.rm=T)/lenp #mean over population including missing data
6128      meanp<-c(meanp,meany)
6129    }
6130    meanps<-cbind(meanps,meanp)
6131    meanp<-NULL
6132  }
6133  meanps
6134
6135  plot(meanps[,1],type="p",pch=1,xlab="SNP_position",ylab="prop het")
6136  for (p in 2:ncol(meanps)){
6137    points(meanps[,p],pch=p)
6138  }
6139
6140  #find het loci (top 99% hets)
6141  hetlocs<-NULL
6142  for (q in 1:length(pop)){
6143    hetloc<-which(meanps[,q]>quantile(meanps,0.99))
6144    hetlocs<-sort(union(hetlocs, hetloc))
6145  }
6146  hetvals<-meanps[hetlocs,]
6147  hetgenos<-data.frame(hetlocs,hetvals)
6148  hetgenos
6149
6150  Appendix 24: Command line to filter by loci location in vcftools

```

```

6151 #vcftools:
6152 #frequency analysis
6153 vcftools --gzvcf /nobackup/mnkz72/new_refmap/vcftools/run3/populations.snps.vcf.gz --freq --out
6154 run3_populations_freqanalysis
6155 #depth:
6156 vcftools --gzvcf /nobackup/mnkz72/new_refmap/vcftools/run3/populations.snps.vcf.gz --depth --out
6157 run3_populations_depthanalysis
6158 #allele counts:
6159 vcftools --gzvcf /nobackup/mnkz72/new_refmap/vcftools/run3/populations.snps.vcf.gz --counts --out
6160 run3_populations_countsanalysis
6161 #heterozygosity:
6162 vcftools --gzvcf /nobackup/mnkz72/new_refmap/vcftools/run3/populations.snps.vcf.gz --het --out
6163 run3_populations_hetanalysis
6164 #site-quality:
6165 vcftools --gzvcf /nobackup/mnkz72/new_refmap/vcftools/run3/populations.snps.vcf.gz --site-quality --out
6166 run3_populations_sitequalityanalysis
6167 #Allele-het-depth counts for raw vcf from ref_map:
6168 vcftools --vcf "/nobackup/mnkz72/new_refmap/bcf_out/populations.snps.vcf" --freq --out
6169 allpopulations_freqanalysis_freq
6170 vcftools --vcf "/nobackup/mnkz72/new_refmap/bcf_out/populations.snps.vcf" --depth --out
6171 allpopulations_freqanalysis_depth
6172 vcftools --vcf "/nobackup/mnkz72/new_refmap/bcf_out/populations.snps.vcf" --het --out
6173 allpopulations_freqanalysis_het
6174 vcftools --vcf "/nobackup/mnkz72/new_refmap/bcf_out/populations.snps.vcf" --counts --out
6175 allpopulations_freqanalysis_counts
6176 vcftools --vcf "/nobackup/mnkz72/new_refmap/bcf_out/populations.snps.vcf" --site-quality --out
6177 allpopulations_freqanalysis_site-quality
6178 vcftools --vcf "/nobackup/mnkz72/new_refmap/bcf_out/populations.snps.vcf" --missing-indv --out
6179 allpopulations_freqanalysis_missingindv
6180 #Filter for bi-allelic alleles:
6181 vcftools --vcf /nobackup/mnkz72/new_refmap/vcftools/allele_filter/allsamples_cp.vcf --min-alleles 2 --max-
6182 alleles 2 --out biallelic_filtered_populations_snps --recode

```

6183

6184

6185 **Appendix 25: Full Individual and Population list for heterozygous filtered analysis**

| name | pop | latitude | region | group |
|---------|-------|----------|---------------|----------|
| 3_11 | A-3 | 36.03633 | South_Spain | Southern |
| 3_13 | A-3 | 36.03633 | South_Spain | Southern |
| 3_14 | A-3 | 36.03633 | South_Spain | Southern |
| 3_15 | A-3 | 36.03633 | South_Spain | Southern |
| 3_27 | A-3 | 36.03633 | South_Spain | Southern |
| 1_1 | B-1 | 36.80044 | South_Spain | Southern |
| 1_26 | B-1 | 36.80044 | South_Spain | Southern |
| 1_6 | B-1 | 36.80044 | South_Spain | Southern |
| 10_15 | C-10 | 37.88211 | South_Spain | Southern |
| 10_26 | C-10 | 37.88211 | South_Spain | Southern |
| 10_27 | C-10 | 37.88211 | South_Spain | Southern |
| 10_30 | C-10 | 37.88211 | South_Spain | Southern |
| 6_1 | D-6 | 37.93551 | South_Spain | Southern |
| 6_29 | D-6 | 37.93551 | South_Spain | Southern |
| 12_10 | E-12 | 42.31008 | Mediterranean | Southern |
| 12_20 | E-12 | 42.31008 | Mediterranean | Southern |
| 12_8 | E-12 | 42.31008 | Mediterranean | Southern |
| 13_1 | F-13 | 43.02902 | North_Spain | Southern |
| 13_12 | F-13 | 43.02902 | North_Spain | Southern |
| 13_19 | F-13 | 43.02902 | North_Spain | Southern |
| 13_22 | F-13 | 43.02902 | North_Spain | Southern |
| 13_23 | F-13 | 43.02902 | North_Spain | Southern |
| 13_8 | F-13 | 43.02902 | North_Spain | Southern |
| Lla_17 | G-Lla | 43.40738 | North_Spain | Southern |
| Lla_20 | G-Lla | 43.40738 | North_Spain | Southern |
| Lla_23 | G-Lla | 43.40738 | North_Spain | Southern |
| Lla_33 | G-Lla | 43.40738 | North_Spain | Southern |
| Lla_A | G-Lla | 43.40738 | North_Spain | Southern |
| Lla_B | G-Lla | 43.40738 | North_Spain | Southern |
| Lla_C | G-Lla | 43.40738 | North_Spain | Southern |
| Vil_1 | H-Vil | 45.09393 | France | Northern |
| Vil_10 | H-Vil | 45.09393 | France | Northern |
| Vil_15 | H-Vil | 45.09393 | France | Northern |
| Vil_2 | H-Vil | 45.09393 | France | Northern |
| Vil_21 | H-Vil | 45.09393 | France | Northern |
| Vil_25p | H-Vil | 45.09393 | France | Northern |
| Vil_27 | H-Vil | 45.09393 | France | Northern |
| Vil_32 | H-Vil | 45.09393 | France | Northern |
| Vil_35 | H-Vil | 45.09393 | France | Northern |
| Vil_36g | H-Vil | 45.09393 | France | Northern |


| | | | | |
|---------|--------|----------|----------|----------|
| Vil_36p | H-Vil | 45.09393 | France | Northern |
| Tal_10 | I-Tal | 47.6997 | France | Northern |
| Tal_11 | I-Tal | 47.6997 | France | Northern |
| Tal_14 | I-Tal | 47.6997 | France | Northern |
| Tal_28 | I-Tal | 47.6997 | France | Northern |
| Tal_3 | I-Tal | 47.6997 | France | Northern |
| Tal_33 | I-Tal | 47.6997 | France | Northern |
| Tal_34 | I-Tal | 47.6997 | France | Northern |
| Tal_38 | I-Tal | 47.6997 | France | Northern |
| Tal_40 | I-Tal | 47.6997 | France | Northern |
| Mat_11 | J-Mat | 48.35697 | France | Northern |
| Mat_12 | J-Mat | 48.35697 | France | Northern |
| Mat_13 | J-Mat | 48.35697 | France | Northern |
| Mat_17 | J-Mat | 48.35697 | France | Northern |
| Mat_2 | J-Mat | 48.35697 | France | Northern |
| Mat_20 | J-Mat | 48.35697 | France | Northern |
| Mat_22 | J-Mat | 48.35697 | France | Northern |
| Mat_24 | J-Mat | 48.35697 | France | Northern |
| Mat_25 | J-Mat | 48.35697 | France | Northern |
| Mat_4 | J-Mat | 48.35697 | France | Northern |
| Mat_7 | J-Mat | 48.35697 | France | Northern |
| Dor_A | K-Dor | 50.6 | South_UK | Northern |
| Dor_B | K-Dor | 50.6 | South_UK | Northern |
| Dor_C | K-Dor | 50.6 | South_UK | Northern |
| Dor_D | K-Dor | 50.6 | South_UK | Northern |
| low2_10 | L-low2 | 50.68183 | South_UK | Northern |
| low2_13 | L-low2 | 50.68183 | South_UK | Northern |
| low2_14 | L-low2 | 50.68183 | South_UK | Northern |
| low2_2 | L-low2 | 50.68183 | South_UK | Northern |
| low2_25 | L-low2 | 50.68183 | South_UK | Northern |
| low2_26 | L-low2 | 50.68183 | South_UK | Northern |
| low2_7 | L-low2 | 50.68183 | South_UK | Northern |
| low2_9 | L-low2 | 50.68183 | South_UK | Northern |
| Tym_16 | M-Tym | 53.30307 | North_UK | Northern |
| Tym_19 | M-Tym | 53.30307 | North_UK | Northern |
| Tym_20 | M-Tym | 53.30307 | North_UK | Northern |
| Tym_26g | M-Tym | 53.30307 | North_UK | Northern |
| Tym_26p | M-Tym | 53.30307 | North_UK | Northern |
| Tym_3 | M-Tym | 53.30307 | North_UK | Northern |
| Tym_30 | M-Tym | 53.30307 | North_UK | Northern |
| Tym_40 | M-Tym | 53.30307 | North_UK | Northern |
| Tym_41 | M-Tym | 53.30307 | North_UK | Northern |
| Tym_5 | M-Tym | 53.30307 | North_UK | Northern |
| Sut_17 | N-Sut | 53.35291 | North_UK | Northern |
| Sut_29 | N-Sut | 53.35291 | North_UK | Northern |
| Sut_33 | N-Sut | 53.35291 | North_UK | Northern |

| | | | | |
|--------|-------|----------|----------|----------|
| Sut_34 | N-Sut | 53.35291 | North_UK | Northern |
| Sut_36 | N-Sut | 53.35291 | North_UK | Northern |
| Sut_4 | N-Sut | 53.35291 | North_UK | Northern |
| Sut_41 | N-Sut | 53.35291 | North_UK | Northern |
| Gis | CUL | NA | Cultivar | Cultivar |
| Lir | CUL | NA | Cultivar | Cultivar |
| Mon | CUL | NA | Cultivar | Cultivar |
| Olg | CUL | NA | Cultivar | Cultivar |
| Olg_1 | CUL | NA | Cultivar | Cultivar |
| Ome | CUL | NA | Cultivar | Cultivar |
| Pri | CUL | NA | Cultivar | Cultivar |
| Rab | CUL | NA | Cultivar | Cultivar |
| Suz | CUL | NA | Cultivar | Cultivar |
| Tin | CUL | NA | Cultivar | Cultivar |


6186

6187

6188



6191 **Appendix 27: DAPC Summary**



6192

6193

6194 **Appendix 28: Tajima D's Summary**

| pop | mean_pi | TajimaD | rare_alleles | demography | selection |
|---|----------|----------|--------------|----------------------------|-----------------|
| A-3 | 82.3 | -0.42 | neutral | neutral | neutral |
| B-1 | 57.58 | -79.44 | many_rare | expansion_after_bottleneck | selective_sweep |
| C-10 | 18.3 | -71.09 | many_rare | expansion_after_bottleneck | selective_sweep |
| D-6 | 63.17 | -182.16 | many_rare | expansion_after_bottleneck | selective_sweep |
| E-12 | 39.25 | -125.65 | many_rare | expansion_after_bottleneck | selective_sweep |
| F-13 | 49.81 | -40.29 | many_rare | expansion_after_bottleneck | selective_sweep |
| G-Lla | 67.41 | -9.04 | many_rare | expansion_after_bottleneck | selective_sweep |
| H-Vil | 9.11 | -15.89 | many_rare | expansion_after_bottleneck | selective_sweep |
| I-Tal | 3.5 | -8.33 | many_rare | expansion_after_bottleneck | selective_sweep |
| J-Mat | 41.01 | -34.55 | many_rare | expansion_after_bottleneck | selective_sweep |
| K-Dor | 41.55 | -18.45 | many_rare | expansion_after_bottleneck | selective_sweep |
| L-low2 | 11.33 | -8.05 | many_rare | expansion_after_bottleneck | selective_sweep |
| M-Tym | 10.56 | -15.02 | many_rare | expansion_after_bottleneck | selective_sweep |
| N-Sut | 21.02 | -14.14 | many_rare | expansion_after_bottleneck | selective_sweep |
| P(T<=t) two-tail between North/South | 0.019298 | 0.023475 | | | |

6195

6196 **Appendix 29: Mean MAF for each population**

| pop | mean_nsites | mean_pi | mean_maf |
|------|-------------|---------|----------|
| A-3 | 811.7 | 82.3 | 0.46 |
| B-1 | 833 | 57.58 | 0.66 |
| C-10 | 858 | 18.3 | 0.86 |
| D-6 | 692 | 63.17 | 0.57 |

| | | | |
|--------|-------|-------|------|
| E-12 | 862 | 39.25 | 0.91 |
| F-13 | 804.9 | 49.81 | 0.64 |
| G-Lla | 857.8 | 67.41 | 0.32 |
| H-Vil | 854.7 | 9.11 | 0.25 |
| I-Tal | 854.2 | 3.5 | 0.55 |
| J-Mat | 832.9 | 41.01 | 0.17 |
| K-Dor | 856 | 41.55 | 0.32 |
| L-low2 | 862 | 11.33 | 0.43 |
| M-Tym | 773.5 | 10.56 | 0.25 |
| N-Sut | 790 | 21.02 | 0.26 |

6197

6198 **Appendix 30: Chapter 5 R Commands**

6199 #Seed Measurements:

6200 setwd("G:/Linum Project/rdir")

6201 library(ggplot2)

6202 library(ggpubr)

6203 library(readr)

6204 seed_measurements_wild_data_pop <- read_csv("seed_measurements_wild_data_pop.csv",

6205 col_types = cols(`Lat_(degrees_N)` = col_number(),

6206 Alt = col_number(), `Seed_length_(Cm)` = col_number(),

6207 `Seed_width_(Cm)` = col_number(),

6208 `Seed_Area_(Cm)` = col_number(),

6209 pc1 = col_number()))

6210 View(seed_measurements_wild_data_pop)

6211

6212 #Modelling for seed area vs latitude:

6213 g <- ggscatter(seed_measurements_wild_data_pop, x = "Lat_(degrees_N)", y = "Seed_Area_(Cm)", size = 1.0,

6214 rug = TRUE, # Add marginal rug

6215 color = "Experiment", palette = "uchicago", facet.by = "Experiment", add = "reg.line", conf.int = TRUE) +

6216 stat_cor(aes(color = Experiment), label.sep = ";", label.x.npc = "left", label.y.npc=0.20, method =

6217 "pearson")+geom_point(aes(color = Experiment))+stat_smooth(method="glm",se=FALSE)+

6218 theme_light()+ggtitle("Seed Area vs Latitude")

6219

```

6220 g +
6221 font("title", size = 22, color = "black", face = "bold")+
6222 font("xlab", size = 18, color = "black", face = "bold")+
6223 font("ylab", size = 18, color = "black", face = "bold")+
6224 font("legend.title", size = 18, color = "black", face = "bold")+
6225 font("legend.text", size = 14, color = "black",)
6226
6227 #Modelling for seed area vs pc1:
6228 h <- ggscatter(seed_measurements_wild_data_pop, x = "pc1", y = "Seed_Area_(Cm)", size = 1.0,
6229               rug = TRUE,                # Add marginal rug
6230               color = "Experiment", palette = "uchicago", facet.by = "Experiment",add = "reg.line", conf.int = TRUE) +
6231               stat_cor(aes(color = Experiment), label.sep = ";", label.x.npc = "left", label.y.npc=0.20, method =
6232 "pearson")+geom_point(aes(color = Experiment))+stat_smooth(method="glm",se=FALSE)+
6233 theme_light()+ggtitle("Seed Area vs Climatic Variables (pc1)")
6234
6235 h +
6236 font("title", size = 22, color = "black", face = "bold")+
6237 font("xlab", size = 18, color = "black", face = "bold")+
6238 font("ylab", size = 18, color = "black", face = "bold")+
6239 font("legend.title", size = 18, color = "black", face = "bold")+
6240 font("legend.text", size = 14, color = "black",)
6241
6242 #Vern/Non-vern on other traits:
6243 library("car")
6244 setwd("G:/Linum Project/rdir")
6245 library(readr)
6246 combined_vern_control_2018_only <- read_csv("combined_vern_control_2018_only.csv",
6247               col_types = cols(Lat = col_number(),
6248               Lon = col_number(), pc1 = col_number(),

```

```

6249         `Height_(Cm)` = col_number(), Stem_no = col_number(),
6250         Bud_no = col_number(), Days_to_fl = col_number())
6251 View(combined_vern_control_2018_only)
6252 scatterplot(Days_to_fl ~ Height_(Cm) | Treatment, data = combined_vern_control_2018_only,
6253             smooth = TRUE, regLine = TRUE, grid = FALSE, frame = FALSE, xlab = "Plant Height(Cm)", ylab = "Days to
6254 Flowering")
6255 abline(lm(Height ~ Days_to_fl data = combined_vern_control_2018_only),col="red")
6256
6257 ggscatter(combined_vern_control_2018_only, x = "Bud_no", y = "Days_to_fl", size = 2.0,
6258           rug = TRUE,                # Add marginal rug
6259           color = "Treatment", palette = "jco", facet.by = "Treatment",add = "reg.line", conf.int = TRUE) +
6260   stat_cor(aes(color = Treatment), method = "pearson")
6261
6262 #Model summary
6263 summary1 <- summary(glm(Stem_no ~ Days_to_fl, data = combined_vern_control_2018_only))
6264 adjRsqr <- summary1$adj.r.squared
6265 fStat <- summary1$statistic
6266 pValue <- pf(fStat[summary1])
6267 summary(summary1)
6268
6269 #Height - Days to fl test:
6270 Heightdaysmod<-cbind(combined_vern_control_2018_only$Days_to_fl,
6271 combined_vern_control_2018_only$'Height_(Cm)')
6272 Heightaysmod2<-glm(Heightdaysmod~Treatment, family=quasibinomial,
6273 data=combined_vern_control_2018_only)
6274 summary(Heightaysmod2)
6275 barcentres<-
6276 barplot(tapply(combined_vern_control_2018_only$'Height_(Cm)',combined_vern_control_2018_only$Treatm
6277 ent,mean),ylim=c(0,max(tapply(combined_vern_control_2018_only$'Height_(Cm)',combined_vern_control_2

```

```

6278 018_only$Treatment,mean)+5)), ylab = "Height (Cm)", xlab = "Treatment", main = "Plant Height vs
6279 Treatments")
6280 means<-
6281 tapply(combined_vern_control_2018_only$'Height_(Cm)',combined_vern_control_2018_only$Treatment,mea
6282 n)
6283 ses<-
6284 tapply(combined_vern_control_2018_only$'Height_(Cm)',combined_vern_control_2018_only$Treatment,sd)/
6285 sqrt(tapply(combined_vern_control_2018_only$'Height_(Cm)',combined_vern_control_2018_only$Treatment
6286 ,length))
6287 arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
6288
6289 #Stem_no - Days to fl test:
6290 Stemdaysmod<-cbind(combined_vern_control_2018_only$Stem_no,
6291 combined_vern_control_2018_only$Days_to_fl)
6292 Stemdaysmod2<-glm(Stemdaysmod~Treatment, family=quasibinomial,
6293 data=combined_vern_control_2018_only)
6294 summary(Stemdaysmod2)
6295 barcentres<-
6296 barplot(tapply(combined_vern_control_2018_only$Stem_no,combined_vern_control_2018_only$Treatment,
6297 mean),ylim=c(0,max(tapply(combined_vern_control_2018_only$Stem_no,combined_vern_control_2018_only
6298 $Treatment,mean)+2)), ylab = "Stem Number", xlab = "Treatment", main = "Stem Number vs Treatments")
6299 means<-
6300 tapply(combined_vern_control_2018_only$Stem_no,combined_vern_control_2018_only$Treatment,mean)
6301 ses<-
6302 tapply(combined_vern_control_2018_only$Stem_no,combined_vern_control_2018_only$Treatment,sd)/sqrt(
6303 tapply(combined_vern_control_2018_only$Stem_no,combined_vern_control_2018_only$Treatment,length))
6304 arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
6305
6306 #Bud_no - Days to fl test:
6307 buddaysmod<-cbind(combined_vern_control_2018_only$Days_to_fl,
6308 combined_vern_control_2018_only$Bud_no)
6309 buddaysmod2<-buddaysmodlm<-lm(buddaysmod~Treatment,data=combined_vern_control_2018_only)

```



```

6310 summary(pc1daysmodlm)
6311 barcentres<-
6312 barplot(tapply(combined_vern_control_2018_only$Bud_no,combined_vern_control_2018_only$Treatment,m
6313 ean),ylim=c(0,max(tapply(combined_vern_control_2018_only$Bud_no,combined_vern_control_2018_only$Tr
6314 eatment,mean)+2)), ylab = "Bud Number", xlab = "Treatment", main = "Bud Number vs Treatments")
6315 means<-
6316 tapply(combined_vern_control_2018_only$Bud_no,combined_vern_control_2018_only$Treatment,mean)
6317 ses<-
6318 tapply(combined_vern_control_2018_only$Bud_no,combined_vern_control_2018_only$Treatment,sd)/sqrt(t
6319 apply(combined_vern_control_2018_only$Bud_no,combined_vern_control_2018_only$Treatment,length))
6320 arrows(barcentres, means-ses, barcentres, means+ses, length=0.05, angle=90, code=3)
6321
6322 #PC1 - Height Correlation:
6323 library(readr)
6324 wilddonly_vern18 <- read_csv("wilddonly_vern18.csv",
6325                             col_types = cols(Lat = col_number(),
6326                                                Lon = col_number(), pc1 = col_number(),
6327                                                `Height_(Cm)` = col_number(), Stem_no = col_number(),
6328                                                Bud_no = col_number(), Days_to_fl = col_number()))
6329 View(wilddonly_vern18)
6330
6331 #pc1 - height correlation:
6332 a <- ggscatter(wilddonly_vern18, x = "pc1", y = "Height_(Cm)", size = 1.0,
6333               rug = TRUE,                # Add marginal rug
6334               color = "Treatment", palette = "uchicago", facet.by = "Treatment",add = "reg.line", conf.int = TRUE) +
6335               stat_cor(aes(color = Treatment), label.sep = ";", label.x.npc = "left", label.y.npc=0.8, method =
6336 "pearson")+geom_point(aes(color = Treatment))+stat_smooth(method="glm",se=FALSE)+
6337 theme_light()+ggtitle("Plant Height (Cm) vs Climate Variable (pc1)")
6338
6339 a +
6340   font("title", size = 22, color = "black", face = "bold")+

```

```

6341 font("xlab", size = 18, color = "blue", face = "bold")+
6342 font("ylab", size = 18, color = "blue", face = "bold")+
6343 font("legend.title", size = 18, color = "black", face = "bold")+
6344 font("legend.text", size = 18, color = "blue", face = "bold")
6345
6346
6347 #Lat - height correlation:
6348 b <- ggscatter(wildonly_vern18, x = "Lat", y = "Height_(Cm)", size = 1.0,
6349             rug = TRUE,                # Add marginal rug
6350             color = "Treatment", palette = "uchicago", facet.by = "Treatment", add = "reg.line", conf.int = TRUE) +
6351             stat_cor(aes(color = Treatment), label.sep = ";", label.x.npc = "left", label.y.npc=0.8, method =
6352 "pearson")+geom_point(aes(color = Treatment))+stat_smooth(method="glm",se=FALSE)+
6353 theme_light()+ggtitle("Plant Height (Cm) vs Latitude (degrees North)")
6354
6355 b +
6356 font("title", size = 20, color = "black", face = "bold")+
6357 font("xlab", size = 18, color = "blue", face = "bold")+
6358 font("ylab", size = 18, color = "blue", face = "bold")+
6359 font("legend.title", size = 18, color = "black", face = "bold")+
6360 font("legend.text", size = 18, color = "blue", face = "bold")
6361
6362 #pc1 - Stem number correlation:
6363 c <- ggscatter(wildonly_vern18, x = "pc1", y = "Stem_no", size = 1.0,
6364             rug = TRUE,                # Add marginal rug
6365             color = "Treatment", palette = "uchicago", facet.by = "Treatment", add = "reg.line", conf.int = TRUE) +
6366             stat_cor(aes(color = Treatment), label.sep = ";", label.x.npc = "left", label.y.npc=0.9, method =
6367 "pearson")+geom_point(aes(color = Treatment))+stat_smooth(method="glm",se=FALSE)+
6368 theme_light()+ggtitle("Stem Number vs Climate Variable (pc1)")
6369
6370 c +

```

```

6371 font("title", size = 22, color = "black", face = "bold")+
6372 font("xlab", size = 18, color = "blue", face = "bold")+
6373 font("ylab", size = 18, color = "blue", face = "bold")+
6374 font("legend.title", size = 18, color = "black", face = "bold")+
6375 font("legend.text", size = 18, color = "blue", face = "bold")
6376
6377 #Lat - Stem number correlation:
6378 d <- ggscatter(wildonly_vern18, x = "Lat", y = "Stem_no", size = 1.0,
6379             rug = TRUE,                # Add marginal rug
6380             color = "Treatment", palette = "uchicago", facet.by = "Treatment", add = "reg.line", conf.int = TRUE) +
6381             stat_cor(aes(color = Treatment), label.sep = ";", label.x.npc = "left", label.y.npc=0.85, method =
6382 "pearson")+geom_point(aes(color = Treatment))+stat_smooth(method="glm",se=FALSE)+
6383 theme_light()+ggtitle("Stem Number vs Latitude (degrees North)")
6384
6385 d +
6386 font("title", size = 22, color = "black", face = "bold")+
6387 font("xlab", size = 18, color = "blue", face = "bold")+
6388 font("ylab", size = 18, color = "blue", face = "bold")+
6389 font("legend.title", size = 18, color = "black", face = "bold")+
6390 font("legend.text", size = 18, color = "blue", face = "bold")
6391
6392 #pc1 - Bud number correlation:
6393 e <- ggscatter(wildonly_vern18, x = "pc1", y = "Bud_no", size = 1.0,
6394             rug = TRUE,                # Add marginal rug
6395             color = "Treatment", palette = "uchicago", facet.by = "Treatment", add = "reg.line", conf.int = TRUE) +
6396             stat_cor(aes(color = Treatment), label.sep = ";", label.x.npc = "left", label.y.npc=0.7, method =
6397 "pearson")+geom_point(aes(color = Treatment))+stat_smooth(method="glm",se=FALSE)+
6398 theme_light()+ggtitle("Bud Number vs Climate Variable (pc1)")
6399
6400 e +

```

```

6401 font("title", size = 22, color = "black", face = "bold")+
6402 font("xlab", size = 18, color = "blue", face = "bold")+
6403 font("ylab", size = 18, color = "blue", face = "bold")+
6404 font("legend.title", size = 18, color = "black", face = "bold")+
6405 font("legend.text", size = 18, color = "blue", face = "bold")
6406
6407 #Lat - Bud number correlation:
6408 f <- ggscatter(wildonly_vern18, x = "Lat", y = "Bud_no", size = 1.0,
6409             rug = TRUE,                # Add marginal rug
6410             color = "Treatment", palette = "uchicago", facet.by = "Treatment", add = "reg.line", conf.int = TRUE) +
6411             stat_cor(aes(color = Treatment), label.sep = ";", label.x.npc = "left", label.y.npc=0.53, method =
6412 "pearson")+geom_point(aes(color = Treatment))+stat_smooth(method="glm",se=FALSE)+
6413 theme_light()+ggtitle("Bud Number vs Latitude (degrees North)")
6414
6415 f +
6416 font("title", size = 22, color = "black", face = "bold")+
6417 font("xlab", size = 18, color = "blue", face = "bold")+
6418 font("ylab", size = 18, color = "blue", face = "bold")+
6419 font("legend.title", size = 18, color = "black", face = "bold")+
6420 font("legend.text", size = 18, color = "blue", face = "bold")
6421 Appendix 31: Plant Height, stem number, and bud number

```

| Pop_ind | Height_(Cm) | Stem_no | Bud_no | Days_to_fl |
|---------|-------------|---------|--------|------------|
| Vil_25 | 76.5 | 10 | 2 | 292 |
| Vil_27 | 21.5 | 6 | 0 | 202 |
| Vil_27 | 21.5 | 6 | 0 | 202 |
| Vil_36 | 39.5 | 27 | 2 | 167 |
| Roc_12 | 48 | 16 | 2 | 247 |
| Roc_12 | 52.5 | 25 | 3 | 205 |
| Saf_16 | 29.5 | 10 | 0 | 212 |
| Saf_17 | 41 | 33 | 5 | 294 |
| Saf_19 | 34 | 48 | 2 | 299 |
| Saf_19 | 42 | 41 | 1 | 198 |
| Tal_10 | 34 | 6 | 1 | 149 |

| | | | | |
|----------|------|----|----|-----|
| Tal_10 | 26 | 3 | 0 | 69 |
| Mat_2 | 32 | 6 | 4 | 201 |
| Mat_2 | 39 | 33 | 13 | 191 |
| Mat_14 | 29 | 5 | 3 | 211 |
| Mat_14 | 24 | 12 | 2 | 177 |
| Mat_17 | 26.3 | 6 | 3 | 200 |
| Mat_17 | 49 | 20 | 16 | 198 |
| Mat_18 | 39 | 17 | 1 | 248 |
| Mat_18 | 36.5 | 21 | 12 | 190 |
| Mat_23 | 47 | 22 | 8 | 199 |
| Mat_23 | 31 | 5 | 5 | 196 |
| 19_26 | 43.5 | 11 | 0 | 184 |
| 19_26 | 43.5 | 11 | 0 | 184 |
| 19_30 | 39 | 9 | 3 | 213 |
| 14_21 | 40 | 36 | 3 | 214 |
| 13_1 | 34.5 | 12 | 1 | 191 |
| Lla_17 | 45 | 6 | 1 | 270 |
| Lla_20 | 55 | 33 | 10 | 300 |
| Lla_23 | 29 | 6 | 2 | 212 |
| Lla_43_A | 58 | 18 | 1 | 243 |
| Lla_B | 33 | 26 | 1 | 212 |
| Lla_C | 43 | 25 | 5 | 202 |
| Lla_C | 28 | 14 | 3 | 207 |
| 15_17 | 26.5 | 7 | 3 | 151 |
| 15_17 | 23 | 8 | 3 | 134 |
| 15_18 | 18 | 10 | 3 | 177 |
| 15_18 | 45 | 30 | 1 | 160 |
| 15_28 | 29.5 | 2 | 0 | 236 |
| 15_30 | 32 | 5 | 1 | 184 |
| 15_30 | 41 | 15 | 1 | 182 |
| 3_11 | 49.5 | 12 | 5 | 190 |
| 3_13 | 41.5 | 10 | 2 | 157 |
| 3_13 | 29.5 | 3 | 1 | 156 |
| 3_13 | 31 | 3 | 2 | 193 |
| 3_14 | 35.5 | 10 | 0 | 160 |
| 3_14 | 43 | 4 | 2 | 118 |
| 3_15 | 27 | 3 | 2 | 122 |
| 3_15 | 40.5 | 4 | 2 | 189 |
| 2_3 | 26.5 | 3 | 1 | 109 |
| 2_3 | 29 | 5 | 2 | 98 |
| 2_29 | 23.5 | 12 | 1 | 216 |
| 4_7 | 28 | 7 | 1 | 76 |
| 4_7 | 18.5 | 3 | 0 | 63 |
| 4_23 | 30 | 11 | 1 | 151 |

| | | | | |
|---------|------|----|----|-----|
| 4_27 | 25 | 2 | 0 | 66 |
| 4_27 | 18.5 | 6 | 0 | 67 |
| 4_28 | 31 | 2 | 1 | 102 |
| 4_28 | 22 | 4 | 2 | 82 |
| 4_A | 33 | 12 | 4 | 186 |
| Tor_4 | 17.5 | 6 | 2 | 59 |
| Tor_4 | 16 | 5 | 1 | 71 |
| 1_1 | 25 | 13 | 0 | 138 |
| 1_26 | 30.5 | 5 | 1 | 152 |
| 1_26 | 46 | 5 | 0 | 178 |
| 1_A | 29.2 | 9 | 1 | 113 |
| 1_C | 39 | 12 | 1 | 128 |
| 1_C | 34.5 | 7 | 2 | 124 |
| 5_4 | 20 | 3 | 1 | 54 |
| 5_10 | 19 | 4 | 0 | 73 |
| 8_12 | 34.6 | 9 | 4 | 202 |
| 8_12 | 47 | 13 | 12 | 184 |
| 8_16 | 35.6 | 10 | 1 | 159 |
| 9_23 | 36.5 | 13 | 0 | 105 |
| 9_27 | 29.3 | 1 | 0 | 55 |
| 9_27 | 33.5 | 5 | 4 | 62 |
| 9_34 | 32 | 2 | 1 | 118 |
| 10_15 | 27.5 | 1 | 0 | 153 |
| 10_15 | 18.5 | 3 | 0 | 84 |
| 10_26 | 38 | 7 | 0 | 115 |
| 10_26 | 29.5 | 6 | 1 | 100 |
| 10_27 | 32 | 5 | 3 | 118 |
| 10_27 | 31 | 5 | 0 | 117 |
| 10_30 | 29 | 8 | 0 | 83 |
| 6_1 | 37.7 | 9 | 0 | 103 |
| 6_26 | 42.5 | 11 | 4 | 198 |
| 6_26 | 50 | 7 | 6 | 181 |
| 6_29 | 47.5 | 5 | 3 | 199 |
| 7_17 | 36 | 8 | 1 | 121 |
| 11_23 | 42 | 15 | 0 | 158 |
| Man_4 | 49 | 17 | 3 | 214 |
| Man_4 | 26 | 8 | 1 | 212 |
| Man_5 | 38 | 8 | 1 | 200 |
| Man_8 | 33 | 30 | 8 | 287 |
| Tym_5 | 33.5 | 22 | 0 | 199 |
| Tym_30 | 31 | 6 | 2 | 157 |
| IOW2_2 | 17.5 | 4 | 2 | 74 |
| IOW2_2 | 25.5 | 8 | 7 | 50 |
| IOW2_25 | 23 | 2 | 0 | 57 |

| | | | | |
|---------|------|----|----|-----|
| IOW2_25 | 26.5 | 5 | 1 | 66 |
| IOW2_26 | 23 | 18 | 1 | 212 |
| IOW1_11 | 36 | 15 | 0 | 307 |
| IOW1_17 | 36 | 18 | 1 | 272 |
| Ara | 78.5 | 1 | 2 | 73 |
| Ara | 93 | 3 | 4 | 77 |
| Ari | 63.5 | 4 | 8 | 57 |
| Ari | 88.5 | 2 | 4 | 45 |
| Ari | 91.5 | 3 | 3 | 66 |
| Ble | 62 | 1 | 2 | 49 |
| Ble | 45 | 3 | 8 | 38 |
| Ble | 58 | 1 | 4 | 47 |
| Ble | 40.5 | 1 | 9 | 50 |
| Bol | 54 | 1 | 2 | 68 |
| Bol | 64 | 1 | 0 | 52 |
| Ede | 73 | 1 | 2 | 65 |
| Ede | 76 | 1 | 0 | 63 |
| Ede | 73 | 1 | 2 | 65 |
| Gis | 53 | 3 | 3 | 83 |
| Gis | 52 | 1 | 1 | 58 |
| Gis | 52 | 3 | 2 | 61 |
| Gis | 52 | 5 | 2 | 63 |
| Lir | 57.5 | 2 | 4 | 49 |
| Lir | 86 | 3 | 8 | 49 |
| Lir | 45 | 3 | 6 | 56 |
| Mar | 56 | 2 | 5 | 49 |
| Mar | 44.5 | 3 | 6 | 63 |
| Mon | 47 | 3 | 6 | 54 |
| Mon | 78.5 | 1 | 3 | 40 |
| Mon | 77 | 1 | 3 | 53 |
| Olg | 16.5 | 1 | 0 | 90 |
| Olg | 58.5 | 1 | 0 | 111 |
| Ome | 36 | 1 | 5 | 49 |
| Ome | 44.5 | 1 | 5 | 40 |
| Ome | 47 | 3 | 8 | 49 |
| Ome | 55 | 1 | 10 | 46 |
| Pri | 49.5 | 3 | 3 | 65 |
| Pri | 38.5 | 1 | 2 | 48 |
| Pri | 37 | 1 | 3 | 76 |
| Pri | 32 | 7 | 1 | 129 |
| Rab | 48 | 8 | 14 | 118 |
| Rab | 48 | 3 | 11 | 38 |
| Rab | 57.5 | 1 | 10 | 45 |
| Rab | 63 | 3 | 3 | 57 |

| | | | | |
|--------|------|---|---|-----|
| Suz | 72 | 3 | 2 | 51 |
| Suz | 72 | 1 | 1 | 26 |
| Suz | 78 | 3 | 3 | 52 |
| Suz | 68.5 | 1 | 3 | 77 |
| Tin | 50 | 1 | 3 | 63 |
| Tin | 64 | 1 | 2 | 55 |
| Tin | 61 | 2 | 6 | 68 |
| Vol | 40 | 3 | 5 | 51 |
| Vol | 27.8 | 2 | 2 | 64 |
| Vil_21 | 27 | 3 | 3 | 98 |
| Vil_21 | 26 | 2 | 0 | 101 |
| Vil_25 | 19.5 | 3 | 0 | 105 |
| Vil_25 | 25.5 | 3 | 3 | 96 |
| Vil_27 | 24.5 | 3 | 2 | 93 |
| Vil_27 | 27.5 | 5 | 1 | 101 |
| Vil_36 | 30.5 | 3 | 1 | 92 |
| Vil_36 | 27.5 | 2 | 0 | 99 |
| Roc_12 | 38 | 4 | 2 | 93 |
| Roc_12 | 46 | 5 | 2 | 90 |
| Saf_9 | 21.5 | 4 | 1 | 95 |
| Saf_9 | 25 | 3 | 3 | 106 |
| Saf_10 | 31 | 3 | 1 | 100 |
| Saf_10 | 8 | 6 | 4 | 92 |
| Saf_16 | 25 | 3 | 2 | 98 |
| Saf_16 | 25.5 | 8 | 2 | 92 |
| Saf_19 | 27.5 | 5 | 0 | 110 |
| Saf_19 | 31 | 4 | 0 | 95 |
| Tal_4 | 26 | 4 | 5 | 95 |
| Tal_4 | 32 | 6 | 6 | 96 |
| Tal_23 | 21 | 3 | 2 | 93 |
| Tal_23 | 32.5 | 3 | 1 | 103 |
| Tal_25 | 31.5 | 3 | 0 | 131 |
| Tal_28 | 24.5 | 6 | 1 | 122 |
| Tal_28 | 30 | 4 | 1 | 117 |
| Mat_2 | 45 | 4 | 6 | 91 |
| Mat_2 | 23 | 7 | 5 | 119 |
| Mat_14 | 30.5 | 5 | 0 | 102 |
| Mat_14 | 34.5 | 2 | 2 | 97 |
| Mat_17 | 23.5 | 3 | 0 | 107 |
| Mat_17 | 19.5 | 4 | 0 | 95 |
| Mat_23 | 27.5 | 5 | 5 | 87 |
| Mat_23 | 34.5 | 1 | 1 | 103 |
| Mat_24 | 31 | 4 | 2 | 101 |
| Mat_24 | 45 | 6 | 5 | 98 |

| | | | | |
|--------|------|----|----|-----|
| 12_20 | 36 | 3 | 1 | 85 |
| 19_30 | 25 | 9 | 0 | 99 |
| 19_30 | 30 | 3 | 2 | 106 |
| 14_6 | 17 | 5 | 0 | 105 |
| 14_21 | 31.5 | 6 | 1 | 96 |
| 14_21 | 42 | 2 | 0 | 99 |
| 13_1 | 39 | 1 | 3 | 98 |
| 13_1 | 22.5 | 6 | 2 | 89 |
| 13_5 | 24 | 8 | 0 | 134 |
| 13_12 | 21.5 | 4 | 0 | 106 |
| Lla_17 | 19 | 1 | 1 | 100 |
| Lla_20 | 32.5 | 3 | 1 | 99 |
| Lla_20 | 30 | 2 | 1 | 95 |
| Lla_23 | 25 | 3 | 2 | 131 |
| Lla_23 | 26 | 3 | 0 | 127 |
| Lla_33 | 30 | 3 | 2 | 106 |
| Lla_33 | 28 | 6 | 4 | 106 |
| Lla_A | 30.5 | 7 | 3 | 110 |
| Lla_A | 29.5 | 4 | 2 | 98 |
| Lla_B | 30 | 10 | 11 | 98 |
| Lla_C | 28.5 | 4 | 2 | 99 |
| 15_17 | 29.5 | 1 | 2 | 92 |
| 15_17 | 23 | 4 | 2 | 89 |
| 15_18 | 30 | 5 | 3 | 88 |
| 15_18 | 24.5 | 8 | 3 | 85 |
| 15_27 | 24.5 | 3 | 2 | 97 |
| 15_27 | 32 | 3 | 1 | 96 |
| 15_28 | 33 | 3 | 1 | 101 |
| 15_28 | 29 | 5 | 2 | 102 |
| 15_30 | 32 | 1 | 2 | 97 |
| 15_30 | 29 | 2 | 0 | 103 |
| 3_11 | 27 | 2 | 1 | 90 |
| 3_11 | 30.5 | 4 | 2 | 93 |
| 3_13 | 24.5 | 4 | 0 | 97 |
| 3_13 | 31 | 4 | 1 | 95 |
| 3_14 | 24.2 | 1 | 1 | 91 |
| 3_14 | 34 | 3 | 2 | 97 |
| 3_16 | 58 | 5 | 1 | 101 |
| 2_3 | 27 | 3 | 1 | 80 |
| 2_3 | 26.5 | 3 | 3 | 88 |
| 2_29 | 30 | 5 | 1 | 91 |
| 2_29 | 24 | 7 | 1 | 102 |
| 4_5 | 23 | 2 | 2 | 87 |
| 4_5 | 28.1 | 3 | 2 | 80 |

| | | | | |
|-------|------|---|---|-----|
| 4_7 | 26 | 3 | 1 | 89 |
| 4_7 | 25 | 4 | 3 | 92 |
| 4_23 | 19 | 1 | 1 | 88 |
| 4_23 | 28 | 1 | 1 | 86 |
| 4_27 | 24.3 | 3 | 2 | 84 |
| 4_28 | 25.3 | 1 | 1 | 82 |
| Tor_4 | 18.5 | 6 | 9 | 77 |
| Tor_4 | 17 | 4 | 3 | 79 |
| 1_26 | 24.5 | 3 | 0 | 90 |
| 1_26 | 25 | 3 | 3 | 88 |
| 1_A | 18.5 | 2 | 0 | 86 |
| 1_A | 28 | 3 | 2 | 91 |
| 5_4 | 28 | 4 | 3 | 79 |
| 5_4 | 24.5 | 3 | 3 | 78 |
| 5_4 | 21 | 3 | 2 | 78 |
| 5_10 | 19.5 | 1 | 2 | 83 |
| 5_10 | 18.7 | 2 | 2 | 84 |
| 8_16 | 32 | 5 | 7 | 87 |
| 8_16 | 21 | 7 | 5 | 85 |
| 8_27 | 24.2 | 3 | 0 | 82 |
| 9_24 | 24.5 | 5 | 5 | 82 |
| 9_27 | 23 | 5 | 2 | 84 |
| 10_15 | 24.5 | 5 | 2 | 74 |
| 10_26 | 28 | 5 | 0 | 77 |
| 10_26 | 19.5 | 1 | 0 | 84 |
| 10_27 | 20 | 5 | 0 | 78 |
| 10_27 | 26 | 5 | 7 | 78 |
| 10_30 | 19.3 | 5 | 1 | 76 |
| 6_26 | 21 | 5 | 0 | 92 |
| 6_26 | 24 | 5 | 1 | 95 |
| 6_29 | 30 | 1 | 0 | 98 |
| 6_29 | 27 | 7 | 3 | 98 |
| 11_23 | 24 | 9 | 4 | 86 |
| Man_4 | 21 | 5 | 3 | 94 |
| Man_4 | 24.5 | 3 | 2 | 99 |
| Man_5 | 29 | 5 | 2 | 98 |
| Man_5 | 25 | 3 | 1 | 98 |
| Man_6 | 27 | 4 | 2 | 98 |
| Man_6 | 29 | 5 | 1 | 93 |
| Man_8 | 27.5 | 5 | 0 | 92 |
| Man_8 | 19.5 | 5 | 1 | 105 |
| Tym_3 | 28.5 | 3 | 1 | 111 |
| Tym_3 | 33.5 | 7 | 1 | 105 |
| Tym_5 | 22.5 | 5 | 0 | 109 |

| | | | | |
|---------|------|----|---|-----|
| Tym_5 | 28.5 | 1 | 0 | 106 |
| Tym_26 | 25.5 | 3 | 0 | 106 |
| Tym_26 | 28.5 | 3 | 1 | 78 |
| Tym_30 | 28 | 1 | 0 | 108 |
| Tym_30 | 27.5 | 4 | 2 | 107 |
| Sut_4 | 27.5 | 3 | 0 | 103 |
| Sut_4 | 30.5 | 4 | 1 | 103 |
| Dor | 35 | 3 | 1 | 100 |
| Dor | 37.5 | 4 | 1 | 106 |
| Dor | 31 | 5 | 0 | 103 |
| Dor | 29.5 | 2 | 0 | 109 |
| Dor_C | 29 | 2 | 5 | 93 |
| Dor_C | 33 | 8 | 5 | 103 |
| IOW2_2 | 19 | 4 | 0 | 92 |
| IOW2_2 | 29 | 7 | 7 | 96 |
| IOW2_25 | 29 | 5 | 3 | 90 |
| IOW2_25 | 23.5 | 5 | 3 | 88 |
| IOW2_26 | 22.5 | 4 | 1 | 111 |
| IOW2_26 | 33 | 11 | 0 | 102 |
| IOW2_30 | 16.5 | 6 | 0 | 110 |
| IOW2_30 | 30 | 15 | 2 | 106 |
| IOW1_10 | 24.5 | 4 | 1 | 97 |
| IOW1_10 | 25 | 6 | 4 | 86 |
| IOW1_11 | 26.5 | 5 | 2 | 104 |
| IOW1_11 | 26.5 | 4 | 1 | 99 |
| IOW1_17 | 25.5 | 4 | 1 | 93 |
| IOW1_17 | 27 | 4 | 0 | 101 |
| Ara | 68 | 1 | 4 | 95 |
| Ara | 72.5 | 1 | 2 | 92 |
| Ari | 84 | 1 | 1 | 90 |
| Ari | 68 | 1 | 3 | 93 |
| Ari | 57.2 | 4 | 3 | 86 |
| Ari | 67 | 1 | 4 | 85 |
| Ble | 51 | 1 | 2 | 100 |
| Ble | 37.5 | 1 | 4 | 94 |
| Ble | 66 | 2 | 5 | 102 |
| Ble | 59 | 1 | 5 | 93 |
| Bol | 65.2 | 1 | 4 | 90 |
| Bol | 58 | 1 | 4 | 92 |
| Ede | 79 | 1 | 2 | 97 |
| Ede | 70.5 | 1 | 2 | 99 |
| Gis | 46 | 2 | 2 | 102 |
| Gis | 56 | 3 | 1 | 93 |
| Gis | 50 | 3 | 3 | 93 |

| | | | | |
|-----|------|---|----|-----|
| Gis | 66 | 3 | 4 | 106 |
| Lir | 18 | 1 | 0 | 77 |
| Lir | 52 | 3 | 5 | 86 |
| Mar | 41 | 1 | 3 | 83 |
| Mar | 43.5 | 1 | 6 | 93 |
| Mon | 52.5 | 1 | 4 | 96 |
| Mon | 81.5 | 2 | 6 | 90 |
| Mon | 71 | 1 | 6 | 82 |
| Mon | 56 | 1 | 3 | 97 |
| Olg | 74 | 1 | 2 | 100 |
| Olg | 74 | 1 | 1 | 92 |
| Ome | 25 | 1 | 2 | 82 |
| Ome | 28.5 | 1 | 1 | 83 |
| Pri | 47 | 3 | 6 | 87 |
| Pri | 41.5 | 3 | 4 | 94 |
| Pri | 42 | 2 | 5 | 97 |
| Pri | 23.5 | 9 | 2 | 123 |
| Rab | 62 | 1 | 2 | 94 |
| Rab | 47.5 | 1 | 4 | 90 |
| Rab | 42 | 3 | 6 | 97 |
| Rab | 49.3 | 3 | 11 | 93 |
| Suz | 56 | 1 | 4 | 97 |
| Suz | 56 | 1 | 3 | 94 |
| Tin | 45 | 2 | 1 | 97 |
| Tin | 46 | 1 | 2 | 92 |
| Tin | 53.5 | 1 | 1 | 104 |
| Tin | 63 | 3 | 3 | 93 |
| Vol | 36.5 | 2 | 3 | 95 |
| Vol | 29 | 1 | 2 | 96 |

6422

6423 **Appendix 32: Macro for ImageJ processing (Chapter 5)**

6424 #Opening file and setting known distance unit:

6425 open("D:\\Linum Project\\Cap and seed photos\\Seed photos Aug 2018\\1_1 Seeds.tif");

6426 makeLine(996, 210, 1014, 324);

6427 run("Set Scale...", "distance=115.41 known=1 pixel=1 unit=mm");

6428 run("Split Channels");

6429 selectWindow("1_1 Seeds.tif (blue)");

6430 run("Color Balance...");

6431 run("Apply LUT");

```
6432 run("Analyze Particles...", "size=25-Infinity pixel circularity=0.00-1.00 show=Outlines display");
6433 setAutoThreshold("Default");
6434 //run("Threshold...");
6435 setThreshold(0, 254);
6436 run("Convert to Mask");
6437 setAutoThreshold("Default");
6438 setThreshold(1, 255);
6439 run("Convert to Mask");
6440 run("Analyze Particles...", "size=25-Infinity pixel circularity=0.00-1.00 show=Outlines display");
6441 saveAs("Results", "D:\\Linum Project\\Cap and seed photos\\Seed photos Aug 2018\\1_1 Seeds.xls");
6442
6443 # _____
6444 _____
6445
6446 #Aplying LUT and making binary:
6447
6448 run("Apply LUT");
6449 run("Make Binary");
6450 run("Analyze Particles...", "size=25-Infinity pixel circularity=0.00-1.00 show=Outlines display");
6451
6452 _____
```

6453 **BIBLIOGRAPHY**

6454

6455 Adams, J., 2009. Vegetation-climate interaction: how plants make the global environment. Springer Science &
6456 Business Media.

6457 Addicott, F.T. (1977) 'Flower Behavior in *Linum lewisii*: Some Ecological and Physiological Factors in Opening
6458 and Abscission of Petals', *American Midland Naturalist*, 97(2), p. 321. Available at:
6459 <https://doi.org/10.2307/2425097>.

6460 Adhikari, K.N., Buirchell, B.J. and Sweetingham, M.W. (2012) 'Length of vernalization period affects flowering
6461 time in three lupin species', *Plant Breeding*, 131(5), pp. 631–636. Available at: <https://doi.org/10.1111/J.1439-0523.2012.01996.X>.

6463 Adhikari, P.B., Liu, X. and Kasahara, R.D. (2020) 'Mechanics of Pollen Tube Elongation: A Perspective', *Frontiers
6464 in Plant Science*, 11, p. 1612. Available at: <https://doi.org/10.3389/FPLS.2020.589712/BIBTEX>.

6465 Alcalá, N. *et al.* (2013) 'Peak and persistent excess of genetic diversity following an abrupt migration increase',
6466 *Genetics*, 193(3), pp. 953–971. Available at: [https://doi.org/10.1534/GENETICS.112.147785/-](https://doi.org/10.1534/GENETICS.112.147785/-/DC1/GENETICS.112.147785-4.PDF)
6467 [/DC1/GENETICS.112.147785-4.PDF](https://doi.org/10.1534/GENETICS.112.147785-4.PDF).

6468 Alexander, D.H., Novembre, J. and Lange, K. (2009) 'Fast model-based estimation of ancestry in unrelated
6469 individuals', *Genome Research*, 19(9), pp. 1655–1664. Available at: <https://doi.org/10.1101/GR.094052.109>.

6470 Allaby, R.G. *et al.* (2005) 'Evidence of the domestication history of flax (*Linum usitatissimum* L.) from genetic
6471 diversity of the *sad2* locus', *Theoretical and Applied Genetics*, 112(1), pp. 58–65. Available at:
6472 <https://doi.org/10.1007/S00122-005-0103-3/TABLES/3>.

6473 Allard, A. *et al.* (2016) 'Detecting QTLs and putative candidate genes involved in budbreak and flowering time
6474 in an apple multiparental population', *Journal of Experimental Botany*, 67(9), pp. 2875–2888. Available at:
6475 <https://doi.org/10.1093/JXB/ERW130>.

6476 Alonso, C. *et al.* (2013) 'Among-species differences in pollen quality and quantity limitation: Implications for
6477 endemics in biodiverse hotspots', *Annals of Botany*, 112(7), pp. 1461–1469. doi:10.1093/aob/mct213.

6478 'An Introduction to Generalized Linear Models' (2018) *An Introduction to Generalized Linear Models, Fourth
6479 Edition* [Preprint]. Available at: <https://doi.org/10.1201/9781315182780>.

6480 Analysis of relative gene expression using different real-time quantitative PCR - Abstract - Europe PMC (no
6481 date). Available at: <https://europepmc.org/article/cba/641544> (Accessed: 10 June 2022).

6482 Anderson, J.T., Willis, J.H. and Mitchell-Olds, T. (2011) 'Evolutionary genetics of plant adaptation', *Trends in
6483 Genetics*, 27(7), pp. 258–266. Available at: <https://doi.org/10.1016/J.TIG.2011.04.001>.

6484 Andrews, K.R. *et al.* (no date) 'Harnessing the power of RADseq for ecological and evolutionary genomics'.
6485 Available at: <https://doi.org/10.1101/028837>.

6486 Arnold, B. *et al.* (2013) 'RADseq underestimates diversity and introduces genealogical biases due to
6487 nonrandom haplotype sampling', *Molecular Ecology*, 22(11), pp. 3179–3190. Available at:
6488 <https://doi.org/10.1111/MEC.12276>.

6489 Asíns, M.J. (2002) 'Present and future of quantitative trait locus analysis in plant breeding', *Plant Breeding*, pp.
6490 281–291. Available at: <https://doi.org/10.1046/j.1439-0523.2002.730285.x>.

6491 Ataii, E. *et al.* (2021) 'Genetic characterization of a diversity panel, selected from IPK linseed (*Linum*
6492 *usitatissimum*) world collection', *Plant Breeding*, 140(5), pp. 919–932. Available at:
6493 <https://doi.org/10.1111/pbr.12962>.

6494 ATES, E. (2012) 'Some chemical and morphological properties of five clover species (*Trifolium* sp.) at different
6495 aspect of pasture in Belovets village (Razgrad), Bulgaria', *International Journal of Plant Production*, 5(3), pp.
6496 255–262. Available at: <https://doi.org/10.22069/IJPP.2012.737>.

6497 Atkins, K.E. and Travis, J.M.J. (2010) 'Local adaptation and the evolution of species' ranges under climate
6498 change', *Journal of Theoretical Biology*, 266(3), pp. 449–457. Available at:
6499 <https://doi.org/10.1016/J.JTBI.2010.07.014>.

6500 Auge, G.A. *et al.* (2017) 'Maternal vernalization and vernalization-pathway genes influence progeny seed
6501 germination', *New Phytologist*, 216(2), pp. 388–400. Available at: <https://doi.org/10.1111/nph.14520>.

6502 Bacilieri, R. *et al.* (2013) 'Genetic structure in cultivated grapevines is linked to geography and human
6503 selection', *BMC Plant Biology*, 13(1), pp. 1–14. Available at: [https://doi.org/10.1186/1471-2229-13-](https://doi.org/10.1186/1471-2229-13-25/TABLES/5)
6504 [25/TABLES/5](https://doi.org/10.1186/1471-2229-13-25/TABLES/5).

6505 Baiakhmetov, E. *et al.* (2021) 'Evidence for extensive hybridisation and past introgression events in feather
6506 grasses using genome-wide SNP genotyping', *BMC Plant Biology*, 21(1), pp. 1–20. Available at:
6507 <https://doi.org/10.1186/S12870-021-03287-W/TABLES/3>.

6508 Ballerini, E.S. and Kramer, E.M. (2011) 'In the light of evolution: A reevaluation of conservation in the CO-FT
6509 regulon and its role in photoperiodic regulation of flowering time', *Frontiers in Plant Science*, 2(NOV), p. 81.
6510 Available at: <https://doi.org/10.3389/FPLS.2011.00081/BIBTEX>.

6511 Barboni, D. *et al.* (2004) 'Relationships between plant traits and climate in the Mediterranean region: A pollen
6512 data analysis', *Journal of Vegetation Science*, 15(5), pp. 635–646. doi:10.1111/j.1654-1103.2004.tb02305.x.

6513 Bay, R.A. *et al.* (2018) *Genomic signals of selection predict climate-driven population declines in a migratory*
6514 *bird*.

6515 Behjati, S. and Tarpey, P.S. (2013) 'What is next generation sequencing?', *Archives of Disease in Childhood*.
6516 *Education and Practice Edition*, 98(6), p. 236. Available at: [https://doi.org/10.1136/ARCHDISCHILD-2013-](https://doi.org/10.1136/ARCHDISCHILD-2013-304340)
6517 [304340](https://doi.org/10.1136/ARCHDISCHILD-2013-304340).

6518 Beltrán, R. *et al.* (2019) 'Effect of temperature on pollen germination for several Rosaceae species: Influence of
6519 freezing conservation time on germination patterns', *PeerJ*, 2019(12). Available at:
6520 <https://doi.org/10.7717/PEERJ.8195/SUPP-1>.

6521 Bigot, S. *et al.* (2018) 'Pivotal roles of environmental sensing and signaling mechanisms in plant responses to
6522 climate change', *Global Change Biology*, 24(12), pp. 5573–5589. Available at:
6523 <https://doi.org/10.1111/GCB.14433>.

6524 Bilton, T.P. *et al.* (2018) 'Linkage Disequilibrium Estimation in Low Coverage High-Throughput Sequencing
6525 Data', *Genetics*, 209(2), p. 389. Available at: <https://doi.org/10.1534/GENETICS.118.300831>.

6526 Bird, K.A. *et al.* (2017) 'Population structure and phylogenetic relationships in a diverse panel of *Brassica rapa*
6527 L.', *Frontiers in Plant Science*, 8. Available at: <https://doi.org/10.3389/fpls.2017.00321>.

6528 Blackman, B.K., Michaels, S.D. and Rieseberg, L.H. (2011) 'Connecting the sun to flowering in sunflower
6529 adaptation', *Molecular ecology*, 20(17), pp. 3503–3512. Available at: [https://doi.org/10.1111/J.1365-
6530 294X.2011.05166.X](https://doi.org/10.1111/J.1365-294X.2011.05166.X).

6531 Blanquart, F. *et al.* (2013) 'A practical guide to measuring local adaptation', *Ecology Letters*, 16(9), pp. 1195–
6532 1205. Available at: <https://doi.org/10.1111/ELE.12150>.

6533 Blázquez, M., Koornneef, M. and Putterill, J. (2001) 'Flowering on time: genes that regulate the floral
6534 transition: Workshop on the molecular basis of flowering time control', *EMBO Reports*, 2(12), p. 1078.
6535 Available at: <https://doi.org/10.1093/EMBO-REPORTS/KVE254>.

6536 Blümel, M., Dally, N. and Jung, C. (2015) 'Flowering time regulation in crops — what did we learn from
6537 *Arabidopsis*?', *Current Opinion in Biotechnology*, 32, pp. 121–129. Available at:
6538 <https://doi.org/10.1016/J.COPBIO.2014.11.023>.

6539 Boavida, L.C. and McCormick, S. (2007) 'TECHNICAL ADVANCE: Temperature as a determinant factor for
6540 increased and reproducible in vitro pollen germination in *Arabidopsis thaliana*', *The Plant Journal*, 52(3), pp.
6541 570–582. Available at: <https://doi.org/10.1111/J.1365-313X.2007.03248.X>.

6542 Bolsheva, N.L. *et al.* (2017) 'Evolution of blue-flowered species of genus *Linum* based on high-throughput
6543 sequencing of ribosomal RNA genes', *BMC Evolutionary Biology*, 17(2), pp. 23–36. Available at:
6544 <https://doi.org/10.1186/S12862-017-1105-X/FIGURES/6>.

6545 Booker, T.R., Jackson, B.C. and Keightley, P.D. (2017) 'Detecting positive selection in the genome', *BMC*
6546 *Biology*, 15(1), pp. 1–10. Available at: <https://doi.org/10.1186/S12915-017-0434-Y/FIGURES/2>.

6547 Boshier, D. *et al.* (2015) 'Is local best? examining the evidence for local adaptation in trees and its scale',
6548 *Environmental Evidence*, 4(1). doi:10.1186/s13750-015-0046-3.

6549 Bougeard, S. and Dray, S. (2018) 'Supervised Multiblock Analysis in R with the *ade4* Package', *Journal of*
6550 *Statistical Software*, 86, pp. 1–17. Available at: <https://doi.org/10.18637/JSS.V086.I01>.

6551 BOX, G.E.P. (1953) 'NON-NORMALITY AND TESTS ON VARIANCES', *Biometrika*, 40(3–4), pp. 318–335. Available
6552 at: <https://doi.org/10.1093/BIOMET/40.3-4.318>.

6553 Božič, A. and Šiber, A. (2022) 'Mechanics of inactive swelling and bursting of porate pollen grains', *Biophysical*
6554 *Journal*, 121(5), pp. 782–792. Available at: <https://doi.org/10.1016/J.BPJ.2022.01.019>.

6555 Brandon, R.N., 2014. *Adaptation and environment* (Vol. 1040). Princeton University Press.

6556 Brennan, A.C. *et al.* (2014) 'The genetic structure of *Arabidopsis thaliana* in the south-western Mediterranean
6557 range reveals a shared history between North Africa and southern Europe', *BMC Plant Biology*, 14(1), pp. 1–14.
6558 Available at: <https://doi.org/10.1186/1471-2229-14-17/FIGURES/4>.

6559 Breygina, M., Klimenko, E. and Schekaleva, O. (2021) 'Pollen Germination and Pollen Tube Growth in
6560 Gymnosperms', *Plants*, 10(7). Available at: <https://doi.org/10.3390/PLANTS10071301>.

6561 Buira, A. *et al.* (2020) 'The contribution of the edaphic factor as a driver of recent plant diversification in a
6562 Mediterranean biodiversity hotspot', *Journal of Ecology*, 109(2), pp. 987–999. doi:10.1111/1365-2745.13527.

6563 Bustin, S.A. *et al.* (2009) 'The MIQE guidelines: minimum information for publication of quantitative real-time
6564 PCR experiments', *Clinical chemistry*, 55(4), pp. 611–622. Available at:
6565 <https://doi.org/10.1373/CLINCHEM.2008.112797>.

6566 Butte, A.J., Dzau, V.J. and Glueck, S.B. (2001) 'Further defining housekeeping, or "maintenance," genes Focus
6567 on "A compendium of gene expression in normal human tissues"',
6568 <https://doi.org/10.1152/physiolgenomics.2001.7.2.95>, 7(2), pp. 95–96. Available at:
6569 <https://doi.org/10.1152/PHYSIOLGENOMICS.2001.7.2.95>.

6570 Calvin, J.A. (1998) 'Regression Models for Categorical and Limited Dependent Variables', *Technometrics*, 40(1),
6571 pp. 80–81. Available at: <https://doi.org/10.1080/00401706.1998.10485496>.

6572 Campetella, G. *et al.* (2020) 'Plant functional traits are correlated with species persistence in the herb layer of
6573 old-growth beech forests', *Scientific Reports*, 10(1). Available at: <https://doi.org/10.1038/S41598-020-76289-7>.

6574 Carmona, R. *et al.* (2017) 'Automated identification of reference genes based on RNA-seq data', *BioMedical*
6575 *Engineering OnLine*, 16(Suppl 1). Available at: <https://doi.org/10.1186/S12938-017-0356-5>.

6576 Catchen, J. *et al.* (2013) 'Stacks: an analysis tool set for population genomics', *Molecular ecology*, 22(11), pp.
6577 3124–3140. Available at: <https://doi.org/10.1111/MEC.12354>.

6578 Cave, R.L. and Johnston, M.E. (2010) "Vernalization promotes flowering of a heat tolerant *Calandrinia* while
6579 long days replace vernalization for early flowering of *brunonia*," *Scientia Horticulturae*, 123(3), pp. 379–384.
6580 Available at: <https://doi.org/10.1016/j.scienta.2009.10.004>.

6581 Cave, R.L. *et al.* (2013) 'Modelling temperature, photoperiod and vernalization responses of *Brunonia australis*
6582 (*Goodeniaceae*) and *Calandrinia* sp. (*Portulacaceae*) to predict flowering time', *Annals of botany*, 111(4), pp.
6583 629–639. Available at: <https://doi.org/10.1093/AOB/MCT028>.

6584 Chambers, J. and Hastie, T., 1992. *Statistical models in S*. London: Chapman & Hall

6585 Chandler, J., Wilson, A. and Dean, C. (1996) 'Arabidopsis mutants showing an altered response to
6586 vernalization', *The Plant Journal*, 10(4), pp. 637–644. Available at: [https://doi.org/10.1046/J.1365-](https://doi.org/10.1046/J.1365-313X.1996.10040637.X)
6587 313X.1996.10040637.X.

6588 Chapman, J.R. *et al.* (2009) 'A quantitative review of heterozygosity–fitness correlations in animal populations',
6589 *Molecular Ecology*, 18(13), pp. 2746–2765. Available at: <https://doi.org/10.1111/J.1365-294X.2009.04247.X>.

6590 Chen, C.-C. and Gibson, P.B. (1973) 'Effect of Temperature on Pollen-Tube Growth in *Trifolium repense* after
6591 Cross- and Self-Pollinations1', *Crop Science*, 13(5), pp. 563–566. Available at:
6592 <https://doi.org/10.2135/CROPSCI1973.0011183X001300050021X>.

6593 Chen, D. *et al.* (2018) 'Architecture of gene regulatory networks controlling flower development in *Arabidopsis*
6594 *thaliana*', *Nature Communications* 2018 9:1, 9(1), pp. 1–13. Available at: [https://doi.org/10.1038/s41467-018-](https://doi.org/10.1038/s41467-018-06772-3)
6595 06772-3.

6596 Chiluwal, A. *et al.* (2020) 'Deterioration of ovary plays a key role in heat stress-induced spikelet sterility in
6597 sorghum', *Plant Cell and Environment*, 43(2), pp. 448–462. Available at: <https://doi.org/10.1111/pce.13673>.

6598 Cho, L.H., Yoon, J. and An, G. (2017) 'The control of flowering time by environmental factors', *The Plant*
6599 *Journal*, 90(4), pp. 708–719. Available at: <https://doi.org/10.1111/TPJ.13461>.

6600 Chouard, P. (1960) 'Vernalization and its Relations to Dormancy', *Annual Review of Plant Physiology*, 11(1), pp.
6601 191–238. Available at: <https://doi.org/10.1146/ANNUREV.PP.11.060160.001203>.

6602 Clarke, J.D. (2009) 'Cetyltrimethyl Ammonium Bromide (CTAB) DNA Miniprep for Plant DNA Isolation', *Cold*
6603 *Spring Harbor Protocols*, 2009(3), p. pdb.prot5177. Available at: <https://doi.org/10.1101/PDB.PROT5177>.

6604 Cloutier, S. *et al.* (2009) 'Development and analysis of EST-SSRs for flax (*Linum usitatissimum* L.)', *Theoretical*
6605 *and Applied Genetics*, 119(1), pp. 53–63. Available at: <https://doi.org/10.1007/S00122-009-1016-3/FIGURES/4>.

6606 Cloutier, S. *et al.* (2012) 'Integrated consensus genetic and physical maps of flax (*Linum usitatissimum* L.)',
6607 *Theoretical and Applied Genetics* 2012 125:8, 125(8), pp. 1783–1795. Available at:
6608 <https://doi.org/10.1007/S00122-012-1953-0>.

6609 Clua, A.A. and Gimenez, D.O. (2003) 'Environmental factors during seed development of narrow-leaved bird's-
6610 foot-trefoil (*Lotus tenuis*) influences subsequent dormancy and germination', *Grass and Forage Science*, 58(4),
6611 pp. 333–338. Available at: <https://doi.org/10.1046/j.1365-2494.2003.00385.x>.

6612 Conner, J.K. (2002) 'Genetic mechanisms of floral trait correlations in a natural population', *Nature*, 420(6914),
6613 pp. 407–410. Available at: <https://doi.org/10.1038/NATURE01105>.

6614 Corbesier, L. and Coupland, G. (2006) 'The quest for florigen: a review of recent progress', *Journal of*
6615 *Experimental Botany*, 57(13), pp. 3395–3403. Available at: <https://doi.org/10.1093/JXB/ERL095>.

6616 Cox, M. and Doudna, J., 2015. *Molecular Biology*:2nd Revised edition. New York: W.H. FREEMAN & CO LTD,
6617 pp.221-238, 369-376, 592-593.

6618 Çopur, O., Gur, M.A., Karakus, M. and Demirel, U., 2006. Determination of correlation and path analysis among
6619 yield components and seed yield in oil flax varieties (*Linum usitatissimum* L.). *Journal of Biological Sciences*,
6620 6(4), pp.738-743.

6621 Craufurd, P.Q. and Wheeler, T.R. (2009) 'Climate change and the flowering time of annual crops', *Journal of*
6622 *experimental botany*, 60(9), pp. 2529–2539. Available at: <https://doi.org/10.1093/JXB/ERP196>.

6623 Crean, A.J. and Immler, S. (2021) 'Evolutionary consequences of environmental effects on gamete
6624 performance', *Philosophical Transactions of the Royal Society B: Biological Sciences*, 376(1826). Available at:
6625 <https://doi.org/10.1098/RSTB.2020.0122>.

6626 Danecek, P. *et al.* (2011) 'The variant call format and VCFtools', *Bioinformatics*, 27(15), pp. 2156–2158.
6627 Available at: <https://doi.org/10.1093/BIOINFORMATICS/BTR330>.

6628 Darapuneni, M.K. *et al.* (2014) 'Effect of vernalization and photoperiod on flax flowering time', *Euphytica*,
6629 195(2), pp. 279–285. Available at: <https://doi.org/10.1007/s10681-013-0996-x>.

6630 Davey, J. W., & Blaxter, M. L. (2010). RADSeq: next-generation population genetics. *Briefings in Functional*
6631 *Genomics*, 9(5-6), 416–423. <https://doi.org/10.1093/bfpg/elq031>

6632 Davey, J.W. *et al.* (2011) 'Genome-wide genetic marker discovery and genotyping using next-generation
6633 sequencing', *Nature Reviews Genetics* 2011 12:7, 12(7), pp. 499–510. Available at:
6634 <https://doi.org/10.1038/nrg3012>.

6635 Debieu, M. *et al.* (2013) 'Co-Variation between Seed Dormancy, Growth Rate and Flowering Time Changes with
6636 Latitude in *Arabidopsis thaliana*', *PLoS ONE*, 8(5). Available at: <https://doi.org/10.1371/journal.pone.0061075>.

6637 Delph, L. and Havens, K., 1998. *Sperm Competition and Sexual Selection*. Elsevier Inc. Pp.149-173.

6638 Dempewolf, H. *et al.* (2017) 'Past and Future Use of Wild Relatives in Crop Breeding', *Crop Science*, 57(3), pp.
6639 1070–1082. Available at: <https://doi.org/10.2135/CROPSCI2016.10.0885>.

6640 Descamps, C., Quinet, M. and Jacquemart, A.L. (2021) 'Climate Change–Induced Stress Reduce Quantity and
6641 Alter Composition of Nectar and Pollen From a Bee-Pollinated Species (*Borago officinalis*, Boraginaceae)',
6642 *Frontiers in Plant Science*, 12, p. 2264. Available at: <https://doi.org/10.3389/FPLS.2021.755843/BIBTEX>.

6643 Devasirvatham, V. *et al.* (2012) 'Effect of high temperature on the reproductive development of chickpea
6644 genotypes under controlled environments', *Functional Plant Biology*, 39(12), pp. 1009–1018. Available at:
6645 <https://doi.org/10.1071/FP12033>.

6646 Diamond, J. (2002) 'Evolution, consequences and future of plant and animal domestication', *Nature*,
6647 418(6898), pp. 700–707. Available at: <https://doi.org/10.1038/NATURE01019>.

6648 Dickinson, H., Rodriguez-Enriquez, J. and Grant-Downton, R. (2018) 'Pollen Germination and Pollen Tube
6649 Growth of *Arabidopsis thaliana*: in vitro and Semi in vivo Methods', *Bio-protocol*, 8(16). Available at:
6650 <https://doi.org/10.21769/BIOPROTOC.2977>.

6651 Diederichsen, A. (2001) 'Comparison of genetic diversity of flax (*Linum usitatissimum* L.) between Canadian
6652 cultivars and a world collection', *Plant Breeding*, 120(4), pp. 360–362. Available at:
6653 <https://doi.org/10.1046/J.1439-0523.2001.00616.X>.

6654 Diederichsen, A. and Raney, J.P. (2006) 'Seed colour, seed weight and seed oil content in *Linum usitatissimum*
6655 accessions held by Plant Gene Resources of Canada', *Plant Breeding*, 125(4), pp. 372–377. Available at:
6656 <https://doi.org/10.1111/J.1439-0523.2006.01231.X>.

6657 Diederichsen, A. and Ulrich, A. (2009) 'Variability in stem fibre content and its association with other
6658 characteristics in 1177 flax (*Linum usitatissimum* L.) genebank accessions', *Industrial Crops and Products*, 30(1),
6659 pp. 33–39. Available at: <https://doi.org/10.1016/j.indcrop.2009.01.002>.

6660 Dmitriev, A.A. *et al.* (2021) 'Genome Sequencing of Fiber Flax Cultivar Atlant Using Oxford Nanopore and
6661 Illumina Platforms', *Frontiers in Genetics*, 11, p. 1487. Available at:
6662 <https://doi.org/10.3389/FGENE.2020.590282/BIBTEX>.

6663 Dole, J.M. and Wilkins, H.F. (1994) "Interaction of bulb vernalization and shoot photoperiod on 'Nellie White'
6664 Easter Lily," *HortScience*, 29(3), pp. 143–145. Available at: <https://doi.org/10.21273/hortsci.29.3.143>.

6665 Donohue, K. *et al.* (2005) 'Environmental and genetic influences on the germination of *Arabidopsis thaliana* in
6666 the field', *Evolution*, 59(4), pp. 740–757. Available at: <https://doi.org/10.1111/J.0014-3820.2005.TB01750.X>.

6667 Donohue, K. *et al.* (2010) 'Germination, postgermination adaptation, and species ecological ranges', *Annual*
6668 *Review of Ecology, Evolution, and Systematics*, 41, pp. 293–319. Available at: [https://doi.org/10.1146/annurev-](https://doi.org/10.1146/annurev-ecolsys-102209-144715)
6669 [ecolsys-102209-144715](https://doi.org/10.1146/annurev-ecolsys-102209-144715).

6670 Dray, S. and Dufour, A.B. (2007) 'The ade4 Package: Implementing the Duality Diagram for Ecologists', *Journal*
6671 *of Statistical Software*, 22(4), pp. 1–20. Available at: <https://doi.org/10.18637/JSS.V022.I04>.

6672 Dudley, S.A. (1996) 'THE RESPONSE TO DIFFERING SELECTION ON PLANT PHYSIOLOGICAL TRAITS: EVIDENCE
6673 FOR LOCAL ADAPTATION', *Evolution*, 50(1), pp. 103–110. Available at: [https://doi.org/10.1111/J.1558-](https://doi.org/10.1111/J.1558-5646.1996.TB04476.X)
6674 [5646.1996.TB04476.X](https://doi.org/10.1111/J.1558-5646.1996.TB04476.X).

6675 Ejsmond, M.J. *et al.* (2011) 'Does climate affect pollen morphology? Optimal size and shape of pollen grains
6676 under various desiccation intensity', *Ecosphere*, 2(10), pp. 1–15. Available at: [https://doi.org/10.1890/ES11-](https://doi.org/10.1890/ES11-00147.1)
6677 [00147.1](https://doi.org/10.1890/ES11-00147.1).

6678 El-Beltagi, H. and Salama, Z. (2007) *Evaluation of fatty acids profile and the content of some secondary*
6679 *metabolites in seeds of different flax cultivars (*Linum usitatissimum* L.) Cytotoxic and antioxidant properties of*
6680 *active principals isolated from water hyacinth against four cancer cells lines View project Ascorbic Acid*

6681 *Enhances Growth and Yield of Sweet Peppers (Capsicum annum) by Mitigating Salinity Stress View project.*
6682 Available at: <https://www.researchgate.net/publication/255606154>.

6683 Elfving, N. *et al.* (2011) 'The Arabidopsis thaliana Med25 mediator subunit integrates environmental cues to
6684 control plant development', *Proceedings of the National Academy of Sciences of the United States of America*,
6685 108(20), pp. 8245–8250. Available at: <https://doi.org/10.1073/PNAS.1002981108>.

6686 Ellstrand, N.C., Prentice, H.C. and Hancock, J.F. (2003) 'Gene Flow and Introgression from Domesticated Plants
6687 into their Wild Relatives', <https://doi.org/10.1146/annurev.ecolsys.30.1.539>, 30, pp. 539–563. Available at:
6688 <https://doi.org/10.1146/ANNUREV.ECOLSYS.30.1.539>.

6689 Essemine, J., Ammar, S. and Bouzid, S. (2010) 'Impact of heat stress on germination and growth in higher
6690 plants: physiological, biochemical and molecular repercussions and mechanisms of defence.', *Journal of*
6691 *Biological Sciences*, 10(6), pp. 565–572.

6692 Fábrián, A. *et al.* (2019) 'Stigma functionality and fertility are reduced by heat and drought co-stress in wheat',
6693 *Frontiers in Plant Science*, 10. Available at: <https://doi.org/10.3389/FPLS.2019.00244/FULL>.

6694 Falconer, D.S., 1996. Introduction to quantitative genetics. Pearson Education India.

6695 FAOSTAT data (2006) Food and Agriculture Organization of the United Nations, FAO Statistical Data-
6696 bases, URL: <http://faostat.fao.org/>

6697 Fedorenko, O.M., Savushkin, A.I. and Olimpienko, G.S. (2001) 'Genetic Diversity in Natural Populations of
6698 Arabidopsis thaliana (L.) Heynh. from Karelia', *Russian Journal of Genetics* 2001 37:2, 37(2), pp. 162–167.
6699 Available at: <https://doi.org/10.1023/A:1009041922186>.

6700 Firon, N., Nepi, M. and Pacini, E. (2012) 'Water status and associated processes mark critical stages in pollen
6701 development and functioning', *Annals of Botany*, 109(7), pp. 1201–1214. Available at:
6702 <https://doi.org/10.1093/AOB/MCS070>.

6703 Flowers, J.M. *et al.* (2009) 'Population Genomics of the Arabidopsis thaliana Flowering Time Gene Network',
6704 *Molecular Biology and Evolution*, 26(11), pp. 2475–2486. Available at:
6705 <https://doi.org/10.1093/MOLBEV/MSP161>.

6706 Fragallah, S.A.D.A. *et al.* (2019) 'Effects of Sucrose, Boric Acid, pH, and Incubation Time on in Vitro Germination
6707 of Pollen and Tube Growth of Chinese fir (Cunninghamia lanceolata L.)', *Forests* 2019, Vol. 10, Page 102, 10(2),
6708 p. 102. Available at: <https://doi.org/10.3390/F10020102>.

6709 Frichot, E. *et al.* (2014) 'Fast and efficient estimation of individual ancestry coefficients', *Genetics*, 196(4), pp.
6710 973–983. Available at: <https://doi.org/10.1534/GENETICS.113.160572/-/DC1>.

6711 Frichot, E. and François, O. (2015) 'LEA: An R package for landscape and ecological association studies',
6712 *Methods in Ecology and Evolution*, 6(8), pp. 925–929. Available at: <https://doi.org/10.1111/2041-210X.12382>.

6713 Fu, Y.-B., Diederichsen, A. and Allaby, R.G. (2011) "Locus-specific view of flax domestication history," *Ecology*
6714 *and Evolution*, 2(1), pp. 139–152. Available at: <https://doi.org/10.1002/ece3.57>.

6715 Fu, Y.B. (2012) 'Population-based resequencing revealed an ancestral winter group of cultivated flax:
6716 implication for flax domestication processes', *Ecology and Evolution*, 2(3), pp. 622–635. Available at:
6717 <https://doi.org/10.1002/ECE3.101>.

6718 Fu, Y.X. (1995) 'Statistical Properties of Segregating Sites', *Theoretical Population Biology*, 48(2), pp. 172–197.
6719 Available at: <https://doi.org/10.1006/TPBI.1995.1025>.

6720 Furness, C.A. and Rudall, P.J. (2004) 'Pollen aperture evolution – a crucial factor for eudicot success?', *Trends in*
6721 *Plant Science*, 9(3), pp. 154–158. Available at: <https://doi.org/10.1016/J.TPLANTS.2004.01.001>.

6722 Galloway, L.F. and Burgess, K.S. (2009) 'Manipulation of flowering time: phenological integration and maternal
6723 effects', *Ecology*, 90(8), pp. 2139–2148. Available at: <https://doi.org/10.1890/08-0948.1>.

6724 Galloway, L.F. and Burgess, K.S. (2012) 'Artificial selection on flowering time: influence on reproductive
6725 phenology across natural light environments', *Journal of Ecology*, 100(4), pp. 852–861. Available at:
6726 <https://doi.org/10.1111/J.1365-2745.2012.01967.X>.

6727 Gao, Y.-B. et al. (2014) "Low temperature inhibits pollen tube growth by disruption of both tip-localized
6728 reactive oxygen species and endocytosis in *Pyrus bretschneideri* Rehd.," *Plant Physiology and Biochemistry*, 74,
6729 pp. 255–262. Available at: <https://doi.org/10.1016/j.plaphy.2013.11.018>.

6730 Gargiulo, R., Kull, T. and Fay, M.F. (2021) 'Effective double-digest RAD sequencing and genotyping despite large
6731 genome size', *Molecular Ecology Resources*, 21(4), pp. 1037–1055. Available at: <https://doi.org/10.1111/1755-0998.13314>.

6733 Gelman, A. and Hill, J., 2006. *Data analysis using regression and multilevel/hierarchical models*. Cambridge
6734 university press.

6735 Gerristen, H., 2018. CRAN - Package mapplots. [online] [Cran.r-project.org](https://cran.r-project.org/web/packages/mapplots/index.html). Available at: <https://cran.r-project.org/web/packages/mapplots/index.html> [Accessed 22 June 2022].

6737 Gilad, Y. et al. (2002) 'Evidence for positive selection and population structure at the human MAO-A gene',
6738 *Proceedings of the National Academy of Sciences*, 99(2), pp. 862–867. Available at:
6739 <https://doi.org/10.1073/PNAS.022614799>.

6740 Giles, B.E. (1990) 'The effects of variation in seed size on growth and reproduction in the wild barley *Hordeum*
6741 *vulgare* ssp. *spontaneum*', *Heredity* 1990 64:2, 64(2), pp. 239–250. Available at:
6742 <https://doi.org/10.1038/hdy.1990.29>.

6743 Gill KS (1966) *Evolutionary relationships among Linum species*. Ph.D. Thesis, University of California, Riverside,
6744 USA

6745 Giménez-Benavides, L., Escudero, A. and Iriondo, J.M. (2007) 'Reproductive limits of a late-flowering high-
6746 mountain Mediterranean plant along an elevational climate gradient', *New Phytologist*, 173(2), pp. 367–382.
6747 Available at: <https://doi.org/10.1111/J.1469-8137.2006.01932.X>.

6748 Ginzinger, D.G. (2002) 'Gene quantification using real-time quantitative PCR: An emerging technology hits the
6749 mainstream', *Experimental Hematology*, 30(6), pp. 503–512. Available at: [https://doi.org/10.1016/S0301-](https://doi.org/10.1016/S0301-472X(02)00806-8)
6750 [472X\(02\)00806-8](https://doi.org/10.1016/S0301-472X(02)00806-8).

6751 Giorno, F. *et al.* (2013) 'Ensuring Reproduction at High Temperatures: The Heat Stress Response during Anther
6752 and Pollen Development', *Plants 2013, Vol. 2, Pages 489-506*, 2(3), pp. 489–506. Available at:
6753 <https://doi.org/10.3390/PLANTS2030489>.

6754 Giriden, E.R., 1992. ANOVA: Repeated measures (No. 84). sage.

6755 Glantz, S., Slinker, B. and Neilands, T., n.d. Primer of applied regression et analysis of variance.

6756 Godwin, I.D., Aitken, E.A.B. and Smith, L.W. (1997) 'Application of inter simple sequence repeat (ISSR) markers
6757 to plant genetics', *ELECTROPHORESIS*, 18(9), pp. 1524–1528. Available at:
6758 <https://doi.org/10.1002/ELPS.1150180906>.

6759 Goudenhoft, C., Bourmaud, A. and Baley, C. (2019) 'Flax (*Linum usitatissimum* L.) fibers for composite
6760 reinforcement: Exploring the link between plant growth, cell walls development, and fiber properties',
6761 *Frontiers in Plant Science*. Frontiers Media S.A. Available at: <https://doi.org/10.3389/fpls.2019.00411>.

6762 Grapin, A. *et al.* (1998) 'Diploid *Musa acuminata* genetic diversity assayed with sequence-tagged microsatellite
6763 sites', *ELECTROPHORESIS*, 19(8–9), pp. 1374–1380. Available at: <https://doi.org/10.1002/ELPS.1150190829>.

6764 Grassein, F., Lavorel, S. and Till-Bottraud, I. (2014) 'The importance of biotic interactions and local adaptation
6765 for plant response to environmental changes: field evidence along an elevational gradient', *Global Change*
6766 *Biology*, 20(5), pp. 1452–1460. Available at: <https://doi.org/10.1111/GCB.12445>.

6767 Greer, S. *et al.* (2010) 'Housekeeping genes; expression levels may change with density of cultured cells',
6768 *Journal of Immunological Methods*, 355(1–2), pp. 76–79. Available at:
6769 <https://doi.org/10.1016/J.JIM.2010.02.006>.

6770 Gregis, V. *et al.* (2009) 'The Arabidopsis floral meristem identity genes AP1, AGL24 and SVP directly repress
6771 class B and C floral homeotic genes', *The Plant Journal*, 60(4), pp. 626–637. Available at:
6772 <https://doi.org/10.1111/J.1365-313X.2009.03985.X>.

6773 Guan, S. *et al.* (2021) 'Genetic analysis and population structure of wild and cultivated wishbone flower
6774 (*Torenia fournieri* Lind.) lines related to specific floral color', *PeerJ*, 7, p. e11702. Available at:
6775 <https://doi.org/10.7717/PEERJ.11702/SUPP-6>.

6776 Günther, T. *et al.* (2016) 'Genomic and phenotypic differentiation of *Arabidopsis thaliana* along altitudinal
6777 gradients in the North Italian Alps', *Molecular Ecology*, 25(15), pp. 3574–3592. Available at:
6778 <https://doi.org/10.1111/MEC.13705>.

6779 Guo, D. *et al.* (2020) 'Resequencing 200 Flax Cultivated Accessions Identifies Candidate Genes Related to Seed
6780 Size and Weight and Reveals Signatures of Artificial Selection', *Frontiers in Plant Science*, 10. Available at:
6781 <https://doi.org/10.3389/fpls.2019.01682>.

6782 Guo, Q. (2012) 'Incorporating latitudinal and central–marginal trends in assessing genetic variation across
6783 species ranges', *Molecular Ecology*, 21(22), pp. 5396–5403. Available at: <https://doi.org/10.1111/MEC.12012>.

6784 Gupta, S. *et al.* (2021) 'Identification and characterization of a novel long juvenile resource AGS 25', *Genetic
6785 Resources and Crop Evolution*, 68(3), pp. 1149–1163. Available at: [https://doi.org/10.1007/S10722-020-01055-
6786 7/TABLES/2](https://doi.org/10.1007/S10722-020-01055-7/TABLES/2).

6787 Gutaker, R.M. *et al.* (2019) 'Flax latitudinal adaptation at LuTFL1 altered architecture and promoted fiber
6788 production', *Scientific Reports 2019 9:1*, 9(1), pp. 1–12. Available at: [https://doi.org/10.1038/s41598-018-
6789 37086-5](https://doi.org/10.1038/s41598-018-37086-5).

6790 Gutiérrez-Valencia, J. *et al.* (2022) 'Genomic analyses of the *Linum distyly* supergene reveal convergent
6791 evolution at the molecular level', *Current Biology*, 32(20), pp. 4360-4371.e6. Available at:
6792 <https://doi.org/10.1016/J.CUB.2022.08.042>.

6793 El Haddad, N. *et al.* (2021) 'Crop wild relatives in durum wheat breeding: Drift or thrift?', *Crop Science*, 61(1),
6794 pp. 37–54. Available at: <https://doi.org/10.1002/CSC2.20223>.

6795 Hall, M.C. and Willis, J.H. (2006) 'DIVERGENT SELECTION ON FLOWERING TIME CONTRIBUTES TO LOCAL
6796 ADAPTATION IN *MIMULUS GUTTATUS* POPULATIONS', *Evolution*, 60(12), pp. 2466–2477. Available at:
6797 <https://doi.org/10.1111/J.0014-3820.2006.TB01882.X>.

6798 Hämälä, T. and Savolainen, O. (2019) 'Genomic Patterns of Local Adaptation under Gene Flow in *Arabidopsis
6799 lyrata*', *Molecular Biology and Evolution*, 36(11), pp. 2557–2571. Available at:
6800 <https://doi.org/10.1093/MOLBEV/MSZ149>.

6801 Hampe, A. and Petit, R.J. (2005) 'Conserving biodiversity under climate change: the rear edge matters', *Ecology
6802 Letters*, 8(5), pp. 461–467. Available at: <https://doi.org/10.1111/J.1461-0248.2005.00739.X>.

6803 Han, Q. *et al.* (2021) 'Association between RsFT, RsFLC and RsCOL5 (A&B) expression and flowering regulation
6804 in Japanese wild radish', *AoB PLANTS*, 13(4). Available at: <https://doi.org/10.1093/aobpla/plab039>.

6805 Hanley, M.E. *et al.* (2008) 'Breeding system, pollinator choice and variation in pollen quality in British
6806 herbaceous plants', *Functional Ecology*, 22(4), pp. 592–598. Available at: [https://doi.org/10.1111/J.1365-
6807 2435.2008.01415.X](https://doi.org/10.1111/J.1365-2435.2008.01415.X).

6808 Harvey, M.G. *et al.* (2016) "Sequence capture versus restriction site associated DNA sequencing for Shallow
6809 Systematics," *Systematic Biology*, 65(5), pp. 910–924. Available at: <https://doi.org/10.1093/sysbio/syw036>.

6810 Hayat, M.J. and Higgins, M. (2014) 'Understanding poisson regression', *Journal of Nursing Education*, 53(4), pp.
6811 207–215. Available at: <https://doi.org/10.3928/01484834-20140325-04>.

6812 He, Y., Chen, T. and Zeng, X. (2020) 'Genetic and Epigenetic Understanding of the Seasonal Timing of
6813 Flowering', *Plant Communications*, 1(1). Available at:
6814 [https://doi.org/10.1016/J.XPLC.2019.100008/GENETIC_AND_EPIGENETIC_UNDERSTANDING_OF_THE_SEASON
6815 AL_TIMING_OF_FLOWERING.PDF](https://doi.org/10.1016/J.XPLC.2019.100008/GENETIC_AND_EPIGENETIC_UNDERSTANDING_OF_THE_SEASONAL_TIMING_OF_FLOWERING.PDF).

6816 Hedhly, A., Hormaza, J.I. and Herrero, M. (2004) 'Effect of temperature on pollen tube kinetics and dynamics in
6817 sweet cherry, *Prunus avium* (Rosaceae)', *American Journal of Botany*, 91(4), pp. 558–564. Available at:
6818 <https://doi.org/10.3732/ajb.91.4.558>.

6819 Hedhly, A., Hormaza, J.I. and Herrero, M. (2005) 'The effect of temperature on pollen germination, pollen tube
6820 growth, and stigmatic receptivity in peach', *Plant Biology*, 7(5), pp. 476–483. Available at:
6821 <https://doi.org/10.1055/S-2005-865850>.

6822 Heisler, M.G. and Jönsson, H. (2007) 'Modelling meristem development in plants', *Current Opinion in Plant*
6823 *Biology*, 10(1), pp. 92–97. Available at: <https://doi.org/10.1016/J.PBI.2006.11.005>.

6824 Herburger, K. and Holzinger, A. (2016) 'Aniline blue and Calcofluor white staining of callose and cellulose in the
6825 streptophyte green algae *Zygnema* and *Klebsormidium*', *Bio-protocol*, 6(20). Available at:
6826 <https://doi.org/10.21769/BIOPROTOCOL.1969>.

6827 Hernández, F. *et al.* (2019) 'Genetic Diversity and Population Structure of Wild Sunflower (*Helianthus annuus*
6828 L.) in Argentina: Reconstructing Its Invasion History', *Journal of Heredity*, 110(6), pp. 746–759. Available at:
6829 <https://doi.org/10.1093/JHERED/ESZ047>.

6830 Hernandez, R.D. *et al.* (2019) 'Ultrarare variants drive substantial cis heritability of human gene expression',
6831 *Nature Genetics* 2019 51:9, 51(9), pp. 1349–1355. Available at: <https://doi.org/10.1038/s41588-019-0487-7>.

6832 Higgins, J.A., Bailey, P.C. and Laurie, D.A. (2010) 'Comparative genomics of flowering time pathways using
6833 brachypodium distachyon as a model for the temperate Grasses', *PLoS ONE*, 5(4). Available at:
6834 <https://doi.org/10.1371/journal.pone.0010065>.

6835 Higuchi, R. *et al.* (1992) 'Simultaneous amplification and detection of specific DNA sequences', *Bio/technology*
6836 (*Nature Publishing Company*), 10(4), pp. 413–417. Available at: <https://doi.org/10.1038/NBT0492-413>.

6837 Hjelmqvist, H. (1950) 'The flax weeds and the origin of cultivated flax.', *Botaniska Notiser*, 2, pp. 257–98.

6838 Hoban, S. *et al.* (2016) 'Finding the genomic basis of local adaptation: Pitfalls, practical solutions, and future
6839 directions', *American Naturalist*, 188(4), pp. 379–397. Available at:
6840 <https://doi.org/10.1086/688018/ASSET/IMAGES/LARGE/FG4.JPEG>.

6841 Hopkins, R., Schmitt, J. and Stinchcombe, J.R. (2008) 'A latitudinal cline and response to vernalization in leaf
6842 angle and morphology in *Arabidopsis thaliana* (Brassicaceae)', *New Phytologist*, 179(1), pp. 155–164. Available
6843 at: <https://doi.org/10.1111/J.1469-8137.2008.02447.X>.

6844 Hothorn, T., Bretz, F. and Westfall, P. (2008a) 'Simultaneous Inference in General Parametric Models',
6845 *Biometrical Journal*, 50(3), pp. 346–363. Available at: <https://doi.org/10.1002/BIMJ.200810425>.

6846 Hothorn, T., Bretz, F. and Westfall, P. (2008b) 'Simultaneous Inference in General Parametric Models',
6847 *Biometrical Journal*, 50(3), pp. 346–363. Available at: <https://doi.org/10.1002/BIMJ.200810425>.

6848 Hu, T. *et al.* (2021) 'Next-generation sequencing technologies: An overview', *Human Immunology*, 82(11), pp.
6849 801–811. Available at: <https://doi.org/10.1016/J.HUMIMM.2021.02.012>.

6850 Huggett, J. *et al.* (2005) 'Real-time RT-PCR normalisation; strategies and considerations', *Genes & Immunity*
6851 2005 6:4, 6(4), pp. 279–284. Available at: <https://doi.org/10.1038/sj.gene.6364190>.

6852 Huis, R., Hawkins, S. and Neutelings, G. (2010) 'Selection of reference genes for quantitative gene expression
6853 normalization in flax (*Linum usitatissimum* L.)', *BMC Plant Biology*, 10(1), pp. 1–14. Available at:
6854 <https://doi.org/10.1186/1471-2229-10-71/FIGURES/6>.

6855 Ibrahim, A. *et al.* (2018) 'A regulator of early flowering in barley (*Hordeum vulgare* L.)', *PLOS ONE*, 13(7), p.
6856 e0200722. Available at: <https://doi.org/10.1371/JOURNAL.PONE.0200722>.

6857 IDT. 2022. Tips for resuspending and diluting your oligonucleotides. [online] Available at:
6858 <[https://eu.idtdna.com/pages/education/decoded/article/tips-for-resuspending-and-diluting-your-](https://eu.idtdna.com/pages/education/decoded/article/tips-for-resuspending-and-diluting-your-oligonucleotides)
6859 [oligonucleotides](https://eu.idtdna.com/pages/education/decoded/article/tips-for-resuspending-and-diluting-your-oligonucleotides)> [Accessed 2 June 2022].

6860 Iguchi, A. *et al.* (2020) 'Deep-sea amphipods around cobalt-rich ferromanganese crusts: Taxonomic diversity
6861 and selection of candidate species for connectivity analysis', *PLOS ONE*, 15(2), p. e0228483. Available at:
6862 <https://doi.org/10.1371/JOURNAL.PONE.0228483>.

6863 Innan, H. *et al.* (1999) 'A Method for Estimating Nucleotide Diversity From AFLP Data', *Genetics*, 151(3), pp.
6864 1157–1164. Available at: <https://doi.org/10.1093/GENETICS/151.3.1157>.

6865 Innes, P. *et al.* (2022) 'Assessment of biogeographic variation in traits of Lewis flax (*Linum lewisii*) for use in
6866 restoration and agriculture', *AoB PLANTS*, 14(2). Available at: <https://doi.org/10.1093/aobpla/plac005>.

6867 Ionkova, I. *et al.* (2013) 'Linum narbonense: A new valuable tool for biotechnological production of a potent
6868 anticancer lignan Justicidine B', *Pharmacognosy Magazine*, 9(33), p. 39. Available at:
6869 <https://doi.org/10.4103/0973-1296.108138>.

6870 Iovane, M. *et al.* (2022) 'High temperature and humidity affect pollen viability and longevity in *olea europaea*
6871 l.', *Agronomy*, 12(1). Available at: <https://doi.org/10.3390/agronomy12010001>.

6872 Iwaizumi, M.G. and Takahashi, M. (2012) 'Effects of pollen supply and quality on seed formation and
6873 maturation in *Pinus densiflora*', *Journal of plant research*, 125(4), pp. 517–525. Available at:
6874 <https://doi.org/10.1007/S10265-011-0463-5>.

6875 Jaakola, L. and Hohtola, A. (2010) 'Effect of latitude on flavonoid biosynthesis in plants', *Plant, Cell &*
6876 *Environment*, 33(8), pp. 1239–1247. Available at: <https://doi.org/10.1111/J.1365-3040.2010.02154.X>.

6877 Jahnke, M.R. and Etterson, J.R. (2019) 'Autonomous self-fertilization in *Linum sulcatum*, a native annual with a
6878 previously unknown mating system', *Journal of plant research*, 132(1), pp. 57–67. Available at:
6879 <https://doi.org/10.1007/S10265-018-1076-Z>.

6880 Jarman, S.N., Ward, R.D. and Elliott, N.G. (2002) 'Oligonucleotide primers for PCR amplification of coelomate
6881 introns', *Marine Biotechnology*, 4(4), pp. 347–355. Available at: <https://doi.org/10.1007/s10126-002-0029-6>.

6882 Jensen, H. *et al.* (2007) 'Multilocus heterozygosity and inbreeding depression in an insular house sparrow
6883 metapopulation', *Molecular ecology*, 16(19), pp. 4066–4078. Available at: <https://doi.org/10.1111/J.1365->
6884 294X.2007.03452.X.

6885 Jhala, A. and Hall, L.M. (2014) *Flax (Linum usitatissimum L.): Current uses and future applications*. Available at:
6886 <https://www.researchgate.net/publication/228952435>.

6887 Jhala, A.J. and Hall, L.M. (2010) 'Flax (*Linum usitatissimum* L.): Current Uses and Future Applications',
6888 *Australian Journal of basic and Applied Sciences*, 4(9), pp. 4304–4312.

6889 Jiang, F., Cadotte, M.W. and Jin, G. (2021) 'Individual-level leaf trait variation and correlation across biological
6890 and spatial scales', *Ecology and Evolution*, 11(10), p. 5344. Available at: <https://doi.org/10.1002/ECE3.7425>.

6891 Johnson, M.A., Harper, J.F. and Palanivelu, R. (1AD) 'A Fruitful Journey: Pollen Tube Navigation from
6892 Germination to Fertilization'. Available at: <https://doi.org/10.1146/annurev-arplant-050718>.

6893 Johnstone, A., 2001. *Biology*. Oxford: Oxford University Press, pp.94-95.

6894 Jombart, T. and Ahmed, I. (2011) 'adeigenet 1.3-1: new tools for the analysis of genome-wide SNP data',
6895 *Bioinformatics*, 27(21), pp. 3070–3071. Available at: <https://doi.org/10.1093/BIOINFORMATICS/BTR521>.

6896 Jombart, T. and Bateman, A. (2008) 'adeigenet: a R package for the multivariate analysis of genetic markers',
6897 *Bioinformatics*, 24(11), pp. 1403–1405. Available at: <https://doi.org/10.1093/BIOINFORMATICS/BTN129>.

6898 Jones, B. and Gliddon, C. (1999) *Reproductive biology and genetic structure in Lloydia serotina*, *Plant Ecology*.

6899 de Jong, M.A. *et al.* (2011) 'Mitochondrial DNA Signature for Range-Wide Populations of *Bicyclus anynana*
6900 Suggests a Rapid Expansion from Recent Refugia', *PLOS ONE*, 6(6), p. e21385. Available at:
6901 <https://doi.org/10.1371/JOURNAL.PONE.0021385>.

6902 de Jong, M.J. *et al.* (2021) 'SambaR: An R package for fast, easy and reproducible population-genetic analyses
6903 of biallelic SNP data sets', *Molecular Ecology Resources*, 21(4), pp. 1369–1379. Available at:
6904 <https://doi.org/10.1111/1755-0998.13339>.

6905 Jordon-Thaden, I.E. *et al.* (2020) 'A basic ddRADseq two-enzyme protocol performs well with herbarium and
6906 silica-dried tissues across four genera', *Applications in Plant Sciences*, 8(4), p. e11344. Available at:
6907 <https://doi.org/10.1002/APS3.11344>.

6908 Jung, J.H. *et al.* (2007) 'The GIGANTEA-Regulated MicroRNA172 Mediates Photoperiodic Flowering
6909 Independent of CONSTANS in Arabidopsis', *The Plant Cell*, 19(9), pp. 2736–2748. Available at:
6910 <https://doi.org/10.1105/TPC.107.054528>.

6911 Kakani, V.G. *et al.* (2002) 'Response of in vitro pollen germination and pollen tube growth of groundnut
6912 (*Arachis hypogaea* L.) genotypes to temperature', *Plant, Cell & Environment*, 25(12), pp. 1651–1661. Available
6913 at: <https://doi.org/10.1046/J.1365-3040.2002.00943.X>.

6914 Kakani, V.G. *et al.* (2005) 'Differences in in vitro Pollen Germination and Pollen Tube Growth of Cotton
6915 Cultivars in Response to High Temperature', *Annals of Botany*, 96(1), pp. 59–67. Available at:
6916 <https://doi.org/10.1093/AOB/MCI149>.

6917 Kamvar, Z.N., Tabima, J.F. and Grünwald, N.J. (2014a) 'Poppr: An R package for genetic analysis of populations
6918 with clonal, partially clonal, and/or sexual reproduction', *PeerJ*, 2014(1), pp. 1–14. Available at:
6919 <https://doi.org/10.7717/PEERJ.281/TABLE-6>.

6920 Kamvar, Z.N., Tabima, J.F. and Grünwald, N.J. (2014b) 'Poppr: An R package for genetic analysis of populations
6921 with clonal, partially clonal, and/or sexual reproduction', *PeerJ*, 2014(1), pp. 1–14. Available at:
6922 <https://doi.org/10.7717/PEERJ.281/TABLE-6>.

6923 Kassambara, A., 2022. 'ggplot2' Based Publication Ready Plots [R package ggpubr version 0.4.0]. [online]
6924 Cran.r-project.org. Available at: <<https://cran.r-project.org/web/packages/ggpubr/index.html>> [Accessed 9
6925 October 2022].

6926 Kayes, J.M. and Clark, S.E. (1998) 'CLAVATA2, a regulator of meristem and organ development in Arabidopsis',
6927 *Development*, 125(19), pp. 3843–3851. Available at: <https://doi.org/10.1242/DEV.125.19.3843>.

6928 Keddy, P.A. and Constabel, P. (1986) 'Germination of Ten Shoreline Plants in Relation to Seed Size, Soil Particle
6929 Size and Water Level: An Experimental Study', *The Journal of Ecology*, 74(1), p. 133. Available at:
6930 <https://doi.org/10.2307/2260354>.

6931 Khan, M.R.G., Ai, X.Y. and Zhang, J.Z. (2014) 'Genetic regulation of flowering time in annual and perennial
6932 plants', *Wiley Interdisciplinary Reviews: RNA*. Blackwell Publishing Ltd, pp. 347–359. Available at:
6933 <https://doi.org/10.1002/wrna.1215>.

6934 Khanday, I. and Sundaresan, V. (2021) 'Plant zygote development: recent insights and applications to clonal
6935 seeds', *Current opinion in plant biology*, 59. Available at: <https://doi.org/10.1016/J.PBI.2020.101993>.

6936 Kim, D.H. *et al.* (2009) 'Vernalization: Winter and the Timing of Flowering in Plants',
6937 <http://dx.doi.org/10.1146/annurev.cellbio.042308.113411>, 25, pp. 277–299. Available at:
6938 <https://doi.org/10.1146/ANNUREV.CELLBIO.042308.113411>.

6939 Kinmonth-Schultz, H. *et al.* (2021a) 'Flowering Times of Wild Arabidopsis Accessions From Across Norway
6940 Correlate With Expression Levels of FT, CO, and FLC Genes', *Frontiers in Plant Science*, 12, p. 2376. Available at:
6941 <https://doi.org/10.3389/FPLS.2021.747740/BIBTEX>.

6942 Kinmonth-Schultz, H. *et al.* (2021b) 'Flowering Times of Wild Arabidopsis Accessions From Across Norway
6943 Correlate With Expression Levels of FT, CO, and FLC Genes', *Frontiers in Plant Science*, 12, p. 2376. Available at:
6944 <https://doi.org/10.3389/FPLS.2021.747740/BIBTEX>.

6945 Koonin, E. V. (2005) 'Orthologs, paralogs, and evolutionary genomics', *Annual review of genetics*, 39, pp. 309–
6946 338. Available at: <https://doi.org/10.1146/ANNUREV.GENET.39.073003.114725>.

6947 Kurokura, T. *et al.* (2017) 'Fragaria vesca CONSTANS controls photoperiodic flowering and vegetative
6948 development', *Journal of Experimental Botany*, 68(17), pp. 4839–4850. Available at:
6949 <https://doi.org/10.1093/JXB/ERX301>.

6950 Kuznetsova, A., Brockhoff, P.B. and Christensen, R.H.B. (2017) 'lmerTest Package: Tests in Linear Mixed Effects
6951 Models', *Journal of Statistical Software*, 82(13), pp. 1–26. Available at: <https://doi.org/10.18637/JSS.V082.I13>.

6952 Kyrkjeeide, M.O. *et al.* (2016) 'Long-distance dispersal and barriers shape genetic structure of peatmosses
6953 (sphagnum) across the Northern Hemisphere', *Journal of Biogeography*, 43(6), pp. 1215–1226.
6954 doi:10.1111/jbi.12716.

6955 Landoni, B. *et al.* (2022) 'Local climate and vernalization requirements explain the latitudinal patterns of
6956 flowering initiation in the crop wild relative *Linum bienne*', *bioRxiv*, p. 2022.01.02.474722. Available at:
6957 <https://doi.org/10.1101/2022.01.02.474722>.

6958 Langin, K.M. *et al.* (2017) 'Partial support for the central-marginal hypothesis within a population: reduced
6959 genetic diversity but not increased differentiation at the range edge of an island endemic bird', *Heredity*,
6960 119(1), pp. 8–15. Available at: <https://doi.org/10.1038/HDY.2017.10>.

6961 Larios, E. *et al.* (2014) 'Natural selection on seed size through the life cycle of a desert annual plant', *Ecology*,
6962 95(11), pp. 3213–3220. Available at: <https://doi.org/10.1890/13-1965.1>.

6963 Lascoux, M., Glémin, S. and Savolainen, O. (2016) 'Local Adaptation in Plants', *eLS*, pp. 1–7. Available at:
6964 <https://doi.org/10.1002/9780470015902.A0025270>.

6965 Lazaro, A., Obeng-Hinne, E. and Albani, M.C. (2018) 'Extended Vernalization Regulates Inflorescence Fate in
6966 *Arabis alpina* by Stably Silencing PERPETUAL FLOWERING1', *Plant Physiology*, 176(4), pp. 2819–2833. Available
6967 at: <https://doi.org/10.1104/PP.17.01754>.

6968 Lee, D.R. *et al.* (2014) 'Developing local adaptation strategies for climate change in agriculture: A priority-
6969 setting approach with application to Latin America', *Global Environmental Change*, 29, pp. 78–91. Available at:
6970 <https://doi.org/10.1016/J.GLOENVCHA.2014.08.002>.

6971 Lee, J. and Lee, I. (2010) 'Regulation and function of SOC1, a flowering pathway integrator', *Journal of*
6972 *Experimental Botany*, 61(9), pp. 2247–2254. Available at: <https://doi.org/10.1093/JXB/ERQ098>.

6973 Lee, J.H. *et al.* (2013) 'Regulation of temperature-responsive flowering by MADS-box transcription factor
6974 repressors', *Science*, 342(6158), pp. 628–632. Available at:
6975 https://doi.org/10.1126/SCIENCE.1241097/SUPPL_FILE/PAPV2.PDF.

6976 Lee, J.H., Kim, J.J. and Ahn, J.H. (2012) 'Role of SEPALLATA3 (SEP3) as a downstream gene of miR156-SPL3-FT
6977 circuitry in ambient temperature-responsive flowering', <https://doi.org/10.4161/psb.21366>, 7(9), pp. 1151–
6978 1154. Available at: <https://doi.org/10.4161/PSB.21366>.

6979 Leijten, W. *et al.* (2018) 'Translating Flowering Time from *Arabidopsis thaliana* to Brassicaceae and Asteraceae
6980 Crop Species', *Plants* 2018, Vol. 7, Page 111, 7(4), p. 111. Available at:
6981 <https://doi.org/10.3390/PLANTS7040111>.

6982 Leimu, R. and Fischer, M. (2008a) 'A meta-analysis of local adaptation in plants', *PLoS ONE*, 3(12). Available at:
6983 <https://doi.org/10.1371/journal.pone.0004010>.

6984 Leimu, R. and Fischer, M. (2008b) 'A Meta-Analysis of Local Adaptation in Plants', *PLOS ONE*, 3(12), p. e4010.
6985 Available at: <https://doi.org/10.1371/JOURNAL.PONE.0004010>.

6986 Lemuth, K. and Rupp, S. (2015) 'Microarrays as Research Tools and Diagnostic Devices', *RNA Technologies*, pp.
6987 259–280. Available at: https://doi.org/10.1007/978-3-319-17305-4_13/FIGURES/2.

6988 Li, H. (2011) 'A statistical framework for SNP calling, mutation discovery, association mapping and population
6989 genetical parameter estimation from sequencing data', *Bioinformatics*, 27(21), p. 2987. Available at:
6990 <https://doi.org/10.1093/BIOINFORMATICS/BTR509>.

6991 Li, H. and Durbin, R. (2009) 'Fast and accurate short read alignment with Burrows-Wheeler transform',
6992 *Bioinformatics (Oxford, England)*, 25(14), pp. 1754–1760. Available at:
6993 <https://doi.org/10.1093/BIOINFORMATICS/BTP324>.

6994 Li, L. *et al.* (2015a) 'Flowering responses to light and temperature', *Science China Life Sciences* 2016 59:4, 59(4),
6995 pp. 403–408. Available at: <https://doi.org/10.1007/S11427-015-4910-8>.

6996 Li, L. *et al.* (2015b) 'Flowering responses to light and temperature', *Science China Life Sciences* 2016 59:4, 59(4),
6997 pp. 403–408. Available at: <https://doi.org/10.1007/S11427-015-4910-8>.

6998 Li, Y.X. *et al.* (2016) 'Identification of genetic variants associated with maize flowering time using an extremely
6999 large multi-genetic background population', *The Plant Journal*, 86(5), pp. 391–402. Available at:
7000 <https://doi.org/10.1111/TPJ.13174>.

7001 Lian, H., Roy, E. and Zheng, H. (2016) 'Microglial Phagocytosis Assay', *BIO-PROTOCOL*, 6(21). Available at:
7002 <https://doi.org/10.21769/BIOPROTOC.1988>.

7003 Liang, W. *et al.* (2018a) 'Selection and evaluation of reference genes for qRT-PCR analysis in *Euscaphis konishii*
7004 Hayata based on transcriptome data', *Plant Methods*, 14(1), pp. 1–9. Available at:
7005 <https://doi.org/10.1186/S13007-018-0311-X/FIGURES/4>.

7006 Liang, W. *et al.* (2018b) 'Selection and evaluation of reference genes for qRT-PCR analysis in *Euscaphis konishii*
7007 Hayata based on transcriptome data', *Plant Methods*, 14(1), pp. 1–9. Available at:
7008 <https://doi.org/10.1186/S13007-018-0311-X/FIGURES/4>.

7009 Linck, E. and Battey, C.J. (2019) 'Minor allele frequency thresholds strongly affect population structure
7010 inference with genomic data sets', *Molecular Ecology Resources*, 19(3), pp. 639–647. Available at:
7011 <https://doi.org/10.1111/1755-0998.12995>.

7012 Lischer, H.E.L. and Excoffier, L. (2012) 'PGDSpider: an automated data conversion tool for connecting
7013 population genetics and genomics programs', *Bioinformatics (Oxford, England)*, 28(2), pp. 298–299. Available
7014 at: <https://doi.org/10.1093/BIOINFORMATICS/BTR642>.

7015 Liu, J. *et al.* (2015) 'Genetic and epigenetic control of plant heat responses', *Frontiers in Plant Science*, 6(APR),
7016 p. 267. Available at: <https://doi.org/10.3389/FPLS.2015.00267/BIBTEX>.

7017 Liu, Y. *et al.* (2019) 'Genome-Wide Single Nucleotide Polymorphisms are Robust in Resolving Fine-Scale
7018 Population Genetic Structure of the Small Brown Planthopper, *Laodelphax striatellus* (Fallén) (Hemiptera:
7019 Delphacidae)', *Journal of Economic Entomology*, 112(5), pp. 2362–2368. Available at:
7020 <https://doi.org/10.1093/JEE/TOZ145>.

7021 Livak, K.J. and Schmittgen, T.D. (2001a) 'Analysis of Relative Gene Expression Data Using Real-Time
7022 Quantitative PCR and the $2^{-\Delta\Delta CT}$ Method', *Methods*, 25(4), pp. 402–408. Available at:
7023 <https://doi.org/10.1006/METH.2001.1262>.

7024 Livak, K.J. and Schmittgen, T.D. (2001b) 'Analysis of Relative Gene Expression Data Using Real-Time
7025 Quantitative PCR and the $2^{-\Delta\Delta CT}$ Method', *Methods*, 25(4), pp. 402–408. Available at:
7026 <https://doi.org/10.1006/METH.2001.1262>.

7027 Loha, A., Tigabu, M. and Fries, A. (2008) 'Genetic variation among and within populations of *Cordia africana* in
7028 seed size and germination responses to constant temperatures', *Euphytica* 2008 165:1, 165(1), pp. 189–196.
7029 Available at: <https://doi.org/10.1007/S10681-008-9806-2>.

7030 Lotz, L.A.P. (1990) 'The Relation Between Age and Size at First Flowering of *Plantago Major* in Various
7031 Habitats', *The Journal of Ecology*, 78(3), p. 757. Available at: <https://doi.org/10.2307/2260897>.

7032 Loveless, M.D. and Hamrick, J.L. (1984a) *Ecological Determinants of Genetic Structure in Plant Populations*,
7033 *Source: Annual Review of Ecology and Systematics*. Available at: <https://about.jstor.org/terms>.

7034 Loveless, M.D. and Hamrick, J.L. (1984b) *Ecological Determinants of Genetic Structure in Plant Populations*,
7035 *Source: Annual Review of Ecology and Systematics*. Available at: <https://about.jstor.org/terms>.

7036 Lovell, J.T. *et al.* (2013) 'Pleiotropy of FRIGIDA enhances the potential for multivariate adaptation', *Proceedings*
7037 *of the Royal Society B: Biological Sciences*, 280(1763). Available at: <https://doi.org/10.1098/rspb.2013.1043>.

7038 Lu, P. *et al.* (2014) 'The Arabidopsis CALLOSE DEFECTIVE MICROSPORE1 Gene Is Required for Male Fertility
7039 through Regulating Callose Metabolism during Microsporogenesis', *Plant Physiology*, 164(4), pp. 1893–1904.
7040 Available at: <https://doi.org/10.1104/PP.113.233387>.

7041 Luo, Z. *et al.* (2019) 'Genetic diversity and population structure of a *Camelina sativa* spring panel', *Frontiers in*
7042 *Plant Science*, 10, p. 184. Available at: <https://doi.org/10.3389/FPLS.2019.00184/BIBTEX>.

7043 Lütfi, P. (2002) *The Effects of Temperature on Pollen Germination and Pollen Tube Growth of Apricot and Sweet*
7044 *Cherry*. Available at: https://www.pubhort.org/ejhs/2002/file_1877.pdf (Accessed: 6 September 2022).

7045 Lyakh, V.A. and Soroka, A.I. (2021) 'Artificial medium for in vitro pollen germination of some ornamental Linum
7046 species', *Advances in Horticultural Science*, 35(3), pp. 329–332. Available at: <https://doi.org/10.36253/AHSC->
7047 10854.

7048 Maisonneuve, B. and Den Nijs, A.P.M. (1984) 'In vitro pollen germination and tube growth of tomato
7049 (*Lycopersicon esculentum* Mill.) and its relation with plant growth', *Euphytica*, 33(3), pp. 833–840. Available at:
7050 <https://doi.org/10.1007/BF00021910/METRICS>.

7051 Manoli, A. *et al.* (2012) 'Evaluation of candidate reference genes for qPCR in maize', *Journal of Plant*
7052 *Physiology*, 169(8), pp. 807–815. Available at: <https://doi.org/10.1016/J.JPLPH.2012.01.019>.

7053 Marth, G.T. *et al.* (2004) 'The Allele Frequency Spectrum in Genome-Wide Human Variation Data Reveals
7054 Signals of Differential Demographic History in Three Large World Populations', *Genetics*, 166(1), pp. 351–372.
7055 Available at: <https://doi.org/10.1534/GENETICS.166.1.351>.

7056 Maruki, T. and Lynch, M. (2014) 'Genome-wide estimation of linkage disequilibrium from population-level
7057 high-throughput sequencing data', *Genetics*, 197(4), pp. 1303–1313. Available at:
7058 <https://doi.org/10.1534/GENETICS.114.165514>.

7059 Mastretta-Yanes, A. *et al.* (2014) 'Gene Duplication, Population Genomics, and Species-Level Differentiation
7060 within a Tropical Mountain Shrub', *Genome Biology and Evolution*, 6(10), p. 2611. Available at:
7061 <https://doi.org/10.1093/GBE/EVU205>.

7062 McDill, J. *et al.* (2009) 'The phylogeny of Linum and Linaceae subfamily Linoideae, with implications for their
7063 systematics, biogeography, and evolution of heterostyly', *Systematic Botany*, 34(2), pp. 386–405. Available at:
7064 <https://doi.org/10.1600/036364409788606244>.

7065 McKinney, G.J. *et al.* (2017) 'Paralogs are revealed by proportion of heterozygotes and deviations in read ratios
7066 in genotyping-by-sequencing data from natural populations', *Molecular Ecology Resources*, 17(4), pp. 656–669.
7067 Available at: <https://doi.org/10.1111/1755-0998.12613>.

7068 McMaster, G.S. *et al.* (2008) 'Simulating the influence of vernalization, photoperiod and optimum temperature
7069 on wheat developmental rates', *Annals of botany*, 102(4), pp. 561–569. Available at:
7070 <https://doi.org/10.1093/AOB/MCN115>.

7071 McNeil, K.A., Newman, I. and Kelly, F.J., 1996. Testing research hypotheses with the general linear model. SIU
7072 Press.

7073 Meek, M.H. *et al.* (2023) 'Understanding Local Adaptation to Prepare Populations for Climate Change',
7074 *BioScience*, 73(1), pp. 36–47. Available at: <https://doi.org/10.1093/biosci/biac101>.

7075 De Mendonça Vilela, M. *et al.* (2017) 'Analysis of Three Sugarcane Homo/Homeologous Regions Suggests
7076 Independent Polyploidization Events of *Saccharum officinarum* and *Saccharum spontaneum*', *Genome Biology*
7077 *and Evolution*, 9(2), pp. 266–278. Available at: <https://doi.org/10.1093/GBE/EVW293>.

7078 Miller, J.M., Cullingham, C.I. and Peery, R.M. (2020) 'The influence of a priori grouping on inference of genetic
7079 clusters: simulation study and literature review of the DAPC method', *Heredity* 2020 125:5, 125(5), pp. 269–
7080 280. Available at: <https://doi.org/10.1038/s41437-020-0348-2>.

7081 Miranda, J.A. and Steward, G.F. (2017) 'Variables influencing the efficiency and interpretation of reverse
7082 transcription quantitative PCR (RT-qPCR): An empirical study using Bacteriophage MS2', *Journal of Virological*
7083 *Methods*, 241, pp. 1–10. Available at: <https://doi.org/10.1016/J.JVIROMET.2016.12.002>.

7084 Mittell, E.A. *et al.* (2020) 'Feral populations of Brassica oleracea along Atlantic coasts in western Europe',
7085 *Ecology and Evolution*, 10(20), p. 11810. Available at: <https://doi.org/10.1002/ECE3.6821>.

7086 MobaXterm., 2022. MobaXterm free Xserver and tabbed SSH client for Windows. [online] Available at:
7087 <<https://mobaxterm.mobatek.net/>> [Accessed 21 September 2022].

7088 MOLES, ANGELA T. *et al.* (2004) "Small-seeded species produce more seeds per square metre of canopy per
7089 year, but not per individual per lifetime," *Journal of Ecology*, 92(3), pp. 384–396. Available at:
7090 <https://doi.org/10.1111/j.0022-0477.2004.00880.x>.

7091 Montalt, R. *et al.* (2019) 'Influence of temperature on the progamic phase in Citrus', *Environmental and*
7092 *Experimental Botany*, 166, p. 103806. Available at: <https://doi.org/10.1016/J.ENVEXPBOT.2019.103806>.

7093 Montesinos-Navarro, A., Picó, F. and Tonsor, S. (2012) 'Clinal variation in seed traits influencing life cycle
7094 timing in *Arabidopsis thaliana*', *Evolution*, 66.

7095 Moran, E. V. (2020) 'Simulating the effects of local adaptation and life history on the ability of plants to track
7096 climate shifts', *AoB PLANTS*, 12(1). Available at: <https://doi.org/10.1093/aobpla/plaa008>.

7097 Mori, A.S. *et al.* (2022) 'Feedbacks Between Biodiversity and Climate Change', *The Ecological and Societal*
7098 *Consequences of Biodiversity Loss*, pp. 281–304. Available at: <https://doi.org/10.1002/9781119902911.CH13>.

7099 Muravenko, O. V. *et al.* (2010a) 'Karyogenomics of species of the genus *Linum* L.', *Russian Journal of Genetics*
7100 2010 46:10, 46(10), pp. 1182–1185. Available at: <https://doi.org/10.1134/S1022795410100091>.

7101 Muravenko, O. V. *et al.* (2010b) 'Karyogenomics of species of the genus *Linum* L.', *Russian Journal of Genetics*
7102 2010 46:10, 46(10), pp. 1182–1185. Available at: <https://doi.org/10.1134/S1022795410100091>.

7103 Murray, B.R. *et al.* (2004) 'Geographical gradients in seed mass in relation to climate', *Journal of Biogeography*,
7104 31(3), pp. 379–388. Available at: <https://doi.org/10.1046/J.0305-0270.2003.00993.X>.

7105 Muthamia, K. (1994) 'Effect of bulb size and duration of vernalization on onion (*Allium cepa* L.) seed yield'.
7106 Available at: <http://erepository.uonbi.ac.ke/handle/11295/23061> (Accessed: 9 October 2022).

7107 Mylne, J. *et al.* (2004) 'Epigenetic Regulation in the Control of Flowering', *Cold Spring Harbor Symposia on*
7108 *Quantitative Biology*, 69, pp. 457–464. Available at: <https://doi.org/10.1101/SQB.2004.69.457>.

7109 Nag, S., Mitra, J. and Karmakar, P.G. (2015) ' An Overview on Flax (*Linum usitatissimum* L.) and its Genetic
7110 Diversity ', *International Journal of Agriculture, Environment and Biotechnology*, 8(4), p. 805. Available at:
7111 <https://doi.org/10.5958/2230-732x.2015.00089.3>.

7112 Nazareno, A.G. et al. (2021) 'By animal, water, or wind: Can dispersal mode predict genetic connectivity in
7113 riverine plant species?', *Frontiers in Plant Science*, 12. doi:10.3389/fpls.2021.626405.

7114 Nei, M. (1972a) *Genetic Distance between Populations*, Source: *The American Naturalist*.

7115 Nei, M. (1972b) *Genetic Distance between Populations*, Source: *The American Naturalist*. Available at:
7116 <https://about.jstor.org/terms>.

7117 Newbigin, E., Anderson, M.A. and Clarke', A.E. (1993) *Gametophytic Self-Incompatibility Systems*, *The Plant*
7118 *Cell*.

7119 Nievola, C.C. et al. (2017) 'Rapid responses of plants to temperature changes', *Temperature: Multidisciplinary*
7120 *Biomedical Journal*, 4(4), p. 371. Available at: <https://doi.org/10.1080/23328940.2017.1377812>.

7121 Normand, S. et al. (2011) 'Postglacial migration supplements climate in determining plant species ranges in
7122 Europe', *Proceedings of the Royal Society B: Biological Sciences*, 278(1725), pp. 3644–3653.
7123 doi:10.1098/rspb.2010.2769.

7124 Nôžková, J., Remeselníková, K. and Bjelková, M. (2014) 'Characterization and evaluation of flax seeds (*Linum*
7125 *usitatissimum* L.) on selected genotypes', *Journal of Central European Agriculture*, 15(1), pp. 193–207.
7126 Available at: <https://doi.org/10.5513/JCEA01/15.1.1434>.

7127 Odong, T.L. et al. (2011) 'Determination of genetic structure of germplasm collections: Are traditional
7128 hierarchical clustering methods appropriate for molecular marker data?', *Theoretical and Applied Genetics*,
7129 123(2), pp. 195–205. Available at: <https://doi.org/10.1007/S00122-011-1576-X/FIGURES/6>.

7130 Öhlund, E., Zurek, K. and Hammer, M. (2015) 'Towards Sustainable Agriculture? The EU framework and local
7131 adaptation in Sweden and Poland', *Environmental Policy and Governance*, 25(4), pp. 270–287. Available at:
7132 <https://doi.org/10.1002/EET.1687>.

7133 Olejniczak, P. et al. (2018) 'Seed size in mountain herbaceous plants changes with elevation in a species-
7134 specific manner', *PLoS ONE*, 13(6). Available at: <https://doi.org/10.1371/journal.pone.0199224>.

7135 Osnato, M. et al. (2022) 'Photoperiod Control of Plant Growth: Flowering Time Genes Beyond Flowering',
7136 *Frontiers in Plant Science*, 12, p. 3359. Available at: <https://doi.org/10.3389/FPLS.2021.805635/BIBTEX>.

7137 Ozhimkova, E., Uschapovsky, I. and Manaenkov, O. (2023) "Study of varietal differences in the composition of
7138 heteropolysaccharides of oil flax and fiber flax," *Polysaccharides*, 4(1), pp. 78–87. Available at:
7139 <https://doi.org/10.3390/polysaccharides4010006>.

7140 Pacini, E. (2008) 'Pollination', *Encyclopedia of Ecology*, pp. 562–565. Available at:
7141 <https://doi.org/10.1016/B978-0-444-63768-0.00859-3>.

7142 Pacini, E. and Dolferus, R. (2016) 'The Trials and Tribulations of the Plant Male Gametophyte — Understanding
7143 Reproductive Stage Stress Tolerance', *Abiotic and Biotic Stress in Plants - Recent Advances and Future
7144 Perspectives* [Preprint]. Available at: <https://doi.org/10.5772/61671>.

7145 Paina, C. *et al.* (2014) 'Vernalization Mediated Changes in the *Lolium perenne* Transcriptome', *PLOS ONE*, 9(9),
7146 p. e107365. Available at: <https://doi.org/10.1371/JOURNAL.PONE.0107365>.

7147 Pali, V. *et al.* (2014) 'Identification of microsatellite markers for fingerprinting popular Indian flax (*Linum
7148 usitatissimum* L.) cultivars and their utilization in seed genetic purity assessments', *Australian Journal of Crop
7149 Sciences*, 8(1).

7150 Pan, C. *et al.* (2018) 'Tomato stigma exertion induced by high temperature is associated with the jasmonate
7151 signalling pathway', *Plant, Cell & Environment*, 42(4), pp. 1205–1221. Available at: [doi:10.1111/pce.13444](https://doi.org/10.1111/pce.13444).

7152 Paradis, E. and Schliep, K. (2019a) 'ape 5.0: an environment for modern phylogenetics and evolutionary
7153 analyses in R', *Bioinformatics*, 35(3), pp. 526–528. Available at:
7154 <https://doi.org/10.1093/BIOINFORMATICS/BTY633>.

7155 Paradis, E. and Schliep, K. (2019b) 'ape 5.0: an environment for modern phylogenetics and evolutionary
7156 analyses in R', *Bioinformatics*, 35(3), pp. 526–528. Available at:
7157 <https://doi.org/10.1093/BIOINFORMATICS/BTY633>.

7158 Park, M.J. *et al.* (2016) 'LATE ELONGATED HYPOCOTYL regulates photoperiodic flowering via the circadian clock
7159 in *Arabidopsis*', *BMC Plant Biology*, 16(1), pp. 1–12. Available at: [https://doi.org/10.1186/S12870-016-0810-
7160 8/FIGURES/5](https://doi.org/10.1186/S12870-016-0810-8/FIGURES/5).

7161 Peet, M.M. and Bartholemew, M. (1996) *Effect of Night Temperature on Pollen Characteristics, Growth, and
7162 Fruit Set in Tomato*, *J. AMER. SOC. HORT. SCI.*

7163 Pembleton, L.W., Cogan, N.O.I. and Forster, J.W. (2013a) 'StAMPP: an R package for calculation of genetic
7164 differentiation and structure of mixed-ploidy level populations', *Molecular Ecology Resources*, 13(5), pp. 946–
7165 952. Available at: <https://doi.org/10.1111/1755-0998.12129>.

7166 Pembleton, L.W., Cogan, N.O.I. and Forster, J.W. (2013b) 'StAMPP: an R package for calculation of genetic
7167 differentiation and structure of mixed-ploidy level populations', *Molecular Ecology Resources*, 13(5), pp. 946–
7168 952. Available at: <https://doi.org/10.1111/1755-0998.12129>.

7169 Peterson, B.K. *et al.* (2012a) 'Double Digest RADseq: An Inexpensive Method for De Novo SNP Discovery and
7170 Genotyping in Model and Non-Model Species', *PLOS ONE*, 7(5), p. e37135. Available at:
7171 <https://doi.org/10.1371/JOURNAL.PONE.0037135>.

7172 Peterson, B.K. *et al.* (2012b) 'Double Digest RADseq: An Inexpensive Method for De Novo SNP Discovery and
7173 Genotyping in Model and Non-Model Species', *PLOS ONE*, 7(5), p. e37135. Available at:
7174 <https://doi.org/10.1371/JOURNAL.PONE.0037135>.

7175 Peterson, B.K. *et al.* (2012c) 'Double Digest RADseq: An Inexpensive Method for De Novo SNP Discovery and
7176 Genotyping in Model and Non-Model Species', *PLOS ONE*, 7(5), p. e37135. Available at:
7177 <https://doi.org/10.1371/JOURNAL.PONE.0037135>.

7178 Pfaffl, M.W. (2001) 'A new mathematical model for relative quantification in real-time RT-PCR', *Nucleic Acids*
7179 *Research*, 29(9), p. e45. Available at: <https://doi.org/10.1093/NAR/29.9.E45>.

7180 Porebski, S., Bailey, L.G. and Baum, B.R. (1997) 'Modification of a CTAB DNA extraction protocol for plants
7181 containing high polysaccharide and polyphenol components', *Plant Molecular Biology Reporter* 1997 15:1,
7182 15(1), pp. 8–15. Available at: <https://doi.org/10.1007/BF02772108>.

7183 Posé, D. *et al.* (2013) 'Temperature-dependent regulation of flowering by antagonistic FLM variants', *Nature*
7184 2013 503:7476, 503(7476), pp. 414–417. Available at: <https://doi.org/10.1038/nature12633>.

7185 Promega.co.uk. 2022. GoTaq® qPCR and RT-qPCR Systems | qPCR Master Mix | RT-qPCR Master Mix. [online]
7186 Available at: <[https://www.promega.co.uk/products/pcr/qpcr-and-rt-qpcr/qpcr-](https://www.promega.co.uk/products/pcr/qpcr-and-rt-qpcr/qpcr-kits/?catNum=A6020#resources)
7187 [kits/?catNum=A6020#resources](https://www.promega.co.uk/products/pcr/qpcr-and-rt-qpcr/qpcr-kits/?catNum=A6020#resources)> [Accessed 30 May 2022].

7188 Przeworski, M. (2002) 'The Signature of Positive Selection at Randomly Chosen Loci', *Genetics*, 160(3), pp.
7189 1179–1189. Available at: <https://doi.org/10.1093/GENETICS/160.3.1179>.

7190 Purcell, S. *et al.* (2007a) 'PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage
7191 Analyses', *The American Journal of Human Genetics*, 81(3), pp. 559–575. Available at:
7192 <https://doi.org/10.1086/519795>.

7193 Purcell, S. *et al.* (2007b) 'PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage
7194 Analyses', *The American Journal of Human Genetics*, 81(3), pp. 559–575. Available at:
7195 <https://doi.org/10.1086/519795>.

7196 Purcell, S. *et al.* (2007c) 'PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage
7197 Analyses', *The American Journal of Human Genetics*, 81(3), pp. 559–575. Available at:
7198 <https://doi.org/10.1086/519795>.

7199 Purugganan, M.D. and Fuller, D.Q. (2009) 'The nature of selection during plant domestication', *Nature*,
7200 457(7231), pp. 843–848. Available at: <https://doi.org/10.1038/NATURE07895>.

7201 Putterill, J., Laurie, R. and Macknight, R. (2004) 'It's time to flower: the genetic control of flowering time',
7202 *BioEssays*, 26(4), pp. 363–373. Available at: <https://doi.org/10.1002/BIES.20021>.

7203 Qaderi, M.M., Martel, A.B. and Dixon, S.L. (2019) 'Environmental Factors Influence Plant Vascular System and
7204 Water Regulation', *Plants*, 8(3), p. 65. Available at: <https://doi.org/10.3390/PLANTS8030065>.

7205 Qin, Y. *et al.* (2009) 'Penetration of the Stigma and Style Elicits a Novel Transcriptome in Pollen Tubes, Pointing
7206 to Genes Critical for Growth in a Pistil', *PLOS Genetics*, 5(8), p. e1000621. Available at:
7207 <https://doi.org/10.1371/JOURNAL.PGEN.1000621>.

7208 *QuantiFluor dsDNA System for High Concentration DNA* (no date). Available at:
7209 <https://www.promega.co.uk/resources/pubhub/quantifluor-dsdna-system-for-high-concentration-dna/>
7210 (Accessed: 3 May 2022).

7211 Rahimi, M.M., Zarei, M.A. and Arminian, A. (2011) 'Selection criteria of flax (*Linum usitatissimum* L.) for seed
7212 yield, yield components and biochemical compositions under various planting dates and nitrogen', *African*
7213 *Journal of Agricultural Research*, 6(13), pp. 3167–3175. Available at: <https://doi.org/10.5897/AJAR11.382>.

7214 Rao, X. *et al.* (2013a) 'An improvement of the $2^{-\Delta\Delta CT}$ method for quantitative real-time
7215 polymerase chain reaction data analysis', *Biostatistics, bioinformatics and biomathematics*, 3(3), p. 71.
7216 Available at: </pmc/articles/PMC4280562/> (Accessed: 13 June 2022).

7217 Rao, X. *et al.* (2013b) 'An improvement of the $2^{-\Delta\Delta CT}$ method for quantitative real-time
7218 polymerase chain reaction data analysis', *Biostatistics, bioinformatics and biomathematics*, 3(3), p. 71.
7219 Available at: </pmc/articles/PMC4280562/> (Accessed: 15 August 2022).

7220 Rastogi, A. *et al.* (2013) *Effect of auxin and gibberellic acid on growth and yield components of linseed (Linum*
7221 *usitatissimum* L.), *Crop Breeding and Applied Biotechnology*.

7222 Rauf, S. *et al.* (2010) *Consequences of Plant Breeding on genetic diversity Bt cotton in Pakistan: Performance*
7223 *and adoption with imperfect information View project Drought tolerant Sunflower breeding material View*
7224 *project International Journal of Plant Breeding Consequences of Plant Breeding on Genetic Diversity*. Available
7225 at: <https://www.researchgate.net/publication/235910237>.

7226 Raven, P., 1982. *Biology of plants*. New York: Worth Publishers, pp.132-137.

7227 Ray, C. (1944) 'CYTOLOGICAL STUDIES ON THE FLAX GENUS, LINUM', *American Journal of Botany*, 31(4), pp.
7228 241–248. Available at: <https://doi.org/10.1002/J.1537-2197.1944.TB08027.X>.

7229 R Core Team, 2022. R: A language and environment for statistical computing. R Foundation for Statistical
7230 Computing, Vienna, Austria. Available online at <https://www.R-project.org/>.

7231 Ream, T.S. *et al.* (2014) 'Interaction of Photoperiod and Vernalization Determines Flowering Time of
7232 *Brachypodium distachyon*', *Plant Physiology*, 164(2), p. 694. Available at:
7233 <https://doi.org/10.1104/PP.113.232678>.

7234 Reeves, P.H. and Coupland, G. (2000) 'Response of plant development to environment: control of flowering by
7235 daylength and temperature', *Current Opinion in Plant Biology*, 3(1), pp. 37–42. Available at:
7236 [https://doi.org/10.1016/S1369-5266\(99\)00041-2](https://doi.org/10.1016/S1369-5266(99)00041-2).

7237 Rehman, H.U. *et al.* (2014) 'Seed Priming Influence on Early Crop Growth, Phenological Development and Yield
7238 Performance of Linola (*Linum usitatissimum* L.)', *Journal of Integrative Agriculture*, 13(5), pp. 990–996.
7239 Available at: [https://doi.org/10.1016/S2095-3119\(13\)60521-3](https://doi.org/10.1016/S2095-3119(13)60521-3).

7240 Remans, T. *et al.* (2008) 'Normalisation of real-time RT-PCR gene expression measurements in Arabidopsis
7241 thaliana exposed to increased metal concentrations', *Planta*, 227(6), pp. 1343–1349. Available at:
7242 <https://doi.org/10.1007/S00425-008-0706-4/FIGURES/1>.

7243 Reynolds, W.W. and Casterlin, M.E. (1980) 'The Role of Temperature in the Environmental Physiology of
7244 Fishes', *Environmental Physiology of Fishes*, pp. 497–518. Available at: https://doi.org/10.1007/978-1-4899-3659-2_19.

7246 Riedel, G. *et al.* (2014a) 'An Extended Δ CT-Method Facilitating Normalisation with Multiple Reference Genes
7247 Suited for Quantitative RT-PCR Analyses of Human Hepatocyte-Like Cells', *PLOS ONE*, 9(3), p. e93031. Available
7248 at: <https://doi.org/10.1371/JOURNAL.PONE.0093031>.

7249 Riedel, G. *et al.* (2014b) 'An Extended Δ CT-Method Facilitating Normalisation with Multiple Reference Genes
7250 Suited for Quantitative RT-PCR Analyses of Human Hepatocyte-Like Cells', *PLOS ONE*, 9(3), p. e93031. Available
7251 at: <https://doi.org/10.1371/JOURNAL.PONE.0093031>.

7252 Riedel, G. *et al.* (2014c) 'An Extended Δ CT-Method Facilitating Normalisation with Multiple Reference Genes
7253 Suited for Quantitative RT-PCR Analyses of Human Hepatocyte-Like Cells', *PLOS ONE*, 9(3), p. e93031. Available
7254 at: <https://doi.org/10.1371/JOURNAL.PONE.0093031>.

7255 Del Río, J.C. *et al.* (2011) 'Structural characterization of guaiacyl-rich lignins in flax (*Linum usitatissimum*) fibers
7256 and shives', *Journal of Agricultural and Food Chemistry*, 59(20), pp. 11088–11099. Available at:
7257 https://doi.org/10.1021/JF201222R/ASSET/IMAGES/LARGE/JF-2011-01222R_0001.JPEG.

7258 Roach, D.A. (1987) *Variation in Seed and Seedling Size in Anthoxanthum odoratum*, Source: *The American*
7259 *Midland Naturalist*.

7260 Rodriguez-Enriquez, M.J. *et al.* (2013) 'A novel method for efficient in vitro germination and tube growth of
7261 Arabidopsis thaliana pollen', *New Phytologist*, 197(2), pp. 668–679. Available at:
7262 <https://doi.org/10.1111/NPH.12037>.

7263 Royo, C. *et al.* (2020) 'Agronomic performance of durum wheat landraces and modern cultivars and its
7264 association with genotypic variation in vernalization response (Vrn-1) and photoperiod sensitivity (Ppd-1)
7265 genes', *European Journal of Agronomy*, 120, p. 126129. Available at:
7266 <https://doi.org/10.1016/J.EJA.2020.126129>.

7267 Rozhmina, T. *et al.* (2021) 'A comprehensive dataset of flax (*Linum uitatissimum* L.) phenotypes', *Data in Brief*,
7268 37. Available at: <https://doi.org/10.1016/j.dib.2021.107224>.

7269 RStudio Team, 2020. RStudio: Integrated Development for R. RStudio, PBC, Boston, MA URL
7270 <http://www.rstudio.com/>.

7271 Rúa, M.A. *et al.* (2016) 'Home-field advantage? evidence of local adaptation among plants, soil, and arbuscular
7272 mycorrhizal fungi through meta-analysis', *BMC Evolutionary Biology*, 16(1). Available at:
7273 <https://doi.org/10.1186/s12862-016-0698-9>.

7274 Ruane, L.G. and Donohue, K. (2007) 'Environmental effects on pollen–pistil compatibility between *Phlox*
7275 *cuspidata* and *P. drummondii* (Polemoniaceae): implications for hybridization dynamics', *American Journal of*
7276 *Botany*, 94(2), pp. 219–227. Available at: <https://doi.org/10.3732/AJB.94.2.219>.

7277 Rychel, S. *et al.* (2019) 'FLOWERING LOCUS T, GIGANTEA, SEPALLATA, and FRIGIDA homologs are candidate
7278 genes involved in white lupin (*Lupinus albus* L.) early flowering', *Molecular Breeding*, 39(3), pp. 1–17. Available
7279 at: <https://doi.org/10.1007/S11032-019-0952-0/FIGURES/2>.

7280 Sa, R. *et al.* (2021) 'Chromosome-Level Genome Assembly and Annotation of the Fiber Flax (*Linum*
7281 *usitatissimum*) Genome', *Frontiers in Genetics*, 12, p. 735690. Available at:
7282 <https://doi.org/10.3389/FGENE.2021.735690/FULL>.

7283 Saini, H.S., Sedgley, M. and Aspinall, D. (1983) 'Effect of Heat Stress During Floral Development on Pollen Tube
7284 Growth and Ovary Anatomy in Wheat (*Triticum aestivum* L.)', *Functional Plant Biology*, 10(2), pp. 137–144.
7285 Available at: <https://doi.org/10.1071/PP9830137>.

7286 Sallam, A.H. *et al.* (2021) "Cold conditioned: Discovery of novel alleles for low-temperature tolerance in the
7287 Vavilov Barley Collection," *Frontiers in Plant Science*, 12. Available at:
7288 <https://doi.org/10.3389/fpls.2021.800284>.

7289 Sand Korneliussen, T. *et al.* (2013) *Calculation of Tajima's D and other neutrality test statistics from low depth*
7290 *next-generation sequencing data*. Available at: <http://www.biomedcentral.com/1471-2105/14/289>.

7291 Sanderson, B.J. *et al.* (2020) "A targeted sequence capture array for phylogenetics and population genomics in
7292 the salicaceae," *Applications in Plant Sciences*, 8(10). Available at: <https://doi.org/10.1002/aps3.11394>.

7293 Santamaría, L. *et al.* (2003) 'PLANT PERFORMANCE ACROSS LATITUDE: THE ROLE OF PLASTICITY AND LOCAL
7294 ADAPTATION IN AN AQUATIC PLANT', *Ecology*, 84(9), pp. 2454–2461. Available at: [https://doi.org/10.1890/02-](https://doi.org/10.1890/02-0431)
7295 [0431](https://doi.org/10.1890/02-0431).

7296 Sasaki, E. *et al.* (2017) 'The genetic architecture of the network underlying flowering time variation in
7297 *Arabidopsis thaliana*', *bioRxiv*, p. 175430. Available at: <https://doi.org/10.1101/175430>.

7298 Schena, M. *et al.* (1995) 'Quantitative monitoring of gene expression patterns with a complementary DNA
7299 microarray', *Science (New York, N.Y.)*, 270(5235), pp. 467–470. Available at:
7300 <https://doi.org/10.1126/SCIENCE.270.5235.467>.

7301 Schliep, K.P. (2011) 'phangorn: phylogenetic analysis in R', *Bioinformatics*, 27(4), pp. 592–593. Available at:
7302 <https://doi.org/10.1093/BIOINFORMATICS/BTQ706>.

7303 Schmittgen, T.D. and Livak, K.J. (2008) 'Analyzing real-time PCR data by the comparative CT method', *Nature*
7304 *Protocols 2008 3:6*, 3(6), pp. 1101–1108. Available at: <https://doi.org/10.1038/nprot.2008.73>.

7305 Schmitz, R.J. and Amasino, R.M. (2007) 'Vernalization: A model for investigating epigenetics and eukaryotic
7306 gene regulation in plants', *Biochimica et Biophysica Acta (BBA) - Gene Structure and Expression*, 1769(5–6), pp.
7307 269–275. Available at: <https://doi.org/10.1016/J.BBAEXP.2007.02.003>.

- 7308 Schneider, C.A., Rasband, W.S. and Eliceiri, K.W. (2012a) 'NIH Image to ImageJ: 25 years of image analysis',
7309 *Nature Methods*, pp. 671–675. Available at: <https://doi.org/10.1038/nmeth.2089>.
- 7310 Schneider, C.A., Rasband, W.S. and Eliceiri, K.W. (2012b) 'NIH Image to ImageJ: 25 years of image analysis',
7311 *Nature Methods* 2012 9:7, 9(7), pp. 671–675. Available at: <https://doi.org/10.1038/nmeth.2089>.
- 7312 Schopf, J.W. *et al.* (1996) 'Activation of floral meristem identity genes in Arabidopsis', *Nature* 1996 384:6604,
7313 384(6604), pp. 59–62. Available at: <https://doi.org/10.1038/384059a0>.
- 7314 Schueler, S., Schlünzen, K.H. and Scholz, F. (2005) 'Viability and sunlight sensitivity of oak pollen and its
7315 implications for pollen-mediated gene flow', *Trees - Structure and Function*, 19(2), pp. 154–161. Available at:
7316 <https://doi.org/10.1007/S00468-004-0376-1/FIGURES/4>.
- 7317 Schuerger, A.C., Brown, C.S. and Stryjewski, E.C. (1997) *Anatomical Features of Pepper Plants (Capsicum*
7318 *annuum L.) Grown under Red Light-emitting Diodes Supplemented with Blue or Far-red Light, Sedgwick Rd.,*
7319 *Bldg.*
- 7320 Schwaegerle, K.E. and Levin, D.A. (1990) 'Quantitative genetics of seed size variation in Phlox', *Evolutionary*
7321 *Ecology* 1990 4:2, 4(2), pp. 143–148. Available at: <https://doi.org/10.1007/BF02270911>.
- 7322 Seppänen, M.M. *et al.* (2010) "Vernalization response of Phleum pratense and its relationships to stem
7323 lignification and floral transition," *Annals of Botany*, 106(5), pp. 697–707. Available at:
7324 <https://doi.org/10.1093/aob/mcq174>.
- 7325 Servedio, M.R. *et al.* (2014) 'Not Just a Theory—The Utility of Mathematical Models in Evolutionary Biology',
7326 *PLoS Biology*, 12(12), p. 1002017. Available at: <https://doi.org/10.1371/JOURNAL.PBIO.1002017>.
- 7327 Shachar-Hill, B. *et al.* (2013) 'Mercury-sensitive water channels as possible sensors of water potentials in
7328 pollen', *Journal of Experimental Botany*, 64(16), pp. 5195–5205. Available at:
7329 <https://doi.org/10.1093/JXB/ERT311>.
- 7330 SHAPIRO, S.S. and WILK, M.B. (1965) 'An analysis of variance test for normality (complete samples)',
7331 *Biometrika*, 52(3–4), pp. 591–611. Available at: <https://doi.org/10.1093/BIOMET/52.3-4.591>.
- 7332 Sharma, N. *et al.* (2020a) 'The molecular mechanism of vernalization in Arabidopsis and cereals: role of
7333 Flowering Locus C and its homologs', *Physiologia Plantarum*, 170(3), pp. 373–383. Available at:
7334 <https://doi.org/10.1111/PPL.13163>.
- 7335 Sharma, N. *et al.* (2020b) 'The molecular mechanism of vernalization in Arabidopsis and cereals: role of
7336 Flowering Locus C and its homologs', *Physiologia Plantarum*, 170(3), pp. 373–383. Available at:
7337 <https://doi.org/10.1111/PPL.13163>.
- 7338 Shavrukov, Y. *et al.* (2017) 'Early flowering as a drought escape mechanism in plants: How can it aid wheat
7339 production?', *Frontiers in Plant Science*, 8, p. 1950. Available at:
7340 <https://doi.org/10.3389/FPLS.2017.01950/BIBTEX>.

7341 Sheldon, C.C. *et al.* (2009) 'Mechanisms of gene repression by vernalization in Arabidopsis', *Plant Journal*,
7342 59(3), pp. 488–498. Available at: <https://doi.org/10.1111/J.1365-313X.2009.03883.X>.

7343 Sherry, R.A. and Lord, E.M. (1996) 'DEVELOPMENTAL STABILITY IN LEAVES OF CLARKIA TEMBLORIENSIS
7344 (ONAGRACEAE) AS RELATED TO POPULATION OUTCROSSING RATES AND HETEROZYGOSITY', *Evolution*, 50(1),
7345 pp. 80–91. Available at: <https://doi.org/10.1111/J.1558-5646.1996.TB04474.X>.

7346 Shindo, C. *et al.* (2005) 'Role of FRIGIDA and FLOWERING LOCUS C in Determining Variation in Flowering Time
7347 of Arabidopsis', *Plant Physiology*, 138(2), pp. 1163–1173. Available at: <https://doi.org/10.1104/PP.105.061309>.

7348 Shirsekar, G. *et al.* (2021) 'Fine-scale Population Structure of North American Arabidopsis thaliana Reveals
7349 Multiple Sources of Introduction from Across Eurasia', *bioRxiv*, p. 2021.01.22.427575. Available at:
7350 <https://doi.org/10.1101/2021.01.22.427575>.

7351 Singh, N. *et al.* (2013) 'Comparison of SSR and SNP Markers in Estimation of Genetic Diversity and Population
7352 Structure of Indian Rice Varieties', *PLOS ONE*, 8(12), p. e84136. Available at:
7353 <https://doi.org/10.1371/JOURNAL.PONE.0084136>.

7354 Singh, N.B. *et al.* (1995) *RELATIONSHIP OF PLANT HEIGHT, DAYS TO FLOWERING AND MATURITY TO GRAIN
7355 YIELD IN SHORT-DURATION DETERMINATE PIGEONPEA* • Available at: www.IndianJournals.com.

7356 Smýkal, P. *et al.* (2018) 'The Impact of Genetic Changes during Crop Domestication', *Agronomy 2018, Vol. 8*,
7357 *Page 119*, 8(7), p. 119. Available at: <https://doi.org/10.3390/AGRONOMY8070119>.

7358 Song, Y.H. *et al.* (2014) 'Distinct roles of FKF1, GIGANTEA, and ZEITLUPE proteins in the regulation of constans
7359 stability in Arabidopsis photoperiodic flowering', *Proceedings of the National Academy of Sciences of the
7360 United States of America*, 111(49), pp. 17672–17677. Available at:
7361 https://doi.org/10.1073/PNAS.1415375111/SUPPL_FILE/PNAS.201415375SI.PDF.

7362 Song, Y.H., Ito, S. and Imaizumi, T. (2013) 'Flowering time regulation: photoperiod- and temperature-sensing in
7363 leaves', *Trends in plant science*, 18(10), p. 575. Available at: <https://doi.org/10.1016/J.TPLANTS.2013.05.003>.

7364 Sork, V.L. (2018) 'Genomic Studies of Local Adaptation in Natural Plant Populations', *Journal of Heredity*,
7365 109(1), pp. 3–15. Available at: <https://doi.org/10.1093/jhered/esx091>.

7366 Soto-Cerda, B.J. *et al.* (2014) 'The potential of pale flax as a source of useful genetic variation for cultivated flax
7367 revealed through molecular diversity and association analyses', *Molecular Breeding*, 34(4), pp. 2091–2107.
7368 Available at: <https://doi.org/10.1007/S11032-014-0165-5/FIGURES/3>.

7369 South, A. (2011) *rworldmap: A New R package for Mapping Global Data*. Available at:
7370 <http://www.un.org/millenniumgoals/bkgd>.

7371 Speck, T. and Burgert, I. (2011) 'Plant Stems: Functional Design and Mechanics',
7372 <http://dx.doi.org/10.1146/annurev-matsci-062910-100425>, 41, pp. 169–193. Available at:
7373 <https://doi.org/10.1146/ANNUREV-MATSCI-062910-100425>.

7374 Sporemex, 2022. Spores and Pollens. [online] Sporomex.co.uk. Available at:
7375 <<https://www.sporomex.co.uk/technology/spores-and-pollens>> [Accessed 8 August 2022].

7376 Sreedharan, S.P., Kumar, A. and Giridhar, P. (2018) 'Primer design and amplification efficiencies are crucial for
7377 reliability of quantitative PCR studies of caffeine biosynthetic N-methyltransferases in coffee', *3 Biotech*, 8(11).
7378 Available at: <https://doi.org/10.1007/s13205-018-1487-5>.

7379 Starlabgroup.com. 2022. 96-Well PCR Plate, Skirted, Low Profile, White - STARLAB. [online] Available at:
7380 <[https://www.starlabgroup.com/GB-en/product/96-well-pcr-plate-skirted-low-profile-white-e1403-
7381 5209.html](https://www.starlabgroup.com/GB-en/product/96-well-pcr-plate-skirted-low-profile-white-e1403-5209.html)> [Accessed 30 May 2022].

7382 Steel, R.G.D. and Torrie, J.H., 1960. Principles and procedures of statistics. Principles and procedures of
7383 statistics.

7384 Stinchcombe, J.R. *et al.* (2004) 'A latitudinal cline in flowering time in *Arabidopsis thaliana* modulated by the
7385 flowering time gene FRIGIDA', *Proceedings of the National Academy of Sciences*, 101(13), pp. 4712–4717.
7386 Available at: <https://doi.org/10.1073/PNAS.0306401101>.

7387 Stöcklin, J., Kuss, P. and Pluess, A.R. (2009) 'Genetische Vielfalt, phänotypische Variabilität und lokale
7388 Adaptation in der Alpenlandschaft: Fallstudien mit vier alpinen Pflanzenarten', *Botanica Helvetica*, 119(2), pp.
7389 125–133. Available at: <https://doi.org/10.1007/s00035-009-0065-1>.

7390 Sultan, S.E. (2000) "Phenotypic plasticity for plant development, function and life history," *Trends in Plant
7391 Science*, 5(12), pp. 537–542. Available at: [https://doi.org/10.1016/s1360-1385\(00\)01797-0](https://doi.org/10.1016/s1360-1385(00)01797-0).

7392 Sun, J. *et al.* (2019) 'Photoperiod sensitivity of Canadian flax cultivars and 5-azacytidine treated early flowering
7393 derivative lines', *BMC Plant Biology*, 19(1). Available at: <https://doi.org/10.1186/s12870-019-1763-5>.

7394 Sun, M. and Wong, K.C. (2001) 'Genetic structure of three orchid species with contrasting breeding systems
7395 using RAPD and allozyme markers', *American Journal of Botany*, 88(12), pp. 2180–2188. Available at:
7396 <https://doi.org/10.2307/3558379>.

7397 Sun, S. and Frelich, L.E. (2011) 'Flowering phenology and height growth pattern are associated with maximum
7398 plant height, relative growth rate and stem tissue mass density in herbaceous grassland species', *Journal of
7399 Ecology*, 99(4), pp. 991–1000. Available at: <https://doi.org/10.1111/J.1365-2745.2011.01830.X>.

7400 Sundaresan, V. (2005) 'Control of seed size in plants', *Proceedings of the National Academy of Sciences of the
7401 United States of America*, 102(50), p. 17887. Available at: <https://doi.org/10.1073/PNAS.0509021102>.

7402 Suter, L. *et al.* (2014) 'Gene Regulatory Variation Mediates Flowering Responses to Vernalization along an
7403 Altitudinal Gradient in *Arabidopsis*', *Plant Physiology*, 166(4), pp. 1928–1942. Available at:
7404 <https://doi.org/10.1104/PP.114.247346>.

7405 Sveinsson, S. *et al.* (2014a) 'Phylogenetic pinpointing of a paleopolyploidy event within the flax genus (*Linum*)
7406 using transcriptomics', *Annals of Botany*, 113(5), pp. 753–761. Available at:
7407 <https://doi.org/10.1093/AOB/MCT306>.

7408 Szakiel, A., Pączkowski, C. and Henry, M. (2010) "Influence of environmental abiotic factors on the content of
7409 saponins in plants," *Phytochemistry Reviews*, 10(4), pp. 471–491. Available at:
7410 <https://doi.org/10.1007/s11101-010-9177-x>.

7411 Szpiech, Z.A. and Rosenberg, N.A. (2011) 'On the size distribution of private microsatellite alleles', *Theoretical*
7412 *Population Biology*, 80(2), pp. 100–113. doi:10.1016/j.tpb.2011.03.006.

7413 Tajima, F. (1989) 'Statistical Method for Testing the Neutral Mutation Hypothesis by DNA Polymorphism',
7414 *Genetics*, 123(3), p. 585. Available at: <https://doi.org/10.1093/GENETICS/123.3.585>.

7415 Tatham, S., 2022. PuTTY User Manual. [online] PuTTY. Available at:
7416 <<https://the.earth.li/~sgtatham/putty/0.77/html/doc/>> [Accessed 21 September 2022].

7417 Teper-Bamnolker, P. and Samach, A. (2005) 'The Flowering Integrator FT Regulates SEPALLATA3 and FRUITFULL
7418 Accumulation in Arabidopsis Leaves', *The Plant Cell*, 17(10), p. 2661. Available at:
7419 <https://doi.org/10.1105/TPC.105.035766>.

7420 *The GNU Bourne-Again Shell* (no date). Available at: <https://tiswww.case.edu/php/chet/bash/bashtop.html>
7421 (Accessed: 5 May 2022).

7422 Thomashow, M.F. (1999) "Plant cold acclimation: Freezing tolerance genes and regulatory mechanisms,"
7423 *Annual Review of Plant Physiology and Plant Molecular Biology*, 50(1), pp. 571–599. Available at:
7424 <https://doi.org/10.1146/annurev.arplant.50.1.571>.

7425 Tigano, A. and Friesen, V.L. (2016) 'Genomics of local adaptation with gene flow', *Molecular Ecology*. Blackwell
7426 Publishing Ltd, pp. 2144–2164. Available at: <https://doi.org/10.1111/mec.13606>.

7427 Toräng, P. *et al.* (2015) 'Large-scale adaptive differentiation in the alpine perennial herb *Arabis alpina*', *The*
7428 *New phytologist*, 206(1), pp. 459–470. Available at: <https://doi.org/10.1111/NPH.13176>.

7429 Tork, D.G. *et al.* (2022) 'Ideotype Selection of Perennial Flax (*Linum* spp.) for Herbaceous Plant Habit Traits',
7430 *Agronomy*, 12(12). Available at: <https://doi.org/10.3390/agronomy12123127>.

7431 Trelease, W., 1931. *Winter botany*. New York: Dover Publications.

7432 Tukey, J., 1949. Comparing Individual Means in the Analysis of Variance. *Biometrics*, 5(2), p.99.

7433 Unacademy, 2022. Draw a Labelled Diagram of the Pollen Grain. [online] Unacademy.com. Available at:
7434 <<https://unacademy.com/content/question-answer/biology/draw-a-labelled-diagram-of-the-pollen-grain/>>
7435 [Accessed 8 August 2022].

7436 Uysal, H. *et al.* (2010) 'Genetic diversity of cultivated flax (*Linum usitatissimum* L.) and its wild progenitor pale
7437 flax (*Linum bienne* Mill.) as revealed by ISSR markers', *Genetic Resources and Crop Evolution*, 57(7), pp. 1109–
7438 1119. Available at: <https://doi.org/10.1007/S10722-010-9551-Y/FIGURES/4>.

7439 Uysal, H. *et al.* (2012) 'Variation in phenotypic characters of pale flax (*Linum bienne* Mill.) from Turkey',
7440 *Genetic Resources and Crop Evolution*, 59(1), pp. 19–30. Available at: <https://doi.org/10.1007/S10722-011->
7441 9663-Z/FIGURES/10.

7442 Valentim, F.L. *et al.* (2015a) 'A quantitative and dynamic model of the Arabidopsis flowering time gene
7443 regulatory network', *PLoS one*, 10(2). Available at: <https://doi.org/10.1371/JOURNAL.PONE.0116973>.

7444 Vandesompele, J. *et al.* (2002) 'Accurate normalization of real-time quantitative RT-PCR data by geometric
7445 averaging of multiple internal control genes', *Genome biology*, 3(7). Available at: <https://doi.org/10.1186/GB->
7446 2002-3-7-RESEARCH0034.

7447 Venables, W.N. and Ripley, B.D. (2002a) 'Random and Mixed Effects', pp. 271–300. Available at:
7448 https://doi.org/10.1007/978-0-387-21706-2_10.

7449 Verdu, C.F. *et al.* (2016) 'Dealing with paralogy in RADseq data: in silico detection and single nucleotide
7450 polymorphism validation in *Robinia pseudoacacia* L.', *Ecology and Evolution*, 6(20), pp. 7323–7333. Available
7451 at: <https://doi.org/10.1002/ECE3.2466>.

7452 Vieira, M.L.C. *et al.* (2016) 'Microsatellite markers: what they mean and why they are so useful', *Genetics and*
7453 *Molecular Biology*, 39(3), p. 312. Available at: <https://doi.org/10.1590/1678-4685-GMB-2016-0027>.

7454 Villanueva, R.A.M. and Chen, Z.J. (2019) 'ggplot2: Elegant Graphics for Data Analysis (2nd ed.)',
7455 <https://doi.org/10.1080/15366367.2019.1565254>, 17(3), pp. 160–167. Available at:
7456 <https://doi.org/10.1080/15366367.2019.1565254>.

7457 Vuletin Selak, G. *et al.* (2013) 'The effect of temperature and genotype on pollen performance in olive (*Olea*
7458 *europaea* L.)', *Scientia Horticulturae*, 156, pp. 38–46. Available at:
7459 <https://doi.org/10.1016/J.SCIENTA.2013.03.029>.

7460 Walker, J.M. (1969) 'One-Degree Increments in Soil Temperatures Affect Maize Seedling Behavior', *Soil Science*
7461 *Society of America Journal*, 33(5), pp. 729–736. Available at:
7462 <https://doi.org/10.2136/SSSAJ1969.03615995003300050031X>.

7463 Walters, D., Keil, D. and Walters, B., 1996. *Vascular plant taxonomy*. Dubuque, Iowa: Kendall/Hunt, p.598.

7464 Wang, J. *et al.* (2018) 'A major locus controls local adaptation and adaptive life history variation in a perennial
7465 plant', *Genome Biology*, 19(1). Available at: <https://doi.org/10.1186/s13059-018-1444-y>.

7466 Wang, Z. *et al.* (2012) 'The genome of flax (*Linum usitatissimum*) assembled de novo from short shotgun
7467 sequence reads', *The Plant journal : for cell and molecular biology*, 72(3), pp. 461–473. Available at:
7468 <https://doi.org/10.1111/J.1365-313X.2012.05093.X>.

7469 Watanabe, S. *et al.* (2011) 'A Map-Based Cloning Strategy Employing a Residual Heterozygous Line Reveals that
7470 the GIGANTEA Gene Is Involved in Soybean Maturity and Flowering', *Genetics*, 188(2), pp. 395–407. Available
7471 at: <https://doi.org/10.1534/GENETICS.110.125062>.

7472 Wee, A.K. et al. (2020) 'Genetic structures across a biogeographical barrier reflect dispersal potential of four
7473 Southeast Asian mangrove plant species', *Journal of Biogeography*, 47(6), pp. 1258–1271.
7474 doi:10.1111/jbi.13813.

7475 Weigel, D. et al. (1992) 'LEAFY controls floral meristem identity in Arabidopsis', *Cell*, 69(5), pp. 843–859.
7476 Available at: [https://doi.org/10.1016/0092-8674\(92\)90295-N](https://doi.org/10.1016/0092-8674(92)90295-N).

7477 Weir, B.S. (2012) 'Estimating F-statistics: A historical view', *Philosophy of science*, 79(5), p. 637. Available at:
7478 <https://doi.org/10.1086/667904>.

7479 Welch, S.M., Dong, Z. and Roe, J.L. (2004) *Modelling gene networks controlling transition to flowering in*
7480 *Arabidopsis*. Available at: www.cropscience.org.au.

7481 Werner, C.R. et al. (2020) 'How Population Structure Impacts Genomic Selection Accuracy in Cross-Validation:
7482 Implications for Practical Breeding', *Frontiers in Plant Science*, 11, p. 2028. Available at:
7483 <https://doi.org/10.3389/FPLS.2020.592977/BIBTEX>.

7484 White, N.J. and Butlin, R.K. (2021) 'Multidimensional divergent selection, local adaptation, and speciation',
7485 *Evolution*, 75(9), pp. 2167–2178. Available at: <https://doi.org/10.1111/evo.14312>.

7486 Whitlock, M.C. (2015) 'Modern approaches to local adaptation', *American Naturalist*, 186, pp. S1–S4. Available
7487 at: <https://doi.org/10.1086/682933>.

7488 Wickham, H., 2016. *ggplot2: elegant graphics for data analysis*. springer.

7489 Wilczek, A.M. et al. (no date) 'Genetic and physiological bases for phenological responses to current and
7490 predicted climates'. Available at: <https://doi.org/10.1098/rstb.2010.0128>.

7491 Williams, G. C. 1966. *Adaptation and natural selection*. Princeton University Press, Princeton, NJ.

7492 Williams, I.J., Martin, A.P. and Clark, S.J. (1990) 'Pollination requirements of linseed (*Linum usitatissimum*)',
7493 *The Journal of Agricultural Science*, 115(3), pp. 347–352. Available at:
7494 <https://doi.org/10.1017/S0021859600075778>.

7495 Williams, J.H. and Reese, J.B. (2019) 'Evolution of development of pollen performance', *Current Topics in*
7496 *Developmental Biology*, pp. 299–336. doi:10.1016/bs.ctdb.2018.11.012.

7497 Wolfe, Lorne M and Wolfe, L M (1995) 'The genetics and ecology of seed size variation in a biennial plant,
7498 *Hydrophyllum appendiculatum* (Hydrophyllaceae)', *Oecologia* 1995 101:3, 101(3), pp. 343–352. Available at:
7499 <https://doi.org/10.1007/BF00328821>.

7500 Wright, S. (1965) 'The Interpretation of Population Structure by F-Statistics with Special Regard to Systems of
7501 Mating', *Evolution*, 19(3), p. 395. Available at: <https://doi.org/10.2307/2406450>.

7502 Wright, S. (1978) 'The Relation of Livestock Breeding to Theories of Evolution', *Journal of Animal Science*, 46(5),
7503 pp. 1192–1200. Available at: <https://doi.org/10.2527/JAS1978.4651192X>.

7504 Wu, C. *et al.* (2016) 'Effect of plant age and vernalization on bolting, plant growth and enzyme activity of garlic
7505 (*Allium sativum* L.)', *Scientia Horticulturae*, 201, pp. 295–305. Available at:
7506 <https://doi.org/10.1016/J.SCIENTA.2016.02.006>.

7507 Xie, D. *et al.* (2018) 'Genome-wide association study identifying candidate genes influencing important
7508 agronomic traits of Flax (*Linum usitatissimum* L.) using SLAF-seq', *Frontiers in Plant Science*, 8, p. 2232.
7509 Available at: <https://doi.org/10.3389/FPLS.2017.02232/BIBTEX>.

7510 Yan, L., Chouw, N. and Jayaraman, K. (2014) 'Flax fibre and its composites - A review', *Composites Part B:
7511 Engineering*, 56, pp. 296–317. Available at: <https://doi.org/10.1016/j.compositesb.2013.08.014>.

7512 Yan, X. and Su, X., 2009. Linear regression analysis. Singapore: World Scientific.

7513 Yang, C.Y. *et al.* (2007) 'Luminescence of aniline blue in hydrophobic cavity of BSA', *Journal of Photochemistry
7514 and Photobiology A: Chemistry*, 188(1), pp. 51–55. Available at:
7515 <https://doi.org/10.1016/J.JPHOTOCHEM.2006.11.017>.

7516 Yang, H. *et al.* (2016) 'Genetic Divergence between *Camellia sinensis* and Its Wild Relatives Revealed via
7517 Genome-Wide SNPs from RAD Sequencing', *PLOS ONE*, 11(3), p. e0151424. Available at:
7518 <https://doi.org/10.1371/JOURNAL.PONE.0151424>.

7519 You, F.M. *et al.* (2017) 'Genetic variability of 27 traits in a core collection of flax (*Linum usitatissimum* L.)',
7520 *Frontiers in Plant Science*, 8. Available at: <https://doi.org/10.3389/fpls.2017.01636>.

7521 You, F.M. *et al.* (2018) 'Chromosome-scale pseudomolecules refined by optical, physical and genetic maps in
7522 flax', *The Plant Journal*, 95(2), pp. 371–384. Available at: <https://doi.org/10.1111/TPJ.13944>.

7523 Zamir, D., Tanksley, S.D. and Jones, R.A. (1981) 'Low temperature effect on selective fertilization by pollen
7524 mixtures of wild and cultivated tomato species', *Theoretical and Applied Genetics*, 59(4), pp. 235–238.
7525 Available at: <https://doi.org/10.1007/BF00265501/METRICS>.

7526 Zhang, J. *et al.* (2020) 'Genomic Comparison and Population Diversity Analysis Provide Insights into the
7527 Domestication and Improvement of Flax', *iScience*, 23(4). Available at:
7528 <https://doi.org/10.1016/j.isci.2020.100967>.

7529 Zhao, P. *et al.* (2022) 'Precipitation and local environment shape the geographic variation of seed size across
7530 natural populations of sand rice (*Agriophyllum squarrosum*)', *Journal of Experimental Botany*, 73(16), pp.
7531 5682–5697. Available at: <https://doi.org/10.1093/JXB/ERAC231>.

7532 Zheng, X. *et al.* (2012) 'A high-performance computing toolset for relatedness and principal component
7533 analysis of SNP data', *Bioinformatics*, 28(24), pp. 3326–3328. Available at:
7534 <https://doi.org/10.1093/BIOINFORMATICS/BTS606>.

7535 Zimmerman, S.J., Aldridge, C.L. and Oyler-McCance, S.J. (2020) 'An empirical comparison of population genetic
7536 analyses using microsatellite and SNP data for a species of conservation concern', *BMC Genomics*, 21(1), pp. 1–
7537 16. Available at: <https://doi.org/10.1186/S12864-020-06783-9/FIGURES/6>.

7538 Zinn, K.E., Tunc-Ozdemir, M. and Harper, J.F. (2010) 'Temperature stress and plant sexual reproduction:
7539 uncovering the weakest links', *Journal of Experimental Botany*, 61(7), p. 1959. Available at:
7540 <https://doi.org/10.1093/JXB/ERQ053>.

7541 Zipper, H. *et al.* (2004) 'Investigations on DNA intercalation and surface binding by SYBR Green I, its structure
7542 determination and methodological implications', *Nucleic Acids Research*, 32(12), p. e103. Available at:
7543 <https://doi.org/10.1093/NAR/GNH101>.

7544 Zohary, D. and Hopf, M. (2000) 'Domestication of plants in the Old World: the origin and spread of cultivated
7545 plants in West Asia, Europe and the Nile Valley.', *Domestication of plants in the Old World: the origin and*
7546 *spread of cultivated plants in West Asia, Europe and the Nile Valley*. [Preprint], (Ed.3).

7547 Zonia, L. and Munnik, T. (2004) 'Osmotically Induced Cell Swelling versus Cell Shrinking Elicits Specific Changes
7548 in Phospholipid Signals in Tobacco Pollen Tubes', *Plant Physiology*, 134(2), p. 813. Available at:
7549 <https://doi.org/10.1104/PP.103.029454>.

7550

7551

7552