



Title	Establishing consensus on key public health indicators for the monitoring and evaluating childhood obesity interventions: a Delphi panel study
Authors(s)	O'Donnell, Shane, Doyle, Gerardine, O'Malley, Grace, Browne, Sarah, O'Connor, James, Mars, Monica, Kechadi, Tahar
Publication date	2020-11-17
Publication information	O'Donnell, Shane, Gerardine Doyle, Grace O'Malley, Sarah Browne, James O'Connor, Monica Mars, and Tahar Kechadi. "Establishing Consensus on Key Public Health Indicators for the Monitoring and Evaluating Childhood Obesity Interventions: A Delphi Panel Study" 20 (November 17, 2020).
Publisher	Springer Nature
Item record/more information	http://hdl.handle.net/10197/24355
Publisher's version (DOI)	10.1186/s12889-020-09814-y

Downloaded 2023-10-31T04:02:18Z

The UCD community has made this article openly available. Please share how this access benefits you. Your story matters! (@ucd_oa)



© Some rights reserved. For more information

RESEARCH ARTICLE

Open Access



Establishing consensus on key public health indicators for the monitoring and evaluating childhood obesity interventions: a Delphi panel study

Shane O'Donnell^{1*}, Gerardine Doyle², Grace O'Malley^{3,4}, Sarah Browne⁵, James O'Connor⁶, Monica Mars⁷ and M-Tahar M. Kechadi⁶

Abstract

Background: Childhood obesity is influenced by myriad individual, societal and environmental factors that are not typically reflected in current interventions. Socio-ecological conditions evolve and require ongoing monitoring in terms of assessing their influence on child health. The aim of this study was to identify and prioritise indicators deemed relevant by public health authorities for monitoring and evaluating childhood obesity interventions.

Method: A three-round Delphi Panel composed of experts from regions across Europe, with a remit in childhood obesity intervention, were asked to identify indicators that were a priority in their efforts to address childhood obesity in their respective jurisdictions. In Round 1, 16 panellists answered a series of open-ended questions to identify the most relevant indicators concerning the evaluation and subsequent monitoring of interventions addressing childhood obesity, focusing on three main domains: built environments, dietary environments, and health inequalities. In Rounds 2 and 3, panellists rated the importance of each of the identified indicators within these domains, and the responses were then analysed quantitatively.

Results: Twenty-seven expert panellists were invited to participate in the study. Of these, 16/27 completed round 1 (59% response rate), 14/16 completed round 2 (87.5% response rate), and 8/14 completed the third and final round (57% response rate). Consensus (defined as > 70% agreement) was reached on a total of 45 of the 87 indicators (49%) across three primary domains (built and dietary environments and health inequalities), with 100% consensus reached for 5 of these indicators (6%).

Conclusion: Forty-five potential indicators were identified, pertaining primarily to the dietary environment, built environment and health inequalities. These results have important implications more widely for evaluating interventions aimed at childhood obesity reduction and prevention.

Keywords: Delphi, Obesity, Intervention, Childhood obesity, Evaluation, Public health, Health policy

* Correspondence: shane.odonnell@ucd.ie

¹School of Sociology, University College Dublin, D04 V1W8, Dublin, Ireland

Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Childhood obesity is considered one of the key public health challenges of the twenty-first century, with worldwide prevalence having increased from < 1% in 1975 to 6–8% in 2016 [1]. Since the publication of the influential *Foresight Tackling Obesities: Future Choices* report, the aim of which was to use the scientific evidence base from multidisciplinary research to identify the broad range of factors that influence obesity [2], policymakers have been increasingly encouraged to adopt a 'system-level approach' in their efforts to design and implement public health interventions to address childhood obesity. This acknowledges that the causes of obesity and overweight are multiple and complex, and the development of effective interventions is dependent on addressing myriad determinants rooted simultaneously in individuals' biology as well as the socio-ecological conditions in which they grow up, learn, play, and work [3–5]. A significant gap in current literature is a socio-ecological inventory of factors important in the aetiology of childhood obesity, that simultaneously provides an indication of the relative weighting of factors in terms of priority for intervention. It is anticipated that such an inventory may be important for the monitoring and evaluation systems that collect multiple sources of data with a systems-level framework. Thus, the aim of this study was to identify indicators pertaining to childhood obesity that should be prioritised in the monitoring and evaluation of childhood obesity interventions. This study was conducted as part of a wider EC H2020 funded project entitled *Big Data Against Childhood Obesity* (BigO) which aims to develop a technology system that leverages the potential of big data to support public health authorities in formulating effective, context-specific policies and interventions addressing childhood obesity [4, 6, 7]. An indicator can be broadly defined as a measure that reveals relative positions in a given area (e.g. health). When evaluated at regular intervals, an indicator can point out the direction of change in different populations and across time [8, 9]. Choosing an appropriate set of indicators relevant to the monitoring and evaluation of health interventions requires a high degree of judgement and consensus-building among potential users and other interested parties [10]. Given the large number of variables that influence obesity at a system level (the Foresight report identifies over 100), and therefore the multiplicity of indicators that could be seen as a priority, a Delphi approach was adopted to reach consensus amongst an international panel of public health experts [11].

Methods

Study design

The Delphi process utilises a range of qualitative and quantitative approaches to reach decision-making among isolated anonymous respondents and is

considered a valid methodology for obtaining a collective view from a group of experts where the only alternative is entirely subjective or based on anecdotal evidence [11, 12]. Delphi guides expert group opinion towards a final decision through triangulation of subjective group judgments [13] and is achieved through the application of **five** core features: (i) anonymity; (ii) iteration (iii) controlled feedback; (iv) statistical group response, and (v) expert input [9, 12]. In this study, we attempt to capture a broad set of indicators in the domains of the dietary environment, built environment and health inequalities. Following other studies that have adopted a Delphi approach for policy planning [14], the research involved five phases: indicator screening and categorisation; recruitment; evaluation; re-evaluation; and final consensus. A summary of the procedural steps taken in this Delphi panel study is as follows: firstly, a list of indicators was identified and categorised by qualitatively analysing initial pilot interviews conducted with experts in the public health sector in Ireland, Greece, and Sweden. Three broad domains of interest were identified, including the built environment, dietary environment and health inequalities. Then, in the first round of the Delphi study, a larger panel of experts was recruited and asked to identify the indicators that were most relevant for the design (and subsequent evaluation) of future policies aimed at reducing childhood obesity in these domains. These indicators were then analysed quantitatively over two rounds. In the second round, participants were asked to give importance ratings for each indicator. Responses were then quantitatively analysed to establish the level of consensus agreement for each indicator. Finally, in the third round, the panel of experts were made aware of the consensus levels of each indicator and asked whether they would like to change their response based on this finding. After importance ratings were given for each indicator, a final level of consensus was established. This process is described in more detail in the following section.

Survey development

This study followed a traditional 3-round decision-making Delphi panel study [9, 15]. In order to identify the domains of interest that would feature in the Delphi panel, an initial pilot study was carried out with PHAs in Ireland, Sweden and Greece. These countries were selected because they were represented within the BigO research consortium. A list of indicators was drafted based on the indicators deemed of particular relevance to childhood obesity outlined in the Foresight report and Systems Map to help guide the discussion [2] (Table 1). Given that the PHAs time to participate was limited, some indicators had to be omitted, so we chose those we felt would be most relevant to PHAs. During

the pilot stage, the PHAs were asked to review and comment on the relative importance of each indicator. We also sought their opinion regarding indicators that were missing. The first pilot interview was conducted face-to-face with a PHA based in Ireland. The same list was then discussed with PHAs in Sweden and Greece via video call. This process helped to further refine the list of indicators by eliminating those that were deemed less relevant to the PHAs. For example, the domain related to individual-level satiety control was deemed not as relevant to PHAs. Three authors (SO'D, GO'M, and SB) analysed the responses to identify themes and propose statements. The outcome of this phase was the identification of a number of areas of interest which could be categorised into three main domains:

- Built environment
- Dietary environment
- Health inequalities

Based on this outcome, a set of questions was developed to be used in the first round of the Delphi survey. These questions were initially piloted amongst a small number of PHAs in Ireland and Sweden ($N=3$) to

ensure that they were congruent, easy to understand and culturally appropriate (Table 2).

Recruitment & Data Collection

A purposeful stratified sampling technique was used to identify potential panellists [16]. Following Novakowski and Wellar (2008), strict criteria for the selection of expert panellists was developed to include only those with:

- Direct influence over policy at both local and national levels
- Indirect impact by shaping policy through scholarly research and public advocacy

Following Keeney et al. [11], we aimed to recruit a total of 15–20 panellists. A panel of experts was identified and 27 invitations to participate were sent to PHAs in Ireland, Sweden, Greece, the Netherlands, Spain and the UK. The countries selected for participation (with the exception of the UK) reflected where the organisations involved in the H2020 BigO project were based and also where the BigO system would initially be rolled out. Thus, we were able to leverage the local knowledge and extensive networks of the BigO research teams in

Table 1 Indicators presented to PHAs as part of initial pilot interviews

Domain	Measurable Influencing Factors	Measurable Influencing Factors (cont.)
Physical Activity	Gender	Walkability
	Age	Urban planning
	BMI Grouping	Sedentary time
	Disability	Accessibility
	Prevalence of inactivity	Time and type
	Inequality	Affordability
Social Environmental	Water quality	Green space
	Air quality	Food waste
	Proximity to motorway	School policy
	Density of food retailers	Organic pollutant
	Location of food retailers	Weather pattern
	Food desserts	Advertising/marketing density
	Psychosocial distress	Stigma
Financial	House price	Disposable income
	Homeownership	Household food spend
	Area deprivation	
Individual level	Medical history	Genetics
	Health service utilisation	Family history

Table 2 Delphi Panellists

Country	Key panellist type	N =
Sweden	Academic	4
	Policy maker	
Greece	Academic	2
	Policy maker	
Ireland	Academic	7
	Policy maker	
	Public health advocate	
Netherlands	Policy maker	1
	Academic	
Spain	Academic	1
	Academic	
United Kingdom	Academic	1
	Academic	

each of these jurisdictions to purposefully identify individuals with the requisite expertise. Of those who were invited, 16 initially agreed to take part. Surveys were then distributed to panellists via the online survey Qualtrics™ in English only. In the first round, panellists were given the freedom to respond to each of four questions in narrative form and encouraged to elaborate on their responses in an in-depth manner.

Data analysis

Round 1

Once the 16 responses were returned content analysis was performed. Each response was analysed line-by-line to identify distinct statements made by panellists that related to measures or indicators [11]. Subsequently, statements similar in nature were grouped together under one 'prototypical' statement to reduce the size of the subsequent questionnaires and ease the burden on panellists in completing later rounds [11]. One issue that emerged at this stage of the analysis was that the term 'measure' in each question was interpreted by some to mean 'actions to be taken' rather than specific indicators or measures of progress. As such, the wording of each statement was changed to avoid further confusion or ambiguity in subsequent rounds. The result was the generation of a list of 87 statements, which remained constant in all subsequent rounds.

Round 2

Each statement was uploaded onto Qualtrics and panellists were invited to rate the relative importance of each statement using a 5-point Likert scale. Panellists were

given 3 weeks to submit their responses. Regular reminders ($N=3$) were sent to those who had not yet completed the survey. A predetermined level of consensus, known as the percentage agreement, was set at $\geq 70\%$. Only indicators rated as either 'very' or 'extremely important', by at least 70% or more of the panellists, were deemed to have reached consensus [6, 17–20]. The results of Round 2 were analysed quantitatively using the Software Package for Social Science (SPSS) [21], to calculate the central tendency (Mean, Median, Mode) and level of dispersion (Standard Deviation), in order to present information concerning the collective judgements of respondents in Round 3.

Round 3

We retained all indicators from Round 2 so that each indicator had an equal opportunity to gain the highest rating of importance possible [11]. Panellists were asked to rate the same statements as presented in Round 2. However, each statement was accompanied by two additional pieces of information: the rating the individual panellist assigned to each statement in the previous round, as well as the average response of the group (mode). Panellists were then invited to consider if they would like to change their response in light of the aggregate opinion of other panellists or stand by their original response. Finally, Round 3 was analysed quantitatively using descriptive statistics as in Round 2.

Results

Of the 27 experts invited to participate in this Delphi study, 16 completed round 1 (59% response rate), 14 completed round 2 (87.5% response rate) and 8 completed round 3 (57% response rate). Table 4 below shows a summary of the statements and the level of consensus achieved in each domain. In Round 1, there was no consensus level, as this round was designed to establish the indicators in each domain. In Round 2, 43 of the 87 indicators (49%) passed the consensus agreement threshold (Table 3). This rose to 45 indicators (52%) in Round 3, with variation in the individual indicators that reached consensus between rounds 2 and 3.

Additionally, in Round 3, 100% consensus was reached for some indicators in the **Built Environment** domain ($n=2$), **Dietary Environment** domain ($n=2$), and **Uncategorised** domain ($n=1$), with no indicator reaching full consensus (consensus range 87.5–0%) in the **Health Inequalities** domain (Table 4). Stability of consensus (< 10% variation) was achieved between rounds 2 and 3 for all four domains [6]. Of the remaining indicators, 12 were just below the percentage agreement level (70%), reaching a consensus level of 62.5%. From the five indicators that reached complete consensus (100%), two were indicators that relate to the school environment,

Table 3 Summary of the number of indicators that reached consensus agreement in each round by domain

Statement domains	Number of statements in each domain			Proportion of statements where consensus was achieved (n)		
	Round 1	Round 2	Round 3	Round 1	Round 2	Round 3
Built environment		34	34	47% (16)		47% (16)
Dietary environment		30	30	40% (12)		50% (15)
Health inequalities		19	19	74% (14)		68% (13)
Uncategories		4	4	25% (1)		25% (1)

one economic (food prices), one governmental (cycle lanes) and one personal health (BMI). For the remainder of indicators that passed consensus threshold, more than half were related to government resources in a given environment (access to facilities or the lack thereof and exposure to unhealthy food sources and their advertisements), economic inequalities, and school structures (access to facilities and healthy food). Indicators that were least likely to reach consensus related to access and affordability of alcoholic beverages (availability of off-licences/liquor stores and minimum alcohol unit pricing). A graph showing the numbers of indicators that reached consensus is also shown [Fig. 1].

Discussion

This study engaged with PHAs and advisors to identify and prioritise indicators deemed important for the monitoring and evaluation of childhood obesity interventions. In our study, Consensus (defined as >70% agreement) was reached on a total of 45 of the 87 indicators (49%) across three primary domains (built and dietary environments and health inequalities), with 100% consensus reached for 5 of these indicators (6%). The consensus reached in a large number of factors underscores the level of complexity involved in obesity intervention and the challenges implementing change in these domains.

With certain exceptions [6, 22–24], few studies have explored indicators relevant to the development and monitoring of childhood obesity-related policies. One study employed the Delphi panel technique and focused on the prioritisation of intervention conditions in childhood obesity [25]. Others have focused on research priorities among clinical and academic experts [22, 26]. For example, the Determinants of Nutrition and Eating framework (DONE) study, which employed a three-round Delphi panel study to examine the priorities of policymakers with respect to healthy eating, identified a similar set of indicators with respect to the dietary environment to those elucidated in this study [22]. Our study further builds on this work by addressing other domains deemed important by PHAs - the built environment and wider inequalities related to childhood obesity and includes a number of PHA from a variety of different countries with contrasting health policies. Interestingly,

given the growing awareness of the role that the social determinants of health play in the aetiology of childhood obesity, it is perhaps somewhat unexpected that there was not 100% consensus for any indicator within the health inequalities domain [4]. However, it is also worth noting that although 100% consensus was not reached, it did have the highest rate of consensus (74% = 14 out of 19 indicators with a range of 87.5 to 0% consensus) compared to other domains and their indicators. Furthermore, those indicators pertaining to health inequalities that reached consensus were more likely to relate to wider patterns of social and economic inequality (e.g. unemployment, local deprivation indices, etc.). In contrast, many of the indicators in this domain that failed to reach consensus were centred on interventions that rely on a greater degree of personal agency and individual-level action (e.g. access to cooking and growing schools programmes, access to community gardens, availability and access to universal primary care services etc.). This is perhaps a reflection of a growing awareness amongst policy makers that while individual-level interventions may be helpful in improving overall population health, they may be less effective in reducing relative health inequalities [5].

This study has a number of important implications. First, the results highlight the variety and range of data that would be relevant to PHAs and the identification of indicators across multiple domains and underscores the system-based focus of PHAs in Europe. As outlined in the Foresight report, there are over 100 factors that contribute to childhood obesity and these are often interdependent (e.g. lower pricing of energy-dense food, excessive marketing of energy-dense food and excessive consumption of energy-dense food by children). Despite high-quality, international and longitudinal research programmes relevant to childhood obesity, integrating and monitoring multi-level system factors still presents a challenge. The Childhood Obesity Surveillance Initiative (COSI) from the World Health Organisation (WHO), for example, collects individual-level data on anthropometry, dietary and physical activity patterns, screen time, and sleep, among others. Recent findings from 6 to 9-year-olds in the WHO European Region demonstrates substantial country-level differences in healthy and unhealthy dietary habits, with patterns that cannot be fully

Table 4 List of indicators and results (Arranged Thematically) - Indicator consensus after 3 rounds of the Delphi Study degree of consensus

Domains	Question for each domain	SD
Built Environment	In the design (and subsequent evaluation) of future policies aimed at improving the built environment to reduce childhood obesity what, in your opinion, are likely to be the most useful measures in your jurisdiction?	
Dietary Environment	In the design (and subsequent evaluation) of future policies aimed at improving the dietary environment to reduce childhood obesity what, in your opinion, are likely to be the most useful measures in your jurisdiction?	
Health Inequalities	In the design (and subsequent evaluation) of future policies aimed at improving inequalities in childhood obesity-related outcomes what, in your opinion, are likely to be the most useful measures in your jurisdiction?	
Uncategorised	Are there any other measures related to childhood obesity prevention and monitoring that you feel are important to capture at community/population levels?	
Measurements	Degree of Consensus (%) Round 3	
Built Environment		
Q1_36 School infrastructure that includes spaces for organized or individual exercise/activity	100%	0.5
Q1_57 Availability of safe cycling paths	100%	0.34
Dietary Environment		
Q2_28 The pricing environment of foods	100%	0.5
Q2_42 Availability of tap water in schools	100%	0.43
Uncategorised		
Q4_24 BMI changing over time in terms of mean, median and shape of distribution	100%	0.43
Built Environment		
Q1_31 Availability of outdoor facilities	87.50%	1
Q1_35 Recreational space within walking space of distance of home	87.50%	0.71
Q1_39 Availability of open spaces in neighbourhood.	87.50%	0.35
Q1_41 Density of public parks	87.50%	0.7
Q1_42 Proximity of green space to home	87.50%	0.71
Q1_47 Design of walkways and physical environment	87.50%	0.71
Dietary Environment		
Q2_29 Range and diversity of food retailers	87.50%	0.66
Q2_30 Number of fast food advertisements within the community	87.50%	0.66
Q2_34 Digital exposure to food advertising	87.50%	1.2
Q2_35 Availability of fresh fruit and vegetables	87.50%	0.7
Q2_36 Retail environment within supermarkets	87.50%	1.2
Q2_38% of taxes on sugar	87.50%	0.66
Q2_39% of taxes imposed on foods high in fat and salt	87.50%	0.66
Q2_41 Availability of healthy meals in school and preschool	87.50%	1
Q2_54 Infant feeding indicators	87.50%	0.73
Inequalities		
Q3_9 Employment status or socio-economic status of family	87.50%	1
Q3_10 Local deprivation indices	87.50%	0.99
Q3_11 Area based food poverty statistics	87.50%	0.97
Q3_12 Number of households experiencing food poverty	87.50%	0.99

Table 4 List of indicators and results (Arranged Thematically) - Indicator consensus after 3 rounds of the Delphi Study degree of consensus (*Continued*)

Domains	Question for each domain	SD	
Q3_13	Unemployment levels	87.50%	0.99
Q3_14	Child and family – Living on public assistance	87.50%	0.97
Q3_16	Ethnicity	87.50%	0.87
Q3_18	Family structure	87.50%	0.5
Q3_22	Relative income poverty in line with government measures on inequality	87.50%	0.7
Q3_23	Consistent poverty in line with government measures on inequality	87.50%	0.7
Q3_24	Deprivation in line with government measures on inequality	87.50%	0.71
Q3_25	Additional metrics of social inequality used both individually and as components of census-derived, weighted, area-level deprivation indices	87.50%	0.7
Built Environment			
Q1_37	Affordability of organized sports: club fees and costs	75%	0.71
Q1_40	Number of public parks	75%	1.1
Q1_44	Availability of public transport to access green spaces	75%	0.71
Q1_49	Sports and physical activity participation levels	75%	1.1
Q1_51	Opening hours of green spaces	75%	0.71
Q1_52	Quality of lighting within green spaces	75%	0.8
Q1_53	Level of reported anti-social behaviour in green spaces / open spaces	75%	1
Q1_56	Accessibility of public transport via foot	75%	1.05
Dietary environment			
Q2_24	Density and type of food retailer in proximity to school	75%	0.83
Q2_31	Advertisements in proximity of schools	75%	2
Q2_40	Availability of High Fat Salt Sugar foods/drinks	75%	0.73
Q2_46	Availability of energy-dense foods in vending machines and cafeterias in the school environment.	75%	1.1
Inequalities			
Q3_8	Education level statistics	75%	1.3
Built Environment			
Q1_27	Availability of indoor facilities	62.50%	0.86
Q1_30	Price of indoor facilities	62.50%	1.1
Q1_34	Price of outdoor facilities	62.50%	1.1
Q1_50	Accessibility of public parks via public transport	62.50%	0.8
Q1_55	Access facilities for fitness training at no cost to the individual	62.50%	1.3
Q1_60	Number of cars located on the road outside home	62.50%	1.3
Dietary environment			
Q2_37	Location where children do their shopping	62.50%	1.2
Q2_45	Data on the range and quality of food served in the workplace settings	62.50%	1.1
Inequalities			
Q3_15	Health literacy	62.50%	1.3
Q3_20	Availability and access to school meals schemes	62.50%	1.05

Table 4 List of indicators and results (Arranged Thematically) - Indicator consensus after 3 rounds of the Delphi Study degree of consensus (*Continued*)

Domains	Question for each domain	SD
Uncategorised		
Q4_26	Monitoring of diets of families/children considered at risk by social workers/social services	62.50% 1.3
Q4_27	Whole production chain needs to be attended to	62.50% 1.12
Built Environment		
Q1_38	Numbers of people who use recreational spaces	50% 1
Q1_43	Proximity of blue space to home	50% 0.5
Q1_48	Child and parental attitudes and knowledge of their built environment	50% 0.87
Q1_62	Number of physical activity referrals / prescriptions in general practice	50% 1.2
Dietary environment		
Q2_26	Density and type of food retailer along school commute	50%
Q2_27	Tracking data on portion sizes in fast-food retailers, other restaurants and single-serving snacks	50% 0.83
Q2_33	Food advertising at specific times	50% 1.1
Q2_50	Exposure to alcohol advertising at sporting events	50% 1
Q2_52	Availability of café/bars	50% 1.05
Q2_44	Access to community gardens	42.90% 0.99
Inequalities		
Q3_17	Gender	42.90% 0.83
Q3_19	Availability and access to universal primary health services	42.90% 1.25
Built Environment		
Q1_29	Density of indoor facilities	37.50% 1.2
Q1_32	Number of outdoor facilities	37.50% 1.3
Q1_33	Density of outdoor facilities	37.50% 1.1
Q1_58	Availability of walk to school groups	37.50% 0.86
Dietary environment		
Q2_25	Density and type of food retailer in proximity to home	37.50% 0.71
Q2_32	% of processed food items with clear and accurate front of pack labelling	37.50% 1
Q2_43	Access to allotments- % of school with allotments	37.50% 1.1
Q2_48	Minimum alcohol unit pricing	37.50% 0.97
Inequalities		
Q3_26	Availability of cooking and growing skills programmes	37.50% 1.05
Uncategorised		
Q4_25	Provide special support to individual cases	37.50% 0.66
Built Environment		
Q1_46	GIS based measures of cyclability	25% 0.6
Dietary environment		
Q2_51	Alcohol use as a contributor to adolescent obesity	25% 0.78
Q2_53	Access to farmers markets	25% 1.17
Built Environment		
Q1_45	GIS based measures of walkability	14.30% 0.64
Q1_28	Number of indoor facilities	12.50% 0.93

Table 4 List of indicators and results (Arranged Thematically) - Indicator consensus after 3 rounds of the Delphi Study degree of consensus (Continued)

Domains	Question for each domain	SD
Q1_59 Availability of Park and Ride schemes	12.50%	1.1
Dietary environment		
Q2_47 Availability of off-licences/liquor stores	12.50%	1.1
Inequalities		
Q3_21 Level of referrals to GP	0%	0.7

explained [27]. With respect to physical activity, the Health Behaviour in School-Aged Children (HBSC) Study reports substantial variation in physical activity participation among school-aged children from 29 countries. National differences in the physical, socio-cultural, economic and policy environment account for individual-level differences [28]. Our study will complement existing knowledge and help to guide researchers who wish to further integrate socio-ecological factors into monitoring systems. Furthermore, our findings are particularly timely as they can help to guide the development of emerging big data solutions for the monitoring and surveillance of ongoing efforts to reduce and prevent childhood obesity through a systems approach [4, 6, 7].

A second implication drawn from our findings is the importance of co-developing monitoring systems in collaboration with policy makers and other relevant stakeholders who have good knowledge of their own local context and the data repositories required or in some cases already available for use. For example, to help evaluate a policy intervention on childhood obesity

prevention, such as regulating the distance a fast-food retailer could be built in proximity to a school, multiple related data sources are needed across the domains identified in this study. To monitor change over time and evaluate the effects of this intervention, data is needed at the level of the child, at the level of the school and at the community level. To successfully source, store, retrieve, analyse, and present the socioeconomic-, health-, dietary, economic- and geospatial data needed in the above example, a number of considerations are required to meet legal, data protection, privacy and ethical requirements in addition to the necessary standards, protocols and technological aspects. Monitoring and surveillance systems need to address these concerns so that public health officials can evaluate and monitor the effectiveness of such interventions on childhood obesity and so that decision-making can be facilitated for scaling up successful interventions. However, it is also important to acknowledge that obesity is the outcome of a complex adaptive system and the success of any intervention is dependent on the wider social context in




Fig. 1 Delphi Panel Study: indicator consensus

which it is deployed and embedded. As one panellist pointed out, exclusive reliance on quantitative analysis of macro-level indicators (regardless of the sophistication of the predictive models used) may be too reductionist to provide the holistic picture needed to understand the nuances of this complex adaptive system. Future research must therefore examine how best to incorporate both quantitatively driven, macro-level indicators and qualitative data, which is more appropriate for capturing the wider context and lived experiences of children or communities for whom obesity interventions may be implemented.

Strengths and weaknesses

Through the use of the Delphi method [12–15, 29], this study captured the collective feedback [$N=16$] from some of the leading obesity experts in Europe across a wide range of European regions with varying health systems. For example, based on the European Core Health Indicator of Expenditure on health care as a percentage of GDP in 2018, Sweden (10.90) and the Netherlands (9.97) are above average while Ireland (6.93) and Greece (7.72) are below average [30].

Consensus was found across a wide range of statements which in turn enabled the research team to delineate a list of indicators which can be used to improve and inform the further development of systems to monitor and evaluate ongoing public health efforts to reduce and prevent childhood obesity. Many of these indicators, particularly pertaining to the built environment, have yet to be prioritised in the extant literature. In addition, the study also highlights where there is perhaps less certainty among policy makers (e.g. indicators pertaining to health inequalities) and therefore areas for further inquiry.

While a key strength of the study is its leveraging of expert knowledge of some of the leading authorities with a remit in childhood obesity prevention in Europe with a wide variation in health policy, recruitment of panellists was nonetheless limited to 6 countries and, as such, their perceptions, understandings and insights may not be generalisable to all European countries or to jurisdictions outside Europe. Furthermore, the recruitment relied on the networks of the BigO research team, and therefore, some element of bias in the selection of participants at both the piloting and main data collection phases cannot be ruled out. Another issue was the diminished response rate between rounds, with half the participants lost between Rounds 1 & 3. Given that Delphi panel studies rely on two or more iterative rounds, the content of which can often be repetitive, an attrition rate of up to 50% is not uncommonly reported in the literature [31]. One possible explanation for the attrition in this context of this study may have been the onerous nature of assessing large numbers of statements in each

round. It has been noted that in instances where Delphi panel studies include a high number of items, panellists are less likely to participate all the way through to study completion [32]. Given many of these participants would have been leading authorities in their respective jurisdictions, their time to complete the survey would have been limited. Whilst the research team were conscious of this risk from the outset of the study, the importance of minimising the burden of participants also had to be balanced against the need to ensure that the final list of indicators was as comprehensive as possible to reflect the complexity of childhood obesity-related policy.

Nevertheless, the final sample size [$N=8$] sits between what the NIHR Health Technology Assessment group [33] identifies as the lower threshold for participation in consensus groups at which point validity begins to decline rapidly [$N=6$], and the upper threshold at which point any improvements in validity may become subject to diminishing returns [$N=12$]. Indeed, there is little existing theoretical or empirical evidence that increasing larger sample sizes in Delphi studies necessarily leads to more reliable or valid results [6, 34–38].

In addition to examining the robustness of the indicators identified in this study among PHAs based in other jurisdictions, future studies should also incorporate the voices of other important stakeholders, such as supra-national organisations (e.g. WHO); industry (e.g. insurance companies, device manufacturers, marketing agencies); the healthcare sector; investors; non-health related government agencies; teachers, children and their parents. Furthermore, the development of co-designed tools and platforms, acceptability and usability studies (particularly focused on privacy and data sharing considerations) should be carried out to further inform how such systems might be used in practice by PHAs, researchers or citizens.

Conclusion

This study contributes to current childhood obesity literature by providing expert consensus on a wide range of key socio-ecological influences and measures that are amenable to policy change, particularly in the areas of the built environment, dietary environment and health inequalities. Factors that should especially be prioritised include the school infrastructure that includes space for organised or individual exercise activity, availability of safe cycling paths, the pricing environment of foods, availability of tap water in schools and BMI changes over time. The volume and complexity of pertinent measures that should be collected require the implementation of smart technology solutions. The findings, therefore, have implications both in informing childhood obesity interventions and in developing systems that can monitor and evaluate those efforts.

Abbreviations

BMI: Body Mass Index; COSI: Childhood Obesity Surveillance Initiative; DONE: Determinants of Nutrition and Eating framework; EU: European Union; GO'M: Grace O'Malley; HSBC: Health Behaviour in School-Aged Children; PHA: Public Health Authority; SB: Sarah Browne; SO'D: Shane O'Donnell; SPSS: Software Package for Social Science; UK: United Kingdom; WHO: World Health Organisation

Acknowledgements

The authors would like to thank the participants who gave their time and expertise to this study as well as the anonymous reviewers for their very helpful comments in previous versions of this manuscript.

Authors' contributions

SO'D, GO'M, GD designed the study and performed the data collection and analysis. All other authors contributed to the interpretation and analysis of results, reviewed and commented on the manuscript. The author(s) read and approved the final manuscript.

Funding

The work leading to these results was part of the EU H2020 project: BigO: Big data against childhood Obesity (Grant No. 727688, <https://bigoprogram.eu>). This project was part of the European Community's Health, demographic change and well-being Programme of EU H2020. GO'M was supported by the Royal College of Surgeons in Ireland StAR Fellowship. The funders had no role in the design of the study and collection, management, analysis, and interpretation of the data and in writing the manuscript. The authors and their contributions to the manuscript are independent of the funder.

Availability of data and materials

The datasets generated and/or analysed during the current study are not publicly available to protect the anonymity of participants but are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

Ethical exemption for the study was granted by the University College Dublin Human Research Ethics Committee (Reference Number: HS-E-18-49).

Consent for publication

Participants consented to their data being used for publication in scientific journals.

Competing interests

The authors declare no competing financial interests.

Author details

¹School of Sociology, University College Dublin, D04 V1W8, Dublin, Ireland. ²UCD College of Business and UCD Geary Institute for Public Policy, University College Dublin, Dublin A94 XF34, Ireland. ³School of Physiotherapy, Division of Population Health Sciences, Royal College of Surgeons Ireland, D02 YN77 Dublin, Ireland. ⁴Children's Health Ireland, Temple Street, D01 XD99 Dublin, Ireland. ⁵School of Public Health, Physiotherapy & Sports Science, Woodview House, Belfield, University College Dublin, Dublin 04V1W8, Ireland. ⁶School of Computer Science, Insight Centre for Data Analytics, University College Dublin, D04 V1W8 Dublin, Ireland. ⁷Division of Human Nutrition and Health, Wageningen University and Research, PO Box 17, NL-6700, AA, Wageningen, The Netherlands.

Received: 16 October 2019 Accepted: 2 November 2020

Published online: 17 November 2020

References

1. Jaacks LM, Vandevijvere S, Pan A, McGowan CJ, Wallace C, Imamura F, et al. The obesity transition: stages of the global epidemic. *Lancet Diabetes Endocrinol*. 2019;7(3):231–40.
2. Butland B, Jebb S, Kopelman P, McPherson K, Thomas S, Mardell J, et al. Tackling obesity: future choices-project report. London: Department of Innovation, Universities and Skills; 2007. Available at: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/287937/07-1184x-tackling-obesity-future-choices-report.pdf [Accessed 05/02/2019].
3. Timmins KA, Green MA, Radley D, Morris MA, Pearce J. How has big data contributed to obesity research? A review of the literature. *Int J Obes*. 2018; 42(12):1951–62.
4. Morris MA, Wilkins E, Timmins KA, Bryant M, Birkin M, Griffiths C. Can big data solve a big problem? Reporting the obesity data landscape in line with the foresight obesity system map. *Int J Obes*. 2018;42(12):1963–95.
5. Link BG, Phelan J. Social conditions as fundamental causes of disease. *J Health Soc Behav*. 1995;1:80–94.
6. Vogel C, Zwolinsky S, Griffiths C, Hobbs M, Henderson E, Wilkins E. A Delphi study to build consensus on the definition and use of big data in obesity research. *Int J Obes*. 2019;17:1.
7. Diou C, Sarafis I, Papapanagiotou V, Alagiadoglou L, Lekka I, Filos D, Stefanopoulos L, Kilintzis V, Maramis C, Karavidopoulou Y, Maglaveras N. BigO: A public health decision support system for measuring obesogenic behaviors of children in relation to their local environment. *arXiv preprint arXiv:2005.02928*. 2020 May 6. Available at: <https://arxiv.org/pdf/2005.02928.pdf> Accessed 29/08/2020.
8. Nardo, M., Saisana, M., Saltelli, A., Tarantola, S., Hoffman, A. & Giovannini, E. Handbook on constructing composite indicators: methodology and user guide, EC Joint Research Centre and OECD Statistics Directorate and the Directorate for Science, Technology and Industry, OECD publishing. 2008. <https://www.oecd.org/sdd/42495745.pdf> Accessed 01/09/2020.
9. Gibney S, Sexton E, Shannon S. Measuring what matters: achieving consensus on a positive aging Indicator set for Ireland. *J Aging Soc Policy*. 2018;8:1–6.
10. Lippeveld T, Sauerborn R, Bodart C. Design and implementation of health information systems. Geneva: World Health Organization; 2000. <https://apps.who.int/iris/handle/10665/42289> Accessed 15/09/2020.
11. Keeney S, Hasson F, McKenna HP. The Delphi technique in nursing and health research. Oxford: Wiley-Blackwell; 2011.
12. Broomfield D, Humphris GM. Using the Delphi technique to identify the cancer education requirements of general practitioners. *Med Educ*. 2001; 35(10):928–37.
13. Cantrill JA, Sibbald B, Buetow S. The Delphi and nominal group techniques in health services research. *Int J Pharm Pract*. 1996;4(2):67–74.
14. Sébastien L, Bauler T, Lehtonen M. Can indicators bridge the gap between science and policy? An exploration into the (non) use and (non) influence of indicators in EU and UK policy making. *Nat Culture*. 2014;9(3):316–43.
15. Iqbal S, Pison-Young L. Methods-the Delphi method—a guide from Susanne Iqbal and Laura Pison-Young. *Psychologist*. 2009;22(7):598.
16. Research methods in anthropology: qualitative and quantitative approaches. London: Rowman & Littlefield; 2017.
17. Williams PL, Webb C. The Delphi technique: a methodological discussion. *J Adv Nurs*. 1994;19(1):180–6.
18. Dajani JS, Sincoff MZ, Talley WK. Stability and agreement criteria for the termination of Delphi studies. *Technol Forecast Soc Chang*. 1979;13(1): 83–90.
19. Green P. The Content of a College-Level Outdoor Leadership Course. 1982. <https://eric.ed.gov/?id=ED276546> Accessed 04/02/2019.
20. Dias A, Mooren T, Kleber RJ. Reducing consequences of child maltreatment during adulthood by public health actions: a Delphi study. *Eur J Pub Health*. 2019;29(3):425–31.
21. IBM Corp. SPSS Statistics for Windows, Version 24.0. Armonk: IBM Corp.; 2016.
22. Stok FM, Hoffmann S, Volkert D, Boeing H, Ensenauer R, Stelmach-Mardas M, et al. The DONE framework: Creation, evaluation, and updating of an interdisciplinary, dynamic framework 2.0 of determinants of nutrition and eating. *PLoS One*. 2017;12(2):e0171077.
23. Haynes E, Palermo C, Reidlinger DP. Modified policy-Delphi study for exploring obesity prevention priorities. *BMJ Open*. 2016;6(9):e011788.
24. Kim MJ, Sung E, Choi EY, Ju YS, Park EW, Cheong YS, et al. Delphi Survey for Designing a Intervention Research Study on Childhood Obesity Prevention. *Korean J Family Med*. 2017;38(5):284 Foster BA, Winkler P, Weinstein K, Parra-Medina D. Developing a patient-centered outcome for targeting early childhood obesity across multiple stakeholders. *BMC obesity*. 2018; 5(1):39.
25. Taylor RW, Robinson A, Espinel PT, Baur LA, Wake M, Sabin MA. Research priorities in 2012 for the effective management of childhood obesity. *Clin Obes*. 2013;3(1–2):3–6.

26. Byrne S, Wake M, Blumberg D, Dibley M. Identifying priority areas for longitudinal research in childhood obesity: Delphi technique survey. *Int J Pediatr Obes.* 2008;3(2):120–9.
27. Williams J, Buoncristiano M, Nardone P, Rito AI, Spinelli A, Hejgaard T, Kierkegaard L, Nurk E, Kunešová M, Musić Milanović S, García-Solano M. A snapshot of European Children's eating habits: results from the fourth round of the WHO European childhood obesity surveillance initiative (COSI). *Nutrients.* 2020;12(8):2481.
28. Weinberg D, Stevens GW, Bucksch J, Inchley J, de Looze M. Do country-level environmental factors explain cross-national variation in adolescent physical activity? A multi-level study in 29 European countries. *BMC Public Health.* 2019;19(1):680.
29. Novakowski N, Wellar B. Using the Delphi technique in normative planning research: methodological design considerations. *Environ Plan A.* 2008;40(6):1485–500.
30. European Core Health Indicators (ECHI) European Commission. Available at :<http://ec.europa.eu/health/dyna/echi/datatool/index.cfm?indlist=77a> [Accessed 22/08/2020].
31. Khodyakov D, Grant S, Denger B, Kinnett K, Martin A, Peay H, Coulter I. Practical considerations in using online modified-Delphi approaches to engage patients and other stakeholders in clinical practice guideline development. *Patient-Patient-Centered Outcomes Res.* 2020;13(1):11–21.
32. Gargon E, Crew R, Burnside G, Williamson PR. Higher number of items associated with significantly lower response rates in COS Delphi surveys. *J Clin Epidemiol.* 2019;108:110–20.
33. Murphy MK, Black NA, Lamping DL, McKee CM, Sanderson CF, Askham J, Marteau T. Consensus development methods, and their use in clinical guideline development. *Health Technol Assessment (Winchester, England).* 1998;2(3):i–88.
34. Powell C. The Delphi technique: myths and realities. *J Adv Nurs.* 2003;41(4):376–82.
35. De Villiers MR, De Villiers PJ, Kent AP. The Delphi technique in health sciences education research. *Med Teacher.* 2005;27(7):639–43.
36. Slade SC, Dionne CE, Underwood M, Buchbinder R. Standardised method for reporting exercise programmes: protocol for a modified Delphi study. *BMJ Open.* 2014;4(12):e006682.
37. Okoli C, Pawlowski SD. The Delphi method as a research tool: an example, design considerations and applications. *Inf Manag.* 2004;42(1):15–29.
38. Hasson F, Keeney S, McKenna H. Research guidelines for the Delphi survey technique. *J Adv Nurs.* 2000;32(4):1008–15.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

