



ANIMAL SCIENCE

Embryonic and post-embryonic development of the spider *Polybetes pythagoricus* (Sparassidae): A biochemical point of view

SOFIA ROMERO, ALDANA LAINO, GABRIEL MOLINA, MONICA CUNNINGHAM & CARLOS FERNANDO GARCIA

Abstract: Analysis of energy expense during development has achieved special interest through time on account of the crucial role of the consumption of resources required for offspring survival. Spider eggs have a fixed composition as well as some initial energy that is supplied by mothers. These resources are necessary to support the metabolic expense not only through the embryonic period but also during the post-embryonic period, as well as for post emerging activities before spiderlings become self-sustaining. Depletion of these resources would be critical for spiders since it could give rise to prey competition as well as filial cannibalism. Even though spiders represent a megadiverse order, information regarding the metabolic requirements during spiders development is very scarce. In this study, we analyse the changes in protein, lipid and carbohydrate content as well as the variation in lipovitellin reserves and hemocyanin content during *Polybetes pythagoricus* development. Our results show that lipovitellins and phospholipids represent the major energy source throughout embryonic and post-embryonic development. Lipovitellin apolipoproteins are gradually consumed but are later depleted after dispersion. Phosphatidylethanolamine is mainly consumed during the post-embryonic period, while triacylglycerides are consumed after juveniles' dispersion. Finally, hemocyanin concentration starts to increase in postembryonic stages.

Key words: development, fatty acids, hemocyanin, lipids, lipovitellin, spider.

INTRODUCTION

With more than 49,000 species described (World Spider Catalog 2021), spiders represent one of the most important and megadiverse groups and are found in different habitats (Foelix 2011, Grismado et al. 2014). They are of great importance to humans both for the harmful effect that the venom present in some species generates on human health (Vetter & Isbister 2008) and for their role in the biological pest control of different agroecosystems owing to their being generalist predators (Saba et al. 2020). Although existing information on the development and reproduction of spiders is very

abundant, biochemical and metabolic studies on these topics are very scarce.

The study of energetic expense during development has achieved significant interest over time, and oviparous animals have become a special model for biological studies on development. Their eggs have a fixed composition and, therefore, a certain amount of energy whose variations can be followed throughout embryo development (Garcia et al. 2008, Trabalon et al. 2018). In the case of spiders, mothers lay their eggs in silk cocoons built by themselves (Foelix 2011) thus determining the amount and quality of the resources they allocate to their offspring (Salomon et al. 2011,

Ruhland et al. 2016). The presence of this cocoon influences the development of spiders due to the fact that some post-embryonic stages also take place inside this structure. Because of this, initial composition and energy resources of the eggs are needed not only to support the metabolic demand throughout the embryonic period but also to bear the metabolic cost during the post-embryonic stage and the post emergent activities before spiderlings become self-sustaining (Anderson 1978, Ruhland et al. 2016, Trabalon et al. 2018).

The different resources accumulated by mothers and transferred to the eggs may influence the offspring's traits (Salomon et al. 2011). The process of synthesis and storage of nutrients in the oocytes is known as vitellogenesis. During vitellogenesis, oocytes size increases by the progressive accumulation of yolk, lipid droplets, as well as various proteins, carbohydrates and lipids, which are required for offspring development (Byrne et al. 1989, Choi & Moon 2003, Fruttero et al. 2011, Romero et al. 2018, Thompson & Russel 1999, Trabalon et al. 1992). The depletion of yolk is a critical time for spiders, since it could give rise to competition for prey and filial cannibalism (Yip & Rayor 2014). Information regarding yolk synthesis in spiders is quite recent and very scarce (Bednarek et al. 2019, Laino et al. 2011b, 2013, Romero et al. 2019). Moreover, the use of energetic resources during development is variable and the energetic demand required for embryonic development can be supplied by lipids, proteins and carbohydrates (Campos et al. 2006, Mohamed 2000, Santana et al. 2014). However, the only contribution about the consumption of energetic resources in spiders was that by Trabalon and collaborators, who observed the key role played by triacylglycerides, carbohydrates, and proteins during the embryonic and post-embryonic

development of the spider *Pardosa saltans* (Trabalon et al. 2018).

Focusing on spiders' diversity, the main aim of this research is to make headway in the understanding of the role played by the different biomolecules during *Polybetes pythagoricus* spider development. *P. pythagoricus* is a South American spider found in Chile, Bolivia, Brazil, Paraguay, Uruguay and in north-central Argentina, where different biochemical and physiological aspects have been analysed (Cunningham & Pollero 1996, Cunningham et al. 1994, 2002, 2007, Laino et al. 2009, 2011a, b, 2015b, Romero et al. 2018, 2019). Copulation and mating, along with other relevant data about *P. pythagoricus* life cycle were previously described (Galiano 1979, Scioscia 1984), as well as post-embryonic development (Galiano 1971). However, there is no information about the different energetic resources and their use during *P. pythagoricus* development.

The aims of this study were: firstly, to determine the quantitative changes in the total content of carbohydrates and proteins. Secondly, to characterize, through immunological techniques, lipovitellins consumption and hemocyanin content, respectively. Lastly, to provide information on the lipid and fatty acids dynamics during *P. pythagoricus* development. We hypothesized that: (1) carbohydrates are gradually consumed throughout the development process; 2) lipovitellins are gradually consumed throughout development; (3) hemocyanin is synthesized in the advanced stages of development; 4) triacylglycerides are the main energy source used throughout the development process, 5) phospholipid content decreases as lipovitellins are consumed.

MATERIALS AND METHODS

Ethics statement

The species used in the present experiments (*P. pythagoricus*) is not endangered or protected. Our research conforms to the legal requirements for the treatment of animals in research facilities using invertebrate species and taking care for them using accepted ethical standards.

Spider collection and rearing

Spiders were captured in a forest of *Eucalyptus* sp. of Martin Rodriguez Park in the city of Ensenada (34°52'56" S, 57°56'07" W) and in Pereyra Iraola Park (Berazategui-La Plata) Argentina (34°50'39" S, 58°10'55" W). Capture permits were No 117/16 Protected Areas, Province of Buenos Aires. They were housed individually in cylindrical terrariums (10 cm in diameter × 5 cm in height) without any substrate, fed once a week with Turkestan cockroach (*Shelfordella tartara*) or Dubia roach (*Blaptica dubia*), and kept at 25 ± 1 °C under a 14-h light – 10-h dark photoperiodic cycle. Vitellogenic females were maintained in these conditions until oviposition. All the egg-sacs were built by the spider females. An ootheca was obtained from each female and was kept in the laboratory at 25 ± 1 °C.

Experimental groups

A total of 32 egg-sacs (1.21 ± 0.48 g/egg-sac) were used and divided into eight experimental groups to analyse the different energy resources and their variation during *P. pythagoricus* development. These experimental stages were numbered from 1 to 8, the first five are stages that are develop inside the cocoon and the last three are stages after emergence (outside the cocoon). To standardize the analysis, stages inside the cocoon (stages 1 to 5) were temporarily separated by 5-day difference between each of them. Since the stages used in this study

started to number from the embryonic period, and in order to allow a comparison between our study with previously reported studies (Galiano 1971, Scioscia 1984), we highlight that the post-embryonic stages 4 and 5 of this study are the non-mobile free juveniles described by Galiano (1971), while the first emergence stage analysed in the present study (stage 6) are the mobile free juveniles characterized by the same author (Galiano 1971).

The stages analysed of the embryonic period (stages after oviposition and before hatching) were: Stage 1 (egg stage, less than 12-h laying); Stage 2 (5 days after oviposition) and Stage 3 (10 days after oviposition). The stages analysed of the post-embryonic period (after hatching) that take place inside the egg-sac were: Stage 4 (15 days after oviposition) and Stage 5 (20 days after oviposition). Stages analysed outside the egg-sac were: Stage 6 (25 ± 1 day after oviposition, juveniles in gregarious stage at the moment of emergence from egg-sac), Stage 7 (35 ± 1 day after oviposition, juveniles after the first moult outside the egg-sac) and Stage 8 (42 ± 1 day after oviposition, juveniles in dispersal stage one week after the first moult outside the egg-sac). Moreover, two complementary stages were processed and analysed: an advanced juvenile stage (AJ, 1.2 ± 0.4 g) and previtellogenic females as an adult stage (A, 2.01 ± 0.6 g), which represent stages that have already been fed.

General measurements

All egg-sacs were weighed in a Mettler-Toledo New Classic MS-204S analytical balance. Before being processed, the individuals were observed using a Leica S8 APO stereomicroscope and photos were taken with a digital camera.

The individuals of each egg-sac were weighed together and frozen-killed at -20 °C, and later on, homogenized in 1.5 ml buffer potassium phosphate regardless of your weight

(50 mM, pH 7.4) with the addition of a protease inhibitor cocktail 1/1000 (v/v) (Sigma Chemical Co., St. Louis, MO, USA). For each sample, 0.75 ml was centrifuged at 10,500 g for 20 min, the pellet discarded, and supernatant stored at -70°C until the time of proteins and carbohydrates analysis. The rest of the sample was used to extract lipids following the procedure of Folch et al. (1957). In the case of AJ and A, the entire body of each spider was homogenized in 3 ml buffer potassium phosphate (50 mM, pH 7.4) with the addition of a protease inhibitor cocktail 1/1000 (v/v) and following the same procedure as the egg-sacs.

Proteins and glycogen analysis

The total protein content of the homogenates was quantified by the method of Lowry et al. (1951) with bovine-serum albumin as standard and expressed in μg of protein/mg of wet weight of individuals. The protein-subunit analyses were performed by polyacrylamide gel electrophoresis (PAGE) under dissociating conditions adding sodium dodecyl sulphate (SDS) on a 4–23% SDS-PAGE (Laemmli 1970) and using β -mercaptoethanol as a reducing agent. Twenty μg of sample was loaded per well; and after resolution, gels were stained with Coomassie Brilliant Blue R-250 (Sigma Chemical Co.). The respective molecular weights were calculated using a lyophilized mixture of ten highly purified well-characterized proteins to use in molecular weight determination in the presence of SDS (Precision Plus Protein™ Dual Color Standards, BIO-RAD Code:1610374). The isolation of lipovitellins (LVs) of eggs and the very-high-density lipoprotein (VHDL) of *P. pythagoricus* hemolymph, the same as the obtaining of rat-serum antibodies against hemocyanin (Hc) and LV, were performed following the same steps described in Romero et al. (2019).

Western blotting analysis initially involved the separation of proteins by SDS-PAGE loaded with 15 μg of protein of each sample per well. The proteins from the resolved unstained gel were transferred to a nitrocellulose membrane following the same steps as Romero et al. (2019). Then, membranes were incubated for 2 h one with the anti-LV polyclonal antibodies (1:5000) and the other one with the anti-Hc polyclonal antibodies (1:3000), both raised in rat. Immune complexes were detected after an incubation with a Goat Anti-Rat IgG H&L (HRP) (1:2000; ABCAM Inc.) and the immunoreactivity signals emitted were detected using Chemidoc Imaging system (Bio-Rad) and analysed with ImageJ software (NIH).

The enzyme-linked immunosorbent assay (ELISA) was used to quantify Hc and LV in the homogenates of the eight development stages previously described according to the assay of Engvall & Perlmann (1972). In order to quantify LV and Hc, standards curves were produced with the isolated LV and purified Hc. The Hc quantification was carried out following the same steps taken by Romero et al. (2019). In LV case, the curve was prepared by loading with 50 μl per well of LV in the micro well plates (Nunc-Immuno™ MicroWell™ PolySorp) at different concentrations of: 0, 5, 10, 50, 100, 200, 400, 600, 800, and 1000 ng dissolved in coating buffer (35 mM NaHCO_3 , mM Na_2CO_3 , pH 9.6). The supernatants from stage 1 to 8 were incubated with anti-LV polyclonal antibodies diluted (1:2000), and then with the Goat Anti-Rat IgG H&L (HRP) diluted (1:5000) following the same steps described in Trabalon et al. (2018). The same detection and readout of data steps were carried out for both LV and Hc. After washes, 50- μl aliquots of substrate solution that contained ABTS (2, 2'-Azinobis [3-ethylbenzothiazoline-6-sulfonic acid]-diammonium salt) and H_2O_2 were added to each well, incubated during 15 min, and

finally the reaction was stopped with 2% oxalic acid (50 μ L). The absorbance reading at 405 nm was performed in a Beckman Coulter DTX 880 Multimode Detector. Triplicate measurements were taken for each sample.

Glycogen of each sample was precipitated using a 66% KOH 1:1 (v/v) solution and heated for 20 minutes over a boiling water bath (100 °C). After cooled at room temperature, a solution of Na₂SO₄ 10% (v/v) and ethanol 96° 0.25% (v/v) was added in each sample, and then were heated for 5 minutes over boil water. Samples were finally centrifuged at medium speed for 10 min, supernatants were discarded and the glycogen precipitates obtained were resuspended in 1 ml H₂O. The glycogen concentration of the different homogenates was determined applying a modified anthrone-based assay (Van Handel 1985) and quantified using a glycogen standard (0.1-40 μ g/ml). The results were expressed as μ g of glycogen/mg of wet weight of individuals.

Lipids and Fatty-acid analysis

After lipid extraction, the total lipid concentration was determined by gravimetry (Cunningham & Pollero 1996) and expressed in μ g of lipids/mg of wet weight of individuals. Then, the quantitative determination of lipid classes was performed by thin layer chromatography (TLC) coupled to a flame ionization detector (FID) in an Iatroscan apparatus model TH-10 (Iatron Laboratories, Tokyo, Japan) after separation on type S-III Chromarods (Ackman et al. 1990). The different lipid classes were separated and quantified using monoacylglycerol as an internal standard and the calibration curves for each lipid presented correlation coefficients greater than 0.95. The different samples were analysed following the procedure as that used in Romero et al. (2018) using also the same kind of lipid standards and solvent systems.

Fatty-acid methyl esters from the total lipids of each sample were prepared using a BF₃-MeOH solution according to the method of Morrison & Smith (1964). The fatty-acid methyl esters analysis was performed by gas-liquid chromatography (GLC) in an HP-6890 capillary chromatograph (Hewlett Packard, Palo Alto, California, USA), equipped with a flame-ionization detector and fitted with an Omegawax 250 fused silica column of dimensions 30 m \times 0.25 mm and with a 0.25 μ m solid phase (Supelco, Bellefonte, California, USA), and under the same conditions that were performed in Romero et al. (2018). Peaks were identified by comparison with retention times of Supelco 37 component fatty acid methyl ester mix.

Statistical analyses

Statistical comparison was performed by a one-way ANOVA after checking for normality and homogeneity of variances. The different values were expressed as the means \pm standard deviations (SDs). Significant differences ($p < 0.05$, $n=4$) were compared by Tukey's post hoc test. Data were analysed using GraphPad InStat 3.01 (GraphPad Software, San Diego, California, USA).

RESULTS

In this study, eight developmental stages of *P. pythagoricus* were analysed (Figure 1): five took place inside the egg-sac and the other three outside the egg-sac. Females laid 209 ± 88 eggs without any substance holding them together. After egg laying, spherical green eggs with 1.91 ± 0.11 mm diameter are observed (Stage 1). Then, after 5 days of embryo development an active organogenesis could be observed where a white cumulus cells are identified in the green eggs (Stage 2), and 10 days after oviposition, the appendages of the embryo begin to be appreciated (Stage 3). The post-embryonic




Figure 1. The analysed stages of *P. pythagoricus* development. Inside the egg-sac: the embryonic period (Stages 1 to 3) and post-embryonic period (Stages 4 and 5); Outside the egg-sac: juveniles in gregarious stage (Stage 6) and juveniles in dispersal stages (Stages 7 and 8). E: embryonic period; Post-E: post-embryonic period; GJ: juveniles in gregarious stage; DJ: juveniles in dispersal stages.

period takes place when the embryo hatches and moults after 14 or 15 days of oviposition. Stage 4 (15 days after oviposition) are juveniles with a transparent and unpigmented cuticle. The yolk shifted to the opisthosoma gives a characteristic green colour to this body area and whose size can double the prosoma size. In stage 5 (20 days after oviposition) juveniles show a lightly pigmented brown cuticle. These juveniles have frequent leg movements but without being able to walk and their opisthosoma remains the largest body area. Between the 22 ± 1 days after oviposition, juveniles moults inside the egg-sac producing a well-defined light pink and more rigid exuvia. The resulting juveniles after this moults present longer legs which enable them to walk and be more active, and both prosoma and opisthosoma are comparable in size and present a darker pigmentation throughout the

body. Between 25 ± 1 days after oviposition, these juveniles emerge from the egg-sac and remain together as juveniles in gregarious stage (Stage 6) moving around the cocoon surface. Spiderling emergence may or may not be carried out in the presence of mothers because it was observed that they can emerge and continue to grow even when their mother is not present. Ten days after emergence, these gregarious spiderlings moults and give rise to dispersed juveniles (Stage 7 and 8). The tough exuvia presents a dark pink pigmentation. In both stage 7 (35 ± 1 days after laying) and 8 (42 ± 1 days after laying), juveniles leave the gregarious life walking and running with a higher speed than juveniles in gregarious stage. The cuticle of these juveniles has a darker pigmentation throughout the body and opisthosoma is slightly smaller than prosoma.

At this time, juveniles have long legs with the characteristic lateral position of *P. pythagoricus*.

Protein concentration remained relative constant until stage 5, where it was observed an increase that reaches $203.01 \pm 6.47 \mu\text{g}$ of proteins/mg wet weight (Figure 2). After emergence, values decreased up to $88.06 \pm 9.84 \mu\text{g}$ of proteins/mg wet weight in stage 8. Then, a significant increase was observed in AJ and A (already fed individuals).

Taking into account the principal polypeptides observed by SDS-PAGE analysis (Figure 3a and 4a), it was observed the same protein profile during the embryonic period (Stages 1 to 3) with the polypeptides 170, 120, 75, 67, 46, and 30 kDa present. After hatching of embryos, 75 kDa polypeptide disappeared almost completely in stage 4, while 170 and 120 kDa polypeptides disappeared completely before the emergence of juveniles (Stage 6). Low-molecular-weight polypeptides were differentially consumed, 30 kDa started to

diminish after embryos' hatching (Stage 4) but 46 kDa polypeptide did not disappear until juveniles dispersed (Stages 7 and 8). Conversely, 67 kDa polypeptide was present in all stages.

Moreover, the western blot analysis using polyclonal antibody against LV revealed the presence of 120, 75, 46, and 30 kDa polypeptides of LV during the embryonic period (Stages 1 to 3) (Figure 3b). During the post-embryonic period, 75 kDa polypeptide disappeared in stage 4 and then also 120 kDa polypeptide disappeared in stage 5, remaining only low-molecular-weight polypeptides (30 and 46 kDa). Finally, juveniles in gregarious stage only preserved the 46 kDa polypeptide (Stage 6) and no polypeptides sharing immunological identity with LV were detected in dispersed juveniles (Stage 7 and 8). Conversely, the presence of 67 kDa polypeptide was observed throughout the all stages studied applying a western blot analysis using polyclonal antibody against Hc (Figure 4b).




Figure 2. Analysis of protein concentration during *P. pythagoricus* development until adult's stage. Values are the means \pm SDs. Different letters above the bars indicate significant differences among the different stages at $p < 0.05$, as determined by Tukey's *post hoc* test. 1 to 8: stages of *P. pythagoricus* development; AJ: advanced juvenile stage; A: adult stage; E: embryonic period; Post-E: post-embryonic period; GJ: juveniles in gregarious stage; DJ: juveniles in dispersal stages.

The content of LV and Hc in the homogenates of the eight stages was quantified using the method of ELISA with polyclonal antibody against LV and Hc (Figure 5). It was observed a gradual LV consumption beginning from protein value of $410 \pm 77.7 \mu\text{g LV/mg}$ of proteins in stage 1 and reaching $296.8 \pm 55.8 \mu\text{g LV/mg}$ of proteins stage 4 (after embryo hatch). Then, the LV amount decreased significantly to $102.89 \pm 27.6 \mu\text{g LV/mg}$ of proteins in stage 6 (juveniles in gregarious stage) and $18.04 \pm 5.53 \mu\text{g LV/mg}$ of proteins in stage 8 (juveniles in dispersal stages). LV consumption between the first stage and the last stage of juveniles in dispersal phase was linear and gradual ($R^2= 0.96$). By contrast, the amount of Hc presented an exponential increase ($R^2= 0.81$). It began with an average of

$159.22 \pm 27.93 \mu\text{g Hc/mg}$ of proteins, and then it reached values of $274.67 \pm 1.86 \mu\text{g Hc/mg}$ of proteins in stage 5. Finally, it reached $641.58 \pm 16.47 \mu\text{g Hc/mg}$ of proteins in the last stage of dispersed juveniles analysed.

During the embryonic period, it was observed an increase in the glycogen concentration in the stage 3 ($725 \pm 0.18 \mu\text{g glycogen/mg}$ wet weight) (Figure 6). After embryo hatch, glycogen concentration decreased gradually until values of $0.18 \pm 0.02 \mu\text{g glycogen/mg}$ wet weight in stage 8. The glycogen concentration was also quantified in AJ and A, where values were strikingly lower in AJ whereas in A were comparable with the first dispersal stage.

The concentration of total lipids in the embryonic period showed a quite variable values




Figure 3. Analysis of soluble proteins during *P. pythagoricus* development. Electrophoresis (SDS-PAGE): 20 μg of soluble proteins per well (a) and immunoblotting with anti-LV polyclonal antibodies: 15 μg of soluble proteins per well (b). std: molecular weight standard (kDa); LV: lipovitellin isolated from egg cytosol; 1 to 8: stages of *P. pythagoricus* development. Numbers with the same colour belong to the same period of development that are delimited below the figure as: E: embryonic period; Post-E: post-embryonic period; GJ: juveniles in gregarious stage; DJ: juveniles in dispersal stages.

with an average of $60.47 \pm 20.2 \mu\text{g}$ of lipids/mg wet weight, and then no significant decreases were observed with the post-embryonic stages ($32.42 \pm 8.2 \mu\text{g}$ of lipids/mg wet weight) (Figure 7). The amount of lipids decreased significantly to $12.3 \pm 1.7 \mu\text{g}$ of lipids/mg of wet weight in the last stage of dispersed juveniles (Stage 8). Comparable values were observed in AJ and A.

Analysing the different lipid classes, the principal concentration of structural lipids in the embryonic stages was represented by phosphatidylethanolamine (PE) with values of $30.25 \pm 8.05 \mu\text{g}$ PE/mg of wet weight (Figure 8). After hatching, PE content decreased until spiderlings emergence from the egg-sac (Stage 6, $2.83 \pm 1.68 \mu\text{g}$ PE/mg of wet weight), values that remained relative constant in advanced

stages. Phosphatidylcholine (PC) presented a similar pattern as that of PE, beginning with a concentration of $12.65 \pm 1.46 \mu\text{g}$ PC/mg of wet weight in the embryonic stages and decreased until spiderlings emergence from the egg-sac (Stage 6). Sphingomyelin (SM) maintained a small proportion of less than $1.2 \mu\text{g}$ SM/mg of wet weight throughout the developmental stages occurring inside and outside the egg-sac (Figure 8). However, in A this value decreased to average concentration of $0.28 \pm 0.02 \mu\text{g}$ SM/mg of wet weight.

Triacylglycerides (TAG) concentration values were relatively constant until the stage 6 (juveniles in gregarious stage), between 12.55 ± 1.8 and $17.8 \pm 3.7 \mu\text{g}$ TAG/mg of wet weight (Figure 9). After dispersion, values decreased reaching




Figure 4. Analysis of soluble proteins during *P. pythagoricus* development. Electrophoresis (SDS-PAGE): 20 µg of soluble proteins per well (a) and immunoblotting with anti-Hc polyclonal antibodies: 15 µg of soluble proteins per well (b). std: molecular weight standard (kDa); VHDL: very-high-density lipoprotein of *P. pythagoricus*; 1 to 8: stages of *P. pythagoricus* development. Numbers with the same colour belong to the same period of development and are delimited below the figure as: E: embryonic period; Post-E: post-embryonic period; GJ: juveniles in gregarious stage; DJ: juveniles in dispersal stages.

0.77 ± 0.2 µg TAG/mg of wet weight in the stage 8. Then, the concentration rose to 2.6 ± 0.2 µg TAG/mg of wet weight in advanced stages. Free fatty acids (FFA) concentration had minimum values in the embryonic and post-embryonic periods (less than 2 µg of FFA/mg of wet weight), but once juveniles emerged the concentration increased to an average of 5.2 ± 0.63 µg of FFA/mg of wet weight (Figure 9). Later on, concentration values reduced to 1.2 ± 0.8 µg of FFA/mg of wet weight in AJ and A. Diacylglycerides (DAG) only appeared in small amounts during the post-embryonic development (Figure 9).

Hydrocarbons (HC) content increased during the embryonic development and remained relatively constant during post-embryonic development with averages of 1.03 ± 0.55µg of HC/mg of wet weight (Figure 10). Outside the egg-sac, dispersed juveniles increased their HC

content to 4.63 ± 1.19 µg of HC/mg of wet weight and 3 ± 0.66 µg of HC/mg of wet weight in stage 7 and 8, respectively. Cholesterol (C) throughout the embryonic development fluctuated between 1.21 ± 0.12 to 1.51 ± 0.25 µg of C/mg of wet weight (Figure 10), but then decreased to 0.89 ± 0.28 µg of C/mg of wet weight when juveniles emerged from the egg-sac (Stage 6). In AJ and A, C concentration average was of 0.65 ± 0.05 µg of C/mg of wet weight. Sterol esters (SE) values were only detected in the post-embryonic stages with small amounts between 0.1 ± 0.06 and 0.96 ± 0.28 µg of SE/mg of wet weight (Figure 10).

Analysing the fatty-acid methyl esters (Figure 11), fatty acids percentages remained constant until the gregarious stage (Stage 6); however, significant changes appeared after dispersion (Stages 7 and 8). Palmitic acid (16:0) decreased from 13.1 ± 2.7% (Stage 6) to 6.9 ±




Figure 5. Analysis of lipovitellin (LV) and hemocyanin (Hc) concentration during *P. pythagoricus* development. Values are the means ± SDs. Different letters above the bars indicate significant differences among the different stages at $p < 0.05$, as determined by Tukey's *post hoc* test. 1 to 8: stages of *P. pythagoricus* development; E: embryonic period; Post-E: post-embryonic period; GJ: juveniles in gregarious stage; DJ: juveniles in dispersal stages.

0.16% (Stage 8), whereas stearic acid (18:0) increased from $7.2 \pm 2.7\%$ to $12.6 \pm 0.17\%$. In the case of monounsaturated fatty acids, oleic acid (18:1) diminished from $45.2 \pm 0.3\%$ (Stage 6) to $24.8 \pm 1.7\%$ (Stage 8), whereas linoleic acid (18:2) began with $23.4 \pm 0.7\%$ and enriched to $38.45 \pm 0.4\%$. Moreover, eicosatetraenoic acid (20:4) and eicosapentaenoic acid (20:5) increased their values from less than 1.5% to 6% in stages 7 and 8. Throughout the development, no changes were observed in the total amount of saturated fatty acids (SFA), while the unsaturated fatty acids underwent changes after juveniles emerged. The monounsaturated fatty acids (MUFA) content decreased from $49.45 \pm 1.35\%$ to $26.04 \pm 2.02\%$, while the polyunsaturated fatty acids (PUFA) increased from $28.6 \pm 1.93\%$ to $52.2 \pm 2.08\%$.

DISCUSSION

The presence or absence of maternal care, gregarious stages and the competition for food and intra-species cannibalism are behaviours that may influence a spider's development. Thus, the study of resource variation during development may provide new information to explain or accompany behavioural studies. Up to the present it has been clear that the identity of the energetic resources that support embryonic and post-embryonic development in spiders and its understanding is limited in this megadiverse group.

Our results showed that *P. pythagoricus* had no variation in total protein concentration when accompanied by gradual consumption of LV during the early stages of development. These findings would suggest there is a balance between new protein synthesis and yolk protein degradation. This has also been described for




Figure 6. Analysis of glycogen concentration during *P. pythagoricus* development until adult's stage. Values are the means \pm SDs. Different letters above the bars indicate significant differences among the different stages at $p < 0.05$, as determined by Tukey's *post hoc* test. 1 to 8: stages of *P. pythagoricus* development; AJ: advanced juvenile stage; A: adult stage; E: embryonic period; Post-E: post-embryonic period; GJ: juveniles in gregarious stage; DJ: juveniles in dispersal stages.

other arthropods (Campos et al. 2006, Kamel & Ragaa 1981). However, something different was reported in *P. saltans* spider, where gradual consumption of proteins was observed, even when LV levels did not decrease during embryonic development (Trabalon et al. 2018), which showed some differences between both species. *P. pythagoricus* showed differential consumption of LV polypeptides: higher-molecular-weight polypeptides (120 and 75 kDa) were consumed in post-embryonic instars, whereas lower-molecular-weight polypeptides (46 and 30 kDa) were consumed after juveniles emerged. Higher-molecular-weight polypeptides might be found to be more exposed to the proteolytic enzyme than lower-molecular-weight polypeptides during development as was previously described for other arthropods where LVs was observed to have a higher subunit (surrounding smaller polypeptides) that was more susceptible to enzymatic attack (Garcia et al. 2008). After juveniles of *P. pythagoricus* emerged from egg-sacs, total protein consumption and LV

reserve exhaustion was probably a consequence of the intense activity of the mobile juveniles, especially during dispersion instars (Stage 7 and 8). Therefore, LV proteolysis would be linked to the energetic and structural needs of the organism (Lu & Warner 1991, Perona & Vallejo 1985, Warner et al. 1995).

Increase in the total amount of proteins in advanced stages of *P. pythagoricus* could be due to prey consumption. It was described that spider growth could be maximized with higher protein content diets (Jensen et al. 2011) since this may be needed to build new tissues and fulfil the basic requirements of metabolism, maintaining protein homeostasis in adults (Jensen et al. 2011, Romero et al. 2019, Wilder 2011). Salomon et al. (2011) showed that a protein-enriched diet in *Stegodyphus lineatus* resulted in higher juveniles' survival in this subsocial spider species.

The importance of Hc in O₂ transport in aqueous medium in arthropods and other invertebrates has been shown in different




Figure 7. Analysis of total lipid concentration during *P. pythagoricus* development until adult's stage. Values are the means ± SDs. Different letters above the bars indicate significant differences among the different stages at $p < 0.05$, as determined by Tukey's *post hoc* test. 1 to 8: stages of *P. pythagoricus* development; AJ: advanced juvenile stage; A: adult stage; E: embryonic period; Post-E: post-embryonic period; GJ: juveniles in gregarious stage; DJ: juveniles in dispersal stages.

research papers (Foelix 2011, Markl & Decker 1992), this being the protein’s main function. The presence of Hc in the eggs of several arthropods has been demonstrated over time (Chen et al. 2017), and it has also been recently described for the present model of study (Romero et al. 2019). This passage of Hc from the mother to the eggs could provide an initial pool of Hc to the embryo until this can be able to synthesize its own Hc; even more if we consider that Hc is not only an oxygen-carrying protein but also a protein storage of this molecule (Foelix 2011, Starrett et al. 2013). Active involvement of Hc in oxygen exchange in embryos of *Locusta migratoria* was previously described, where the authors proposed that hemocyanin plays an important role in oxygen transport during embryogenesis (Chen et al. 2017). By means of a process called inversion, during the embryonic development of Araneomorphae spiders, it has been observed

that there is a great change in the organization of the body that results in the incorporation of yolk mass in the embryo (Wolff & Hilbrant 2011). Thus, the embryo would have a pool of Hc to supply the demand for oxygen which cannot be met by simple diffusion produced by the high rate of aerobic metabolism involved in embryonic development (Ingrisch 1987, Rakshpal 1962). Moreover, this could be an important fact to be highlighted if we consider that the eggs are small and isothermal in their local microclimate, and they are often challenged by oxygen shortage in fluctuating environments, even at mildly high temperatures (Woods et al. 2005, Woods & Hill 2004).

Although we cannot rule out the early synthesis of embryonic Hc, we might consider the beginning of Hc expression during development as occurring in late stages, the same as what was described for other arthropods (Chen




Figure 8. Concentration of the main phospholipids during *P. pythagoricus* development until adult’s stage: phosphatidylethanolamine (PE), phosphatidylcholine (PC) and sphingomyelin (SM). Values are the means \pm SDs. Different letters above the bars indicate significant differences among the different stages at $p < 0.05$, as determined by Tukey’s *post hoc* test. 1 to 8: stages of *P. pythagoricus* development; AJ: advanced juvenile stage; A: adult stage; E: embryonic period; Post-E: post-embryonic period; GJ: juveniles in gregarious stage; DJ: juveniles in dispersal stages.

et al. 2017, Pick et al. 2010), even in members of Subphylum Chelicerata where an increase in Hc content was observed after the first embryonic moulting (Sugita & Sekiguchi 1979). The Hc increase observed during the post-embryonic stages of *P. pythagoricus* would allow juveniles to deal with the new challenges of the environment after hatching, since Hc has a great variety of functions such as the defence against pathogens (Coates & Nairn 2014, Riciluca et al. 2012), transporting hormones of the moulting (Jaenicke et al. 1999), acting as lipid transporter (Cunningham et al. 2007, Cunningham & Pollero 1996, Laino et al. 2009, 2015a), being part of the cuticle component (Paul et al. 1994) and having phenoloxidase activity (Jaenicke & Decker 2008). This last function is associated with the sclerotization process, a remarkable function if we consider that the pigmentation of the cuticle begins to show in this stage (Stage 5).

In various arthropods, carbohydrate mobilization has been described as being responsible for obtaining the energy which will finally be used for the growing embryo (Campos et al. 2006, Ludwig & Ramazzotto 1965, Mohamed 2000, Santana et al. 2014, Santos et al. 2008, Tralalon et al. 2018). Although glycogen is stored in eggs and is used in the early stages of development, its content has been observed to increase in late embryonic stages in *Drosophila melanogaster*, when organogenesis takes place (Waltero et al. 2020, Yamazaki & Nusse 2002). This is similar to what was observed in *P. pythagoricus* whose glycogen level increased at the end of the embryonic development (Stage 3). In advanced stages of development, different sources of carbohydrates, such as glycogen, may be important for chitin biosynthesis, which requires great amounts of glucose (Chippendale 1978). Our results showed a gradual and constant consumption of carbohydrates after




Figure 9. Concentration of energetic lipids during *P. pythagoricus* development until adult's stage: triacylglycerides (TAG), free fatty acids (FFA), and diacylglycerides (DAG). Values are the means \pm SDs. Different letters above the bars indicate significant differences among the different stages at $p < 0.05$, as determined by Tukey's *post hoc* test. 1 to 8: stages of *P. pythagoricus* development; AJ: advanced juvenile stage; A: adult stage; E: embryonic period; Post-E: post-embryonic period; GJ: juveniles in gregarious stage; DJ: juveniles in dispersal stages.

hatching, similar to that observed in *P. saltans* spider (Trabalon et al. 2018). However, these carbohydrates represented less than 1% of the total reserves in *P. pythagoricus* eggs.

In *P. pythagoricus*, lipid content during development was greater in the embryonic and non-dispersed juvenile stages than in dispersed juveniles and adults. The same was observed in other arachnids and in various holometabolous insects, where this energy reserve would offset the partial or total energy required for adult life stages (Lease & Wolf 2011). Throughout the time, it has been highlighted that energetic lipids represent an important resource for the embryonic development of many insects, mites and crustaceans (Campos et al. 2006, Geister et al. 2009, Santos et al. 2011, Van Handel 1993).

However, in most spiders this does not seem to be true. *P. pythagoricus* eggs initially have 72.2% of structural lipids, and only 24.3% of energetic lipids. This agrees with the analysis performed in *Schizocosa malitiosa* (Lycosidae) where it was observed that eggs had 67 ± 27% of structural lipids (PE + PC + SM), only 26.6 ± 18.5% of energetic lipids (TAG + FFA) and 5.5 ± 3.7% of C + SE + HC (Laino et al. 2013). Nevertheless, this does not coincide with the values observed in *P. saltans* (Lycosidae), where the reported figures were 33.1 ± 1.2% for structural lipids, 51.5 ± 8.1% for energetic lipids, and 12.8 ± 4% for C + SE + HC + LPL (Lysophosphatidylcholine) (Trabalon et al. 2018). According to the authors, the difference found in the lipid composition of eggs of these Lycosidae spiders could be due to the difference




Figure 10. Concentration of modulating or signaling lipids during *P. pythagoricus* development until adult's stage: hydrocarbons (HC), cholesterol (C) and sterol esters (SE). Values are the means ± SDs. Different letters above the bars indicate significant differences among the different stages at $p < 0.05$, as determined by Tukey's *post hoc* test. 1 to 8: stages of *P. pythagoricus* development; AJ: advanced juvenile stage; A: adult stage; E: embryonic period; Post-E: post-embryonic period; GJ: juveniles in gregarious stage; DJ: juveniles in dispersal stages.

in the type of food mothers received during the test, as it was observed in the hemolymph of *Eurypelma californicum* and *Brachypelma albopilosum* that were fed with different animals (Schartau & Leidescher 1983, Trabalon 2011, Trabalon et al. 2018).

Oocytes of different spider families show the presence of aggregates of organelles and inclusions usually arranged into orderly concentric zones called Balbiani bodies or yolk nucleus, with concentric lamellae corresponding to the endoplasmic reticulum (Andre & Rouiller 1957, Jędrzejowska & Kubrakiewicz 2010, Sotelo

& Trujillo-Cenoz 1957), which might show a high content of structural lipids. Moreover, the major presence of these structural lipids in yolk lipoproteins was previously described in *P. pythagoricus* (Laino et al. 2011b), where their percentages were $49 \pm 1\%$ for LV1 and $37.6 \pm 1\%$ for LV2, while energetic lipids represented only $9.5 \pm 0.4\%$. During vitellogenesis, the presence of structural lipids was highlighted as a key component of the ovary in *P. pythagoricus* females; this organ became charged with 63% of these lipids (especially with PE) and only with 23% of energetic lipids (Romero et al. 2018).




Figure 11. Composition of the major fatty acids during *P. pythagoricus* development: palmitate (16:0), stearate (18:0), oleate (18:1), linoleate (18:2), eicosatetraenoic acid (20:4) and eicosapentaenoic acid (20:5). The values are the percentage means \pm SDs. Different letters indicate significant differences among the different stages at a $p < 0.05$, as determined by Tukey's *post hoc* test. 1 to 8: stages of *P. pythagoricus* development; E: embryonic period; Post-E: post-embryonic period; GJ: juveniles in gregarious stage; DJ: juveniles in dispersal stages.

Likewise, the eggs of this species have a great percentage of PE, which coincides with the composition reported for their own LVs (Laino et al. 2011b) and with that reported for some embryos of crustaceans (Wang et al. 2015). Up to the present, it has been proposed that PE needs a unique structure to form the inner mitochondrial membranes, thus improving the functioning of the electron transport chain (Ikou & Ryan 2017, Romero et al. 2018, Tasseva et al. 2012, Teague et al. 2013). However, its partial consumption in the post-embryonic period (over 50%) and its great consumption in the emerged juveniles (over 90%), represent considerable evidence that PE could also have an energetic function as some authors suggest for structural lipids (Hagen et al. 1996, Mayzaud et al. 2003). Furthermore, it has been described that phospholipids can act as energetic resource during the early development of some crustaceans (Sibert et al. 2004). This hypothesis could be reinforced by the scarce content of this phospholipid in dispersed juveniles and adults (0.4% and 0.21% of the initial content of PE). PE would be hydrolysed mainly in the post-embryonic stages, thus providing an ethanolamine moiety that could covalently modify several proteins (Vance & Tasseva 2013). Great consumption of phospholipids during the post-embryonic stages would reveal that phospholipase activity may probably be greater than lipase activity.

TAG are found in yolk forming lipid droplets (Schie et al. 2013, Walther & Farese 2012), and they are the most important energetic molecules because of their great caloric capacity and the way they get stored (Laino et al. 2011b, 2013). TAG were consumed in the more active stages analysed in *P. pythagoricus*: 50% in stage 7 and 96% in stage 8 (juveniles in dispersion instar). This kind of late hydrolysis of TAG was also observed in *P. saltans* spider (Trabalon et al. 2018). Although the *de-novo* synthesis of FFA

cannot be discarded, in the present work it can be inferred that the increase of fatty acids in juveniles in the gregarious stage (Stages 6) is probably due to phospholipid hydrolysis (mainly PE), whereas in the dispersed juveniles (Stage 7 and 8) it could be due to TAG hydrolysis.

HC in spiders could have several functions such as that of being the first chemical barrier to prevent the entry of pathogens, serving as kairomones for entomopathogenic fungi and bacteria (Lecuona et al. 1991), participating as controllers of hydric homeostasis (Gibbs 1998, Hadley 1981) and having an active role in the chemical communication among congeners (Blomquist & Bagnères 2010). HC content varies to a great extent in *P. pythagoricus*, mainly in embryonic stages and in juveniles in dispersion. This may be due to the *de novo* synthesis that was previously described for arthropods (Fan et al. 2008). On the other hand, both C and SE are membrane components and moulting hormone precursors (Andersen 1979, Martin-Creuzburg et al. 2007). Therefore, it is clear that, although HC, C, and SE are minor in adults, they play an important biological role such as the formation of ecdysteroids (Andersen 1979, Martin-Creuzburg et al. 2007). However, the lack of information with relation to the subject hinders the adequate interpretation of the role played by these lipids in spiders' development.

In *P. pythagoricus*, egg reserve mostly contains unsaturated fatty acids 18:1 and 18:2, and saturated fatty acids 16:0 and 18:0. These major fatty acids were reserved in the ovaries during vitellogenesis (Romero et al. 2018), to later become part of yolk, thus coinciding with the general pattern of fatty acids of their LVs that are components of yolk (Laino et al. 2011b). This pattern of fatty acids was also observed in two members of the family Lycosidae, *S. malitiosa* and *P. saltans*, with 18:2, 18:1, 16:0 and 18:0 as major fatty acids (Laino et al. 2013, Trabalon et al. 2018).

Although it is described that in some insects the composition of fatty acids remains constant throughout embryonic development (Hoppe et al. 1975), in other arthropods consumption does not seem to be equal, since some fatty acids are consumed and others produced during embryogenesis (Figueiredo et al. 2012). In the case of spiders, the total percentage of SFA, MUFA, and PUFA remained constant until the gregarious stage. Then, the percentage of MUFA decreased with the concomitant increase of PUFA during dispersion, along with a specific change in fatty acid composition inside these groups. The impoverishment of 16:0 was offset by the enrichment of 18:0, which explains the absence of SFA, mainly in dispersion stages where the consumption of TAG was evident. Apart from being an important source of energy, it was observed that derivatives of 16:0—as 14-methylhexadecanoic acid—can be found in the silk and have important antimicrobial activity provided by the protective effect of the methyl group (Heimer 1988, Tahir et al. 2017), which is an important fact considering that active silk production already takes place in these dispersed stages. In the case of unsaturated fatty acids, there was an impoverishment of 18:1 and a major enrichment of 18:2, which play an important role in functions such as cell physiology, immunity, and reproduction (Malcicka et al. 2018). Fatty acid species present in phospholipids and TAG appear to be similar in both types of lipids. It is assumed that energetic lipids mostly contain SFA and that phospholipids have a higher amount of PUFA (Innis 1991). In *P. pythagoricus*, phospholipids do not seem to have the majority of polyunsaturated fatty acids or else these do not correlate with the amount of PE used. Even so, 20:4 and 20:5 fatty acids in the dispersed juveniles increased six times more than in embryonic stages, which are considered to be essential fatty acids for arthropods. The

importance of fatty acids with C20 lies in the fact that they are precursors of eicosanoids (prostaglandins, leukotrienes) (Meijer et al. 1986, Petzel 1993, Stanley-Samuelson et al. 1991, Stanley-Samuelson & Pedibhotla 1996). Several studies on arthropods describe the function of eicosanoids in the defence against microorganisms through immune reactions (Morishima et al. 1997, Park et al. 2003, Stanley-Samuelson et al. 1991), something that would be of vital importance for the survival of just dispersed juveniles.

As a conclusion, we can highlight in the development of *P. pythagoricus* spider that: 1) carbohydrates are gradually consumed after hatching, but they represent less than 1% of the reserves; 2) apolipoproteins of the LV in these spider species are gradually and differentially consumed, being first consumed the higher-molecular-weight polypeptides (120 and 75 kDa) and then the lower-molecular-weights polypeptides (46 and 30 kDa); 3) while LV amount and the total protein content decrease, there is an important increase of Hc content from post-embryonic instars to dispersion instars, which allows us to consider that the Hc synthesis of the offspring during development is late; 4) the main energy resources during development are represented by LV and phospholipids, but TAG surprisingly do not represent the majority energy source being consumed in advanced stages; and finally 5) the use of phospholipids (PE and PC) and unsaturated fatty acids (especially MUFA) gains great importance in the post-embryonic and emerged juveniles respectively. It is clear that, although there is a need to generate new research to better understand the metabolism of the development of spiders, the present work is the first one to have obtained a continuous and complete perspective of the biochemical changes generated during spider development.

Acknowledgments

This work was supported by Grants from Agencia Nacional de Promoción Científica y Tecnológica (PICT-2017-0684), and UNLP Argentina N794 and N7207. L. A., G. F. and C. M. are members of CONICET, Argentina. R. S. and M. G. are a CONICET scholarship fellow. The authors are grateful to Rosana del Cid and Laura Cristina Prezioso for the review of the English, Sebastian Romero and Mario Ramos for the figure design, to Romina Becerra (INIBIOLP, Argentina) for her technical assistance, and to the arachnology laboratory of CEPAVE for providing the service and equipment necessary to take the pictures for the current work.

REFERENCES

- ACKMAN RG, MCLEOD C & BANERJEE AK. 1990. An overview of analyses by chromarod iatroskan TLC-FID. *J Planar Chromatogr* 3: 450-490.
- ANDERSEN SO. 1979. Biochemistry of insect cuticle. *Annu Rev Entomol* 24: 29-59.
- ANDERSON JF. 1978. Energy content of spider eggs. *Oecologia* 37: 41-57.
- ANDRE J & ROUILLER C. 1957. The ultrastructure of the vitelline body in the oocyte of the spider *Tegenaria parietina*. *J Biophys Biochem Cytol* 3: 977-984.
- BEDNAREKAW, SAWADRO MK, NICEWICZ L & BABCZYŃSKA AI. 2019. Vitellogenins in the spider *Parasteatoda tepidariorum* - expression profile and putative hormonal regulation of vitellogenesis. *BMC Dev Biol* 19: 4.
- BLOMQUIST GJ & BAGNÈRES AG. 2010. Insect hydrocarbons: biology, biochemistry, and chemical ecology. Cambridge: Cambridge University Press, 506 p.
- BYRNE BM, GRUBER M & AB G. 1989. The evolution of egg yolk proteins. *Prog Biophys Mol Biol* 53: 33-69.
- CAMPOS ET AL. 2006. Kinetics of energy source utilization in *Boophilus microplus* (Canestrini, 1887) (Acari: Ixodidae) embryonic development. *Vet Parasitol* 138: 349-357.
- CHEN B, MA R, DING D, WEIL & KANG L. 2017. Aerobic respiration by haemocyanin in the embryo of the migratory locust. *Insect Mol Biol* 26: 461-468.
- CHIPPENDALE GM. 1978. Carbohydrates in reproduction and embryonic development. In: Rockstein M (Ed), *Biochemistry of Insects*, New York: Academic Press, New York, USA, p. 42-45.
- CHOI YS & MOON MJ. 2003. Fine structure of the ovarian development in the Orb-web Spider, *Nephila clavata*. *Entomol Res* 33: 25-32.
- COATES CJ & NAIRN J. 2014. Diverse immune functions of hemocyanins. *Dev Comp Immunol* 45: 43-55.
- CUNNINGHAM M, GARCIA F, GONZALEZ BARO MR, GARDA H & POLLERO R. 2002. Organophosphorous insecticides fenitrothion alters the lipid dynamics in the spider *Polybetes pythagoricus* high density lipoproteins. *Pestic Biochem Phys* 73: 37-47.
- CUNNINGHAM M, GARCIA F & POLLERO RJ. 2007. Arachnid lipoproteins: Comparative aspects. *Comp Biochem Phys C* 146: 79-87.
- CUNNINGHAM M & POLLERO R. 1996. Characterization of lipoprotein fraction with high content of hemocyanin in the hemolymphatic plasma of *Polybetes pythagoricus*. *J Exp Zool* 274: 275-280.
- CUNNINGHAM M, POLLERO R & GONZALEZ A. 1994. Lipid circulation in spiders. Transport of phospholipids, free acids and triacylglycerols as the major lipid classes by a high-density lipoprotein fraction isolated from plasma of *Polybetes pythagoricus*. *Comp Biochem Physiol B* 109: 333- 338.
- ENGVALL E & PERLMANN P. 1972. Enzyme-linked immunosorbent assay, ELISA: III. Quantitation of specific antibodies by enzyme-labeled anti-immunoglobulin in antigen-coated tubes. *J Immunol* 109: 129-135.
- FAN Y, ELIYAHU D & SCHAL C. 2008. Cuticular hydrocarbons as maternal provisions in embryos and nymphs of the cockroach *Blattella germanica*. *J Exp Biol* 211: 548-554.
- FIGUEIREDO J, LIN J, ANTO J & NARCISO L. 2012. The consumption of DHA during embryogenesis as an indicative of the need to supply DHA during early larval development: a review. *J Aquacult Res Dev* 3: 140.
- FOELIX RF. 2011. *Biology of spiders*, 3rd ed., New York: Oxford University Press, 432 p.
- FOLCH J, LEES M & SLOANE STANLEY GH. 1957. A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 226: 497-509.
- FRUTTERO LL, FREDE S, RUBIOLO ER & CANAVOSO LE. 2011. The storage of nutritional resources during vitellogenesis of *Panstrongylus megistus* (Hemiptera: Reduviidae): the pathways of lipophorin in lipid delivery to developing oocytes. *J Insect Physiol* 57: 475-486.
- GALIANO ME. 1971. El desarrollo postembrionario larval en especies de género *Polybetes* Simon, 1897 (Araneae, Sparassidae). *Acta Zool Lillo* 28: 221-225.

- GALIANO ME. 1979. Datos adicionales sobre el ciclo vital de *Polybetes pythagoricus* (Holmberg, 1874) (Araneae, Eusparassidae). *Acta Zool Lillo* 35: 75-86.
- GARCIA F, CUNNINGHAM ML, GARDA H & HERAS H. 2008. Embryo lipoproteins and yolk lipovitellin consumption during embryogenesis in *Macrobrachium borellii* (Crustacea: Palaemonidae). *Comp Biochem Physiol B* 151: 317-322.
- GEISTER TL, LORENZ MW, HOFFMANN KH & FISCHER K. 2009. Energetics of embryonic development: effects of temperature on egg and hatchling composition in a butterfly. *J Comp Physiol B* 179: 87-98.
- GIBBS AG. 1998. Water-proofing properties of cuticular lipids. *Am Zool* 38: 471-482.
- GRISMADO CJ, RAMÍREZ MJ & IZQUIERDO MA. 2014. Araneae: Taxonomía, diversidad y clave de identificación de familias de la Argentina. In: Roig-Juñent S, Claps LE, Morrone JJ (Eds), Biodiversidad de artrópodos argentinos, volumen 3. Buenos Aires: División Aracnología, Museo Argentino de Ciencias Naturales "Bernardino Rivadavia", Buenos Aires, Argentina, p. 55-94.
- HADLEY NF. 1981. Cuticular lipids of terrestrial plants and arthropods: a comparison of their structure, composition, and waterproofing function. *Biol Rev* 56: 23-47.
- HAGEN W, VLEET E & KATTNER G. 1996. Seasonal lipid storage as overwintering strategy of Antarctic krill. *Mar Ecol Prog Ser* 134: 85-89.
- HEIMER S. 1988. *Wunderbare Welt der Spinnen*. Leipzig: Urania-Verlag.
- HOPPE KT, HADLEY NF & TRELEASE RN. 1975. Changes in lipid and fatty acid composition of eggs during development of the beet armyworm, *Spodoptera exigua*. *J Insect Physiol* 21: 1427-1430.
- IKON N & RYAN RO. 2017. Cardiolipin and mitochondrial cristae organization. *Biochim Biophys Acta*, 1859: 1156-1163.
- INGRISCH S. 1987. Oxygen consumption by developing and diapausing eggs of *Eupholidoptera smyrnensis* (Orthoptera: Tettigoniidae). *J Insect Physiol* 33: 861-865.
- INNIS SM. 1991. Essential fatty acids in growth and development. *Prog. Lipid Res* 30:39-103.
- JAENICKE E & DECKER H. 2008. Kinetic properties of catecholoxidase activity of tarantula hemocyanin. *FEBS J* 275: 1518-1528.
- JAENICKE E, FOLL R & DECKER H. 1999. Spider hemocyanin binds ecdysone and 20-OH-ecdysone. *J Biol Chem* 274: 34267-34271.
- JĘDRZEJOWSKA I & KUBRAKIEWICZ J. 2010. Yolk nucleus-The complex assemblage of cytoskeleton and ER is a site of lipid droplet formation in spider oocytes. *Arthropod Struct Dev* 39: 350-359.
- JENSEN K, MAYNTZ D, TOFT S, RAUBENHEIMER & DSIMPSON SJ. 2011. Prey nutrient composition has different effects on *Pardosa* wolf spiders with dissimilar life histories. *Oecologia* 165: 577-583.
- KAMEL MY & RAGAA RH. 1981. Purification and characterization of pyrophosphatase from developing embryos of *Hyalomma dromedarii*. *Insect Biochem* 11: 691-698.
- LAEMMLI UK. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680-685.
- LAINO A, CUNNINGHAM M, COSTA FG & GARCIA CF. 2013. Energy sources from the eggs of the wolf spider *Schizocosa malitiosa*: isolation and characterization of lipovitellins. *Comp Biochem Phys B* 165: 172-180.
- LAINO A, CUNNINGHAM ML, GARCIA F & HERAS H. 2009. First insight into the lipid uptake, storage and mobilization in arachnids: role of midgut diverticula and lipoproteins. *J Insect Physiol* 55: 1118-1124.
- LAINO A, CUNNINGHAM ML, HERAS H & GARCIA F. 2011a. In vitro lipid transfer between lipoproteins and midgut-diverticula in the spider *Polybetes pythagoricus*. *Comp Biochem Phys B* 160: 181-186.
- LAINO A, CUNNINGHAM ML, HERAS H & GARCIA F. 2011b. Isolation and characterization of two vitellins from eggs of the spider *Polybetes pythagoricus* (Araneae: Sparassidae). *Comp Biochem Physiol B* 158: 142-148.
- LAINO A, CUNNINGHAM M, SUAREZ G & GARCIA CF. 2015a. Identification and characterization of the lipid transport system in the tarantula *Grammostola rosea*. *OJAS* 5: 9-20.
- LAINO A, GARCIA CF & CUNNINGHAM M. 2015b. Protein characterization and fatty acid composition of VHDL subfraction II of the spider *Polybetes pythagoricus*. *Biocell* 39: 33-40.
- LEASE HM & WOLF BO. 2011. Lipid content of terrestrial arthropods in relation to body size, phylogeny, ontogeny and sex. *Physiol Entomol* 36: 29-38.
- LECUONA R, RIBA G, CASSIER P & CLEMENT JL. 1991. Alterations of insect epicuticular hydrocarbons during infection with *Beauveria bassiana* or *B. brongniartii*. *J Invertebr Pathol* 58: 10-18.

- LOWRY OH, ROSENBOUGH NJ, FARR AL & RANDALL R. 1951. Protein measurement with the Folin phenol reagent. *J Biol Chem* 193: 265-275.
- LUDWIG D & RAMAZZOTTO LJ. 1965. Energy sources during embryogenesis of the yellow mealworm, *Tenebrio molitor*. *Ann Entomol Soc Am* 58: 543-546.
- LU J & WARNER AH. 1991. Immunodetection of thiol proteinase levels in various populations of *Artemia* cysts and during development. *Biochem Cell Biol* 69: 96-101.
- MALCICKA M, VISSER B, & ELLERS J. 2018. An evolutionary perspective on linoleic acid synthesis in animals. *Evol Biol* 45: 15-26.
- MARKL J & DECKER H. 1992. Molecular structure of the arthropod hemocyanins. In: Mangum CP (Ed), *Advances in Comparative and Environmental Physiology*. Berlin: Springer-Verlag Berlin Heidelberg, Berlin, Germany, p. 326-363.
- MARTIN-CREUZBURG D, WESTERLUND SA & HOFFMANN KH. 2007. Ecdysteroid levels in *Daphnia magna* during a molt cycle: determination by radioimmunoassay (RIA) and liquid chromatography-mass spectrometry (LC-MS). *Gen Comp Endocr* 151: 66-71.
- MAYZAUD P, BOUTOUTE M & ALONZO F. 2003. Lipid composition of the euphausiids *Euphausia vallentini* and *Thysanoes samacrura* during summer in the Southern Indian Ocean. *Antarc Sci* 15: 463-475.
- MEIJER L, BRASH AR, BRYANT RW, NG K, MACLOUF J & SPRECHER H. 1986. Stereospecific induction of starfish oocyte maturation by (8R)-hydroxyeicosatetraenoic acid. *J Biol Chem* 261: 17040-17047.
- MOHAMED SA. 2000. α -Amylase from developing embryos of the camel tick *Hyalomma dromedarii*. *Comp Biochem Phys B* 126: 99-108.
- MORISHIMA I, YAMANO Y, INOUE K & MATSUO N. 1997. Eicosanoids mediate induction of immune genes in the fat body of the silkworm, *Bombyx mori*. *FEBS Lett* 419: 83-86.
- MORRISON WR & SMITH LM. 1964. Preparation of fatty acid methyl esters and dimethylacetals from lipids with boron fluoride-methanol. *J Lipid Res* 5: 600-608.
- PAUL R, BERGNER B, PFEFFER-SEIDL A, DECKER H, EFINGER R & STORZ H. 1994. Gas transport in the haemolymph of arachnids - oxygen transport and the physiological role of haemocyanin. *J Exp Biol* 188: 25-46.
- PARK Y, KIM Y, PUTNAM SM & STANLEY DW. 2003. The bacterium *Xenorhabdus nematophilus* depresses nodulation reactions to infection by inhibiting eicosanoid biosynthesis in tobacco hornworms, *Manduca sexta*. *Arch Insect Biochem* 52: 71-80.
- PERONA R & VALLEJO CG. 1985. Acid hydrolases during *Artemia* development: A role in yolk degradation. *Comp Biochem Physiol B* 81: 993-1000.
- PETZEL DH. 1993. Prostanoids and Fluid Balance in Insects. In: Stanley-Samuelson DW & Nelson DR (Eds), *Insect Lipids: Chemistry, Biochemistry, and Biology*. Lincoln: University of Nebraska Press, Lincoln, USA, p. 139-178.
- PICK C, SCHNEUER M & BURMESTER T. 2010. Ontogeny of hemocyanin in the ovoviviparous cockroach *Blattella germanica* suggests an embryo-specific role in oxygen supply. *J Insect Physiol* 56: 455-460.
- RAKSHPAL R. 1962. Diapause in the eggs of *Gryllus pennsylvanicus* Burmeister (Orthoptera: Gryllidae). *Can J Zool* 40: 179-194.
- RICILUCA KCT, SAYEGH RSR, MELO RL & SILVA PI. 2012. Rondonin an antifungal peptide from spider (*Acanthoscurria rondoniae*) haemolymph. *Res Immunol* 2: 66-71.
- ROMERO S, LAINO A, ARRIGHETTI F, CUNNINGHAM M & GARCIA CF. 2018. First study on lipid dynamics during the female reproductive cycle of *Polybetes pythagoricus* (Araneae Saporassidae). *Can J Zool* 96: 847-858.
- ROMERO S, LAINO A, ARRIGHETTI F, GARCÍA CF & CUNNINGHAM M. 2019. Vitellogenesis in spiders: first analysis of protein changes in different reproductive stages of *Polybetes pythagoricus*. *J Comp Physiol B* 189: 335-350.
- RUHLAND F, PETILLON J & TRABALON M. 2016. Physiological costs during the first maternal care in the wolf spider *Pardosa saltans* (Araneae, Lycosidae). *J Insect Physiol* 95:42-50.
- SABA M, AWAN DS & YOUSAF S. 2020. Spider as a biological agent in pest control-A Review. *J Wildlife Manag* 4: 27-34.
- SALOMON M, MAYNTZ D, TOFT S & LUBIN Y. 2011. Maternal nutrition affects offspring performance via maternal care in a subsocial spider. *Behav Ecol Sociobiol* 65: 1191-1202.
- SANTANA CC, DO NASCIMENTO JS, COSTA MM, DA SILVA AT, DORNELAS CB & GRILLO LA. 2014. Embryonic development of *Rhynchophorus palmarum* (Coleoptera: Curculionidae): dynamics of energy source utilization. *J Insect Sci* 14: 280.
- SANTOS R, MARIANO AC, ROSAS-OLIVEIRA R, PASCARELLI B, MACHADO EA, MEYER-FERNANDES JR & GONDIM KC. 2008. Carbohydrate accumulation and utilization by oocytes of *Rhodnius prolixus*. *Arch Insect Biochem Physiol* 67: 55-62.

- SANTOS R, ROSAS-OLIVEIRA R, SARAIVA FB, MAJEROWICZ D & GONDIM KC. 2011. Lipid accumulation and utilization by oocytes and eggs of *Rhodnius prolixus*. Arch Insect Biochem Physiol 77: 1-16.
- SCHARTAU W & LEIDESCHER T. 1983. Composition of the hemolymph of the tarantula *Eurypelma californicum*. J Comp Physiol 152: 73-77.
- SCHIE IW, NOLTE L, PEDERSEN TL, SMITH Z, WU J, YAHIAÏÈNE I, NEWMAN JW & HUSER T. 2013. Direct comparison of fatty acid ratios in single cellular lipid droplets as determined by comparative Raman spectroscopy and gas chromatography. Analyst 138: 6662-6670.
- SCIOSCIA CL. 1984. Análisis del crecimiento en *Polybetes pythagoricus* (Holmberg, 1874) (Araneae, Sparassidae). Rev Soc Entomol Argent 43: 1-4.
- SIBERT V, OUELLETP & BRÊTHES JC. 2004. Changes in yolk total proteins and lipid components and embryonic growth rates during lobster (*Homarus americanus*) egg development under a simulated seasonal temperature cycle. Mar Biol 144: 1075-1086.
- SMITH DM, HUNTER BJ, ALLAN GL, ROBERTS DCK, BOOTH MA & GLENCROSS BD. 2004. Essential fatty acids in the diet of silver perch (*Bidyanus bidyanus*): effect of linolenic and linoleic acid on growth and survival. Aquaculture 236: 377-390.
- SOTELO JR & TRUJILLO-CENOZ O. 1957. Electron microscope study of the vitelline body of some spider oocytes. J Biophys Biochem Cytol 3: 301-310.
- STANLEY-SAMUELSON DW, JENSEN E, NICKERSON KW, TIEBEL K, OGG CL & HOWARD RW. 1991. Insect immune response to bacterial infection is mediated by eicosanoids. P Natl Acad Sci 88: 1064-1068.
- STANLEY-SAMUELSON DW & PEDIBHOTLA VK. 1996. What can we learn from prostaglandins and related eicosanoids in insects? Insect Biochem Mol Biol 26: 223-234.
- STARRETT J, HEDIN M, AYOUB N & HAYASHI CY. 2013. Hemocyanin gene family evolution in spiders (Araneae), with implications for phylogenetic relationships and divergence times in the infraorder Mygalomorphae. Gene 524: 175-186.
- SUGITA H & SEKIGUCHI K. 1979. Protein components in the perivitelline fluid of the embryo of the horseshoe crab, *Tachypleus tridentatus*. Dev Biol 73: 183-192.
- TAHIR HM, ZAHRA K, ZAHEER A & SAMIULLAH K. 2017. Spider silk: An excellent biomaterial for medical science and industry. Punjab Univ J Zool 32: 143-154.
- TASSEVA G, BAI HD, DAVIDESCU M, HAROMY A, MICHELAKIS E & VANCE JE. 2012. Phosphatidylethanolamine deficiency in mammalian mitochondria impairs oxidative phosphorylation and alters mitochondrial morphology. J Biol Chem 288: 4158-4173.
- TEAGUE WE JR, SOUBIAS O, PETRACHE H, FULLER N, HINES KG, RAND RP & GAWRISCH K. 2013. Elastic properties of polyunsaturated phosphatidylethanolamines influence rhodopsin function. Faraday Discuss 161: 383-339.
- THOMPSON MB & RUSSEL KJ. 1999. Embryonic energetics in eggs of two species of Australian skink, *Morethia boulengeri* and *Morethia adelaidensis*. J Herpetol 33: 291-297.
- TRABALON M. 2011. Agonistic interactions, cuticular and hemolymphatic lipid variations during the foraging period in spider females *Brachypelma albopilosa* (Theraphosidae). J Insect Physiol 57: 735-743.
- TRABALON M, BAUTZ AM, MORINIERE M & PORCHERON P. 1992. Ovarian development and correlated changes in hemolymphatic ecdysteroid levels in two spiders, *Coelotes terrestris* and *Tegenaria domestica* (Araneae, Agelenidae). Gen Comp Endocr 88: 128-136.
- TRABALON M, RUHLAND F, LAINO A, CUNNINGHAM M & GARCIA F. 2018. Embryonic and post-embryonic development inside wolf spiders' egg sac with special emphasis on the vitellus. J Comp Physiol B 188: 211-224.
- VAN HANDEL E. 1985. Rapid determination of glycogen and sugars in mosquitoes. J Am Mosq Control Assoc 1: 299-301.
- VAN HANDEL E. 1993. Fuel metabolism of the mosquito (*Culex quinquefasciatus*) embryo. J Insect Physiol 39: 831-833.
- VANCE JE & TASSEVA G. 2013. Formation and function of phosphatidylserine and phosphatidylethanolamine in mammalian cells. Biochim Biophys Acta 1831: 543-554.
- VETTER RS & ISBISTER GK. 2008. Medical aspects of spider bites. Annu Rev Entomol 53: 409-429.
- WALTHER TC & FARESE JR RV. 2012. Lipid droplets and cellular lipid metabolism. Annu Rev Biochem 81: 687-714.
- WALTERO C, MARTINS R, CALIXTO C, DA FONSECA RN, ABREU LA, DA SILVA VAZ I JR & LOGULLO C. 2020. The hallmarks of GSK-3 in morphogenesis and embryonic development metabolism in arthropods. Insect Biochem Molec Biol 118: 103307.
- WANG LH, LIN C, HUANG CY & TSAI S. 2015. Studies on lipid content and composition in banded coral shrimp

(*Stenopus hispidus*) embryos. *J Crustacean Biol* 35: 622-626.

WARNER AH, PERZ MJ, OSAHAN JK & ZIELINSKI BS. 1995. Potential role in development of the major cysteine protease in larvae of the brine shrimp *Artemia franciscana*. *Cell Tissue Res* 282: 21-31.

WILDER SM. 2011. Spider nutrition: An integrative perspective. *Adv Insect Physiol* 40: 87-136.

WOODS HA, BONNECAZE RT & ZRUBEK B. 2005. Oxygen and water flux across eggshells of *Manduca sexta*. *J Exp Biol* 208: 1297-1308.

WOODS HA & HILL RI. 2004. Temperature-dependent oxygen limitation in insect eggs. *J Exp Biol* 207: 2267-2276.

WOLFF C & HILBRANT M. 2011. The embryonic development of the central American wandering spider *Cupiennius salei*. *Front Zool* 8: 15.

WORLD SPIDER CATALOG. 2021. World Spider Catalog. Version 22.0. Natural History Museum Bern, online at <http://wsc.nmbe.ch>, accessed on March 2021. doi: 10.24436/2

YAMAZAKI H & NUSSE RI. 2002. Identification of DCAP, a *Drosophila* homolog of a glucose transport regulatory complex. *Mech Dev* 119: 115-119.

YIP EC & RAYOR LS. 2014. Maternal care and subsocial behaviour in spiders. *Biol Rev* 89: 427-449.

¹Universidad Nacional de La Plata/Consejo Nacional de Investigaciones Científicas y Técnicas (UNLP-CONICET), Instituto de Investigaciones Bioquímicas de La Plata "Prof. Dr. Rodolfo R. Brenner" (INIBIOLP), FCM, Calle 60 y 120, La Plata (1900), Buenos Aires, Argentina

Correspondence to: **Carlos Fernando Garcia, Monica Cunningham**
E-mail: cfgarcia1123@yahoo.com.ar,
cunninghammoni@gmail.com

Author contributions

Romero, S. Conceptualization, Methodology, Formal analysis, Investigation, Visualization, Writing - Original Draft, Writing - Review & Editing. Laino A., Investigation, Methodology, Formal analysis, Funding acquisition, Writing - Review & Editing. Molina G., Investigation, Formal analysis, Writing - Review & Editing. Cunningham M. Conceptualization, Funding acquisition, Writing - Original Draft, Writing- Review & Editing. Garcia C. Conceptualization, Funding acquisition, Writing - Original Draft, Writing- Review & Editing.



How to cite

ROMERO S, LAINO A, MOLINA G, CUNNINGHAM M & GARCIA CF. 2021. Embryonic and post-embryonic development of the spider *Polybetes pythagoricus* (Sparassidae): A biochemical point of view. *An Acad Bras Cienc* 94: e20210159. DOI 10.1590/0001-3765202220210159.

Manuscript received on February 2, 2021;
accepted for publication on November 18, 2021

SOFIA ROMERO¹

<https://orcid.org/0000-0002-4032-2771>

ALDANA LAINO¹

<https://orcid.org/0000-0002-7299-6439>

GABRIEL MOLINA¹

<https://orcid.org/0000-0002-8677-9347>

MONICA CUNNINGHAM¹

<https://orcid.org/0000-0001-5413-2015>

CARLOS FERNANDO GARCIA¹

<https://orcid.org/0000-0001-5934-6899>