

Survey of potential parasitoids (Hymenoptera) of *Drosophila suzukii* (Diptera: Drosophilidae) in Buenos Aires province, Argentina

GALLARDO, Fabiana E.^{1,2,*}, RECHE, Vanina A.¹, MARGARÍA, Cecilia B.³, AQUINO, Daniel A.^{1,3,4}, ANSA, María A.^{5,6}, DETTLER, María A.^{5,6}, VAZQUEZ, Florencia⁵, BARRIENTOS, Gualterio^{5,6}, SANTADINO, Marina^{5,6}, MARTÍNEZ, Emilia⁶ & RIQUELME VIRGALA, María^{5,6}

¹ División Entomología. Facultad de Ciencias Naturales y Museo. Universidad Nacional de La Plata. Paseo del Bosque s/n, B1900FWA, La Plata, Argentina.

² Comisión de Investigaciones Científicas de la provincia de Buenos Aires (CICPBA), Argentina.

³ Zoología Agrícola, Centro de Investigación en Sanidad Vegetal, Facultad de Ciencias Agrarias y Forestales. Universidad Nacional de La Plata, Avenida 60 y 119, B1900FWA, La Plata, Argentina.

⁴ CEPAVE (CONICET-UNLP). Boulevard 120 entre 60 y 64, La Plata, Buenos Aires, Argentina.

⁵ Departamento de Tecnología, Universidad Nacional de Luján (UNLu). Av. Constitución y Ruta 5, (6700) Luján, Buenos Aires, Argentina.

⁶ Instituto de Ecología y Desarrollo Sustentable (UNLu-CONICET). Av. Constitución y Ruta 5, (6700) Luján, Buenos Aires, Argentina.

* E-mail: gallardo@fcnym.unlp.edu.ar

Received 24 - VI - 2021 | Accepted 30 - XII - 2021 | Published 31 - III - 2022

<https://doi.org/10.25085/rsea.810107>

Estudio de potenciales parasitoides (Hymenoptera) de *Drosophila suzukii* (Diptera: Drosophilidae) en la provincia de Buenos Aires, Argentina

RESUMEN. *Drosophila suzukii* (Matsumura) es una plaga que afecta a una gran variedad de cultivos frutales. El objetivo de este trabajo fue conocer la diversidad de parasitoides (Hymenoptera) que podrían estar asociados con SWD en cultivos frutales ubicados en el noreste de la provincia de Buenos Aires, Argentina. Las especies identificadas fueron *Euxestophaga argentinensis* Gallardo, *Ganaspis pelleranoi* (Brèthes), *Leptopilina bouvardi* (Barbotin, Carton & Kelner-Pillault), *Hexacola bonaerensis* Reche (Cynipoidea, Figitidae), *Trichopria anastrephae* (Costa Lima) (Diaprioidea, Diapriidae) y *Pachycrepoideus vindemmiae* (Rondani) (Chalcidoidea, Pteromalidae). Se registró por primera vez a *L. bouvardi*, *E. argentinensis* y *T. anastrephae* en la provincia de Buenos Aires.

PALABRAS CLAVE. Cultivos frutales. *Drosophila* de alas manchadas. Parasitoides. Provincia biogeográfica Pampeana.

ABSTRACT. *Drosophila suzukii* (Matsumura) is a pest that affects a great variety of fruit crops. The objective of this work was to learn about the diversity of parasitoids (Hymenoptera) that might be associated with SWD in fruit crops located in northeastern Buenos Aires province, Argentina. Species were identified as *Euxestophaga argentinensis* Gallardo, *Ganaspis pelleranoi* (Brèthes), *Leptopilina bouvardi* (Barbotin, Carton & Kelner-Pillault), *Hexacola bonaerensis* Reche (Cynipoidea, Figitidae), *Trichopria anastrephae* (Costa Lima) (Diaprioidea, Diapriidae) and *Pachycrepoideus vindemmiae* (Rondani) (Chalcidoidea, Pteromalidae). *Leptopilina bouvardi*, *E. argentinensis* and *T. anastrephae* were reported for the first time in the Buenos Aires province.

KEYWORDS. Fruit crops. Parasitoids. Pampean biogeographic province. Spotted-wing *Drosophila*.

Drosophila suzukii (Matsumura), commonly known as spotted wing drosophila (SWD) is a pest that affects a great variety of fruit crops (Cini et al., 2012). SWD female oviposits under the skin of healthy and mature fruit by piercing the epicarp with a serrated ovipositor. This fact together with softening due to the larval activity, exposes fruits to different pathogens, leading to their decay and the consequent economic losses (Cini et al., 2012). Despite SWD is native to Southeast Asia, it has been recently registered in the European and American continents as a result of its high dispersion capacity (Bolda et al., 2010; Calabria et al., 2012). In Argentina, SWD was detected for the first time in 2014 in raspberry crops of the province of Río Negro (Cichón et al., 2015). Almost simultaneously, it was reported in blueberry crops from the province of Buenos Aires (Santadino et al., 2015) and traps located in orange plantations in Entre Ríos (Díaz et al., 2015). Thereafter, SWD was recorded from different fruit-producing regions, such as NOA (Lue et al., 2017; Funes et al., 2018; Lavagnino et al., 2018), Cuyo (Dagatti et al., 2018), Patagonia (Garrido et al., 2018; Lavagnino et al., 2018; de la Vega et al., 2019), NEA (SENASA, 2016) and Pampeana (Gonsebatt et al., 2017; Lavagnino et al., 2018). To date, there are no registered chemical products for SWD control in Argentina, being cultural and biological control two of the main tools for its management (Escudero-Colomar, 2016). In this context, the search for potential natural enemies of this pest is essential for integrated control programs (García, 2020; Wang et al., 2020). A recent worldwide review of SWD natural enemy species was published by Kirschbaum et al. (2020). Parasitoids were the most numerous and most studied group of SWD's natural enemies. The known species of parasitoid wasps that attack *D. suzukii* belong to five genera. Two of them are pupal parasitoids, namely, *Pachycrepoideus* Ashmead (Chalcidoidea, Pteromalidae) and *Trichopria* Ashmead (Diaprioidea, Diapriidae), and the remaining three are larval parasitoids: *Leptopilina* Förster, *Ganaspis* Förster (Cynipoidea, Figitidae) and *Asobara* Förster (Ichneumonoidea, Braconidae) (Asplen et al., 2015). Species of the genus *Ganaspis* are among the most efficient SWD parasitoids, showing the highest rates of parasitoidism (Cini et

al., 2012). In this sense, Giorgini et al. (2019) suggested that *Ganaspis brasiliensis* (von Ihering) was a promising classical biological control agent for release in invaded regions because of its demonstrated high specificity. Recent studies in Argentina are encouraging, since they have reported *G. brasiliensis* (Gallardo et al., 2021), *Ganaspis hookeri* Crawford (Lue et al., 2017), *Leptopilina boulandi* (Barbotin, Carton & Kelner-Pillault) (Garrido et al., 2018), *Leptopilina clavipes* (Hartig) (Lue et al., 2017), *Pachycrepoideus vindemmiae* (Rondani) (Funes et al., 2020), *Spalangia endius* Walker (Gómez Segade et al., 2021) and more recently a new species of *Dieucoila* Ashmead, *D. octoflagella* Reche (Reche et al., 2021) in association with SWD. Thus, the objective of this work was to find out the diversity of parasitoids that might be associated with SWD in fruit crops located in northeastern Buenos Aires province, Argentina.

The study was conducted in fruit orchards in three localities: Luján (34° 35' S; 59° 04' W), Lobos (35° 11' S; 59° 16' W) and Mercedes (34° 41' S; 59° 24' W). Samplings were carried out during 2017 and 2018 in SWD-infested crops of strawberries, blueberries, raspberries, cherries, peaches and blackberries. The specimens were captured using traps baited with 200 cc of apple vinegar diluted to 50% with water. Traps were placed in the central part of each fruit crop along the windbreaks and were renewed every fifteen days. The material collected was taken to the laboratory to count adult SWD and potential parasitoids. Fruit samples with no evident fruit damage collected from the field during the harvest period. In the laboratory, each fruit was individually placed in ventilated containers inside a GC-300 growth chamber (Lab. Companion®) under controlled conditions (23 ± 2 ° C, Relative humidity 60 ± 10%, Photoperiod 16L: 8D), until adult emergence. The proportion of infested fruit and the number of adults were used to estimate Incidence and Severity.

The wasp specimens recovered (n = 26) were sent for identification to the Sección Hymenoptera Parasítica of the División Entomología, Museo de La Plata, Buenos Aires, Argentina. The specimens were identified using taxonomic keys (Risbec, 1950; Gibson, et al., 1997; Guimarães et al., 2003) and specialized bibliography. The dipterans specimens

were identified using taxonomic keys (Vlach, 2010). The wasp specimens examined were deposited in the entomological collection of the Museo de La Plata. The number of SWD and adult wasps recovered per crop sample is summarized in Table I. Besides documenting *Drosophila suzukii*, six species of Hymenoptera parasitoids belonging to Figitidae, Diapriidae and Pteromalidae were also

collected: *Euxestophaga argentinensis* Gallardo, *Ganaspis pelleranoi* (Brèthes), *L. boulandi*, *Hexacola bonaerensis* Reche (Cynipoidea, Figitidae), *Trichopria anastrephae* (Costa Lima) (Diaprioidea, Diapriidae) and *P. vindemmiae* (Chalcidoidea, Pteromalidae). Fruit infestation was variable between species and no parasitoids were recorded (Table II).

Table I. Total number of captures of female and male *Drosophila suzukii* and wasps from June 2017 to June 2018.

Locality	Crop	<i>D. suzukii</i>	<i>E. argentinensis</i>	<i>G. pelleranoi</i>	<i>H. bonaerensis</i>	<i>L. boulandi</i>	<i>T. anastrephae</i>	<i>P. vindemmiae</i>
Lobos	Cherry	246			3 females	1 female		1 female
	Raspberry	31				1 female		
	Blueberry	171				2 females		
Luján	Strawberry	5					1 female	
	Blackberry	51		3 females		1 female	1 female 1 male	
	<i>Morus</i> spp.	134				1 female	1 female	
	Peach	5	1 female	1 female			1 female	
Mercedes	<i>Morus</i> spp.	3				4 females		
	Blackberry	22		1 female	1 female			

***Euxestophaga argentinensis* Gallardo**

Distribution. Neotropical region. Argentina. It is distributed throughout the provinces of Santa Fe, Tucumán (Gallardo et al., 2017), and Buenos Aires (new record).

Biology. Primary larval-pupal solitary koinobiont endoparasitoid of Diptera Otitidae: *Euxesta eluta* Loew (Gallardo et al., 2017).

Material examined. Argentina. Buenos Aires. Luján. 1 female, crop of *Prunus persica* (Rosaceae) (peaches), experimental field UNLu, 15-XI-2017, Dettler coll.

***Ganaspis pelleranoi* (Brèthes)**

Distribution. Neotropical region: Mexico, Belize, Guatemala, El Salvador, Costa Rica, Panama, Colombia, Venezuela, Brazil, Peru, Bolivia and Argentina (Díaz & Gallardo 2001). In Argentina, it is distributed throughout the provinces of Buenos Aires, Corrientes, Jujuy, Misiones, Salta, Tucumán (Díaz & Gallardo, 2014) and La Rioja, (Nuñez-Campero & Ovruski, 2013).

Biology. Primary larval-pupal solitary koinobiont

endoparasitoid of Diptera Tephritidae: *Anastrepha fraterculus*, *Anastrepha schultzi*, *Anastrepha* sp., *Ceratitis capitata*, and Lonchaeidae: *Lonchaea* sp. (Guimarães et al., 2003; Díaz & Gallardo, 2014).

Material examined. Argentina. Buenos Aires. Luján. 1 female, crop of *Prunus persica* (Rosaceae) (peaches), experimental field UNLu, 18-I-2018, Ansa coll.; 2 females, crop of *Rubus ulmifolius* (Rosaceae) (blackberries), experimental field UNLu, 26-XII-2017, Ansa coll.; 1 female, crop of *Rubus ulmifolius* (Rosaceae) (blackberries), experimental field UNLu, 18-I-2018, Ansa coll. **Mercedes.** 1 female, crop of *Rubus ulmifolius* (Rosaceae) (blackberries), family farm, 06-XI-2017, Riquelme Virgala coll.

***Hexacola bonaerensis* Reche**

Distribution. Nearctic and Neotropical regions: US, Cuba and Argentina, where it is distributed throughout the province of Buenos Aires (Reche & Gallardo, 2015).

Biology. Primary larval-pupal solitary koinobiont endoparasitoid of Diptera Ephydriidae (Reche & Gallardo, 2015).

Material examined. Argentina. Buenos Aires. Lobos. 2 females, crop of *Prunus avium* (Rosaceae) (cherries), commercial field, 15-V-2018, Dettler coll.; 1 female, crop of *Prunus avium* (Rosaceae) (cherries), commercial field, 21-VI-2018, Dettler coll. Mercedes. 1 female, crop of *Rubus ulmifolius* (Rosaceae) (blackberries), family farm, 8-I-2018, Dettler coll.

***Leptopilina boulandi* (Barbotin, Carton & Kelner-Pillault)**

Distribution. Cosmopolitan. In Argentina, it has been reported in the provinces of Tucumán (Díaz & Gallardo, 2014), Neuquén, Río Negro (Garrido et al., 2018) and Buenos Aires (new record).

Biology. Primary larval-pupal solitary koinobiont endoparasitoid of Diptera Drosophilidae: *Drosophila melanogaster* (Nordlander, 1980), *Zaprionus indianus* (Marchiori et al., 2003), *Drosophila suzukii* (Cuch-Arguimbau et al., 2013; García Cancino et al., 2015; Wollmann et al., 2016; Knoll et al., 2017; Garrido et al., 2018).

Material examined. Argentina. Buenos Aires. Lobos. 1 female, crop of *Prunus avium* (Rosaceae) (cherries), commercial field, 8-I-2018, Dettler coll.; 1 female, crop of *Rubus idaeus* (Rosaceae) (raspberries), commercial field, 8-I-2018, Dettler coll.; 1 female, crop of *Vaccinium corymbosum* (Ericaceae) (blueberries), commercial field, 8-I-2018, Dettler coll.; 1 female, crop of *Vaccinium corymbosum* (Ericaceae) (blueberries), commercial field, 21-VI-2017, Dettler coll. Luján. 1 female, crop of *Morus* spp. (Moraceae), experimental field UNLu, 26-XII-2017, Ansa coll.; 1 female, crop of *Rubus ulmifolius* (Rosaceae) (blackberries), experimental field UNLu, 26-XII-2017, Ansa coll. Mercedes. 1 female, windbreak of *Morus* spp. (Moraceae), family farm, 06-XI-2017, Dettler coll.; 3 females, windbreak of *Morus* spp. (Moraceae), family farm, 15-V-2018, Dettler coll.

***Trichopria anastrephae* (Costa Lima)**

Distribution. Neotropical region: Brazil (Costa Lima, 1940), Argentina (Turica & Mallo, 1961) and

Venezuela (Boscán & Godoy, 1996). In Argentina, this species is distributed throughout the provinces of Tucumán (Turica & Mallo, 1961) and Buenos Aires (new record).

Biology. Primary pupal solitary idiobiont endoparasitoid of Diptera Tephritidae: *Anastrepha* sp., *A. serpentine* (Costa Lima, 1940), *Anastrepha fraterculus*, *Ceratitis capitata* (Ovruski et al., 2006); Drosophilidae: *Drosophila* spp. (Turica & Mallo, 1961) and *D. suzukii* (Wollmann et al., 2016).

Material examined. Argentina. Buenos Aires. Luján, 1 female, crop of *Fragaria x ananassa* (Rosaceae) (strawberries), commercial field, 27-XII-2017, Martínez coll.; 1 female and 1 male, crop of *Rubus ulmifolius* (Rosaceae) (blackberries), experimental field UNLu, 18-I-2018, Ansa coll.; 1 female, windbreak of *Morus* spp. (Moraceae), experimental field UNLu, 26-XII-2017, Ansa coll.; 1 female, crop of *Prunus persica* (Rosaceae) (peaches), experimental field UNLu, 06-XI-2017, Dettler coll.

***Pachycrepoideus vindemmiae* (Rondani)**

Distribution. Cosmopolitan. In Argentina, it is distributed throughout the provinces of Córdoba (Fischetti et al., 1978), Tucumán (Nasca, 1976), Mendoza (De Santis, 1960), Catamarca (De Santis & Gallego de Sureda, 1988), Buenos Aires, Entre Ríos (Turica et al., 1971) and La Rioja (Ovruski, 2002).

Biology. Polyphagous pupal ectoparasitoid of various families of Diptera Muscomorpha although its members are primary parasitoids, they can also develop as facultative hyperparasitoids (Van Alphen & Thunnissen, 1983; Wang & Messing, 2004). A wide range of hosts from diverse families of dipterous have been recorded: Anthomyiidae, Calliphoridae, Cecidomyiidae, Drosophilidae, Lonchaeidae, Muscidae, Phoridae, Piophilidae, Sarcophagidae, Sphaeroceridae, Stratiomyidae, Syrphidae and Tephritidae (Noyes, 2019).

Material examined. Argentina. Buenos Aires. Lobos. 1 female, crop *Prunus avium* (Rosaceae), commercial field, 27-XI-2017, Vazquez coll.

Table II. Mean of Incidence (% of fruit damaged) and Severity (mean flies developed per fruit) of damage of *Drosophila suzukii*

Locality	Crop	Incidence (%)	Severity (flies per fruit)
Lobos	Cherry	3	0,03
	Raspberry	44	2,38
	Blueberry	0	0
Luján	Strawberry	11	0,35
	Blackberry	40	2,17
	<i>Morus</i> spp.	21	1,24
	Peach	3	0,07
Mercedes	<i>Morus</i> spp.	0	0
	Blackberry	14	0,29

ACKNOWLEDGEMENTS

We thank Cecilia Gorreta (CICPBA) for technical assistance and Comisión de Investigaciones Científicas de la Provincia de Buenos Aires (CICPBA) and Universidad Nacional de La Plata (UNLP) for their constant support.

LITERATURE CITED

- Asplen, M.K., Anfora, G., Biondi, A., Choi, D.S., Chu, D., Daane, K.M., Gibert, P., Gutierrez, A.P., Hoelmer, K.A., Hutchison, W.D., Isaacs, R., Jiang, Z.L., Kárpáti, Z., Kimura, M.T., Pascual, M., Philips, C.R., Plantamp, C., Ponti, L., Véték, G., Vogt, H., Walton, V.M., Yu, Y., Zappalà, L., & Desneux, N. (2015). Invasion biology of spotted wing *Drosophila* (*Drosophila suzukii*): a global perspective and future priorities. *Journal of Pest Science*, **88**, 469–494. <http://dx.doi.org/10.1007/s10340-015-0681-z>
- Boscán, N. de M., & Godoy, F. (1996). Nuevos parasitoides de moscas de las frutas de los géneros *Anastrepha* y *Ceratitis* en Venezuela. *Agronomía Tropical*, **4(4)**, 465-471.
- Bolda, M.P., Goodhue, R.E., & Zalom, F.G. (2010). Spotted Wing *Drosophila*: potential economic impact of a newly established pest. *Agricultural Resource Economics. ARE Update* **13(3)**, 5-8. University of California Giannini -Foundation of Agricultural Economics.
- Calabria, G., Maca, J., Bachili, G., Serra, L., & Pascual, M. (2012). First records of the potential pest species *Drosophila suzukii* (Diptera: Drosophilidae) in Europe. *Journal of Applied Entomology*, **136**, 139-147. <https://doi.org/10.1111/j.1439-0418.2010.01583.x>
- Cichón, L., Garrido, S., & Lago, J. (2015). Primera detección de *Drosophila suzukii* (Matsumura, 1939) (Diptera: Drosophilidae) en frambuesas del Valle de Río Negro, Argentina. In: Libro de Resúmenes del IX Congreso Argentino de Entomología. Posadas, Misiones, Argentina, pp. 270.
- Cini, A., Loriatti, C., & Anafora, A. G. (2012). A review of the invasion of *Drosophila suzukii* in Europe and a draft research agenda for integrated pest management. *Bulletin of Insectology*, **65**, 149-160.
- Costa Lima, A. (1940). Alguns parasitos de moscas de frutas. *Anais da Academia Brasileira de Ciência*, **12(1)**, 17-20.
- Cuch-Arguimbau, N., Escudero-Colomar, L.A., Forshage, M., & Pujade-Villar, J. (2013). Identificadas dos especies de Hymenoptera como probables parasitoides de *Drosophila suzukii* (Matsumura, 1931) en una plantación ecológica de cerezos en Begues (Barcelona, España). *Phytoma-Esp.* **247**, 42-47.
- Dagatti, C.V., Marcucci, B., Herrera, M.E., & Becerra, V.C. (2018). Primera detección de *Drosophila suzukii* (Diptera: Drosophilidae) en frutos de zarzamora en Mendoza, Argentina. *Revista de la Sociedad Entomológica Argentina*, **77(3)**, 26-29. <https://doi.org/10.25085/rsea.770304>
- De la Vega, G.J., Corley, J.C., & Soliani, G. (2019). Genetic assessment of invasion history of *Drosophila suzukii* in Argentina. *Journal of Pest Science*, **93**, 63-75. <http://dx.doi.org/10.1007/s10340-019-01149-x>
- De Santis, L. (1960). Anotaciones sobre calcidoideos Argentinos. II (Hymenoptera). *Revista de la Facultad de Agronomía de La Plata*, **36(2)**, 109-119.
- De Santis, L., & Gallego de Sureda, A.E. (1988). Himenópteros Calcidoideos parasitoides de *Musca domestica* (Dipt.) en América del sur. *Anales de la Academia Nacional de Ciencias de Buenos Aires*, 5-9.
- Díaz, N.B., & Gallardo, F.E. (2001). *Aganaspis* Lin 1987. Generic enlargement and a key for species present in a neotropical region (Cynipoidea: Figitidae: Eucoilinae). *Physis*, **58**, 91-95.

- Díaz, N.B., & Gallardo, F.E. (2014). Figitidae. *Biodiversidad de Artrópodos Argentinos Vol. 4.* (ed. Roig-Juñent, S., Claps, L.E., & Morrone J.J.), pp. 117-125. Editorial INSUE-UNT, San Miguel de Tucumán, Argentina.
- Díaz, B.M., Lavagnino, H.J., Garrán, S., Hochmaier, V., & Fanara, J.J. (2015). Detección de la mosca plaga *Drosophila suzukii* (Matsumura) en la región de Concordia (Entre Ríos). In: Libro de resúmenes de las XV Jornadas Fitosanitarias Argentinas. Santa Fe, Argentina, pp. 287.
- Escudero-Colomar L.A. (2016). Métodos de control para *Drosophila suzukii* (Matsumura) (Diptera: Drosophilidae), una nueva plaga de frutales que se está extendiendo mundialmente. *Revista de Agronomía del Noroeste argentino*, **36**, 19-31.
- Fischetti, D.L., Sosa, H.M., & Cavallo, A. (1978). Biocenosis de la mosca de los frutos en Córdoba. *Memorias de la Tercera Jornada Fitosanitaria Argentina*, **1**, 265-278.
- Funes, C.F., Escobar, L.I., Heredia, A.M., Segade, G., Agostini, J.P., & Kirschbaum, D. S. (2018). Primera detección e incidencia de *Drosophila suzukii* (Diptera: Drosophilidae) en cultivos de berries de Tucumán, Argentina. In: Libro de Resúmenes del X Congreso Argentino de Entomología. Mendoza, Argentina, pp. 242.
- Funes, C.F., Garrido, S.A., Aquino, D.A., Escobar, L.I., Gómez Segade, C., Cichón, L. I., Ovruski, S.M., & Kirschbaum, D.S. (2020). New records of *Pachycrepoideus vindemmiae* (Hymenoptera: Pteromalidae) associated with *Drosophila suzukii* (Diptera: Drosophilidae) in cherry and berry crops in Argentina. *Revista de la Sociedad Entomológica Argentina*, **79(4)**, 39-43. <https://doi.org/10.25085/rsea.790406>
- Gallardo, F.E., Reche, V.A., Bertolaccini, I., Zarate, B., & Curis, C. (2017). A new genus and species of Eucoilinae (Hymenoptera, Cynipoidea, Figitidae) parasitoid of *Euxesta eluta* Loew (Diptera, Otitidae) attacked *Bt* sweet corn in Argentina. *Journal of Hymenoptera Research*, **54**, 57-70. <https://doi.org/10.3897/jhr.54.10096>
- Gallardo, F.E., Funes, C., Reche, V., Kirschbaum, D., Ovruski, S.M., & Buffington, M. (2021). First Record and Distribution of *Ganaspis brasiliensis* (Hymenoptera: Figitidae: Eucoilinae), a parasitoid of *Drosophila suzukii* (Diptera: Drosophilidae) in Argentina. *Neotropical Entomology*, **50(5)** <https://doi.org/10.1007/s13744-021-00912-z>
- García, F.R.M. (2020). Introduction to *Drosophila suzukii* management. *Drosophila suzukii Management* (ed. García, F.R.M.), pp 1-9. Springer Nature, Switzerland AG., pp 1-9. https://doi.org/10.1007/978-3-030-62692-1_1.
- García Cancino, M.D., González Hernández, A., González Cabrera, J., Moreno Carrillo, G., Sánchez González, J.A., & Arredondo Bernal, H.C. (2015). Parasitoides de *Drosophila suzukii* (Matsumura) (Diptera: Drosophilidae) en Colima, México. *Southwestern Entomologist*, **40(4)**, 854-858.
- Garrido, S.A., Cichón, L.I., Lago, J.D., Gallardo, F.E., & Navarro, M.D. (2018). Primer registro de *Leptopilina bouvardi* (Hymenoptera: Figitidae) asociado a *Drosophila suzukii* (Diptera: Drosophilidae) en el Alto Valle de Río Negro y Neuquén, Patagonia, Argentina. *Revista de la Sociedad Entomológica Argentina*, **77(2)**, 22-27. <https://doi.org/10.25085/rsea.770202>
- Gibson, G.A. P., Huber, J.T., & Woolley, J.B. (1997). *Annotated keys to the genera of Nearctic Chalcidoidea (Hymenoptera)*. National Research Council Research Press, Ottawa, Canada.
- Giorgini, M., Wang X-G., Wang, Y., Chen, F-S., Hougardy, E., Zhang, H-M., Chen, Z-Q., Chen, H-Y., Liu, C-X., Cascone, P., Formisano, G., Carvalho, G.A., Biondi, A., Buffington, M., Daane, K.M., Hoelmer K.A., & Guerrier, E. (2019). Exploration for native parasitoids of *Drosophila suzukii* in China reveals a diversity of parasitoid species and narrow host range of the dominant parasitoid. *Journal of Pest Science*, **92(2)**, 509-522. <https://doi.org/10.1007/s10340-018-01068-3>
- Gómez Segade, C.B., Garrido, S.A., Aquino, D.A., Corley, J.C., & Cichón, L.I. (2021). Primer registro de *Spalangia endius* (Hymenoptera: Pteromalidae) asociado a *Drosophila suzukii* (Diptera: Drosophilidae) en cultivos de fruta fina y de carozo de la Patagonia Norte (Argentina). *Revista de la Sociedad Entomológica Argentina*, **80(2)**: 48-52 <https://doi.org/10.25085/rsea.800209>
- Gonsebatt, G., San Pedro, P., Fanara, J.J., Seta, S., Leone A., & Díaz, B.M. (2017). *Drosophila suzukii* (Matsumura) (Diptera: Drosophilidae) en durazneros (*Prunus persica*) en el sur de la provincia de Santa Fe: primer registro. In

- XIX Congreso and XXXVII Reunión Anual de la Sociedad de Biología de Rosario. Zavalla, Santa Fe, Argentina, pp. 73.
- Guimarães, J.A., Gallardo, F.E., Díaz, N.B., & Zucchi, R.A. (2003). Eucoilinae species (Hymenoptera: Cynipoidea: Figitidae) parasitoids of fruit-infesting dipterous larvae in Brazil: identity, geographical distribution and host associations. *Zootaxa*, **278**, 1-23. <https://doi.org/10.11646/zootaxa.278.1.1>
- Kirschbaum, D.S., Funes, C.F., Buonocore-Biancheri M.J., Suárez, L. & Ovruski, S.M. (2020). The biology and ecology of *Drosophila suzukii* (Diptera: Drosophilidae). *Drosophila suzukii Management* (ed. García, F.R.M.), pp. 41-91. Springer Nature, Switzerland AG. <https://doi.org/10.1007/978-3-030-62692-1>.
- Knoll, V., Ellenbroek, T., Romeis, J., & Collatz, J. (2017). Seasonal and regional presence of hymenopteran parasitoids of *Drosophila* in Switzerland and their ability to parasitize the invasive *Drosophila suzukii*. *Scientific Reports*, **7**, 40697. <https://doi.org/10.1038/srep40697>
- Lavagnino, N.J., Díaz, B.M., Cichón, L.I., de la Vega, G.J., Garrido, S.A., Lago, J.D., & Fanara, J.J. (2018). New records of the invasive pest *Drosophila suzukii* (Matsumura) (Diptera: Drosophilidae) in the South American continent. *Revista de la Sociedad Entomológica Argentina*, **77** (1), 27-31. <https://doi.org/10.25085/rsea.770105>
- Lue, C.H., Mottern, J.L., Walsh, G.C., & Buffington, M.L. (2017). New record for the invasive Spotted Wing *Drosophila*, *Drosophila suzukii* (Matsumura, 1931) (Diptera: Drosophilidae) in Anillaco, western Argentina. *Proceedings of the Entomological Society of Washington*, **119**, 146-150. <http://doi.org/10.4289/0013-8797.119.1.146>
- Marchiori, C.H., Arantes, S.B., Pereira, L.A., Moreira Silva Filho, O., & Rodrigues Borges, V. (2003). First record of *Leptopilina boulandi* Barbotin et al. (Hymenoptera: Figitidae: Eucoilinae) parasitizing of *Zaprionus indianus* Gupta (Diptera: drosophilidae) in Brazil. *Ciências Agrárias*, **24**(2), 321-324. <http://dx.doi.org/10.5433/1679-0359.2003v24n2p321>
- Nasca, A.J. (1976). Tiempo óptimo de exposición de las pupas de *Ceratitis capitata* (Wied.) y *Drosophila melanogaster* Meig. en el parasitismo de *Pachycrepoideus vindemniae* (Rond.). *Revista Agronómica del Noroeste Argentino*, **13**(1-4), 77-84.
- Nordlander, G. (1980). Revision of the genus *Leptopilina* Förster, 1869, with notes on the status of some other genera (Hymenoptera, Cynipoidea: Eucoilidae). *Entomologica Scandinavica*, **1**, 428-453. <http://doi.org/10.1163/1876631280794710024>
- Noyes, J.S. (2019). Universal Chalcidoidea Database. World Wide Web electronic publication. <http://www.nhm.ac.uk/chalcidooids>
- Núñez-Campero, S.R., & Ovruski, S.M. (2013). Registro de dos especies de himenópteros parasitoides neotropicales de mosca de la fruta en la localidad de Los Molinos, Departamento Castro Barros (La Rioja, Argentina). *Acta Zoológica Lilloana*, **57**, 7577.
- Ovruski, S.M. (2002). New records of fruit fly parasitoids (Hymenoptera: Braconidae, Figitidae, Pteromalidae) in La Rioja Province, northwestern Argentina. *Proceedings of the Entomological Society of Washington*, **104**(4), 1055-1057.
- Ovruski, S.M., Oroño, L.E., Núñez-Campero, S., Schliserman, P., Albornoz-Medina, P., Bezdjian, L.P., Van Nieuwenhove, G.A., & Martin, C.B. (2008). A review of hymenopterous parasitoid guilds attacking *Anastrepha* spp. and *Ceratitis capitata* (Diptera: Tephritidae) in Argentina. Fruit Flies of Economic Importance: From Basic to Applied Knowledge. Proceedings of the 7th International Symposium on Fruit Flies of Economic Importance. Salvador, Brazil 10-15 September 2006. (ed. Sugayama, R., Zucchi, R.A., Ovruski, S.M. & Sivinski, J.) pp. 113-125.
- Reche, V.A., & Gallardo, F.E. (2015). Revision of Argentinian species of *Hexacola* Foerster (Hymenoptera: Figitidae: Eucoilinae). *Journal of Insect, Biodiversity and Systematics*, **1**(1), 77-86.
- Reche, V.A., Gallardo, F.E., Funes, C., Escobar, L., Ovruski, S., & Kirschbaum, D. (2021). A new species of *Dieucoila* (Hymenoptera Figitidae) associated with *Drosophila suzukii* (Diptera Drosophilidae) in Argentina. *Bulletin of Insectology*, **74**(1): 11-18.
- Risbec, J. (1950). La faune entomologique des cultures au Sénégal et au Soudan: II. Contribution à l'étude des Proctotrupidae (Serphiidae). Gouvernement général de l'Afrique occidentale française, [s.l.], 509-638.
- Santadino, M.V., Riquelme Virgala, M.B., Ansa, M.A., Bruno, M., Di Silvestro, G., & Lunazzi, E.G. (2015). Primer registro de *Drosophila suzukii* (Diptera: Drosophilidae) asociado al cultivo de arándanos (*Vaccinium* spp.) de

- Argentina. *Revista de la Sociedad Entomológica Argentina*, **74** (3-4), 183-185.
- SENASA. (2016). Retrieved from <http://www.senasa.gob.ar/senasa-comunica/noticias/deteccion-de-drosophila-suzukii-en-la-argentina> (Accessed 03, 01, 2021)
- Turica, A., & Mallo, R.G. (1961). Observaciones sobre la población de las "Tephritidae" y sus endoparásitos en algunas regiones citrícolas argentinas. *Revista IDIA Suplement*, **6**, 145-161.
- Turica, A., Vergani, A.R., Quintanilla, R.H., Zerbino M. C., & Ceruso, H.E. (1971). Las moscas de los frutos. Instituto Nacional de Tecnología Agropecuaria. Serie Formación Técnica Agrícola, **7**, 1-17.
- Van Alphen, J.J. M., & Thunnissen, J. (1983). Host selection and allocation by *Pachycrepoideus vindemiae* Rondani (Pteromalidae) as a facultative hyperparasitoid of *Asobara tabida* (Braconidae: Alysiinae) and *Leptopilina heterotoma* (Cynipoidea: Eucolidae). *Netherlands Journal of Zoology*, **33**, 497-514. <http://doi.org/10.1163/002829683X00228>
- Vlach, J. (2010). Identifying *Drosophila suzukii*. Version from June 2, 2010. Oregon Department of Agriculture. agrireseau.qc.ca/lab/documents/clé_identification_drosophila_suzukii.pdf
- Wang, X.G., & Messing, R.H. (2004). The ectoparasitic pupal parasitoid, *Pachycrepoideus vindemiae* (Hymenoptera: Pteromalidae), attacks other primary tephritid fruit fly parasitoids: host expansion and potential non-target impact. *Biological Control*, **31**, 227-236. <http://doi.org/10.1016/j.biocontrol.2004.04.019>
- Wang, X., Daane, K.M., Hoelmer, K.A., & Lee, J.C. (2020). Biological Control of Spotted-Wing Drosophila: An Update on Promising Agents. *Drosophila suzukii Management* (ed. García, F.R.M.), pp 143-167. Springer Nature, Switzerland AG. https://doi.org/10.1007/978-3-030-62692-1_8.
- Wollmann, J., Hoffmann Schlesener, D.C., Soares Ferreira, M., Silveira Garcia, M., Costa, V.A., & Mello Garcia, F.R. (2016). Parasitoids of Drosophilidae with potential for parasitism on *Drosophila suzukii* in Brazil. *Drosophila Information Service*, **99**, 38-42. https://doi.org/10.1007/978-3-030-62692-1_8.