

Chemical Protein Unfolding – A Simple Cooperative Model

Joachim Seelig* and Anna Seelig




Cite This: *J. Phys. Chem. B* 2023, 127, 8296–8304




Read Online

ACCESS |

 Metrics & More

 Article Recommendations

ABSTRACT: Chemical unfolding with guanidineHCl or urea is a common method to study the conformational stability of proteins. The analysis of unfolding isotherms is usually performed with an empirical linear extrapolation method (LEM). A large positive free energy is assigned to the native protein, which is usually considered to be a minimum of the free energy. The method thus contradicts common expectations. Here, we present a multistate cooperative model that addresses specifically the binding of the denaturant to the protein and the cooperativity of the protein unfolding equilibrium. The model is based on a molecular statistical–mechanical partition function of the ensemble, but simple solutions for the calculation of the binding isotherm and the associated free energy are presented. The model is applied to 23 published unfolding isotherms of small and large proteins. For a given denaturant, the binding constant depends on temperature and pH but shows little protein specificity. Chemical unfolding is less cooperative than thermal unfolding. The cooperativity parameter σ is at least 2 orders of magnitude larger than that of thermal unfolding. The multistate cooperative model predicts zero free energy for the native protein, which becomes strongly negative beyond the midpoint concentration of unfolding. The free energy to unfold a cooperative unit corresponds exactly to the diffusive energy of the denaturant concentration gradient necessary for unfolding. The temperature dependence of unfolding isotherms yields the denaturant-induced unfolding entropy and, in turn, the unfolding enthalpy. The multistate cooperative model provides molecular insight and is as simple to apply as the LEM but avoids the conceptual difficulties of the latter.



INTRODUCTION

Chemical denaturation is a process by which the protein conformation is unfolded via addition of denaturants such as guanidineHCl, urea, or SDS (sodium dodecyl sulfate). Chemical denaturation is a common method for determining a protein's conformational stability, relative to its functional properties.^{1,2} Identifying the conditions that maximize the structural stability of a protein is crucial during the development of biologics for therapeutic treatments. Several complementary techniques should be applied to provide a systematic analysis of protein stability.³

Chemical unfolding of proteins is analyzed almost exclusively with a 2-state model, the linear extrapolation method (LEM).⁴ The LEM is an empirical approximation, and its conceptual difficulties have been realized since its initial proposal.^{4–6} In particular, the LEM assigns a large positive Gibbs free energy, $\Delta G_0^{\text{H}_2\text{O}}$ to the native protein. However, “the general understanding in the protein folding field has been that proteins fold into their native conformations driven by the decrease in Gibbs free energy (negative ΔG).”⁷ This thermodynamic hypothesis has become the default physical description of protein folding. In this view, the native state is the most stable one, that is, the global G minimum, not a maximum.

The 2-state model is a noncooperative model. It has no energy parameter for the interaction between neighboring amino acid residues. All amino acid residues unfold simultaneously (all-or-none model).⁸ However, the conforma-

tional change of all amino acid residues at the same time is physically unrealistic. Instead, “peptides that form helices in solution do not show a simple two-state equilibrium between a fully folded and fully unfolded structure. Instead, they form a complex mixture of all helix, all coil, or, most frequently, central helices with frayed coil ends”.⁹ A sequential cooperative unfolding of protein domains is, therefore, a physically more realistic alternative.


We have recently proposed a semiempirical model that describes a cooperative protein unfolding.¹⁰ The model assumes explicitly the binding of the denaturant D to the protein with the binding constant K_D and the cooperative unfolding of the protein with the cooperativity parameter σ .¹⁰ Here, we provide a modification of this model, based on a statistical-mechanical partition function leading to simple expressions for the chemical unfolding isotherm and the associated free energy.

Published chemical unfolding isotherms obtained with spectroscopic techniques and with calorimetry are analyzed. GuanidineHCl, urea, and SDS were studied as denaturants. The protein size ranged from 30 to ~ 1600 amino acid residues,

Received: May 26, 2023

Revised: August 8, 2023

Published: September 22, 2023



including a monoclonal antibody. The present model yields excellent simulations of all unfolding isotherms. The native protein is the reference state with a zero free energy. The free energy becomes negative upon unfolding and decreases with the logarithm of the denaturant concentration. The temperature dependence of the free energy provides the unfolding entropy and, in turn, the unfolding enthalpy. The latter agrees with the calorimetric measurements. The binding constant K_D depends essentially on the type of denaturant and varies little with the nature of the protein. The cooperativity parameter for chemical unfolding is compared to that for thermal unfolding.¹¹ The multistate cooperative model is firmly grounded in statistical mechanical thermodynamics. With a multistate cooperative approximation, we provide a simple expression for cooperative chemical denaturation analysis, which is equally easy to apply as the LEM.

MATERIALS AND METHODS

Chemical unfolding experiments with guanidine HCl, urea, and SDS (subsequently performed with different spectroscopic and calorimetric techniques) are selected from the available literature. The focus is on the chemical unfolding of lysozyme with guanidineHCl, urea, and SDS. Altogether, 23 published chemical denaturation isotherms of proteins of different structure and size have been investigated.

Chemical Unfolding Models. Chemical denaturants such as guanidineHCl and urea are commonly used to study protein stability. They change the polarity of the environment, bind to backbone and amino acid residues, and thus change the native protein conformation (N) into the unfolded conformation (U) (all-or-none folding). In contrast, the transition of individual amino acids from their native state (n) to their unfolded state (u) denotes multistate unfolding.

Multistate Cooperative Unfolding Model. The multistate cooperative model considers the individual amino acid residues of the protein. The amino acid residues in the native or in the unstructured confirmation are designated as “n” and “u”. The initial step of this model is the binding of a denaturant D to an amino acid “n” in a structured protein segment, inducing a conformational transition to an unstructured state “u”:



This chemical equilibrium is described by a simple equation:

$$K_D(T) = \frac{c_u}{c_n c_D} \quad (2)$$

The concentrations c_n and c_u are the concentrations of the amino acid residues participating in unfolding. The binding constant $K_D(T)$ is a function of the temperature only. Unfolding is a dynamic equilibrium between many different protein conformations.

The statistical interpretation of a chemical equilibrium requires a grand canonical partition function. To make the connection to the textbook literature, we consider a system of only two types of particles. “In a grand canonical ensemble, the number of type A particles, N_A , and type B particles, N_B , are both variable. Let μ_A and μ_B be the respective chemical potentials of the two components. The grand partition function is

$$Z(\mu_A, \mu_B, V, T) = \sum_{N_A, N_B, i} e^{\beta(N_A \mu_A + N_B \mu_B - E_{N_A, N_B, i})} \quad (3)$$

(ref 12, chapter 13.9). Equation 2 defines a three-component system measuring the ensemble size in concentration units c_i (mol/L). The chemical potential is given by

$$\sum_i \mu_i = \sum_i (\mu_i^0 + RT \ln c_i) = \Delta G(c_D) \quad (4)$$

$$\Delta G(c_D) = -RT(\ln K_D + \ln c_D + \ln c_u - \ln c_n) \quad (5)$$

In chemical unfolding experiments, the protein concentration is typically $\sim 10 \mu\text{M}$, whereas the concentration of the denaturant is 1–8 M. The chemical potential of the equilibrium is dominated by the denaturant concentration, leading to the following approximation.

$$\Delta G(c_D) = \sum_i \mu_i \cong -RT \ln (K_D c_D) \quad (6)$$

The conditional probability, $q(c_D)$ of a residue “n” in a structured protein segment is defined as 1, the conditional probability of the unfolded residue “u” is

$$q(c_D) = e^{-\Delta G/RT} = K_D c_D \quad (7)$$

provided the residue is located at the end of a native stretch of amino acid residues. If unfolding occurs in the middle of a native segment, unfolding is more difficult, and the corresponding conditional probability is $\sigma q(c_D)$ ($\sigma \ll 1$). The conditional probability $q(c_D)$ is inserted into the Zimm–Bragg partition function $Z(c_D)$.¹²

$$\begin{aligned} Z(c_D) &= (1 \quad 0) \begin{pmatrix} 1 & \sigma q(c_D) \\ 1 & q(c_D) \end{pmatrix}^\nu \begin{pmatrix} 1 \\ 1 \end{pmatrix} \\ &\approx (1 \quad 0) \begin{pmatrix} 1 & \sigma K_D c_D \\ 1 & K_D c_D \end{pmatrix}^\nu \begin{pmatrix} 1 \\ 1 \end{pmatrix} \end{aligned} \quad (8)$$

ν denotes the number of amino acid residues participating in the unfolding reaction. The cooperativity parameter σ determines the steepness of the unfolding transition. A small σ value leads to a sharp transition. The cooperativity parameter in chemical unfolding is typically $\sigma \approx 10^{-2}$ – 10^{-3} and is 10 to a hundred times larger than that of thermal unfolding. The fraction of unfolded protein is

$$\Theta_U(c_D) = \frac{q(c_D)}{\nu} \frac{d(\ln Z(c_D))}{dc_D} \left(\frac{dq(c_D)}{dc_D} \right)^{-1} \quad (9)$$

For a noncooperative ensemble with $\sigma = 1$, the partition function eq 8 reduces to $Z(q) = (1 + q)^\nu$. The fraction of unfolded protein becomes independent of ν and is, $\Theta_U(q) = \frac{\partial \ln Z(q)}{\partial \ln q} = \frac{q}{1 + q}$, which is identical to the Langmuir adsorption isotherm.

Figure 1 shows the unfolding isotherm $\Theta_U(c_D)$ (eq 9) for different cooperativity parameters σ . The binding constant is $K_D = 0.25 \text{ M}^{-1}$, which is typical for binding of guanidineHCl to most proteins. The binding constant is too small to induce complete protein unfolding for a noncooperative ensemble ($\sigma = 1$) as demonstrated by the Langmuir isotherm in Figure 1A. A dramatic change in the binding isotherm is induced by including even small cooperative interactions (red to green lines in Figure 1A).

The extent of unfolding (eq 9) is the result of the partition function $Z(c_D)$. Likewise, the free energy is also related to the




Figure 1. Multistate cooperative unfolding model. Variation of the cooperativity parameter σ . (A) Extent of unfolding $\Theta_U(c_D)$ (eq 9). Black: $\sigma = 1$. Red: $\sigma = 10^{-1}$. Blue: $\sigma = 10^{-2}$. Green: $\sigma = 10^{-3}$ (B) Free energy as a function of cooperativity parameter σ and denaturant concentration (eq 10). Binding constant, $K_D = 0.25 \text{ M}^{-1}$. Midpoint concentration $c_0 = 4.0 \text{ M}$. Number of amino acid residues participating in transition $\nu = 129$. Temperature $T = 298 \text{ K}$. The simulation parameters K_D , c_0 , ν , T and $\sigma = 10^{-3}$ correspond to the unfolding of lysozyme in the guanidineHCl solution.

partition function according to standard statistical thermodynamics.¹³ The free energy change of unfolding is

$$\Delta F(c_D) = -RT_0 \ln(Z(c_D)) \quad (10)$$

$\Delta F(c_D)$ upon addition of guanidineHCl is displayed in Figure 1B. The native protein is the reference state with zero free energy. Upon addition of denaturant, the free energy decreases. In the case of no cooperativity ($\sigma = 1$), the free energy decreases already at low concentrations of denaturant. For a cooperative ensemble, the free energy remains close to zero up to the midpoint of unfolding c_0 and decreases rapidly beyond this concentration. Compared to noncooperative denaturation, the free energy change for a cooperative system is distinctly smaller.

The temperature dependence of the free energy is

$$\frac{\partial \Delta F}{\partial T} = -\Delta S \quad (11)$$

The unfolding enthalpy can then be calculated as

$$\Delta H = \Delta F + T\Delta S \quad (12)$$

Simple Multistate Cooperative Approximation. For the biochemical practitioner, the above formalism may act as a deterrent. Fortunately, the matrix eq 8 can be replaced by a simple linear expression, which can easily be calculated. The cooperativity parameter in chemical unfolding experiments is always $\sigma \geq 10^{-3}$, and the largest eigenvalue λ_0 of the above matrix is a sufficient approximation, resulting in a simpler partition function¹²

$$Z_\lambda(c_D) = (\lambda_0(c_D))^\nu \quad (13)$$

with

$$\lambda_0(c_D) = \frac{1 + K_D c_D + [(1 - K_D c_D)^2 + 4\sigma K_D c_D]^{1/2}}{2} \quad (14)$$

The fraction of unfolded protein is given by

$$\Theta_{U,\lambda}(c_D) \cong \frac{1}{2} \left(1 + \frac{K_D c_D - 1 + 2\sigma}{[(1 - K_D c_D)^2 + 4\sigma K_D c_D]^{1/2}} \right) \quad (15)$$

At the midpoint of unfolding with $c_D = c_0$ and $\Theta(c_0) = 1/2$ follows

$$K_D c_0 = 1 \text{ or } K_D = 1/c_0 \quad (16)$$

In the multistate cooperative model, the binding constant of the denaturant is simply the reciprocal of the midpoint concentration of unfolding. Equation 15 can also be written as

$$\Theta_{U,\lambda}(c_D) \cong \frac{1}{2} \left(1 + \frac{c_D/c_0 - 1 + 2\sigma}{[(1 - c_D/c_0)^2 + 4\sigma c_D/c_0]^{1/2}} \right) \quad (17)$$

Equations 15 and 17 are equivalent to a cooperative sorption isotherm, which is based on a statistical-mechanical partition function. They are in line with many other sorption isotherms.¹⁴ The only fit parameter in the simulation of chemical unfolding isotherms is, therefore, the cooperativity parameter σ .

The free energy of the system is

$$\Delta F_\lambda(c_D) = -\nu RT_0 \ln \lambda_0(c_D) \quad (18)$$

For denaturant concentrations $c_D \gg c_0$, the following approximation is valid

$$\Delta F_\lambda(c_D) \approx -\nu RT_0 \ln(K_D c_D) \text{ for } c_D \gg c_0 \quad (19)$$

Concentration Gradient $\Delta c = c_{\text{end}} - c_{\text{ini}}$ and the Free Energy of the $n \rightarrow u$ transition. The multistate cooperative model predicts a one-to-one relationship between the free energy of the concentration gradient $\Delta c = c_{\text{end}} - c_{\text{ini}}$ and the energy required to induce the $n \rightarrow u$ transition. Unfolding takes place in the concentration interval $c_{\text{ini}} \leq c_D \leq c_{\text{end}}$. The concentration gradient $\Delta c = c_{\text{end}} - c_{\text{ini}}$ constitutes a diffusive or osmotic free energy

$$\Delta G_{\text{Diff}} = -RT_0 \ln \frac{c_{\text{end}}}{c_{\text{ini}}} \quad (20)$$

The free energy at ambient temperature is typically $\Delta G_{\text{Diff}} \approx -0.4$ to -1.0 kcal/mol. In parallel, the binding of D is associated with the free energy of the $n \rightarrow u$ transition per amino acid residue Δg_{nu} .

$$\Delta g_{\text{nu}} = -RT_0 \ln \frac{q(c_{\text{ini}})}{q(c_{\text{end}})} = RT_0 \ln \frac{c_{\text{end}}}{c_{\text{ini}}} \quad (21)$$

The predicted unfolding free energy per amino acid residue is thus identical to the concentration gradient (with opposite sign), that is, $\Delta g_{\text{nu}} + \Delta G_{\text{diff}} = 0$. The multistate cooperative model is thus consistent with the thermodynamic expectation of a reversible equilibrium.

Chemical Equilibrium Two-State Unfolding. The common model to describe chemical unfolding isotherms is a non-cooperative two-state model, which has dominated the field for the last 30–40 years. Only two types of protein conformations are assumed in solution, the native protein (N) and the unfolded protein (U). No intermediate structures and no specific interaction between denaturant D and protein are considered. The equilibrium $N \rightleftharpoons U$ is simply

$$K_{\text{NU}}(c_{\text{D}}) = \frac{[\text{U}]}{[\text{N}]} \quad (22)$$

The equilibrium constant $K(c_{\text{D}})$ varies with the concentration of denaturant. To calculate $K_{\text{NU}}(c_{\text{D}})$, the model makes an unconventional assumption about the free energy.

$$\Delta G_{\text{NU}}(c_{\text{D}}) = \Delta G_0^{\text{H}_2\text{O}} - mc_{\text{D}} \quad (23)$$

Equation 23 is difficult to understand in thermodynamic terms for two reasons. First, a large positive free energy $\Delta G_0^{\text{H}_2\text{O}}$ is assigned to the stable native protein. Second, the free energy is linear, not the usual logarithmic function of the concentration c_{D} . The so-called linear extrapolation method allows the calculation of spectroscopic unfolding transitions. The fraction of unfolded protein $\Theta_{\text{U}}(c_{\text{D}})$ is

$$\begin{aligned} \Theta_{\text{U}}(c_{\text{D}}) &= K_{\text{NU}}(c_{\text{D}})/(1 + K_{\text{NU}}(c_{\text{D}})) \\ &= e^{-\Delta G_{\text{NU}}(c_{\text{D}})/RT_0}/(1 + e^{-\Delta G_{\text{NU}}(c_{\text{D}})/RT_0}) \end{aligned} \quad (24)$$

The disadvantage of this model is its conceptual simplicity of just two protein conformations and the rather questionable linear extrapolation method. It is generally believed that the native protein sits at a free energy minimum. In contrast, the LEM predicts a large positive free energy for a stable native protein.

By definition the midpoint concentration c_0 is the concentration where native and unfolded protein occur at equal concentrations and the free energy is consequently zero in the LEM. It thus follows that

$$m = \Delta G_0^{\text{H}_2\text{O}}/c_0 \quad (25)$$

m is not an independent variable, which is however often ignored in the analysis of spectroscopic data.

RESULTS

Chemical Unfolding of Lysozyme with GuanidineHCl at 30 °C. Figure 2 displays the chemical unfolding of lysozyme in guanidine HCl at 30 °C (experimental data from ref 15).




Figure 2. Unfolding of lysozyme in guanidineHCl solution at 30 °C. Red lines: multistate cooperative model. Blue lines: linear extrapolation method. Vertical lines: midpoint concentration $c_0 = 3.9$ M. (A) (Box solid) data taken from ref 15. Simulation parameters of the multistate cooperative mode: $K_{\text{D}} = 1/c_0 = 0.26 \text{ M}^{-1}$, $\sigma = 1 \times 10^{-3}$. LEM parameters: $\Delta G_0^{\text{H}_2\text{O}} = 8.837 \text{ kcal/mol}$, $m = \Delta G_0^{\text{H}_2\text{O}}/c_0 = 2.34 \text{ kcal/mol}^2$. (B) Temperature profiles of the free energy.

The midpoint concentration of unfolding is $c_0 = 3.9$ M. Chemical unfolding takes place in the concentration range of $2.9 \text{ M} \leq c_{\text{D}} \leq 5.3 \text{ M}$

The solid lines in Figure 2A simulate the experimental data with the parameters given in the legend of Figure 2. Both methods simulate the unfolding transitions equally well. Both models predict exactly 50% unfolding at the midpoint concentration c_0 , but the concentration profiles of the free energy are quite different (Figure 2B). The LEM assigns a large positive free energy, $\Delta G_0^{\text{H}_2\text{O}} = 8.93 \text{ kcal/mol}$, to the native protein. At the midpoint concentration c_0 , the free energy is exactly zero ($\Delta G(c_0) = 0$). The multistate cooperative model predicts zero free energy for the native protein. The free energy change is slightly negative up to c_0 , ($\Delta F = -2.0 \text{ kcal/mol}$) and decreases rapidly at $c_{\text{D}} > c_0$ (eq 18). The change in free energy upon unfolding (in the concentration range $2.9 \text{ M} \leq c_{\text{D}} \leq 5.3 \text{ M}$) is $\Delta F = -21.7 \text{ kcal/mol}$ for the multistate cooperative model but only $\Delta G = -11.8 \text{ kcal/mol}$ for the LEM.

Lysozyme Unfolding in Urea. Urea is less effective in chemical denaturation than guanidineHCl as is obvious in Figure 3. The urea midpoint concentration is 6.7 M compared to 4.1 M for guanidine HCl. All thermodynamic data are summarized in Table 2, see ref 16.




Figure 3. Chemical unfolding of lysozyme with guanidineHCl (box solid) and urea (circle solid) at 25 °C and pH 7. Data taken from ref 16. The midpoint concentrations $c_0 = 4.1$ M for guanidine HCl cooperativity parameter is $\sigma = 2 \times 10^{-3}$ for both denaturants. The figure also compares the exact and approximate solutions for the binding isotherm and free energy. (A) Extent of unfolding. Red line: matrix solution eq 9. Black line: simplified isotherm eq 17. (B) Free energy of unfolding. Red line: eq 10. Black line: eq 18.

The urea-induced transition (unfolding concentration range $\Delta c = 4.8 \text{ M}$) is broader than that of guanidineHCl ($\Delta c = 3 \text{ M}$). However, in spite of these large differences, the corresponding diffusive or osmotic free energies ΔG_{diff} (eq 20) constituted by these gradients are identical with $\Delta G_{\text{diff}} = -0.431 \text{ kcal/mol}$. Likewise, the free energy changes $\Delta F \approx -24 \text{ kcal/mol}$ and the cooperativity parameter $\sigma = 2 \times 10^{-3}$ are also identical for the two denaturants. The only significant thermodynamic difference is that the urea binding constant $K_{\text{D}} = 0.15 \text{ M}^{-1}$ is distinctly smaller than the guanidineHCl binding constant $K_{\text{D}} = 0.245 \text{ M}^{-1}$. The broadening of the transition region in the urea solution is thus caused by the low urea binding constant and not by a change in the protein cooperativity.

Figure 3 compares the exact and the approximate solutions for the extent of unfolding and (eqs 9 and 10 versus eqs 17 and

18). The approximate solutions show almost complete overlap with the exact solutions.

Temperature-Dependence of Chemical Unfolding of Lysozyme. Lysozyme unfolding depends on the pH and temperature. A decrease in pH or an increase in temperature shift the unfolding transition toward lower c_0 values. This is illustrated in Figure 4 for chemical denaturation with guanidineHCl at three different temperatures.




Figure 4. Temperature dependence of lysozyme unfolding in guanidineHCl solution. Data points taken from Figure 2 of ref 17. Solid lines are the simulations of the multistate cooperative model. Blue: 60 °C, $K_D = 0.9 \text{ M}^{-1}$, $\sigma = 3 \times 10^{-2}$. Red: 40 °C, $K_D = 0.34 \text{ M}^{-1}$, $\sigma = 2.5 \times 10^{-3}$. Black: 10 °C, $K_D = 0.26 \text{ M}^{-1}$, $\sigma = 1 \times 10^{-3}$. (A) Extent of unfolding $\Theta_U(c_D)$. Simulations with eq 9. (B) Free energy $F(c_D)$, calculated with eq 10. (pink star) Free energies at 95% unfolding, calculated with eq 19.

Figure 4A displays unfolding isotherms obtained with UV and CD spectroscopy.¹⁷ The solid lines are simulations with the multistate cooperative model. The binding constants K_D are exactly the reciprocal of the midpoint concentrations c_0 . The binding constants increase with temperature and the transition regions broaden. The cooperative interactions are reduced, leading to a larger cooperativity parameter σ . A temperature increase from 10 to 60 °C increases the cooperativity parameter by a factor of about 10.

Figure 4B displays the predicted concentration dependence of the free energy. The free energy is zero for the native protein

and decreases rapidly beyond the midpoint of unfolding c_0 . The solid lines are calculated with the exact solution eq 9. The three stars-marked data points were calculated with eq 19, corresponding to 95% unfolding. The comparison with the exact solution demonstrates that the simple eq 19 is an excellent approximation for the free energy if $c_d \gg c_0$.

We have analyzed published denaturation experiments of lysozyme in guanidineHCl solutions at different temperatures, and the results are summarized in Table 1.

Polypeptides and proteins of different sizes and structures were also analyzed. The fits of the experimental data with the multistate cooperative model were all of comparable quality as shown for lysozyme in Figures 2–4. The data are summarized in Table 2.

Chemical Unfolding with Sodium Dodecyl Sulfate (SDS). SDS is a much stronger denaturant than guanidine HCl or urea as shown in Figure 5 (data taken from ref 18). Unfolding was measured with calorimetry. At 25 °C, the midpoint concentration of unfolding is only 4.35 mM, resulting in a large binding constant of $K_D = 230 \text{ M}^{-1}$. The cooperativity of SDS-induced unfolding is relatively low with $\sigma = 7 \times 10^{-2}$.

The enthalpy of unfolding, ΔH_0 , is the most important parameter in the thermal unfolding studies. Spectroscopic measurements of chemical unfolding isotherms cannot provide this thermodynamic property. It is obtained, however, by a direct calorimetric measurement, as shown in Figure 5. Lysozyme is titrated with low concentrations of SDS, and the heat of reaction is measured in a calorimeter. Each data point in Figure 5 corresponds to an independent measurement.¹⁸ The total unfolding enthalpy is $\Delta H_n = 55 \text{ kcal/mol}$ at 25 °C. The analogous measurement at 35 °C results in 61.3 kcal/mol, and the heat capacity change is $\Delta C_p = 0.65 \text{ kcal/molK}$. Likewise, the result of an early isothermal enthalpimetric titration of lysozyme with guanidineHCl at pH 2.5 and 40 °C yielded 56 kcal/mol.¹⁹

Altogether, 22 chemical unfolding isotherms of polypeptides and proteins of different size and structure were analyzed. The fits of the experimental data with the multistate cooperative model were all excellent. The results are summarized in Table 2.

DISCUSSION

Characteristics of the Multistate Cooperative Model.

The multistate cooperative model is based on a statistical–mechanical partition function that contains two molecular

Table 1. Chemical Unfolding of Lysozyme with Guanidine HCl at Different Temperatures

temp °C	c_{mid} M	K_D M^{-1}	σ	c_{ini} M	c_{end} M	ΔF kcal/mol	g_{den} kcal/mol	$\Delta G_{\text{H}_2\text{O}}$ kcal/mol	m kcal L/mol ²
10 ²⁰	3.8	0.263	1.50×10^{-03}	2.7	5.3	−21.8	−0.379	9.55	2.512
15 ²⁰	4.0	0.25	1.80×10^{-03}	2.7	5.6	−22.7	−0.417	8.598	2.150
20 ²⁰	3.9	0.257	2.00×10^{-03}	2.6	5.5	−23.8	−0.436	8.598	2.210
25 ²⁰	3.8	0.263	2.50×10^{-03}	2.4	5.5	−26.1	−0.491	9.554	2.513
35 ²⁰	3.3	0.303	5.00×10^{-03}	1.75	5.1	−32.6	−0.654	8.598	2.605
10 ¹⁷	3.8	0.26	1.00×10^{-03}	2.9	5.25	−20.1	−0.334	9.55	2.483
40 ¹⁷	2.9	0.34	2.50×10^{-03}	1.9	4.1	−26.41	−0.493	8.36	2.842
60 ¹⁷	1.1	0.9	3.00×10^{-02}	0.25	2.4	−65.5	−1.5	4.299	3.869
25 ²²	3.9	0.256	1.00×10^{-03}	2.8	5.15	−21.2	−0.361	10.03	2.568
10 ²¹	4.1	0.243	2.00×10^{-03}	2.7	5.8	−22.55	−0.43	8.837	2.147391
30 ¹⁵	3.9	0.258	1.00×10^{-03}	2.9	5.3	−21.7	−0.363	8.837	2.236
45 ¹⁵	3.2	0.31	1.00×10^{-02}	1.3	5.3	−39.3	−0.888	4.777	2.236

Table 2. Chemical Unfolding of Peptides and Proteins of Different Sizes and Structures in Guanidine HCl and Urea

protein, peptide	N_{aa}	pH	denaturant	K_D M^{-1}	c_{mid} M	σ	c_{ini} M	c_{end} M	Δc M	g_{den} kcal/mol	ΔF kcal/mol	lit.	figure
BBL	41	7	GndHCl	0.33	3.03	3.00×10^{-02}	0.2	7.5	7.3	-2.14	-20.9	31	Figure 2B
ubiquitin WT	76		GndHCl	0.34	2.94	7.00×10^{-03}	1.4	5	3.6	-0.753	-22.1	32	Figure 1
ubiquitin L67S	76		GndHCl	0.72	1.39	1.30×10^{-02}	0.5	2.7	2.2	-0.998	-28.4	32	Figure 1
ubiquitin	76	2	GnddHCl	0.395	2.53	8.00×10^{-03}	1.2	4.5	3.3	-0.782	-24.1	33	Figure 4
ubiquitin	76	5.5	GndHCl	0.295	3.39	1.20×10^{-02}	1.3	6.3	5	-0.934	-26.4	33	
Apo lipoprotein A1	120	8	GndHCl	1	1.00	2.00×10^{-02}	0.3	2	1.7	-1.12	-49.7	34	Figure 1
Apo lipoprotein A1	122	7.4	GndHCl	0.952	1.05	5.00×10^{-03}	0.6	1.7	1.1	-0.61	-32.8	35	Figure 7
Apo lipoprotein A1	122	7.4	GndHCl	0.95	1.05	3.00×10^{-02}	0.3	2.3	2	-1.21	-55.9	36	
lysozyme	129	7	GndHCl	0.245	4.08	2.00×10^{-03}	2.8	5.8	3	-0.431	-24.6	16	Figure 1
lysozyme	129	2	GndHCl	0.42	2.38	2.00×10^{-02}	0.8	4.6	3.8	-1.03	-49.6	37	Figure 2A
lysozyme	129	2.5	GndHCl	0.48	2.08	5.00×10^{-03}	1.1	3.2	2.1	-0.632	-31.1	37	Figure 1
PMS-Ct	145	4	GndHCl	0.52	1.92	1.00×10^{-03}	1.1	2.5	1.4	-0.486	-22.7	38	Figure 4
PMS-Ct	241	4	GndHCl	0.513	1.95	4.00×10^{-03}	1.1	2.7	1.6	-0.531	-45.3	38	Figure 4
average				0.551	2.22	0.012			2.931	-0.90			
STDV				0.256	0.93	0.0096			1.659	0.43			
Ac-tyr-(ala-glu-ala-ala-lys-ala) ₅ -phe-NH ₂	32	7	urea	0.25	4.00	8.00×10^{-02}						6	Figure 1
Ac-tyr-(ala-glu-ala-ala-lys-ala) ₈ -phe-NH ₂	50	7.0	urea	0.2	5.00	1.00×10^{-02}	2.2	9.8	7.6	-0.884	-18.1	39	Figure 1 273 K
HPr protein	85	7	urea	0.215	4.65	8.00×10^{-03}	2	8	6	-0.793	-26.2	40	Figure 1B 288 K
cytochrome c	106	5	urea	0.2	5.00	4.00×10^{-03}	2.9	7.7	4.8	-0.578	-25.1	3	
cytochrome c	106	7	urea	0.155	6.46	4.00×10^{-03}	3.6	9.5	5.9	-0.574	-22.5	3	Figure 5
P22 1-domain	123	6	urea	0.3	3.33	9.00×10^{-03}	1.5	5.6	4.1	-0.753	-35.18	41	Figure 2
lysozyme	129	2	urea	0.303	3.30	2.00×10^{-02}	2.2	6.4	4.2	-0.632	-47.9	37	Figure 2B
lysozyme	129	5.5	urea	0.15	6.67	2.00×10^{-03}	4.5	9.3	4.8	-0.43	-23.3	16	Figure 1
anti-EF-R mab	1600	7	urea	0.18	5.56	2.00×10^{-02}	2	8.3	6.3	-0.85	-407	3	Figure 2
average				0.22	4.80	1.74×10^{-02}			5.46	-0.69			
STDV				0.05	1.15	0.0229			1.12	0.15			
lysozyme	129		SDS	230.00	0.00	7.00×10^{-02}	0.001	0.012	0.011	-1.47	-77.7	18	Table 1 298 K




Figure 5. Calorimetric titration of 68 μ M lysozyme with a SDS solution at 25 $^{\circ}$ C. (A) Experimental data taken from ref 18, Table 1. The published data were normalized to 1 mol lysozyme. Each data point is a separate measurement. Red line: multistate cooperative model. $K_D = 230 M^{-1}$, $\sigma = 7 \times 10^{-2}$, enthalpy of unfolding $\Delta H_0 = 55$ kcal/mol. (B) Unfolding isotherm. Red line: multistate cooperative model, calculated with the same parameters as listed in panel (A). Blue line: K_D as in panel (A), but $\sigma = 1$ (noncooperative Langmuir isotherm).

parameters: the binding constant $K_D = 1/c_0$ and the protein cooperativity σ . The cooperativity parameter determines the steepness of the unfolding transition (Figure 1). The model

takes into account the number ν of amino acid residues participating in the unfolding reaction and makes the following predictions. (i) The free energy of the $n \rightarrow u$ transition Δg_{nu} is identical to the free energy provided by the concentration gradient ΔG_{diff} of complete unfolding (eqs 20 and 21). (ii) The free energy of the native protein is the reference state with a zero free energy. This is the minimum free energy of a stable protein. Unfolding requires energy, which is stored in the unfolded protein and can be delivered if the unfolded protein returns to its ground state. (iii) A simple approximation can be given (eqs 17–19), which fits the experimental data extremely well and is as easy to apply as the LEM.

Temperature Dependence of Unfolding. Denaturant binding is a complex process. The denaturant binds to the protein backbone and to amino acid side chains. Binding can be electrostatic or hydrophobic. In addition, the denaturant can change the hydrophobicity of the hydration layer. The binding affinity will be different for different amino acid residues. The binding constant and the cooperativity parameter reported in this paper are average values of all possible interactions. An interpretation in terms of a specific binding model is not attempted.

The temperature dependence of the binding constant K_D and the cooperativity parameter σ together with that of the free energy is displayed in Figure 6 for lysozyme (Table 1). The simulation of the unfolding isotherms yields excellent fits in each case. Nevertheless, the scatter of the data in Figure 6 is considerable, as the isotherms are obtained by different authors




Figure 6. Lysozyme unfolding experiments in guanidineHCl at different temperatures and by different authors. (box solid),²⁰ (circle solid),¹⁷ (pink diamond),^{15,21} (blue triangle)²². The solid lines are introduced to guide the eye.

under different experimental conditions. Two temperature regions can be discerned. Between 10 and 35 °C, the effect of temperature is small, above 35 °C all properties change rapidly.

The midpoint concentration and the free energy decrease with temperature, and the cooperativity parameter and the binding constant increase. At higher temperatures, the protein cooperativity decreases. The cooperativity parameter of chemical unfolding is at least 2 orders of magnitude larger than that of thermal unfolding.^{11,23–25} For example, the σ -parameter of lysozyme for chemical unfolding at 25 °C is $\sigma = 1 \times 10^{-3}$ (see Figure 2) but is only $\sigma = 10^{-6}$ for temperature-induced unfolding (see Figure 1 in ref 11). σ -Parameters of thermal unfolding of other proteins are found in refs 11,23–25. Chemical unfolding is less cooperative than thermal unfolding.

A detailed analysis of the temperature dependence of the free energy is given in Figure 7. $\Delta H = \Delta F + T\Delta S$. Figure 7A displays data obtained from a consistent set of experiments for the temperature range 10 to 25 °C (Figure 1 in ref 20).

From the negative slope of the straight line in Figure 7A, the entropy $-\Delta S = \frac{\partial \Delta F}{\partial T}$ is calculated as $\Delta S = 0.280 \pm 0.047$ kcal/mol K. Figure 7B then displays the entropy term $T\Delta S$ and the predicted enthalpy $\Delta H = \Delta F + T\Delta S$. The average enthalpy is $\Delta H_0 = 58 \pm 1$ kcal/mol. An early isothermal titration study of lysozyme with guanidineHCl at pH 2.5 and 40 °C yielded $\Delta H = 56$ kcal/mol.²⁶ The present result is also in agreement with the SDS isothermal calorimetry shown in Figure 5, resulting in an enthalpy of 55 kcal/mol at 25 °C.

The general understanding in the protein folding field has been that proteins fold into their native conformations driven by a decrease in free energy (negative ΔG).⁷ The native protein is thermodynamically the most stable conformation. In the so-called funnel hypothesis, the native protein sits at the bottom of a rough-edged funnel, representing the minimum free energy. The multistate cooperative model is fully consistent with this hypothesis. The native protein is the




Figure 7. Chemical unfolding of lysozyme in guanidineHCl solution at different temperatures. (A) Free energy $\Delta F(T)$ calculated with the multistate cooperative model. Evaluation of unfolding isotherms of ref 20. Straight line: linear regression analysis with slope $-\Delta S = -0.28 \pm 0.046$ kcal/molK ($R^2 = 0.947$). (B) (pink diamond) Enthalpy ΔH and (blue triangle) entropy $T\Delta S$.

reference state with zero free energy, and unfolding requires energy. In returning reversibly to the native state, the unfolded protein loses its free energy.

GuanidineHCl and Urea Binding Constants. Different molecular mechanism of denaturation have been proposed.^{5,27} An indirect mechanism postulates changes in the water structure and hydrophobic effect. The alternative view is the direct interaction of the chemical denaturant with the protein. Strong support for the latter mechanism comes from isothermal titration calorimetry (ITC) of guanidineHCl and urea with various proteins.²⁸ X-ray studies also demonstrate a close interaction between guanidineHCl and the protein backbone of lysozyme.²⁹

The guanidineHCl binding constants deduced from the lysozyme isotherms are $0.2 \text{ M}^{-1} \leq K_D \leq 0.9 \text{ M}^{-1}$ in the temperature range 10–60 °C (Table 1). Urea binds with a lower affinity of $K_D = 0.15 \text{ M}^{-1}$ (at 25 °C). Isothermal titration calorimetry was used to measure guanidineHCl and urea binding to different proteins.²⁸ The data were analyzed by assuming a set of independent noncooperative binding sites.²⁸ The binding constants deduced for lysozyme were $K_D \approx 0.4\text{--}0.8 \text{ M}^{-1}$ for guanidine HCl and 0.06 M^{-1} for urea. This is in broad agreement with the results of the multistate cooperative model.

The binding constants of guanidineHCl and urea are small, and the binding of a single guanidineHCl molecule is not sufficient to induce the $n \rightarrow u$ transition of an amino acid residue. Only the cooperative binding of many denaturants induces protein unfolding.

The adsorption isotherm (eq 15) is also applicable to denaturation with organic solvents. Lysozyme denaturation isotherms have been reported for ethanol and DMSO.³⁰ The midpoint concentrations and corresponding binding constants are 5 and 0.25 M^{-1} for ethanol and 7 and 0.17 M^{-1} for DMSO. In contrast, anionic SDS has a high affinity for overall cationic lysozyme.¹⁸ Based on Figure 1 of ref 18, the midpoint concentration is $\sim 4.3 \text{ mM}$ and the binding constant $K_D \approx 230 \text{ M}^{-1}$.

free energy of the $N \rightleftharpoons U$ equilibrium

The multistate cooperative model provides a simple adsorption isotherm to analyze chemical unfolding (eq 15). The only unknown parameter is the cooperativity σ . The free energy follows from the partition function (eqs 10 or 18). The free energy of the native protein is zero, and the concentration profile is shown in Figure 1. Addition of denaturant initially leads to only a small negative free energy. The free energy becomes distinctly more negative at concentrations near and above the midpoint concentration c_0 . For large denaturant concentrations $c_D \gg c_0$, a simple approximation of $\Delta F(c_D) \approx -RT_0 N \ln(K_D c_D)$ (eq 19) is valid. Numerical values of the free energy change $\Delta F = \Delta F(c_{\text{end}}) - \Delta F(c_{\text{ini}})$, calculated with eq 10, are given in Table 1. The extent of unfolding in these calculations is $0.01 \leq \Theta_U \leq 0.95$. It should also be noted that the shape of the free energy in chemical unfolding (Figure 1) is identical to the free energy temperature profile in temperature-induced unfolding.¹¹ However, the free energy change ΔF in chemical unfolding is about 3-fold higher more negative than the free energy of thermal unfolding.¹¹ This must be traced back to the binding of the denaturants.

Cooperativity in Chemical and Thermal Unfolding.

Chemical σ parameters are plotted as a function of the temperature in Figure 6. Cooperativity parameter is $\sigma \approx 3 \times 10^{-3}$ for temperatures below 30 °C and strongly increases at higher temperature. The increase in temperature facilitates chemical unfolding by decreasing the cooperativity of the protein. The free energy to start a new folded sequence within an unfolded region is given by $\Delta F_\sigma = -RT \ln \sigma$.

CONCLUSIONS

It is obvious that the linear extrapolation method is a poor model to describe chemical unfolding. The two fit parameters have no well-defined thermodynamic or molecular basis. It is therefore attractive to modify an existing multistate cooperative model for thermal protein unfolding to address chemical unfolding. The new two-parameter model includes the binding

constant of the denaturant and the cooperativity of protein unfolding. It is simple and easy to apply, as the binding constant is the reciprocal of the midpoint concentration of unfolding. Simple equations can be given for the binding isotherm and free energy. The free energy profile of chemical unfolding parallels the experimental results for thermal unfolding. The free energy of the native protein is zero and becomes distinctly negative upon unfolding only after reaching the midpoint concentration. The new model is satisfying as it provides an excellent description of chemical denaturation isotherms and for the first time allows comparing cooperative, chemical, and thermal unfolding.

AUTHOR INFORMATION

Corresponding Author

Joachim Seelig – Biozentrum, University of Basel, CH-4056 Basel, Switzerland; orcid.org/0000-0001-8564-104X; Phone: +41-61-207 2190; Email: joachim.seelig@unibas.ch

Author

Anna Seelig – Biozentrum, University of Basel, CH-4056 Basel, Switzerland; orcid.org/0000-0003-3509-2569

Complete contact information is available at:

<https://pubs.acs.org/10.1021/acs.jpcb.3c03558>

Notes

The authors declare no competing financial interest.

REFERENCES

- Myers, J. K.; Bell, E.; Myers, J. K. *Chemical Denaturation. In Molecular Life Sciences*; Springer: New York, NY, 2014.
- Grimsley, G. R.; Trevino, S. R.; Thurlkill, R. L.; Scholtz, J. M. Determining the conformational stability of a protein from urea and thermal unfolding curves. *Curr. Protoc Protein Sci.* **2013**, *28*, 28.4.1–28.4.1.
- Freire, E.; Schon, A.; Hutchins, B. M.; Brown, R. K. Chemical denaturation as a tool in the formulation optimization of biologics. *Drug Discovery Today* **2013**, *18*, 1007–1013.
- Pace, C. N. Determination and analysis of urea and guanidine hydrochloride denaturation curves. *Methods Enzymol.* **1986**, *131*, 266–280.
- Makhatadze, G. I. Thermodynamics of protein interactions with urea and guanidinium hydrochloride. *J. Phys. Chem. B* **1999**, *103*, 4781–4785.
- Bolen, D. W.; Yang, M. Effects of guanidine hydrochloride on the proton inventory of proteins: implications on interpretations of protein stability. *Biochemistry* **2000**, *39*, 15208–15216.
- Sorokina, I.; Mushegian, A. R.; Koonin, E. V. Is Protein Folding a Thermodynamically Unfavorable, Active, Energy-Dependent Process? *Int. J. Mol. Sci.* **2022**, *23*, 521.
- Zhou, Y.; Hall, C. K.; Karplus, M. The calorimetric criterion for a two-state process revisited. *Protein Sci.* **1999**, *8*, 1064–1074.
- Doig, A. J. Recent advances in helix-coil theory. *Biophys. Chem.* **2002**, *101–102*, 281–293.
- Li-Blatter, X.; Seelig, J. Thermal and Chemical Unfolding of Lysozyme. Multistate Zimm-Bragg Theory Versus Two-State Model. *J. Phys. Chem. B* **2019**, *123*, 10181–10191.
- Seelig, J.; Seelig, A. Molecular Understanding of Calorimetric Protein Unfolding Experiments. *Biophys. Rep.* **2022**, *2*, 100037.
- Davidson, N. *Statistical Mechanics*; Mac Graw-Hill: New York, 1962.
- Baumann, R. P. *Evaluation of thermodynamic properties In Modern Thermodynamics with Statistical Mechanics*; McConnin, R. A., Eds.; Macmillan Publishing Company: New York, USA; 1992; pp. 341.

- (14) Adamson, A. W. *Physical chemistry of surfaces*; John Wiley & Sons: New ork, N.Y., USA; 1976.
- (15) Ibarra-Molero, B.; Sanchez-Ruiz, J. M. A model-independent, nonlinear extrapolation procedure for the characterization of protein folding energetics from solvent-denaturation data. *Biochemistry* **1996**, *35*, 14689–14702.
- (16) Ahmad, F.; Bigelow, C. C. Estimation of the free energy of stabilization of ribonuclease A, lysozyme, alpha-lactalbumin, and myoglobin. *J. Biol. Chem.* **1982**, *257*, 12935–12938.
- (17) Sasahara, K.; Nitta, K. Pressure-induced unfolding of lysozyme in aqueous guanidinium chloride solution. *Protein Sci.* **1999**, *8*, 1469–1474.
- (18) Behbeheni, G. R.; Ramazani, S.; Gonbadi, K. A Thermodynamic Investigation on the Binding of Lysozyme with Sodium Dodecyl Sulfate. *Chem. Soc. Pak.* **2013**, *35*, 1427–1431.
- (19) Pfeil, W.; Privalov, P. L. Thermodynamic investigations of proteins. II. Calorimetric study of lysozyme denaturation by guanidine hydrochloride. *Biophys. Chem.* **1976**, *4*, 33–40.
- (20) Sasahara, K.; Sakurai, M.; Nitta, K. Pressure effect on denaturant-induced unfolding of hen egg white lysozyme. *Proteins* **2001**, *44*, 180–187.
- (21) Laurents, D. V.; Baldwin, R. L. Characterization of the unfolding pathway of hen egg white lysozyme. *Biochemistry* **1997**, *36*, 1496–1504.
- (22) Sasahara, K.; Sakurai, N.; Nitta, K. The volume and compressibility changes of lysozyme associated with guanidinium chloride and pressure-assisted unfolding. *J. Mol. Biol.* **1999**, *291*, 693–701.
- (23) Seelig, J.; Schönfeld, H.-J. Thermal protein unfolding by differential scanning calorimetry and circular dichroism spectroscopy Two-state model versus sequential unfolding. *Q. Rev. Biophys.* **2016**, *49*, No. e9.
- (24) Seelig, J.; Seelig, A. Protein Stability. Analysis of Heat and Cold Denaturation without and with Unfolding Models. *J. Phys. Chem. B* **2023**, *127*, 3352–3363.
- (25) Seelig, J.; Seelig, A. Protein Unfolding-Thermodynamic Perspectives and Unfolding Models. *Int. J. Mol. Sci.* **2023**, *24*, 5457.
- (26) Pace, C. N.; Vajdos, F.; Fee, L.; Grimsley, G.; Gray, T. How to measure and predict the molar absorption coefficient of a protein. *Protein Sci.* **1995**, *4*, 2411–2423.
- (27) Konermann, L. Protein unfolding and denaturants. In *eLSJohn*; Wiley & Sons, Ltd.: Chichester; 2012; 1–7.
- (28) Makhatadze, G. I.; Privalov, P. L. Protein interactions with urea and guanidinium chloride. A calorimetric study. *J. Mol. Biol.* **1992**, *226*, 491–505.
- (29) Raskar, T.; Koh, C. Y.; Niebling, S.; Kini, R. M.; Hosur, M. V. X-ray crystallographic analysis of time-dependent binding of guanidine hydrochloride to HEWL: First steps during protein unfolding. *Int. J. Biol. Macromol.* **2019**, *122*, 903–913.
- (30) Sirotkin, V. A.; Winter, R. Volume Changes Associated with Guanidine Hydrochloride, Temperature, and Ethanol Induced Unfolding of Lysozyme. *J. Phys. Chem. B* **2010**, *114*, 16881–16886.
- (31) Neuweiler, H.; Sharpe, T. D.; Johnson, C. M.; Teufel, D. P.; Ferguson, N.; Fersht, A. R. Downhill versus barrier-limited folding of BBL 2: mechanistic insights from kinetics of folding monitored by independent tryptophan probes. *J. Mol. Biol.* **2009**, *387*, 975–985.
- (32) Haririnia, A.; Verma, R.; Purohit, N.; Twarog, M. Z.; Deshaies, R. J.; Bolon, D.; Fushman, D. Mutations in the hydrophobic core of ubiquitin differentially affect its recognition by receptor proteins. *J. Mol. Biol.* **2008**, *375*, 979–996.
- (33) Ibarra-Molero, B.; Loladze, V. V.; Makhatadze, G. I.; Sanchez-Ruiz, J. M. Thermal versus guanidine-induced unfolding of ubiquitin. An analysis in terms of the contributions from charge-charge interactions to protein stability. *Biochemistry* **1999**, *38*, 8138–8149.
- (34) Edelstein, C.; Scanu, A. M. Effect of guanidine hydrochloride on the hydrodynamic and thermodynamic properties of human apolipoprotein A-I in solution. *J. Biol. Chem.* **1980**, *255*, 5747–5754.
- (35) Gwynne, J.; Brewer, B.; Edelhofer, H. The molecular properties of ApoA-I from human high density lipoprotein. *J. Biol. Chem.* **1974**, *249*, 2411–2416.
- (36) Mantulin, W. W.; Pownall, H. J. Reversible folding reactions of human apolipoprotein A-I: pressure and guanidinium chloride effects. *Biochim. Biophys. Acta* **1985**, *836*, 215–221.
- (37) Singh, R.; Hassan, M. I.; Islam, A.; Ahmad, F.; Khan, R. H. Cooperative Unfolding of Residual Structure in Heat Denatured Proteins by Urea and Guanidinium Chloride. *PLoS One* **2015**, *10*, No. e0128740.
- (38) Santoro, M. M.; Bolen, D. W. Unfolding free energy changes determined by the linear extrapolation method. I. Unfolding of phenylmethanesulfonyl alpha-chymotrypsin using different denaturants. *Biochemistry* **1988**, *27*, 8063–8068.
- (39) Scholtz, J. M.; Barrick, D.; York, E. J.; Stewart, J. M.; Baldwin, R. L. Urea unfolding of peptide helices as a model for interpreting protein unfolding. *Proc. Natl. Acad. Sci.* **1995**, *92*, 185–189.
- (40) Nicholson, E. M.; Scholtz, J. M. Conformational stability of the Escherichia coli HPr protein: test of the linear extrapolation method and a thermodynamic characterization of cold denaturation. *Biochemistry* **1996**, *35*, 11369–11378.
- (41) Newcomer, R. L.; Fraser, L. C. R.; Teschke, C. M.; Alexandrescu, A. T. Mechanism of Protein Denaturation: Partial Unfolding of the P22 Coat Protein I-Domain by Urea Binding. *Biophys. J.* **2015**, *109*, 2666–2677.