



Sex differences in brain structures throughout the lifetime

Mohammadamin Parsaei^a, Hossein Sanjari Moghaddam^{b,*},
 Mohammad Hadi Aarabi^c

^a School of Medicine, Tehran University of Medical Science, Tehran, Iran

^b Psychiatry and Psychology Research Center, Roozbeh Hospital, Tehran University of Medical Sciences, Tehran, Iran

^c Department of Neuroscience (DNS), Padova Neuroscience Center, University of Padova, Padua, Italy

The neuroanatomical characteristics of the brain exhibit variations between females and males, encompassing both healthy and pathological conditions [1]. Bethlehem et al. (2022) recently developed a human brain chart based on Magnetic Resonance Imaging (MRI) data from over 100,000 participants ranging in age from 115 days post-conception to 100 years [2]. This study discovered that males and females have significantly different brain tissue volumes throughout their lives and that these differences can also be seen in the brain growth patterns of people with psychiatric and neurologic conditions. Based on this growth chart trajectory, males have larger brain tissue volumes and more significant variance across MRI phenotypes, compared to females. It is critical to understand the effects of biological sex on brain development as it can significantly affect the physical and mental health of different psychiatric and neurologic patients.

Numerous studies have been conducted to compare brain volumes between healthy males and females. These studies have consistently found that males tend to have larger Total Cerebral Volume (TCV), total Grey Matter Volume (GMV), subcortical Grey Matter Volume (sGMV), total White Matter Volume (WMV), and Cerebrospinal Fluid Volume (CSFV). On the other hand, females generally demonstrate a higher mean Cortical Thickness (CTh) [3]. Abe et al. (2010) examined a population of individuals ranging in age from 21 to 71, revealing that males possess larger total GMV and WMV [4]. Additionally, males exhibited higher rates of decline in GMV and WMV with aging compared to females. This suggests that males and females have distinct brain structures, and the patterns of age-related brain changes are also different.

The precise etiology behind the lifelong disparities in brain structure between males and females remains incompletely understood. Emerging evidence suggests that these variations may have developmental origins. Certain investigations have identified different phenotypes in male and female brain cells, indicating a critical role of sex chromosomes in the emergence of sex-based brain differences. For instance, in vitro research has revealed that embryonal brain cells with XX and XY chromosomal configurations exhibit divergent behaviors [5]. XY cell cultures demonstrate a propensity to yield a higher number of dopamine neurons compared to their XX counterparts [5]. This emphasizes the genetic origin of the sex differences in brain structure. Consistent with this perspective, male infants exhibit larger total GMV and WMV compared to their female counterparts [6]. Moreover, these discrepancies persist into puberty, with males exhibiting greater total GMV, WMV, and TCV [7]. These findings underscore the significance of sex chromosomes in the development and perpetuation of brain differences between males and females.

While numerous studies have provided substantial evidence supporting the genetic origin of brain differences between males and females, a considerable body of research has demonstrated significant effects of sex hormones on brain development. Previous research revealed that testosterone acts as a human fetal brain programming system, leading to fetal dimorphism in brain structure and function

* Corresponding author at: Psychiatric Research Center, Roozbeh Psychiatric Hospital, Tehran University of Medical Sciences, South Kargar Street, Tehran 13337, Iran.

E-mail address: g.h.sanjarimoghaddam@gmail.com (H. Sanjari Moghaddam).

<https://doi.org/10.1016/j.nbas.2023.100098>

Received 18 April 2023; Received in revised form 8 September 2023; Accepted 19 September 2023

Available online 25 September 2023

2589-9589/© 2023 Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

[8,9]. Also, noteworthy sex-by-age interaction between testosterone levels and mean CTh across various brain regions was identified [10]. Furthermore, women with complete androgen insensitivity syndrome, characterized by the absence of androgen action despite a 46XY karyotype, exhibit comparable white matter structure to typical females [11]. This highlights the more prominent influence of sex hormones compared to genetics on brain development.

According to Bethlehem et al. (2022) growth charts for total GMV, WMV, sGMV, and CSFV, the disparities between males and females exhibit the smallest magnitude during the neonatal stage [2]. However, these differences tend to increase notably during post-pubertal periods and subsequently decrease during advanced stages of life, particularly in the postmenopausal periods for females. These observations suggest that although sexual dimorphism is evident at birth, the magnitude of these differences undergo dynamic changes throughout the lifespan, likely influenced by the impact of sex hormones.

Hagemann et al. (2011) revealed that GMV and CSFV undergo alterations throughout the menstrual cycle in women, highlighting the dynamic nature of these brain measures in relation to hormonal fluctuations [12]. Also, postmenopausal women experience a steeper decline in TCV and GMV compared to pre/perimenopausal individuals, implying that sex hormones may have a direct effect on brain growth patterns [13]. These findings collectively contribute to our understanding of the role of sex hormones in brain development, complementing the existing evidence on the developmental origin of brain differences between males and females.

Bethlehem et al. (2022) also investigated the sex differences in the brain changes observed in patients with psychiatric and neurologic disorders [2]. Their study reveals that sex differences are apparent in Anxiety Disorders (ANX) and Major Depressive Disorders (MDD), with males demonstrating a significant increase in global measures such as total GMV, sGMV, WMV, and TCV, while females tend to display non-significant decline in these measures [2]. This study marked the first exploration of sex differences in alterations in total GMV, sGMV, WMV, and TCV in ANX and MDD patients, aligning with prior research showing structural brain differences in specific regions between males and females. den Braber et al. (2013) found that male subjects with obsessive-compulsive disorders exhibited larger GMV in the left middle temporal gyrus, while females displayed larger GMV in the right precuneus [14]. Also, Yang et al. (2017) explored patients with MDD and discovered that males exhibited decreased GMV in the bilateral temporal gyri, whereas females demonstrated decreased GMV in the left lingual gyrus [15]. This suggests that different brain regions may contribute to the development of ANX and MDD in each biological sex, potentially influenced by sex hormones affecting brain structures and functions in these patients.

Based on the Bethlehem et al. (2022) article, male patients with Autism Spectrum Disorders (ASD) exhibit a more pronounced decline in WMV and a greater increase in mean CTh, whereas females demonstrate a more notable increase in CSFV [2]. Interestingly, total GMV and TCV experience a significant decrease in males, while females show a non-significant increase. Conversely, sGMV increases in males but significantly decreases in females [2]. Nevertheless, a systematic review presents contradictory evidence, suggesting that the age-related GMV decline in male ASD patients is less pronounced [16]. This review reveals significant sex differences at different life stages. Specifically, in youth, males have higher CTh in the parahippocampus, whereas in adults, females display higher CTh [16]. Additionally, Zhang et al. (2018) reported a significant sex-age interaction for total WMV in ASD patients, with males exhibiting greater values, especially in adulthood [17]. However, this interaction was not evident for total GMV. In summary, these findings emphasize sex-related discrepancies in lifelong brain alterations among ASD patients, though precise distinctions in each brain region remain a subject of debate.

Regarding patients with Schizophrenia (SCZ), females tend to show a more prominent decline in total GMV, WMV, sGMV, and TCV, while males demonstrate a steeper CTh decline [2]. Research on adolescents and adults with SCZ highlights distinct GMV patterns between sexes [18,19]. However, a study on children with SCZ found no significant sex differences in GMV across brain regions, suggesting that sex hormones, rather than genetics, likely underlie these variations in SCZ patients' brain regions [20].

In the realm of neurological disorders, sex-related patterns in brain metrics in Alzheimer's Disease (AD) patients resembled those seen in SCZ patients, according to Bethlehem et al. [2]. Among individuals with Mild Cognitive Impairment (MCI), both males and females exhibited brain metric alterations similar to AD, although males showed a non-significant increase in total WMV, while females displayed a non-significant decline [2]. However, some studies did not report significant sex-age interactions in GMV, sGMV, and CTh alterations among AD patients [21–23]. Sangha et al. (2021) found that in AD and MCI patients, CTh was thinner in elderly males, but these changes paralleled those observed in healthy controls, suggesting they may reflect broader population alterations rather than disorder-specific effects [24].

Reviewing the existing literature emphasizes the significance of comprehending lifelong brain differences between males and females. These disparities offer critical insights into brain development, aging, and neuropsychiatric disorders. Our discussion highlighted significant variations in brain structures among patients with psychiatric disorders across their lifespans, distinct from sex-specific patterns in healthy individuals. This implies a pivotal role for the sex hormones in psychiatric disorders. Conversely, in neurologic disorder patients like AD and MCI, brain structure sex differences largely mirrored those in the general population, suggesting a diminished impact of sex hormones on these conditions. Some studies have unveiled sex-specific genetic contributors to AD, potentially explaining distinctions between biological sexes [25]. Future research can further illuminate the interplay of sex hormones and genetics in neurologic disorder development, shedding light on brain changes in these patients compared to healthy controls.

Various studies have investigated the effects of sex on brain structure in healthy individuals, while there is a lack of research on individuals with neuropsychiatric disorders in this regard. It is important for future research to focus on the impact of sex on brain structure, particularly in the context of psychiatric and neurological conditions. This will help to further our understanding of these disorders and potentially lead to more targeted and effective treatments.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

- [1] McCarthy MM, Arnold AP, Ball GF, Blaustein JD, De Vries GJ. Sex differences in the brain: the not so inconvenient truth. *The Journal of Neuroscience* 2012;32(7):2241–7.
- [2] Bethlehem RAI, Seidlitz J, White SR, Vogel JW, Anderson KM, Adamson C, et al. Brain charts for the human lifespan. *Nature* 2022;604(7906):525–33.
- [3] Barnes J, et al. Head size, age and gender adjustment in MRI studies: a necessary nuisance? *NeuroImage* 2010;53(4):1244–55.
- [4] Abe O, Yamasue H, Yamada H, Masutani Y, Kabasawa H, Sasaki H, et al. Sex dimorphism in gray/white matter volume and diffusion tensor during normal aging. *NMR in Biomedicine* 2010;23(5):446–58.
- [5] Arnold AP. Sex chromosomes and brain gender. *Nature Reviews. Neuroscience* 2004;5(9):701–8.
- [6] Gilmore JH, Lin W, Prastawa MW, Looney CB, Vetsa YSK, Knickmeyer RC, et al. Regional gray matter growth, sexual dimorphism, and cerebral asymmetry in the neonatal brain. *The Journal of Neuroscience* 2007;27(6):1255–60.
- [7] Group, B.D.C., Total and regional brain volumes in a population-based normative sample from 4 to 18 years: the NIH MRI Study of Normal Brain Development. *Cerebral Cortex*, 2012. 22(1): p. 1-12.
- [8] Lombardo MV, Ashwin E, Auyeung B, Chakrabarti B, Taylor K, Hackett G, et al. Fetal testosterone influences sexually dimorphic gray matter in the human brain. *The Journal of Neuroscience* 2012;32(2):674–80.
- [9] Lombardo MV, Ashwin E, Auyeung B, Chakrabarti B, Lai M-C, Taylor K, et al. Fetal programming effects of testosterone on the reward system and behavioral approach tendencies in humans. *Biological Psychiatry* 2012;72(10):839–47.
- [10] Nguyen T-V, McCracken J, Ducharme S, Botteron KN, Mahabir M, Johnson W, et al. Testosterone-related cortical maturation across childhood and adolescence. *Cerebral Cortex* 2013;23(6):1424–32.
- [11] van Hemmen J, et al. Sex Differences in White Matter Microstructure in the Human Brain Predominantly Reflect Differences in Sex Hormone Exposure. *Cerebral Cortex* 2017;27(5):2994–3001.
- [12] Hagemann G, Ugur T, Schleussner E, Mentzel H-J, Fitzek C, Witte OW, et al. Changes in brain size during the menstrual cycle. *PLoS One* 2011;6(2):e14655.
- [13] Than S, Moran C, Beare R, Vincent AJ, Collyer TA, Wang W, et al. Interactions Between Age, Sex, Menopause, and Brain Structure at Midlife: A UK Biobank Study. *The Journal of Clinical Endocrinology and Metabolism* 2021;106(2):410–20.
- [14] den Braber A, de Geus EJC, Boomsma DI, van 't Ent D. Obsessive-compulsive symptoms and related sex differences in brain structure: an MRI study in Dutch twins. *Twin Research and Human Genetics* 2013;16(2):516–24.
- [15] Yang X, Peng Z, Ma X, Meng Y, Li M, Zhang J, et al. Sex differences in the clinical characteristics and brain gray matter volume alterations in unmedicated patients with major depressive disorder. *Scientific Reports* 2017;7(1).
- [16] Walsh MJM, et al. Brain-based sex differences in autism spectrum disorder across the lifespan: A systematic review of structural MRI, fMRI, and DTI findings. *NeuroImage Clin* 2021;31:102719.
- [17] Zhang W, Groen W, Mennes M, Greven C, Buitelaar J, Rommelse N. Revisiting subcortical brain volume correlates of autism in the ABIDE dataset: effects of age and sex. *Psychological Medicine* 2018;48(4):654–68.
- [18] Rapado-Castro M, Bartholomeusz CF, Castro-Fornieles J, González-Pinto A, Otero S, Baeza I, et al. Gender effects on brain changes in early-onset psychosis. *European Child & Adolescent Psychiatry* 2015;24(10):1193–205.
- [19] Mancini-Marie A, Yoon U, Jimenez J, Fahim C, Potvin S, Grant JA, et al. Sex, Age, Symptoms and Illness Duration and Their Relation with Gyrification Index in Schizophrenia. *Clinical Schizophrenia & Related Psychoses* 2018;12(2):57–68.
- [20] Weisinger B, Greenstein D, Mattai A, Clasen L, Lalonde F, Feldman S, et al. Lack of gender influence on cortical and subcortical gray matter development in childhood-onset schizophrenia. *Schizophrenia Bulletin* 2013;39(1):52–8.
- [21] Lee J, Cho H, Jeon S, Kim HJ, Kim YJ, Lee J, et al. Sex-Related Reserve Hypothesis in Alzheimer's Disease: Changes in Cortical Thickness with a Five-Year Longitudinal Follow-Up. *Journal of Alzheimer's Disease* 2018;65(2):641–9.
- [22] Fjell AM, Westlye LT, Amlien I, Espeseth T, Reinvang I, Raz N, et al. Minute effects of sex on the aging brain: a multisample magnetic resonance imaging study of healthy aging and Alzheimer's disease. *The Journal of Neuroscience* 2009;29(27):8774–83.
- [23] Salat DH, Stangl PA, Kaye JA, Janowsky JS. Sex differences in prefrontal volume with aging and Alzheimer's disease. *Neurobiology of Aging* 1999;20(6):591–6.
- [24] Sangha O, Ma Da, Popuri K, Stocks J, Wang L, Beg MF. Structural volume and cortical thickness differences between males and females in cognitively normal, cognitively impaired and Alzheimer's dementia population. *Neurobiology of Aging* 2021;106:1–11.
- [25] Dumitrescu L, Mayeda ER, Sharman K, Moore AM, Hohman TJ. Sex Differences in the Genetic Architecture of Alzheimer's Disease. *Curr Genet Med Rep* 2019;7(1):13–21.