

Please cite the Published Version

Sarawi, Wedad S, Alhusaini, Ahlam M, Fadda, Laila M, Alomar, Hatun A, Albaker, Awatif B, Alghibiwi, Hanan K, Aljrboa, Amjad S, Alotaibi, Areej M, Hasan, Iman H and Mahmoud, Ayman M (2022) Nano-curcumin prevents copper reproductive toxicity by attenuating oxidative stress and inflammation and improving Nrf2/HO-1 signaling and pituitary-gonadal axis in male rats. *Toxics*, 10 (7). 356 ISSN 2305-6304

DOI: <https://doi.org/10.3390/toxics10070356>

Publisher: MDPI AG

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/633044/>

Usage rights:  [Creative Commons: Attribution 4.0](https://creativecommons.org/licenses/by/4.0/)

Additional Information: This is an open access article which originally appeared in *Toxics*, published by MDPI





Data Access Statement: Data analyzed or generated during this study are included in this manuscript.

Enquiries:

If you have questions about this document, contact rsl@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Article

Nano-Curcumin Prevents Copper Reproductive Toxicity by Attenuating Oxidative Stress and Inflammation and Improving Nrf2/HO-1 Signaling and Pituitary-Gonadal Axis in Male Rats

Wedad S. Sarawi ^{1,*}, Ahlam M. Alhusaini ¹, Laila M. Fadda ¹, Hatun A. Alomar ¹, Awatif B. Albaker ¹, Hanan K. Alghibiwi ¹, Amjad S. Aljrboa ¹, Areej M. Alotaibi ^{1,2}, Iman H. Hasan ¹ and Ayman M. Mahmoud ^{3,4,*}

- ¹ Pharmacology and Toxicology Department, Faculty of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia; aelhusaini@ksu.edu.sa (A.M.A.); lfadda@ksu.edu.sa (L.M.F.); hetalomar@ksu.edu.sa (H.A.A.); abaker@ksu.edu.sa (A.B.A.); halghibiwi@ksu.edu.sa (H.K.A.); 441203053@student.ksu.edu.sa (A.S.A.); 442204129@student.ksu.edu.sa (A.M.A.); ihasan@ksu.edu.sa (I.H.H.)
- ² Department of Pharmacology and Toxicology, College of Pharmacy, Umm Al-Qura University, Mecca 21955, Saudi Arabia
- ³ Physiology Division, Zoology Department, Faculty of Science, Beni-Suef University, Beni-Suef 62514, Egypt
- ⁴ Department of Life Sciences, Faculty of Science & Engineering, Manchester Metropolitan University, Manchester M1 5GD, UK
- * Correspondence: wsarawi@ksu.edu.sa (W.S.S.); a.mahmoud@mmu.ac.uk or ayman.mahmoud@science.bsu.edu.eg (A.M.M.)



Citation: Sarawi, W.S.; Alhusaini, A.M.; Fadda, L.M.; Alomar, H.A.; Albaker, A.B.; Alghibiwi, H.K.; Aljrboa, A.S.; Alotaibi, A.M.; Hasan, I.H.; Mahmoud, A.M. Nano-Curcumin Prevents Copper Reproductive Toxicity by Attenuating Oxidative Stress and Inflammation and Improving Nrf2/HO-1 Signaling and Pituitary-Gonadal Axis in Male Rats. *Toxics* **2022**, *10*, 356. <https://doi.org/10.3390/toxics10070356>

Received: 10 June 2022

Accepted: 28 June 2022

Published: 30 June 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Copper is essential for several cellular processes and is an important catalytic factor for many proteins. However, excess copper can provoke oxidative stress and reproductive toxicity. This study evaluated the effect of liposomal nano-curcumin (N-CUR) and CUR on testicular oxidative injury, inflammation, and apoptosis, and altered steroidogenesis and Nrf2/HO-1 signaling induced by copper sulfate (CuSO₄). Rats received CuSO₄ and N-CUR or CUR via oral gavage for 7 days. CuSO₄ induced histopathological changes and altered pituitary-gonadal axis manifested by decreased serum gonadotropins and testosterone. Testicular steroidogenesis genes (StAR, 3β-HSD, CYP17A1, and 17β-HSD) and androgen receptor (AR) were downregulated in rats that received CuSO₄. N-CUR and CUR prevented testicular tissue injury, increased circulating FSH, LH, and testosterone, and upregulated testicular steroidogenesis genes and AR. Additionally, N-CUR and CUR decreased testicular MDA, NO, NF-κB, iNOS, TNF-α, Bax, and caspase-3 while enhanced Bcl-2, Nrf2, and the antioxidants GSH, HO-1, SOD, and catalase. In conclusion, N-CUR and CUR prevented CuSO₄-induced reproductive toxicity in male rats by suppressing oxidative injury and inflammatory response and boosting steroidogenesis, sex hormones, and Nrf2/HO-1 signaling. N-CUR was more effective in ameliorating tissue injury, oxidative stress, inflammation, and apoptosis and enhancing steroidogenesis and Nrf2/HO-1 than the native form.

Keywords: curcumin; reproductive toxicity; copper sulfate; steroidogenesis; oxidative stress

1. Introduction

Copper (Cu) is an essential element for maintaining several physiological functions such as energy metabolism, protein homeostasis, blood clotting, and the synthesis of neurotransmitters [1,2]. The ion of this redox-active metal is essential for many proteins and enzymes, including ceruloplasmin and superoxide dismutase (SOD), and has a key role in the development of the reproductive system, skin, hair, and bone [3]. The body Cu is maintained within normal levels through the regulation of its absorption and excretion via precise regulatory mechanisms [4]. Cu as a metal is widely used in several industries, including electronics, building materials, wood preservatives, and intrauterine contraceptive devices [5,6]. Owing to its widespread use, exposure to Cu can occur frequently and lead to toxic effects upon exceeding physiological levels. Cu could be absorbed through the

skin, lungs, and intestine resulting in toxic effects in different organs, including the liver, kidney, and nervous system [7,8]. Hepatotoxicity, rhabdomyolysis, arrhythmia, seizures, kidney injury, and neurological disorders are excess Cu-linked clinical manifestations [9,10]. Cu was detected in several tissues of aquatic animals and hence has been graded as an environmental pollutant [11,12]. Copper sulfate (CuSO_4) is one of the Cu compounds that possess fungicidal and bactericidal properties. CuSO_4 is used to minimize the contamination risk in tissue culture and is employed in agriculture as a pesticide to repel pests and improve crop yield. However, its intentional or accidental intoxication could result in multiorgan toxicity that might be fatal [13]. Accordingly, exposure of rats to CuSO_4 resulted in increased Cu levels in the liver and kidney and subsequently impairment and damage of these organs [14]. Recent work from our lab has demonstrated neurotoxicity and cardiac injury in rats exposed to CuSO_4 [15,16].

Reproductive toxicity is one of the manifestations of Cu as reported in several *in vivo* and *in vitro* experiments [17–20]. The effects of Cu on male fertility include significant decrease in the number, motility, and viability of spermatozoa [21]. The intake of Cu, even at low doses, induced adverse effects on the morphology of testis in male mice [22]. Reduced levels of circulating testosterone, follicle-stimulating hormone (FSH), and luteinizing hormone (LH) have been demonstrated in immature male rats exposed to Cu for 26 days [23]. Although the effects of Cu on male fertility are not fully understood, excessive reactive oxygen species (ROS) and oxidative stress could play a significant role [19,24]. Given its rich content of highly unsaturated fatty acids, the testis is vulnerable to oxidative injury [25]. Accordingly, studies have demonstrated that Cu upregulates genes of the oxidative phosphorylation pathway and the generation of ROS and decreases antioxidant defenses and spermatogenesis [17,18]. Therefore, mitigating oxidative stress might attenuate Cu-induced testicular injury.

Curcumin (CUR), the principal curcuminoid of turmeric, is a non-polar polyphenol with potent antioxidant and anti-inflammatory properties [26,27]. CUR is used in dietary supplements, cosmetics, flavored beverages and other applications [28]. It has been virtually studied for several diseases and showed antimicrobial [29], anti-diabetic [30], nephroprotective [31], hepatoprotective [32–36], and antitumor efficacies [37]. It is a bis- α,β -unsaturated β -diketone with aromatic rings functionalized with methoxy and hydroxy groups. The presence of conjugated double bonds in CUR permits effective electron donation and counteracting ROS formation [38]. Therefore, CUR can effectively suppress ROS generation and prevent oxidative injury. CUR has been evidenced to act in a multi-targeted fashion through its ability to target bioactive proteins in various signaling pathways or its epigenetic modulating power [39,40]. The protective efficacy of CUR against testicular injury induced by titanium dioxide [41], lead acetate [42], and doxorubicin (DOX) [43] has been reported; however, its protective efficacy against testicular injury caused by CuSO_4 has not been investigated yet. In addition, the rapid metabolism and poor systemic bioavailability of CUR are the main drawbacks that can limit its therapeutic applications [44]. Nanoformulations greatly improved the properties of CUR by amending its cellular uptake, permeability, and plasma bioavailability [45]. Liposomal CUR nanoformulation exhibited a better bioavailability in different diseases, highlighting that liposomes are optimal delivery vehicles [46]. Liposomes are spherical vesicles constituted by phospholipid bilayers where the polar groups are oriented to the outer and inner aqueous phases. Liposomes have a high propensity to carry both hydrophobic and hydrophilic drugs and improve their accumulation in tissues and therapeutic actions and decrease side effects [46,47]. This study explored the efficacy of liposomal nano-CUR (N-CUR) against oxidative stress, inflammation, and testicular injury induced by CuSO_4 , pointing to the involvement of nuclear factor erythroid 2-related factor 2 (Nrf2)/heme oxygenase-1 (HO-1) signaling. Nrf2 is a cytoprotective factor that controls the transcription of antioxidant defenses and prevents oxidative damage and inflammation [48].

2. Materials and Methods

2.1. Chemicals and Reagents

Liposomal N-CUR and deferoxamine (DFO) were supplied by Lipolife (Essex, UK) and Novartis Pharma AG (Rotkreuz, Switzerland), respectively. CUR, CuSO₄, thiobarbituric acid, sodium dodecyl sulfate (SDS), primers, Griess reagent, pyrogallol, agarose, reduced glutathione (GSH), hematoxylin and eosin (H&E), bisacrylamide, hydrogen peroxide (H₂O₂) and carboxymethylcellulose (CMC) were purchased from Sigma (St. Louis, MO, USA). Protein assay kit was obtained from BioBasic (Markham, Canada), tumor necrosis factor (TNF)- α ELISA kit was supplied by R&D Systems (Minneapolis, MN, USA), and testosterone ELISA kit was purchased from Cusabio (Wuhan, China). FSH and LH ELISA kits and antibodies against Nrf2, HO-1, androgen receptor (AR), and β -actin were provided by Novus Biologicals (Centennial, CO, USA). Other chemicals were obtained from standard suppliers.

2.2. Experimental Animals and Treatments

Forty male Wistar rats (180–200 g) were supplied by the Animals Care Centre (King Saud University). The rats were housed in standard cages under a 12-h light/dark cycle and standard temperature and humidity. The animals had free access to food and water and were allocated into 5 groups ($n = 8$) after acclimatization for one week. Rats in group I served as control and received 1% CMC. Rats in groups II–V received 100 mg/kg CuSO₄ [14], and groups III, IV and V were supplemented with 23 mg/kg DFO [49], 80 mg/kg CUR [13,49], and 80 mg/kg N-CUR [13,49], respectively. All treatments were suspended in 1% CMC and given orally for 7 days. On day 8, the rats were anesthetized with ketamine/xylazine, and blood was collected via cardiac puncture for serum preparation. The rats were sacrificed and the testes were removed and washed in cold phosphate-buffered saline (PBS). Pieces from the testes were fixed in Bouin's solution while other samples were homogenized (10% w/v) in Tris-HCl buffer (pH 7.4). The homogenate was centrifuged, and the supernatant was collected for the determination of malondialdehyde (MDA), nitric oxide (NO), GSH, SOD, catalase (CAT), and TNF- α . Other samples were kept frozen at -80°C for RNA isolation and western blotting.

2.3. Assay of Testosterone, Gonadotropins, and TNF- α

Testosterone, FSH, and LH were assayed in the serum and TNF- α was determined in the testicular tissue supernatant using specific ELISA kits according to instructions of the suppliers.

2.4. Assay of MDA, NO and Antioxidants

MDA and NO were assayed in the testicular tissue supernatant following the methods of Ohkawa et al. [50] and Grisham et al. [51], respectively. GSH was determined according to the method of Ellman [52], and the activities of SOD and CAT were determined according to the methods described by Marklund and Marklund [53] and Cohen et al. [54], respectively.

2.5. Histopathological Examination

The testes samples were fixed in Bouin's solution for 24 h, and then processed for paraffin wax embedding. Five- μm sections were cut using a rotary microtome. The sections were stained with H&E for microscopic examination.

2.6. Gene Expression

Changes in the expression of nuclear factor-kappaB (NF- κB) p65, inducible NO synthase (iNOS), B cell lymphoma-2 (BCL-2), BCL-2-associated X protein (BAX), caspase-3, and steroidogenic acute regulatory protein (StAR), 3 β -Hydroxysteroid dehydrogenase (HSD), 17 β -HSD, and cytochrome P450 17A1 (CYP17A1) were evaluated using qRT-PCR [55]. TRIzol (ThermoFisher Scientific, Waltham, MA, USA) was used to isolate RNA which was treated with RNase-free DNase (Qiagen, Hilden, Germany) and quantified on

a nanodrop. RNA samples with A260/A280 nm ratio of ≥ 1.8 were selected for cDNA synthesis which was amplified using Maxima SYBR Green/ROX qPCR master mix (ThermoFisher Scientific, Waltham, MA, USA) and the primers in Table 1. The obtained data were analyzed using the $2^{-\Delta\Delta C_t}$ method [56] and normalized to β -actin.

Table 1. Primers used for qRT-PCR.

Gene	GenBank Accession Number	Primers (5'-3')	Amplicon Size (bp)
StAR	NM_031558.3	F: CCCAGATAGAGTTCGCCAGC R: TGGTGGGCAGTCCTAACAC	84
CYP17A1	XM_006231434.3	F: TCAGTGAAGTGTGACCTGGGA R: GTGGAGCGGAGCAACTTCAA	122
3 β -HSD	NM_001007719.3	F: TGTGCCAGCCTTCATCTAC R: CTCTCGGCCATCCTTTT	145
17 β -HSD	NM_054007.1	F: GACCGCCGATGAGTTTGT R: TTTGGGTGGTGCTGCTGT	140
NF- κ B p65	NM_199267.2	F: CCTCATCTTTCCCTCAGAGCC R: GGTCCCGTGTAGCCATTGAT	189
iNOS	NM_012611.3	F: ATTCCCAGCCCAACAACACA R: GCAGCTTGTCCAGGGATTCT	112
TNF- α	NM_012675.3	F: AAATGGGCTCCCTCTCATCAGTTC R: TCTGCTTGGTGGTTTGCTACGAC	111
Caspase-3	NM_012922.2	F: GGAGCTTGAACCGGAAGAA R: ACACAAGCCCATTCAGGGT	169
BAX	NM_017059.2	F: AGGACGCATCCACCAAGAAG R: CAGTTGAAGTTGCCGTCTGC	166
BCL-2	NM_016993.1	F: ACTCTTCAGGGATGGGGTGA R: TGACATCTCCCTGTTGACGC	94
β -actin	NM_031144.3	F: AGGAGTACGATGAGTCCGGC R: CGCAGCTCAGTAACAGTCCG	71

2.7. Western Blotting

To evaluate changes in the expression of Nrf2, HO-1, and proliferating cell nuclear antigen (PCNA), the tissue was homogenized in RIPA buffer, and the homogenate was centrifuged at 10,000 rpm. The concentration of protein was assayed in the supernatant using a Bradford protein assay kit. Fifty μ g protein was subjected to 10% SDS/PAGE and the separated protein bands were transferred to nitrocellulose membranes. After blocking in 5% milk in tris-buffered saline tween (TBST), the membranes were incubated overnight at 4 °C with anti-Nrf2, anti-HO-1, anti-AR, or anti- β -actin. The membranes were washed three times in TBST and probed with the secondary antibodies for 1 h at room temperature. After washing with TBST, the bands were developed with Clarity™ Western ECL Substrate from BIO-RAD (Hercules, CA, USA) and visualized using ImageQuant LAS 4000. The band intensity was quantified using ImageJ (version 1.32j, NIH, Bethesda, MD, USA).

2.8. Statistical Analysis

The results are expressed as mean \pm standard error of the mean (SEM). Statistical analysis and multiple comparisons were performed by one-way ANOVA and Tukey's post-hoc test using GraphPad Prism 8. A p value < 0.05 was considered significant.

3. Results

3.1. N-CUR and CUR Enhance the Pituitary-Gonadal Axis and Prevent Testicular Injury in CuSO₄-Administered Rats

The circulating levels of LH, FSH, and testosterone were assayed to evaluate the impact of CuSO₄ on the pituitary-gonadal axis and the protective effect of N-CUR and CUR (Figure 1). CuSO₄-treated rats exhibited a decrease in LH (Figure 1A), FSH (Figure 1B), and testosterone (Figure 1C) as compared to the control ($p < 0.001$). DFO, CUR, or N-CUR alleviated serum LH, FSH, and testosterone in CuSO₄-administered rats. CUR increased

serum FSH significantly ($p < 0.05$) when compared to DFO, while N-CUR alleviated both LH and FSH levels.




Figure 1. N-CUR and CUR enhance the pituitary-gonadal axis in Cu-intoxicated rats. CUR, N-CUR, and DFO increased serum (A) LH, (B) FSH, and (C) testosterone in rats that received CuSO₄. Data are mean \pm SEM, ($n = 8$). * $p < 0.05$ and *** $p < 0.001$ versus Control. # $p < 0.05$, ## $p < 0.01$, and ### $p < 0.001$ versus CuSO₄.

Examination of the testis of control rats revealed normal seminiferous tubules with normal spermatogonia, spermatocytes, spermatids and a large number of spermatozoa (Figure 2A,B). Oral supplementation of CuSO₄ induced degeneration of the seminiferous tubules and loss of spermatogenic series and complete lack of mature spermatozoa as shown in Figure 2C,D. Concurrent treatment of the rats with DFO (Figure 2E,F), CUR (Figure 2G,H), or N-CUR (Figure 2I,J) prevented CuSO₄-induced testicular tissue injury and the seminiferous tubules appeared to possess the normal histological architecture with spermatogonia, primary spermatocytes, spermatids, and spermatozoa.

3.2. N-CUR and CUR Upregulate Steroidogenesis and AR in CuSO₄-Administered Rats

The effect of CuSO₄ on steroidogenesis and the ameliorative potential of N-CUR and CUR were evaluated by determining the mRNA abundance of StAR (Figure 3A), 3 β -HSD (Figure 3B), CYP17A1 (Figure 3C), and 17 β -HSD (Figure 3D), and the protein expression of AR (Figure 3E). CuSO₄ downregulated the mRNA abundance of StAR, 3 β -HSD, CYP17A1, and 17 β -HSD, as well as AR protein in the testis of rats ($p < 0.001$). Both forms of CUR as well as DFO upregulated the expression of steroidogenesis genes and AR ($p < 0.001$). N-CUR upregulated StAR, 3 β -HSD, CYP17A1, 17 β -HSD, and AR significantly when compared to either DFO or CUR (Figure 3).




Figure 2. N-CUR and CUR prevent Cu-induced testicular tissue injury. Photomicrographs of H&E-stained sections in the testis of (A,B) control rats showing normal histological architecture, (C,D) Cu-intoxicated rats showing degenerative changes and loss of spermatogenic series, and spermatozoa, and (E–J) Cu-intoxicated rats treated with DFO (E,F), CUR (G,H), and N-CUR (I,J) showing normal seminiferous tubules with spermatogonia, primary spermatocytes, spermatids, and spermatozoa. ($\times 200$: A,C,G,E,I. $\times 400$: B,D,F,H,J).




Figure 3. N-CUR and CUR upregulate steroidogenesis and AR in Cu-intoxicated rats. Treatment with DFO, CUR, and N-CUR increased (A) StAR, (B) 3β-HSD, (C) CYP17A1, and (D) 17β-HSD mRNA abundance, and (E) AR protein expression. Data are mean ± SEM, (n = 8). *** p < 0.001 versus Control and ### p < 0.001 versus CuSO₄.

3.3. N-CUR and CUR Attenuate Testicular Oxidative Stress in CuSO₄-Administered Rats

Both MDA (Figure 4A) and NO (Figure 4B) levels showed a significant elevation in the testis of CuSO₄-administered rats (p < 0.001). In contrast, testicular GSH (Figure 4C), SOD (Figure 4D), and CAT (Figure 4E) were significantly decreased following the administration of CuSO₄ (p < 0.001). All treatments markedly decreased testicular MDA and NO, and increased GSH, SOD and CAT (p < 0.001). N-CUR was more effective in ameliorating oxidants, GSH and CAT than DFO and CUR, and increasing SOD activity than CUR.




Figure 4. N-CUR and CUR attenuate testicular oxidative stress in Cu-intoxicated rats. Treatment with DFO, CUR, and N-CUR decreased (A) MDA and (B) NO and increased (C) GSH, (D) SOD, and (E) CAT. Data are mean \pm SEM, ($n = 8$). ** $p < 0.01$ and *** $p < 0.001$ versus Control, and ### $p < 0.001$ versus CuSO₄.

3.4. N-CUR and CUR Upregulate Nrf2/HO-1 Signaling in CuSO₄-Administered Rats

Nrf2 and HO-1 in the testis of rats were determined using western blotting that revealed significant downregulation of both proteins in CuSO₄-administered rats ($p < 0.001$; Figure 5A–C). Treatment of these rats with CUR and N-CUR markedly increased the expression of testicular Nrf2 and HO-1. Of note, N-CUR upregulated Nrf2 and HO-1 significantly ($p < 0.001$) when compared with either DFO or CUR.




Figure 5. N-CUR and CUR upregulate Nrf2/HO-1 signaling in Cu-intoxicated rats. (A–C) Treatment with DFO, CUR, and N-CUR increased testicular Nrf2 and HO-1 protein levels. Data are mean ± SEM, (n = 8). ** p < 0.01 and *** p < 0.001 versus Control, and ### p < 0.001 versus CuSO₄.

3.5. N-CUR and CUR Mitigate Testicular Inflammation in CuSO₄-Administered Rats

The inflammatory response provoked in the testis was evaluated through the assessment of NF-κB p65, iNOS, and TNF-α expression as depicted in Figure 6. CuSO₄-administered rats exhibited a significant upregulation of NF-κB p65 (Figure 6A) and iNOS (Figure 6B) mRNA abundance and TNF-α protein (Figure 6C) (p < 0.001). CUR, DFO, and N-CUR prevented the increase in NF-κB p65, iNOS, and TNF-α in the testis of CuSO₄-administered rats (p < 0.001). Notably, N-CUR was more effective in decreasing NF-κB p65 than CUR and iNOS, and TNF-α than DFO and CUR.




Figure 6. N-CUR and CUR mitigate testicular inflammation in Cu-intoxicated rats. CUR, N-CUR, and DFO decreased (A) NF-κB p65 and (B) iNOS mRNA abundance, and (C) TNF-α protein. Data are mean ± SEM, (n = 8). * p < 0.05, ** p < 0.01 and *** p < 0.001 versus Control, and ### p < 0.001 versus CuSO₄.

3.6. N-CUR and CUR Prevent Testicular Apoptosis in CuSO₄-Administered Rats

To evaluate the protective effect of N-CUR and CUR against Cu-induced testicular apoptosis, we assayed the mRNA abundance of Bax, caspase-3, and Bcl-2 along with the protein expression levels of PCNA. Administration of CuSO₄ provoked upregulation of testicular Bax (Figure 7A) and caspase-3 (Figure 7B) and downregulated Bcl2 (Figure 7C) and PCNA (Figure 7D) as compared to the control rats ($p < 0.001$). All treatment agents ameliorated the expression of all assayed markers significantly. In comparison with either CUR or DFO, N-CUR was more effective in ameliorating the expression of apoptosis markers and PCNA.




Figure 7. N-CUR and CUR prevent testicular apoptosis in Cu-intoxicated rats. CUR, N-CUR, and DFO downregulated (A) Bax and (B) caspase-3, and upregulated (C) Bcl-2 and (D) PCNA in the testis of rats. Data are mean \pm SEM, ($n = 8$). * $p < 0.05$, ** $p < 0.01$ and *** $p < 0.001$ versus Control, and ### $p < 0.001$ versus CuSO₄.

4. Discussion

Copper is an essential cofactor for antioxidant enzymes, immune function and a plethora of vital cellular processes [2]. However, its extensive usage in many enterprises, such as, electronics, pesticides, building and transportation materials, water pipes and others incurred a notable eco-environmental contamination by polluting land and water

resources [5,6,57]. Exposure to Cu compounds has been associated with adverse effects, including testicular oxidative stress, and apoptosis, along with declined sperm quality [19,24]. CUR has shown potent antioxidant properties and protected the testis against injury induced by various agents [26,27,41–43]. Herein, we evaluated the role of CUR and N-CUR in attenuating CuSO₄-induced testicular injury in rats, pointing to changes in steroidogenesis, Nrf2/HO-1 signaling, and inflammatory and apoptosis markers.

Exposure to CuSO₄ caused testicular tissue damage and disrupted hormones of the pituitary-gonadal axis. CuSO₄ ingestion resulted in degenerative changes, germ cells depletion and significantly reduced the number of spermatozoa in the seminiferous tubules and decreased the circulating levels of gonadotropins and testosterone. Accordingly, recent studies demonstrated declined serum LH and FSH in pigs [24] and testosterone in rats [19] exposed to Cu compounds, including CuSO₄. FSH and LH are important for the proliferation and normal function of Sertoli cells and Leydig cells, respectively. FSH is necessary for the proliferation, maturation, and function of Sertoli cells which generate signals for the initiation and maintenance of spermatogonia [58]. Therefore, normal secretion of FSH is essential for maintaining spermatogenesis [58]. LH induces the proliferation of Leydig cells and stimulates the secretion of testosterone which maintain the structural and functional integrity of reproductive organs and male accessory glands [58]. The observed decrease in serum testosterone in rats exposed to CuSO₄ could be directly ascribed to insufficient LH secretion and/or reduced responsiveness of Leydig cells to LH.

Moreover, our findings introduced new information on the effect of Cu exposure on steroidogenesis. Rats that received CuSO₄ exhibited a significant decrease in testicular StAR, 3 β -HSD, CYP17A1 and 17 β -HSD mRNA abundance. These enzymes are essential for maintaining steroidogenesis and male fertility. StAR controls the rate-limiting step of testosterone biosynthesis by promoting the transport of cholesterol into the mitochondria [59], and its inhibition can stop the steroidogenic machinery in Leydig cells [60]. Within the mitochondria, CYP11A1 converts cholesterol to pregnenolone which is then hydroxylated by CYP17A1. 3 β -HSD and 17 β -HSD catalyze the subsequent steps of testosterone biosynthesis [61]. The reduced levels of FSH, LH, and testosterone along with the downregulated steroidogenic genes confirmed the endocrine and reproductive toxic effect of CuSO₄.

CUR and N-CUR prevented testicular injury, ameliorated sex hormone levels, and enhanced steroidogenesis in rats that received Cu. In addition, both CUR and N-CUR upregulated the expression of AR in the testis of rats. ARs, ligand-activated nuclear receptors, are highly expressed in the testes, particularly Sertoli cells, and play a central role in maintaining spermatogenesis. Upon binding of testosterone or its metabolite, AR dissociates from heat-shock proteins, dimerizes, and binds to its response element in the nucleus to promote the expression of genes responsible for the maintenance, growth and maturation of male gametes [62]. The total or conditional knockout of AR caused severe defects in reproductive development and spermatogenesis in mice [63]. Although the effect of CuSO₄ on spermatogenesis has been previously reported [19], we introduced evidence that it downregulates AR in the testis and that CUR and its nanoform can increase testosterone biosynthesis and AR. The ability of CUR to prevent testicular damage caused by various toxicants, including titanium dioxide [41], lead [42], and DOX [43] has been previously demonstrated. In these studies, the protective effect of CUR prevented histopathological alterations, increased testosterone release and enhanced sperm parameters. Here, N-CUR showed a more potent protective effect when compared to the native form, an effect that is directly attributed to the improved properties by nanoformulation. It is noteworthy pointing out that the chelating agent DFO exerted a protective effect against CuSO₄-induced testicular toxicity. Hence, the chelating property of CUR might be involved in its protective effect against Cu toxicity. In support of this notion, the chelate complex of CUR with Cu(II) was studied by employing quantum chemical computations as reported by Balasubramanian [64]. The primary site for chelating Cu(II) is the β -diketone bridge of CUR and the chelation mechanism involves the loss of an enolic proton of CUR [64].

Oxidative stress has been suggested to play a key role in Cu toxicity [15,16], and some researchers have demonstrated that its testicular toxicity occurs via provoking oxidative damage [19,24]. Cu is able to cycle easily between unstable reduced and stable oxidized states and facilitate redox reactions. However, this renders it toxic because of the surplus release of hydroxyl radicals [65,66]. Also, Cu can upregulate genes of the oxidative phosphorylation pathway and enhance the generation of mitochondrial ROS, decrease antioxidants, and suppress spermatogenesis [17,18]. ROS are potent oxidizers that cause oxidative damage of lipids, DNA and proteins, and provoke cell death [66]. Accordingly, Cu increased testicular MDA and NO and suppressed the antioxidants GSH, SOD and CAT, demonstrating oxidative stress. The well-documented antioxidant and radical-scavenging properties of CUR [32,33,38,67] are involved in its protective effect against CuSO₄-induced testicular toxicity. The antioxidant potency of CUR was demonstrated in animal models of testicular toxicity induced by lead acetate [42], and DOX [43] where it decreased MDA and enhanced GSH and antioxidant enzymes. In this study, N-CUR decreased MDA and NO, and boosted antioxidant defenses in the testis of CuSO₄-administered rats significantly as compared to the native form, demonstrating its enhanced radical-scavenging activity.

Besides its radical scavenging properties, activation of Nrf2/HO-1 signaling might be involved in the protective effect of CUR and N-CUR against CuSO₄-induced testicular toxicity. The results revealed that exposure to Cu downregulated testicular Nrf2 and HO-1, an effect that was prevented in rats supplemented either CUR or N-CUR. Nrf2 is a redox-sensitive factor located in the cytoplasm sequestered by Keap-1. Upon activation by ROS or electrophiles, it dissociates from Keap-1 and translocates into the nucleus where it binds to ARE and stimulate the expression of detoxifying, cytoprotective and antioxidant genes, including HO-1 [48]. Activation of Nrf2/HO-1 signaling protected the testis against heavy metal-induced oxidative stress, inflammation, and apoptosis [55,68]. CUR upregulated Nrf2/HO-1 and prevented oxidative stress in the testis of cadmium-intoxicated mice [68]. The ability of CUR to upregulate Nrf2 signaling has been investigated in different in vitro and in vivo models (reviewed in [69]). In consistence with the results of antioxidant enzymes, N-CUR was more effective in upregulating testicular Nrf2/HO-1 signaling in CuSO₄-administered rats.

Inflammation and apoptosis are implicated in the toxic effects of Cu, including neurotoxicity [15], cardiotoxicity [16] and testicular toxicity [24]. Here, NF- κ B, iNOS, and TNF- α were significantly upregulated in the testes of rats that received CuSO₄, denoting an inflammatory response. The generation of ROS by Cu is the main culprit behind the developed inflammation. Excess ROS can activate many signaling molecules, including NF- κ B which subsequently elicits the transcription of iNOS and inflammatory cytokines [70]. Recently, we have reported the activation of TLR4/NF- κ B and MAPK signaling and the release of pro-inflammatory cytokines in the cardiac tissue of Cu-administered rodents [16]. In conjunction with inflammation, testicular apoptosis was observed in CuSO₄-administered rats. CuSO₄ increased Bax and caspase-3 and decreased the anti-apoptotic marker Bcl-2. Similar findings have been previously reported in the brain and heart of rats treated with CuSO₄ [15,16]. ROS and pro-inflammatory mediators can activate Bax which promotes the loss of mitochondrial membrane potential and subsequently cytochrome c is released to the cytosol where it activates the executioner caspase-3 [71]. Caspase-3 provokes DNA fragmentation, degradation of cytoskeletal proteins and further release of cytochrome c, leading to cell death [72]. Furthermore, CuSO₄ decreased the expression of PCNA in the testes of rats. PCNA is related to DNA synthesis, chromosome recombination and RNA transcription, and is involved in the proliferation and differentiation of spermatogonia [73]. The positive expression of PCNA is conducive to DNA replication, cell proliferation and smooth spermatogenesis [73]. CUR and N-CUR downregulated NF- κ B, iNOS, TNF- α , Bax and caspase-3, and increased Bcl-2 and PCNA in the testes of Cu-administered rats. These findings demonstrated the anti-inflammatory and anti-apoptosis properties of CUR and pinpointed the superior activity of N-CUR in attenuating inflammation and apoptosis. CUR prevented testicular inflammation and apoptosis as reported in previous studies.

For instance, CUR prevented apoptosis of testicular Leydig cells challenged with palmitic acid [74], and attenuated inflammation in the testes of methotrexate-intoxicated rats [75]. Similar to its antioxidant effect, the anti-inflammatory activity of CUR could be linked to the upregulation of Nrf2/HO-1 signaling. This notion is supported by the study of Boyanapalli et al. who reported that the anti-inflammatory activity of CUR was abolished in Nrf2^{-/-} mice and macrophages [76].

N-CUR showed a significant modulatory effect on LH, genes of steroidogenesis, AR, LPO, cellular antioxidants, Nrf2/HO-1 signaling, and inflammatory, proliferation and apoptosis markers when compared with CUR. This superior activity could be directly attributed to the improvement of its properties upon nanoformulation. When compared with the native form, N-CUR exhibited stronger antioxidant and anti-LPO efficacies than CUR in hepatoma cells lines [77], and attenuated inflammatory response in LPS-treated macrophages [78].

5. Conclusions

These results introduce compelling evidence for the deleterious effect of CuSO₄ on testicular tissue, pituitary-gonadal axis, and steroidogenesis and the protective efficacy of CUR and N-CUR. Treatment with N-CUR and the native form prevented Cu-induced testicular tissue damage, oxidative stress, inflammation, and apoptotic cell death, and ameliorated cell proliferation, circulating levels of sex hormones, and steroidogenesis in rats. These effects were associated with activation of Nrf2/HO-1 signaling and enhanced antioxidants. N-CUR showed stronger protection when compared with CUR, an effect that is ascribed to the improved properties. Therefore, N-CUR is a potent protective agent against testicular toxicity caused by CuSO₄, pending further investigations to explore other mechanisms.

Author Contributions: Conceptualization, W.S.S.; A.M.A. (Ahlam M. Alhusaini) and A.M.M.; methodology, W.S.S.; A.M.A. (Areej M. Alotaibi); L.M.F.; H.K.A.; H.A.A.; A.B.A.; A.S.A.; I.H.H. and A.M.M.; validation, W.S.S. and A.M.M.; formal analysis, W.S.S.; A.M.A. (Ahlam M. Alhusaini) and A.M.M.; investigation, W.S.S.; A.M.A.; H.K.A.; L.M.F.; H.A.A.; A.B.A.; I.H.H. and A.M.M.; resources, W.S.S.; A.M.A. (Ahlam M. Alhusaini); A.M.A. (Areej M. Alotaibi) and A.M.M.; data curation, W.S.S. and A.M.M.; writing—original draft preparation, A.M.M.; writing—review and editing, A.M.M.; visualization, W.S.S.; A.M.A. (Ahlam M. Alhusaini); I.H.H. and A.M.M.; supervision, A.M.A. (Ahlam M. Alhusaini) and A.M.M.; project administration, A.M.A. (Ahlam M. Alhusaini); I.H.H. and W.S.S.; funding acquisition, W.S.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the Deanship of Scientific Research at King Saud University, grant number RG-1441-546.

Institutional Review Board Statement: The experiment was conducted according to the guidelines of the National Institutes of Health (NIH publication No. 85–23, revised 2011) and was approved by the research ethics Committee at King Saud University (Ethical reference no. SE-19-129, approved on 20 November 2019).

Informed Consent Statement: Not applicable.

Data Availability Statement: Data analyzed or generated during this study are included in this manuscript.

Acknowledgments: The authors extend their appreciation to the Deanship of Scientific Research at King Saud University for funding this work through research group number RG-1441-546.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Scheiber, I.F.; Mercer, J.F.; Dringen, R. Metabolism and functions of copper in brain. *Prog. Neurobiol.* **2014**, *116*, 33–57. [[CrossRef](#)]
2. Uriu-Adams, J.Y.; Keen, C.L. Copper, oxidative stress, and human health. *Mol. Asp. Med.* **2005**, *26*, 268–298. [[CrossRef](#)]

3. Bost, M.; Houdart, S.; Oberli, M.; Kalonji, E.; Huneau, J.-F.; Margaritis, I. Dietary copper and human health: Current evidence and unresolved issues. *J. Trace Elem. Med. Biol. Organ Soc. Miner. Trace Elem. (GMS)* **2016**, *35*, 107–115. [[CrossRef](#)]
4. Denoyer, D.; Masaldan, S.; La Fontaine, S.; Cater, M.A. Targeting copper in cancer therapy: ‘Copper that cancer’. *Met. Integr. Biometal. Sci.* **2015**, *7*, 1459–1476. [[CrossRef](#)]
5. Roychoudhury, S.; Nath, S.; Massanyi, P.; Stawarz, R.; Kačániová, M.; Kolesarova, A. Copper-Induced Changes in Reproductive Functions: In Vivo and In Vitro Effects. *Physiol. Res.* **2016**, *65*, 11–22. [[CrossRef](#)]
6. Zhang, L.; Yuan, Z.; Bi, J. Estimation of Copper In-use Stocks in Nanjing, China. *J. Ind. Ecol.* **2012**, *16*, 191–202. [[CrossRef](#)]
7. Committee on Copper in Drinking. *Copper in Drinking Water*; National Academy Press: Washington, DC, USA, 2001.
8. Lamtai, M.; Zghari, O.; Ouakki, S.; Marmouzi, I.; Mesfioui, A.; El Hessni, A.; Ouichou, A. Chronic copper exposure leads to hippocampus oxidative stress and impaired learning and memory in male and female rats. *Toxicol. Res.* **2020**, *36*, 359–366. [[CrossRef](#)]
9. Brewer, G.J. Alzheimer’s disease causation by copper toxicity and treatment with zinc. *Front. Aging Neurosci.* **2014**, *6*, 92. [[CrossRef](#)]
10. Montes, S.; Rivera-Mancia, S.; Diaz-Ruiz, A.; Tristan-Lopez, L.; Rios, C. Copper and Copper Proteins in Parkinson’s Disease. *Oxidative Med. Cell. Longev.* **2014**, *2014*, 147251. [[CrossRef](#)]
11. Zietz, B.; De Vergara, J.D.; Schneider, H.; Kessler, B.; Dunkelberg, H. Prevalence of elevated copper concentrations in tap water in two areas of Germany used for infant feeding and possible health implications. *Eur. J. Med. Res.* **1999**, *4*, 298.
12. Hsiao, C.-D.; Wu, H.-H.; Malhotra, N.; Liu, Y.-C.; Wu, Y.-H.; Lin, Y.-N.; Saputra, F.; Santoso, F.; Chen, K. Expression and Purification of Recombinant GHK Tripeptides Are Able to Protect against Acute Cardiotoxicity from Exposure to Waterborne-Copper in Zebrafish. *Biomolecules* **2020**, *10*, 1202. [[CrossRef](#)]
13. Hashish, E.A.; Elgaml, S.A. Hepatoprotective and Nephroprotective Effect of Curcumin Against Copper Toxicity in Rats. *Indian J. Clin. Biochem.* **2016**, *31*, 270–277. [[CrossRef](#)]
14. Kumar, V.; Kalita, J.; Misra, U.; Bora, H. A study of dose response and organ susceptibility of copper toxicity in a rat model. *J. Trace Elem. Med. Biol.* **2015**, *29*, 269–274. [[CrossRef](#)]
15. Sarawi, W.S.; Alhusaini, A.M.; Fadda, L.M.; Alomar, H.A.; Albaker, A.B.; Aljrboa, A.S.; Alotaibi, A.M.; Hasan, I.H.; Mahmoud, A.M. Curcumin and Nano-Curcumin Mitigate Copper Neurotoxicity by Modulating Oxidative Stress, Inflammation, and Akt/GSK-3 β Signaling. *Molecules* **2021**, *26*, 5591. [[CrossRef](#)]
16. Sarawi, W.S.; Alhusaini, A.M.; Fadda, L.M.; Alomar, H.A.; Albaker, A.B.; Aljrboa, A.S.; Alotaibi, A.M.; Hasan, I.H.; Mahmoud, A.M. Nano-curcumin prevents cardiac injury, oxidative stress and inflammation, and modulates TLR4/NF- κ B and MAPK signaling in copper sulfate-intoxicated rats. *Antioxidants* **2021**, *10*, 1414. [[CrossRef](#)]
17. Zhao, D.; Zhang, X.; Li, X.; Ru, S.; Wang, Y.; Yin, J.; Liu, D. Oxidative damage induced by copper in testis of the red swamp crayfish *Procambarus clarkii* and its underlying mechanisms. *Aquat. Toxicol.* **2019**, *207*, 120–131. [[CrossRef](#)]
18. Lin, S.; Qiao, N.; Chen, H.; Tang, Z.; Han, Q.; Mehmood, K.; Fazlani, S.A.; Hameed, S.; Li, Y.; Zhang, H. Integration of transcriptomic and metabolomic data reveals metabolic pathway alteration in mouse spermatogonia with the effect of copper exposure. *Chemosphere* **2020**, *256*, 126974. [[CrossRef](#)]
19. Khushboo, M.; Murthy, M.K.; Devi, M.S.; Sanjeev, S.; Ibrahim, K.S.; Kumar, N.S.; Roy, V.K.; Gurusubramanian, G. Testicular toxicity and sperm quality following copper exposure in Wistar albino rats: Ameliorative potentials of L-carnitine. *Environ. Sci. Pollut. Res.* **2018**, *25*, 1837–1862. [[CrossRef](#)]
20. Chen, H.; Kang, Z.; Qiao, N.; Liu, G.; Huang, K.; Wang, X.; Pang, C.; Zeng, Q.; Tang, Z.; Li, Y. Chronic Copper Exposure Induces Hypospermatogenesis in Mice by Increasing Apoptosis Without Affecting Testosterone Secretion. *Biol. Trace Elem. Res.* **2020**, *195*, 472–480. [[CrossRef](#)]
21. Sakhaee, E.; Emadi, L.; Abshenas, J.; Kheirandish, R.; Azari, O.; Amiri, E. Evaluation of epididymal sperm quality following experimentally induced copper poisoning in male rats. *Andrologia* **2012**, *44*, 110–116. [[CrossRef](#)]
22. Babaei, H.; Roshangar, L.; Sakhaee, E.; Abshenas, J.; Kheirandish, R.; Dehghani, R. Ultrastructural and morphometrical changes of mice ovaries following experimentally induced copper poisoning. *Iran. Red Crescent Med. J.* **2012**, *14*, 558–568.
23. Chattopadhyay, A.; Sarkar, M.; Sengupta, R.; Roychowdhury, G.; Biswas, N.M. Antitesticular effect of copper chloride in albino rats. *J. Toxicol. Sci.* **1999**, *24*, 393–397. [[CrossRef](#)]
24. Li, Y.; Chen, H.; Liao, J.; Chen, K.; Javed, M.T.; Qiao, N.; Zeng, Q.; Liu, B.; Yi, J.; Tang, Z.; et al. Long-term copper exposure promotes apoptosis and autophagy by inducing oxidative stress in pig testis. *Environ. Sci. Pollut. Res.* **2021**, *28*, 55140–55153. [[CrossRef](#)]
25. Aitken, R.J.; Roman, S. Antioxidant Systems and Oxidative Stress in the Testes. *Oxidative Med. Cell. Longev.* **2008**, *1*, 616821. [[CrossRef](#)]
26. Saghir, S.A.M.; Alharbi, S.A.; Al-Garadi, M.A.; Al-Gabri, N.; Rady, H.Y.; Olama, N.K.; Abdulghani, M.A.M.; Al Hroob, A.M.; Almainan, A.A.; Bin-Jumah, M.; et al. Curcumin Prevents Cyclophosphamide-Induced Lung Injury in Rats by Suppressing Oxidative Stress and Apoptosis. *Processes* **2020**, *8*, 127. [[CrossRef](#)]
27. Pizzo, P.; Scapin, C.; Vitadello, M.; Florean, C.; Gorza, L. Grp94 acts as a mediator of curcumin-induced antioxidant defence in myogenic cells. *J. Cell. Mol. Med.* **2010**, *14*, 970–981. [[CrossRef](#)]
28. Esatbeyoglu, T.; Huebbe, P.; Ernst, I.M.A.; Chin, D.; Wagner, A.E.; Rimbach, G. Curcumin-From Molecule to Biological Function. *Angew. Chem. Int. Ed.* **2012**, *51*, 5308–5332. [[CrossRef](#)]

29. De, R.; Kundu, P.; Swarnakar, S.; Ramamurthy, T.; Chowdhury, A.; Nair, G.B.; Mukhopadhyay, A.K. Antimicrobial Activity of Curcumin against *Helicobacter pylori* Isolates from India and during Infections in Mice. *Antimicrob. Agents Chemother.* **2009**, *53*, 1592–1597. [[CrossRef](#)]
30. Tsuda, T. Curcumin as a functional food-derived factor: Degradation products, metabolites, bioactivity, and future perspectives. *Food Funct.* **2018**, *9*, 705–714. [[CrossRef](#)]
31. Mahmoud, A.M.; Ahmed, O.M.; Galaly, S.R. Thymoquinone and curcumin attenuate gentamicin-induced renal oxidative stress, inflammation and apoptosis in rats. *EXCLI J.* **2014**, *13*, 98–110.
32. Alhusaini, A.; Fadda, L.; Hasan, I.H.; Zakaria, E.; Alenazi, A.M.; Mahmoud, A.M. Curcumin Ameliorates Lead-Induced Hepatotoxicity by Suppressing Oxidative Stress and Inflammation, and Modulating Akt/GSK-3 β Signaling Pathway. *Biomolecules* **2019**, *9*, 703. [[CrossRef](#)]
33. Al-Dossari, M.H.; Fadda, L.M.; Attia, H.A.; Hasan, I.H.; Mahmoud, A.M. Curcumin and Selenium Prevent Lipopolysaccharide/Diclofenac-Induced Liver Injury by Suppressing Inflammation and Oxidative Stress. *Biol. Trace Elem. Res.* **2019**, *196*, 173–183. [[CrossRef](#)]
34. Galaly, S.R.; Ahmed, O.M.; Mahmoud, A.M. Thymoquinone and curcumin prevent gentamicin-induced liver injury by attenuating oxidative stress, inflammation and apoptosis. *J. Physiol. Pharmacol. Off. J. Pol. Physiol. Soc.* **2014**, *65*, 823–832.
35. Damiano, S.; Longobardi, C.; Andretta, E.; Prisco, F.; Piegari, G.; Squillacioti, C.; Montagnaro, S.; Pagnini, F.; Badino, P.; Florio, S.; et al. Antioxidative Effects of Curcumin on the Hepatotoxicity Induced by Ochratoxin A in Rats. *Antioxidants* **2021**, *10*, 125. [[CrossRef](#)]
36. Longobardi, C.; Damiano, S.; Andretta, E.; Prisco, F.; Russo, V.; Pagnini, F.; Florio, S.; Ciarcia, R. Curcumin Modulates Nitrosative Stress, Inflammation, and DNA Damage and Protects against Ochratoxin A-Induced Hepatotoxicity and Nephrotoxicity in Rats. *Antioxidants* **2021**, *10*, 1239. [[CrossRef](#)]
37. Aggarwal, B.B.; Harikumar, K.B. Potential therapeutic effects of curcumin, the anti-inflammatory agent, against neurodegenerative, cardiovascular, pulmonary, metabolic, autoimmune and neoplastic diseases. *Int. J. Biochem. Cell Biol.* **2009**, *41*, 40–59. [[CrossRef](#)]
38. Chikara, S.; Nagaprashantha, L.D.; Singhal, J.; Horne, D.; Awasthi, S.; Singhal, S.S. Oxidative stress and dietary phytochemicals: Role in cancer chemoprevention and treatment. *Cancer Lett.* **2018**, *413*, 122–134. [[CrossRef](#)]
39. Mirzaei, H.; Masoudifar, A.; Sahebkar, A.; Zare, N.; Sadri Nahand, J.; Rashidi, B.; Mehrabian, E.; Mohammadi, M.; Mirzaei, H.R.; Jaafari, M.R. MicroRNA: A novel target of curcumin in cancer therapy. *J. Cell. Physiol.* **2018**, *233*, 3004–3015. [[CrossRef](#)]
40. Carlos-Reyes, Á.; López-González, J.S.; Meneses-Flores, M.; Gallardo-Rincón, D.; Ruíz-García, E.; Marchat, L.; La Vega, H.A.-D.; De La Cruz, O.N.H.; López-Camarillo, C. Dietary Compounds as Epigenetic Modulating Agents in Cancer. *Front. Genet.* **2019**, *10*, 79. [[CrossRef](#)]
41. Karimi, S.; Khorsandi, L.; Nejaddehbashi, F. Protective effects of Curcumin on testicular toxicity induced by titanium dioxide nanoparticles in mice. *JBRA Assist. Reprod.* **2019**, *23*, 344–351. [[CrossRef](#)]
42. Sudjarwo, S.A.; Sudjarwo, G.W. Koerniasari Protective effect of curcumin on lead acetate-induced testicular toxicity in Wistar rats. *Res. Pharm. Sci.* **2017**, *12*, 381–390. [[CrossRef](#)] [[PubMed](#)]
43. Huyut, Z.; Alp, H.H.; Yaman, T.; Keleş, Ö.F.; Yener, Z.; Türkan, F.; Ayengin, K. Comparison of the protective effects of curcumin and caffeic acid phenethyl ester against doxorubicin-induced testicular toxicity. *Andrologia* **2020**, *53*, e13919. [[CrossRef](#)] [[PubMed](#)]
44. Flora, G.; Gupta, D.; Tiwari, A. Nanocurcumin: A Promising Therapeutic Advancement over Native Curcumin. *Crit. Rev. Ther. Drug Carrier Syst.* **2013**, *30*, 331–368. [[CrossRef](#)] [[PubMed](#)]
45. Gera, M.; Sharma, N.; Ghosh, M.; Huynh, D.L.; Lee, S.J.; Min, T.; Kwon, T.; Jeong, D.K. Nanoformulations of curcumin: An emerging paradigm for improved remedial application. *Oncotarget* **2017**, *8*, 66680–66698. [[CrossRef](#)] [[PubMed](#)]
46. Feng, T.; Wei, Y.; Lee, R.J.; Zhao, L. Liposomal curcumin and its application in cancer. *Int. J. Nanomed.* **2017**, *12*, 6027–6044. [[CrossRef](#)]
47. Nisini, R.; Poerio, N.; Mariotti, S.; De Santis, F.; Fraziano, M. The Multirole of Liposomes in Therapy and Prevention of Infectious Diseases. *Front. Immunol.* **2018**, *9*, 155. [[CrossRef](#)]
48. Satta, S.; Mahmoud, A.M.; Wilkinson, F.L.; Alexander, M.Y.; White, S.J. The role of Nrf2 in cardiovascular function and disease. *Oxid. Med. Cell. Longev.* **2017**, *2017*, 9237263. [[CrossRef](#)]
49. Alhusaini, A.; Fadda, L.; Hassan, I.; Ali, H.M.; Alsaadan, N.; Aldowsari, N.; Aldosari, A.; Alharbi, B. Liposomal curcumin attenuates the incidence of oxidative stress, inflammation, and DNA damage induced by copper sulfate in rat liver. *Dose Response A Publ. Int. Hormesis Soc.* **2018**, *16*, 1559325818790869. [[CrossRef](#)]
50. Ohkawa, H.; Ohishi, N.; Yagi, K. Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Anal. Biochem.* **1979**, *95*, 351–358. [[CrossRef](#)]
51. Grisham, M.B.; Johnson, G.G.; Lancaster, J.R., Jr. Quantitation of nitrate and nitrite in extracellular fluids. *Methods Enzymol.* **1996**, *268*, 237–246.
52. Elluman, G.L. Tissue sulfhydryl groups. *Arch. Biochem. Biophys.* **1959**, *82*, 70–77. [[CrossRef](#)]
53. Marklund, S.L. Superoxide dismutase isoenzymes in tissues and plasma from New Zealand black mice, nude mice and normal BALB/c mice. *Mutat. Res.* **1985**, *148*, 129–134. [[CrossRef](#)]
54. Cohen, G.; Dembiec, D.; Marcus, J. Measurement of catalase activity in tissue extracts. *Anal. Biochem.* **1970**, *34*, 30–38. [[CrossRef](#)]

55. Alotaibi, M.F.; Al-Joufi, F.; Seif, H.S.A.; Alzoghaibi, M.A.; Djouhri, L.; Ahmeda, A.F.; Mahmoud, A.M. Umbelliferone Inhibits Spermatogenic Defects and Testicular Injury in Lead-Intoxicated Rats by Suppressing Oxidative Stress and Inflammation, and Improving Nrf2/HO-1 Signaling. *Drug Des. Dev. Ther.* **2020**, *14*, 4003–4019. [[CrossRef](#)] [[PubMed](#)]
56. Livak, K.J.; Schmittgen, T.D. Analysis of relative gene expression data using real-time quantitative pcr and the 2⁻ $\Delta\Delta$ CT method. *Methods* **2001**, *25*, 402–408. [[CrossRef](#)]
57. Vázquez-Blanco, R.; Arias-Estévez, M.; Bååth, E.; Fernández-Calviño, D. Comparison of Cu salts and commercial Cu based fungicides on toxicity towards microorganisms in soil. *Environ. Pollut.* **2020**, *257*, 113585. [[CrossRef](#)]
58. Oduwole, O.O.; Peltoketo, H.; Huhtaniemi, I.T. Role of Follicle-Stimulating Hormone in Spermatogenesis. *Front. Endocrinol.* **2018**, *9*, 763. [[CrossRef](#)]
59. Miller, W.L.; Bose, H.S. Early steps in steroidogenesis: Intracellular cholesterol trafficking. *J. Lipid Res.* **2011**, *52*, 2111–2135. [[CrossRef](#)] [[PubMed](#)]
60. Caron, K.M.; Soo, S.-C.; Wetsel, W.C.; Stocco, D.M.; Clark, B.J.; Parker, K.L. Targeted disruption of the mouse gene encoding steroidogenic acute regulatory protein provides insights into congenital lipid adrenal hyperplasia. *Proc. Natl. Acad. Sci. USA* **1997**, *94*, 11540–11545. [[CrossRef](#)]
61. Bremer, A.A.; Miller, W.L. Chapter 13—Regulation of steroidogenesis. In *Cellular Endocrinology in Health and Disease*; Ulloa-Aguirre, A., Conn, P.M., Eds.; Academic Press: Boston, MA, USA, 2014; pp. 207–227.
62. Dušková, M.; Pospíšilová, H. The Role of Non-Aromatizable Testosterone Metabolite in Metabolic Pathways. *Physiol. Res.* **2011**, *60*, 253–261. [[CrossRef](#)]
63. Wang, R.-S.; Yeh, S.; Tzeng, C.-R.; Chang, C. Androgen Receptor Roles in Spermatogenesis and Fertility: Lessons from Testicular Cell-Specific Androgen Receptor Knockout Mice. *Endocr. Rev.* **2009**, *30*, 119–132. [[CrossRef](#)] [[PubMed](#)]
64. Balasubramanian, K. Quantum chemical insights into alzheimer’s disease: Curcumin’s chelation with Cu(ii), Zn(ii), and Pd(ii) as a mechanism for its prevention. *Int. J. Quantum Chem.* **2016**, *116*, 1107–1119. [[CrossRef](#)]
65. Gunther, M.R.; Hanna, P.; Mason, R.P.; Cohen, M. Hydroxyl Radical Formation from Cuprous Ion and Hydrogen Peroxide: A Spin-Trapping Study. *Arch. Biochem. Biophys.* **1995**, *316*, 515–522. [[CrossRef](#)]
66. Gaetke, L.M.; Chow-Johnson, H.S.; Chow, C.K. Copper: Toxicological relevance and mechanisms. *Arch. Toxicol.* **2014**, *88*, 1929–1938. [[CrossRef](#)] [[PubMed](#)]
67. Mahmoud, A.M.; Sayed, A.M.; Ahmed, O.S.; Abdel-Daim, M.M.; Hassanein, E.H.M. The role of flavonoids in inhibiting IL-6 and inflammatory arthritis. *Curr. Top. Med. Chem.* **2022**, *22*, 746–768. [[CrossRef](#)]
68. Yang, S.-H.; He, J.-B.; Yu, L.-H.; Li, L.; Long, M.; Liu, M.-D.; Li, P. Protective role of curcumin in cadmium-induced testicular injury in mice by attenuating oxidative stress via Nrf2/ARE pathway. *Environ. Sci. Pollut. Res.* **2019**, *26*, 34575–34583. [[CrossRef](#)]
69. Shahcheraghi, S.H.; Salemi, F.; Peirovi, N.; Ayatollahi, J.; Alam, W.; Khan, H.; Saso, L. Nrf2 Regulation by Curcumin: Molecular Aspects for Therapeutic Prospects. *Molecules* **2021**, *27*, 167. [[CrossRef](#)]
70. Bonizzi, G.; Karin, M. The two NF- κ B activation pathways and their role in innate and adaptive immunity. *Trends Immunol.* **2004**, *25*, 280–288. [[CrossRef](#)]
71. Shi, Y.; Chen, J.; Weng, C.; Chen, R.; Zheng, Y.; Chen, Q.; Tang, H. Identification of the protein–protein contact site and interaction mode of human VDAC1 with Bcl-2 family proteins. *Biochem. Biophys. Res. Commun.* **2003**, *305*, 989–996. [[CrossRef](#)]
72. Redza-Dutordoir, M.; Averill-Bates, D.A. Activation of apoptosis signalling pathways by reactive oxygen species. *Biochim. Biophys. Acta (BBA)—Mol. Cell Res.* **2016**, *1863*, 2977–2992. [[CrossRef](#)]
73. Wrobel, K.-H.; Bickel, D.; Kujat, R. Immunohistochemical study of seminiferous epithelium in adult bovine testis using monoclonal antibodies against Ki-67 protein and proliferating cell nuclear antigen (PCNA). *Cell Tissue Res.* **1996**, *283*, 191–201. [[CrossRef](#)]
74. Chen, Z.; Wen, D.; Wang, F.; Wang, C.; Yang, L. Curcumin protects against palmitic acid-induced apoptosis via the inhibition of endoplasmic reticulum stress in testicular Leydig cells. *Reprod. Biol. Endocrinol.* **2019**, *17*, 71. [[CrossRef](#)]
75. Kilinc, L.; Uz, Y.H. Protective effects of curcumin against methotrexate-induced testicular damage in rats by suppression of the p38-MAPK and nuclear factor-kappa B pathways. *Clin. Exp. Reprod. Med.* **2021**, *48*, 211–220. [[CrossRef](#)]
76. Boyanapalli, S.S.S.; Paredes-Gonzalez, X.; Fuentes, F.; Zhang, C.; Guo, Y.; Pung, D.; Saw, C.L.L.; Kong, A.-N.T. Nrf2 Knockout Attenuates the Anti-Inflammatory Effects of Phenethyl Isothiocyanate and Curcumin. *Chem. Res. Toxicol.* **2014**, *27*, 2036–2043. [[CrossRef](#)]
77. Yen, F.-L.; Wu, T.-H.; Tzeng, C.-W.; Lin, L.-T.; Lin, C.-C. Curcumin Nanoparticles Improve the Physicochemical Properties of Curcumin and Effectively Enhance Its Antioxidant and Antihepatoma Activities. *J. Agric. Food Chem.* **2010**, *58*, 7376–7382. [[CrossRef](#)]
78. Nahar, P.P.; Slitt, A.L.; Seeram, N.P. Anti-Inflammatory Effects of Novel Standardized Solid Lipid Curcumin Formulations. *J. Med. Food* **2015**, *18*, 786–792. [[CrossRef](#)]