

Dufour Dominique (Orcid ID: 0000-0002-7794-8671)
 ABOLORE Bello (Orcid ID: 0000-0002-8871-6163)
 Olaosebikan Deborah Olamide (Orcid ID: 0000-0003-1470-1150)
 Fotso Kuate Apollin (Orcid ID: 0000-0002-5247-7519)
 Adinsi Laurent (Orcid ID: 0000-0001-8853-5445)
 Forsythe Lora (Orcid ID: 0000-0001-9931-4453)

Varietal impact on women's labour, workload and related drudgery in processing root, tuber and banana crops. Focus on cassava in sub-Saharan Africa

Alexandre Bouniol^{1,2,3*§}, Hernan Ceballos⁴, Bello Abolore⁵, Béla Teeken⁵, Deborah Olamide Olaosebikan⁵, Durodola Owoade⁵, Agbona Afolabi^{5&6}, Fotso Kuate Apollin⁷, Tessy Madu⁸, Benjamin Okoye⁸, Miriam Ofoeze⁸, Solomon Nwafor⁸, Nnaemeka Onyemauwa⁸, Laurent Adinsi¹, Lora Forsythe¹⁰, **Dominique Dufour**^{3&9§*}

§ These authors contributed equally to the work

† *In Memoriam*

* Corresponding author

- 1) Laboratoire de Sciences des Aliments, Faculté des Sciences Agronomiques, Université d' Abomey-Calavi, Jéricho 03 BP 2819, Benin.
- 2) CIRAD, UMR QUALISUD, Cotonou 01 BP 52, Benin.
- 3) QualiSud, Univ Montpellier, Avignon Université, CIRAD, Institut Agro, IRD, Université de La Réunion, Montpellier, France.
- 4) International Consultant, Malaga, Spain.
- 5) International Institute of Tropical Agriculture (IITA), Ibadan PMB 5320, Nigeria.
- 6) Department of Soil and Crop Science, Molecular & Environmental Plant Sciences, Texas A & M University, College Station, TX 77843, USA.
- 7) International Institute of Tropical Agriculture, BP 2008, Messa, Yaoundé, Cameroon.
- 8) International National Root Crops Research Institute (NRCRI), Umudike, Umuahia PMB 7006, Nigeria.
- 9) CIRAD, UMR QualiSud, F-34398 Montpellier, France.
- 10) Natural Resources Institute, University of Greenwich, Central Avenue, Chatham Maritime, Kent ME4 4TB, UK.

Author	Email	ORCID ID
Alexandre Bouniol	alexandre.bouniol@cirad.fr	0000-0002-6140-424X
Hernan Ceballos	hernanceballos154@gmail.com	0000-0002-8744-7918
Bello Abolore	A.Bello@cgiar.org	0000-0002-8871-6163
Béla Teeken	B.Teeken@cgiar.org	0000-0002-3150-1532
Deborah Olamide Olaosebikan	D.Nwanze@cgiar.org	0000-0003-1470-1150
Durodola Owoade	d.owoade@cgiar.org	0000-0001-8711-0138
Agbona Afolabi	aafolabi@tamu.edu	0000-0002-9756-5432
Fotso Kuate Apollin	a.fotso@cgiar.org	0000-0002-5247-7519
Tessy Madu	tessmadu@gmail.com	0000-0002-0098-3567
Benjamin Okoye	okoyebenjamen@yahoo.com	0000-0001-7407-3032
Miriam Ofoeze	mimiofoeze@gmail.com	0000-0003-0839-7607
Solomon Nwafor	solomonnwafor8@gmail.com	0000-0001-7499-1757
Nnaemeka Onyemauwa	emeka.visco@gmail.com	0000-0002-8099-998X
Laurent Adinsi	adinsil2003@yahoo.fr	0000-0001-8853-5445
Lora Forsythe	l.forsythe@greenwich.ac.uk	0000-0001-9931-4453

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the [Version of Record](#). Please cite this article as doi: [10.1002/jsfa.12936](https://doi.org/10.1002/jsfa.12936)

Abstract

Roots, tubers and cooking bananas are bulky and highly perishable. In Africa, except for yams, their consumption is mainly after transport, peeling and cooking in the form of boiled pieces or dough, a few days after harvest. To stabilize, better preserve the products and, in the case of cassava, release toxic cyanogenic glucosides, a range of intermediate products have been developed, mainly for cassava, related to fermentation and drying after numerous processing operations. This review highlights, for the first time, the impact of genotypes on labour requirements, productivity, and the associated drudgery in processing operations primarily carried out by women processors. Peeling, soaking/grinding/fermentation, dewatering, sieving, and toasting steps were evaluated on a wide range of new hybrids and traditional landraces. The review highlights case studies of gari production from cassava. Results show that, depending on the genotypes used, women's required labour can be more than doubled and even the sum of the weights transported along the process can be up to four times higher for the same quantity of end product. Productivity and loads carried between each processing operation are highly influenced by root shape, ease of peeling, dry matter content and/or fiber content. Productivity and the often related experienced drudgery are key factors to be considered for a better acceptance of new genotypes by actors in the value-addition chain, leading to enhanced adoption, and ultimately to improved livelihoods for women processors.

Keywords: Breeding pipeline, Gender, Varietal Adoption, Technological Characteristics, Food Product Profile, End-product Productivity, Value Chain, Market Segment.

Introduction

Drudgery in processing of root, tuber and banana crops (RTBs) has been recognized as a major, complex social, economic and health problem¹. However, little attention has been paid to the influence of varietal differences on drudgery, and the potential to exploit breeding of appropriate varieties (genotypes) as a partial solution. Although it can have a significant impact on the livelihoods and wellbeing of women (who perform the majority of processing labour), addressing labour requirements, productivity and related drudgery in food processing is often ignored in the development of improved RTB genotypes. Throughout sub-Saharan Africa, women play a vitally important role in agriculture and post-harvest activities, 50% of agricultural work is done by women, with significant variations within and among regions and countries². Agri-food processing at artisanal or small scale is mainly carried out by women with the help of family labour (often young and/or elderly people and neighbors). For processing of RTBs, these operations are often carried out entirely by women -- from the peeling to the elaboration of the end-products ready to be marketed or consumed at the household level^{3,4,5,6,7}. The Collaborative Study of Cassava in Africa (COSCA), conducted in six African countries, showed that women lead in root transportation (68%) and processing operations (76%)⁸. On average, cassava processing was carried out mostly by women in about 75% of the surveyed villages, mostly by men in less than 5%, and by both equally in about 20%¹. Women in sub-Saharan Africa have the highest average agricultural labour-force participation rates in the world⁹.

Assessment of processing productivity and the drudgery associated with processing RTB-based foods is limited. Genotype acceptability and drudgery in processing appear to be strongly linked, i.e. women are more likely to prefer RTB varieties with traits that reduce the drudgery of processing^{10,11,12,13}. Furthermore, cassava processing, for example, is associated with challenging working conditions and serious health hazards^{14,15,16} which increase the likelihood that such operations are perceived as drudgery. Prolonged labour associated with all the operations significantly impinges on the productivity and wellbeing of the (mostly) women operators^{17, 18}. Drudgery has been defined as the dissatisfactory experiences that constrain work performance in any activity¹⁹ and is often related to time-consuming, repetitive and menial work. Physical and mental strain, agony, monotony and hardship have been linked to the drudgery often experienced in farm operations²⁰

Whether a task is experienced as drudgery depends on many different factors which include specific working conditions (which include the quality and type of tools used), the extent of the task but also how the work is culturally validated and looked upon which in turn determines the type of meaning executors of the task attribute to the task²¹.

RTBs have several important features of note in relation to labour input. They tend to be bulky and perishable, making them difficult to transport, and as a result most are often eaten fresh, after cooking, soon after harvest. Alternatively, RTBs are processed soon after harvest to convert them into less perishable and more easily transported products. Cassava, in particular, is extremely sensitive (3 to 5 days) to post-harvest deterioration²². Post-harvest handling and storage offer many challenges for RTBs, eliciting development of a wide variety of food products^{23,24,25, 26}.

Cassava is by far the most widely grown and consumed root in sub-Saharan Africa²⁷. The starchy roots contain toxic cyanogenetic compounds at various levels^{28,29}. Typically consumers perceive bitterness from cyanogens beyond a cyanogenic potential of about 80-100 ppm³⁰. These toxic compounds are only very partially (20%) removed by boiling or frying³¹ so genotypes below 50 ppm are recommended for consumption as boiled or fried pieces (genotypes above 100 ppm become toxic for mammals and must be detoxified to avoid health risks³². The detoxification is done mainly by grinding or rasping, often with fermentation before or after that process. Fermentation is done either by soaking the whole

or peeled roots in containers (retting), or by resting the pulp in sacks or containers for several days. During these operations the volatile cyanogenic compounds are released³³. These double-purpose operations of fermentation (detoxification and softening) of the roots in water are essential for the fiber removal and preparation of safe traditional staple foods in Africa: fufu, lafun, batôn, Chikwangué, agbelima, gari, attieke³⁴. Varying the length of fermentation allows formulation of products according to the preferred tastes of consumers, which are very diversified for a range of products, are specific to production zones of origin, and may vary according to availability in the market^{35,36}.

The current study is an output of the project *Breeding RTB products for end user preferences* (RTBfoods <https://rtbfoods.cirad.fr/>). This project developed a new five-step methodology for developing food product profiles, through mobilizing a multidisciplinary team of breeders, social scientists, and food technologists to capture the preferred traits of farmers, end-users and consumers^{37,38}. The data were produced following a standardized participatory processing diagnosis procedure common to all RTBs³⁹ and mainly published in 2021 as a special issue of the *International Journal of Food Science and Technology*: Consumers have their say: assessing preferred quality traits of roots, tubers and cooking bananas, and implications for breeding <https://ifst.onlinelibrary.wiley.com/toc/13652621/2021/56/3>.

The main objectives of the current study were *i)* to dissect the individual processing steps involved in different RTB food products; *ii)* to evaluate effects of genetic differences among varieties on processor workload, particularly comparing improved and traditional varieties; and *iii)* to guide breeders by highlighting the traits responsible for varietal differences in required labour inputs, which in turn influence acceptability of new varieties.

This review, therefore, does not directly evaluate the level of drudgery experienced based on different genotypes, but rather examines the impact of varieties on productivity and the associated labour requirements in processing operations. Regardless of how an operation is perceived, a decrease in productivity or an increase in labour requirements can potentially contribute to the drudgery experienced by processors. Therefore, this review examined the influence of different varieties on the productivity of processors individually for various processing operations. The study measured processing operator productivity by the amount of mass produced per unit of time per operator within each operation. Additionally, as a secondary measure related to productivity, we evaluated the weights of products that processors had to transport between operations. This assessment served as an indicator, albeit productivity-related, to quantify the labour involved in processing different varieties.

Materials and Methods

In rural or peri-urban areas of East and West Africa, RTB processing sites were selected according to a predefined sampling methodology. The processing trials were participatory and carried out with experienced *champion processors*. Both preferred (locally commonly used/popular varieties) and non-preferred genotypes (locally known as non-adequate for expected product profile) were included to provide a wide range of technological and physico-chemical characteristics (fixed effects from the statistical point of view). Processors provided feedback on the varieties before, during each step, and after processing to identify relevant characteristics of the crop and product. Processing parameters were measured at each step. The specific traits related to the processing ability were included in the different Food Product Profiles.

Proper and representative quantity of product was harvested from at least four genotypes (more if possible). Each processing step was conducted in duplicate (averages of two women, who collaborated by processing half of each batch of material for each genotype).

- Processors were invited to observe each raw genotype and give their views, as an operation unit, on its quality characteristics.
- Dry matter content (DMC) of all the collected samples was measured.
- The processing was carried out in real/normal conditions in processors' own communities.
- Each processor started with the first operation unit stage, processing one variety at a time.

For boiled or pounded products (cassava, plantain, yam), the process is relatively simple, including steps such as peeling, washing, boiling, steaming or frying, and (in some cases), mashing and pounding. The parameters measured were: peeling yield, duration of peeling, cooking (or pounding), and productivity of each operation unit. These products were processed by only 2-3 operation units.

Processing operations of fermented products, on the other hand, are more complex and involve more steps. To standardize the data collected in different countries and on different production lines, all the data were scaled to obtain 1000 kg of the end product. Each process was broken down into separate operation units.

- *Gari* (granulated product): peeling; grating; washing; fermentation & draining; mechanical dewatering; wet pulp sieving; mash toasting; and dry gari sieving⁴⁰.
- *Dry Fufu* (soaked cassava product: Couscous, Lafun): peeling; washing; soaking; fiber & non-softened material removal; draining in bags; sun drying; and dry fufu grinding⁴¹.

A representative sample of approximately 20-50 kg of roots from each genotype were processed to perform the technological diagnosis, and used to measure the yield/productivity of each operation unit (The coding used for each genotype is available in supplementary table S1.). All trials were replicated twice. For each operation unit the average weight was measured before and after processing. The yields of each operation unit were thus calculated according to the genotypes studied and reported as ***Yield by operation unit (%)***. The time needed to perform each operation, on all the available material, was measured with a chronometer. ***Operator productivity by operation unit (kg/h/Operator)*** was calculated for each operation and per genotype studied. The calculation table allowed to visualize the incoming and outgoing quantities for each operation unit for 1000kg of final product: ***Processed material by operation unit (kg /1000 kg final product)***; ***Operator time by operation (h)*** and ***Time distribution by operation unit (%)*** were thus obtained. Three global values allowed characterizing genotypes by:

- ***Gari Yield (%)*** expressed by weight of initial root/weight of finished product.
- ***Operator Time by kg of final product (h/kg)***, is the time needed to produce one kilogram of end product per genotype. It is calculated as the sum of the time spent to finalize each operation unit, reported to one kilogram of final product.
- ***Weight carried by operator per kg of final product (kg)*** calculated as the sum of all intermediate products carried or moved between each operation unit to obtain one kilogram of final product.

The methodology described above became a processing diagnostics and mass balances for the main RTBs consumed in subtropical Africa. Yam in Benin & Nigeria; Sweetpotato & Potato in Uganda, Plantain in Cameroon, Matooke in Uganda, Cassava in Benin, Cameroon, Nigeria, Uganda.

Results

Productivity and peeling yields of RTBs strongly influence the labour required from the processors. Specifically for cassava processing, multiple operation units and teamwork make it difficult to evaluate

the labour required to produce one kilogram of end-product. New data on cassava, sweetpotato, yam, potato and cooking bananas (plantain and matooke) collected following Fliedel *et al.*, (2018) methodology were used³⁹. The data collected allowed estimating the average productivity of each operation according to the genotypes used.

RTB peeling yield related to shape and ease of peeling.

Table 1 reports the RTB peeling data (available in open access) in each RTB processing diagnostic summary for the main product profiles. The flowsheet of each RTB final product has been established and are fully available in each crop report on Table 1 links.

The lowest yields were for bananas, with 55% yield measured for the matooke peeling operation and a productivity of 28 kg/h/operator. For plantains a yield of 50% was observed and a productivity of 43 kg/h/operator of pulp obtained after peeling. These results are in general agreement with previous reports^{42,43}, although yields were slightly lower. For potatoes 75% of peeling yield and 9 kg/h/operator were found for manual peeling in Uganda. In the industry, using steam peeling for potato, losses range from 6 to 10% of fresh weight^{44, 45}. In Uganda, sweetpotato studies reported a peeling yield of 79% and a peeling productivity of 14 kg/h/operator. In India, losses ranged from 3 to 21% among 18 sweetpotato genotypes studied, with an average peel loss of 11%⁴⁶. Yam peeling yield of 80% has been reported in Nigeria and Benin, with an average productivity of 34 kg/h/operator.

For cassava, where the largest number of trials were conducted, the average peeling yield was 74% and the productivity 57 kg/hour/operator. However, there was a high variability among locations and end products (Table 1). The physical characteristics of the peel could influence both peeling labour and the amount of product lost at the peeling stage. These differences in the properties of the peel can include variation in thickness, texture and strength of adhesion to the root flesh⁴⁷. The peel consists of two basic components: the phelloderm, or the *bark*, is the thin, usually rough, outer layer; the cortex is the fibrous layer that attaches directly to the pulp. Low peel adhesion strength to cassava flesh allow easy removal of the external phelloderm by making an incision within the cortex with a knife, followed by pulling and removing the peel around the flesh or by rubbing it off in mechanized systems (figure 1). Cassava genotypes with easy peel removal reduced product waste and labour.

Table 1. Peeling yield and productivity and RTB shape reported on RTBfoods RTB processing diagnostic summary.

Product by institution and country	Link to published flowsheets	Average ± standard deviation					Number of genotypes		Age of plants at harvest (months)
		RTB Shape			Peeling yield (% w.b)	Productivity (kg/h/Operator)	Landraces	Improved	
		Weight (g)	Girth (cm)	Length (cm)					
Boiled Yam, UAC-FSA-Bénin (48)	http://agritrop.cirad.fr/602027/	1441±589	***	35±9	75 ± 11	39 ± 23	6	0	9
Boiled Yam, NRCRI-Nigeria (49)	https://agritrop.cirad.fr/602029/	***	***	***	87 ± 3	38 ± 9	0	4	***
Pounded Yam, Bowen Univ.-Nigeria (50)	https://agritrop.cirad.fr/602026/	978±340	31±6	36±5	79 ± 10	24 ± 11	10	***	***
Boiled sweetpotato, CIP-Uganda (51)	https://agritrop.cirad.fr/602026/	184±95	***	***	79 ± 5	14 ± 3	6	1	***
Boiled potato, NARL-CIP-Uganda (52)	https://agritrop.cirad.fr/602025/	68±42	***	***	75 ± 6	9 ± 4	4	2	***
Boiled plantain, CARBAP-Cameroon (53)	https://agritrop.cirad.fr/602023/	222±57	13,7±1,3	24,3±4,5	50 ± 5	43 ± 4	3	1	***
Matooke, NARL-Uganda (25)	https://agritrop.cirad.fr/602041/	26±9	13,7±1,4	19,3±1,4	55 ± 4	28 ± 5	3	3	4
Bâton, CIRAD/IITA-Cameroon (54)	https://agritrop.cirad.fr/595635/	875±482	21±6	27±4	70 ± 5	54 ± 13	0	8	13
Gari, CIRAD/UAC-FSA-Benin (55)	https://agritrop.cirad.fr/597596/	509±178	19±7	28±3	66 ± 3	66 ± 18	2	13	12
Dry Fufu, CIRAD/IITA-Cameroon (56)	https://agritrop.cirad.fr/597597/	***	***	***	73 ± 6	40 ± 10	3	18	13
Dry Fufu, CIRAD/UAC-FSA-Benin (57)	https://agritrop.cirad.fr/597594/	1068±707	23±6	34±11	73 ± 3	64 ± 13	3	13	12
Boiled cassava, NaCRRI-Uganda (58)	https://agritrop.cirad.fr/602019/	***	***	***	70 ± 8	60 ± 27	4	5	*
Boiled cassava, UAC-FSA-Bénin (59)	https://agritrop.cirad.fr/602013/	388±345	16±5	29±9	77 ± 6	16 ± 8	6	0	12
Water Fufu, IITA –Nigeria (60)	https://doi.org/10.1111/ijfs.14862	577±585	22±6	26±9	80 ± 3	54 ± 7	5	15	13
Gari/Eba, IITA-Nigeria (60)	https://doi.org/10.1111/ijfs.14862	581±453	25±9	21±8	78 ± 3	63 ± 12	5	15	13
Gari/Eba, NRCRI-Nigeria (61)	https://agritrop.cirad.fr/602035/	***	***	***	75 ± 3	***	***	***	***
Water Fufu, NRCRI-Nigeria (62)	https://agritrop.cirad.fr/602033/	***	***	***	83 ± 7	23 ± 4	1	3	12
Attieke, CNRA-Cote d'Ivoire (63)	https://agritrop.cirad.fr/603470/	1254±559	24±4	37±10	70±5	34±7	2	4	15

*** Not evaluated



Figure 1. Peeling cassava roots. **A.** Illustration of the occurrence of constrictions (left) and variation of root shape and size (right). **B.** Traditional peeling by slashing in Africa. **C.** Illustration of roots where the peel can be easily removed.

Roots of the majority of existing cassava genotypes in Africa can only be peeled (with current technologies) by slashing it off the flesh of the root with a sharp knife or machete, increasing enormously the losses and the women processor fatigue⁶⁴. Some authors used a large number of cassava genotypes in Colombia⁶⁵, Uganda⁶⁶ or India⁶⁷ for evaluating phelloderm and root cortex thickness: (0.79 to 5.14 mm in Colombia on 64 genotypes; 0.3–4.9 mm in East Africa on 825 genotypes; 0.2 to 0.5 mm in India on 10 genotypes, with cortex thickness at the proximal, middle, and distal varying from 1.2 to 3.1 mm). This large genetic variation in peel thickness is associated with the difficulty of peeling, which strongly affects peeling yields and productivity. Industrial or small-scale processors prefer genotypes that are easier to peel and for which labour and drudgery are reduced. For hand peeling, mainly in Africa, yield and productivity are also strongly related to the size and shape of the roots and the presence of constrictions^{68,69}.

The data collected by the RTBfoods project teams allowed to visualize the peeling yield and productivity as a function of root girth, root weight and root length (Figure 2). Although there was a strong increase in peeling yields as a function of increasing root diameter, weight and length, the dispersion of the data highlights other factors such as irregular shape, peel thickness, and/or the presence of constrictions affecting peeling efficiency. Processors know how to predict accurately the ease of peeling. It is mainly the women processors who set the purchase prices of the genotypes to be processed and thereby contribute directly through their processing preferences to the varietal adoption by cassava farmers. These gendered decisions are primarily influenced by the labor-intensive

nature of the tasks, particularly in relation to product yield, processing productivity, and the perceived level of drudgery.




Figure 2. The impact of root morphology on peeling yield (%) and productivity (kg/h/operator) in cassava. **1A)** Root weight (g); **1B)** Root length (cm); **1C)** Root girth (cm). Based on Gari UAC/FSA⁵⁵, Lafun UAC/FSA⁵⁷, and Bâton IITA-Cirad Cameroun⁵⁴ data.




Figure 3. General description of the process to produce gari and average amounts of roots and intermediate products involved in the production of one ton of dry gari in three different regions of West Africa represented by the three different institutes that lead the work in the different regions. Lines on top of each rectangle represent the respective standard deviations.

Gari processing: diagnosis and genotype differences

Accepted Article

Gari processing experiments were conducted by three different institutes in three different locations: Savalou, Colline, UAC/FSA, Benin (Supplementary Table S2), IITA, Osun State, Nigeria (Supplementary Table S3) and Umudike, Abia State, NRCRI, Nigeria (Supplementary Table S4). A varying number of genotypes was evaluated in each location (Supplementary Table S1). Only one genotypes (TME B419) was evaluated in the three locations.

Figure 3 illustrates the entire process to produce one ton of gari and the relative losses through the different stages. Yields were obtained for each operation unit on three different sites -- UAC/FSA in Benin, and NRCRI (Umudike) and IITA (Osun State) in Nigeria. There were large differences in the average tons of fresh roots required to produce one ton of gari: 12.9 (NRCRI), 7.0 (UAC/FSA) and 5.1 t (IITA). To a large extent, as illustrated later, the variation in the amounts of roots required was closely related to the average DMC of the genotypes processed in each location (31.4, 37.4 and 39.2%, respectively).

Root peeling is the first stage (operation unit) in gari processing. About 30-35% of the initial root weight is lost in the (manual) peeling. While shape and size of the roots play an important role in defining the losses during peeling (noted in previous section), this study did not characterize those parameters. As shown later in the article, there is also a remarkably variable required labour from the female processors, depending on each genotype and location. The average time to obtain 100Kg of peeled roots was 3.3 ± 2.0 hours by operator, but the variation was very large ranging from 1.3 (easy to peel and high DMC) to 11.5 hours for the most difficult genotypes (small size, very adherent peel and/or low DMC).

A second stage in which weights are drastically reduced (by about 40%) is during the fermentation, draining and mechanical dewatering (Figure 3). The third important reduction in weights (around 50%) takes place during pulp toasting^{70, 71, 72}.

For each processing site, the operators (mainly women) must carry fresh roots and the residual material after each operation unit in order to proceed to the next one. For the three trials, depending on the genotypes used, to obtain 1000kg of gari, a woman operator had to carry the accumulated weight of 25.7, 48.0 and 19.3 tons respectively in UAC/FSA, NRCRI and IITA.

Figure 3 presents the information for each location, dissecting for each genotype the losses through the individual processing operations in the gari production. The three graphs show a great influence of the location and edaphoclimatic parameters on the quantity of roots necessary to produce one ton of gari. There were large genetic differences in the losses at different stages of gari production by location. The DMCs of local varieties as well as the new hybrids tested in NRCRI were much lower than those observed in UAC/FSA or IITA ($31.4\% \pm 2.3$; $37.4\% \pm 5.9$ and $39.2\% \pm 2.9$ respectively).

The weight of discarded peel was much lower at IITA (954 kg) compared to NRCRI and UAC/FSA (2732 and 1879 kg, respectively). The low DMC of the trial at NRCRI would explain the important losses observed during the dewatering operation in that trial. In this region, locally grown genotypes (Agric, Nwacho and Mgboto) on the right of the plot, have a higher overall processing yield than the new hybrids evaluated. For the IITA and UAC/FSA trials, on the other hand, many of the new hybrids introduced showed higher gari yields than the locally grown landraces.

The standard deviations provided in Figure 4 demonstrate important differences among genotypes at the different stages of gari production. The coefficient of variation ($\sigma/\mu * 100$) can provide an insight into these stages during gari production where there is relatively more variation among genotypes regarding losses. (Supplementary Table S5)

Influence of root DMC on the amount of raw cassava roots required to produce 1000 kg of gari.

Figure 5 depicts the relationship between DMC of raw roots from 57 cassava genotypes and the respective amounts of raw roots required to produce one ton of gari. There was a clear negative correlation between DMC and the amount of fresh roots required ($\text{tons of fresh roots} = 26,078 - 0,525 * \text{DMC}$; $R^2=0.674$). The red dots identify commercial checks often used to produce gari in the regions (Dale, Kati Kati, Salome, AGRIC, Nwageri, Chigazu, Durungwo, Nwocha, Mgboto Umuahia, Honourable 1 and 2, Omoh Local 1 and 2 and Akpu). Commercial checks are scattered from the top left down to the bottom right. TME B419 in green was processed in the three trials.




Figure 6 shows the data generated at Savalou, colline, UAC/FSA, Benin and Osun state, IITA Nigeria on the time required to complete each key step of the process for the gari production for 1000kg of final gari. By far the most time-demanding activities in the production of gari are pulp toasting (53.5%) and root peeling (31.0%). Other activities remained relatively minor in terms of labour demand (6% for pulp sieving, and root washing and dry gari sieving with about 3.7% each). The standard deviations indicate that there is considerable variation among genotypes in relation of their demand of labour to produce one ton of gari. The graph reveals that toasting was less efficient at IITA but, on the other hand, peeling was less time consuming there as compared with UAC/FSA (Figure 6A).




Figure 6. Time invested to produce one ton of gari. **(6A)** Partitioning the total labour required to produce one ton of gari into the different activities of the process. **(6B)** Time invested to produce one ton of gari in the UAC/FSA trial (6B) and IITA trial **(6C)** discriminated by genotypes and activities.

The variety Dale has been ignored in Figure 6 because of its outlying performance. As indicated, there were marked differences in the time invested for each operation among the genotypes evaluated, suggested by the standard deviations in Figure 6A. The plots of Figures 6B and 6C confirm large genetic variation for labour requirements to produce one ton of gari.

The coefficients of variation ($CV = \sigma/\mu * 100$) from UAC/SFA data, for the different operations, were: overall time (hours required to produce 1 ton of gari): 19.3%; peeling: 22.7%; washing: 33.2%; grating: 18.0%; dewatering: 6.8%; pulp sieving: 22.9%; pulp toasting: 23.6 % and dry gari sieving: 27.2% In the case of IITA trials CVs were: overall time: 16.0%; peeling= 15.6%; dewatering: 7.4%; pulp sieving: 27.3% and pulp toasting: 19.7 %

The CVs provide a general appreciation of where variation among genotypes is particularly important for a given activity and point out results that require further exploration. For example, in the IITA trial (Figure 6B), root peeling was relatively more uniform ($CV = 15.6\%$) than pulp sieving ($CV = 27.3\%$). I1 and Honourable 2 required more than 40 hours in pulp sieving whereas Akpu and I9 required less than 20 hours. Understanding why these genotypes are so contrasting for pulp sieving would help breeders to use more efficient selection approaches.

Figures 6B and 6C also provide information on DMC for each genotype. U3 has low DMC and requires considerable labour input. RB 8509 (or U9) had the highest DMC and was second lowest clone in labour requirement (Figure 6C). There is some association between DMC and labour requirements. On the other hand, U2 with an intermediate DMC (37.7%), required considerably less labour than U1, which had excellent DMC (42.3%). This last clone required 80 more hours than the average of 523 hours (excluding Dale) for UAC/FSA.




Figure 7. UAC/FSA trial Relationship between dry matter content of activities (Figure 6A). Therefore, the relationship between DMC and the time required to carry the roots and the total labour requirements to produce one ton of gari.

Influence of DMC on workload to process one ton of gari.

Labour hours required to produce one ton of gari versus DMC of raw roots (UAC/FSA data) was plotted in Figure 7. Dale is a locally grown variety in Benin with low DMC and small roots. If this genotype (clearly outlying) is included, the regression analysis comparing DMC and labour hours results in a $R^2 = 0.55$. However, when Dale was ignored the coefficient of determination was drastically reduced ($R^2 = 0.26$). The large range of time required to produce one ton of gari (ignoring Dale) goes from 400 to 800 hours. Relatively little of that variation is explained by DMC, as also indicated by data from Figures 6B and 6C. There are, therefore, other factors that influence the variation in the labour required for processing one ton of gari such as shape and size of roots, which has a huge impact on the peeling operation.

Root peeling, pulp sieving and pulp toasting are the most labour-demanding out these activities was further analyzed and depicted in Supplementary Figure S1. Data for this figure included trials at UAC/FSA and IITA. The most important influence of DMC was found to be for pulp sieving. However, there is a large dispersion of data (Supplementary Figure S1). The negligible influence of DMC on labour requirements for pulp toasting is not surprising. Most of the effect of DMC would be on stages prior to the toasting. When the material being processed reaches that stage, they already have a much more uniform (and increased) DMC because of the mechanical de-watering that takes place after fermentation.

Summary of the influence of DMC on gari production.

A summary of the relative importance of DMC in gari yield and labour requirements has been consolidated in Figure 8. In addition, this figure provides another important parameter related to the drudgery of gari production: the total amount of mass that women had to move throughout the entire process. The highest coefficient of determination was observed for the regression of kg of mass movement per kg of gari produced, on DMC ($R^2 = 0.65$).




Figure 8: Impact of raw material dry matter content (%) on gari yield (8A), operator time (8B) and drudgery (8C)

Figure 8A shows a positive correlation between gari yield and the raw root dry matter content. However, the dry matter content doesn't impact significantly the time spent for processing (Figure 8B). Other parameters may have to be considered for their impact on process productivity, such as: root peeling ability, draining behavior during fermentation, and draining, sieving ability linked to fiber content. A clear negative correlation can be observed between mass movement and dry matter content (Figure 8C). The dry matter content is thus an important trait to be considered in order to reduce labour requirements and thus the often related drudgery in gari processing.

Key findings and Conclusions

This review shows a strong genetic influence on processing yield, operators' efficiency, workload and fatigue of RTB processors (mainly women, young children and elderly). Processing yield depends not only on varieties but also on the level of complexity of the processes. Indeed, low processing yields are the result of complicated operations with many steps to reduce water content, enhance the shelf-life of products and, in the case of cassava, detoxify it if necessary.

Thus, more complex processes tend to increase the likelihood of perceived drudgery, reduce global processing yields but, on the other hand, lengthen the shelf life of food products. In that sense, complexity, and concomitantly some degree of drudgery, is the price to pay to increase the shelf life of the end-product coming from the same raw material.

Reduced processing yields do not only have a direct negative economic impact on the value addition, but also have an indirect (but strongly correlated) effect on the accumulated weight carried through

the process. By combining all the data sets for the different food product profiles, Figure 9 illustrated the clear association between weights carried out and processing yields.




Figure 9: Weight (kg) carried out by operator per kg of end product according processing yield (% w.b). Based on Gari CIRAD/UAC-FSA⁵⁵, Gari IITA⁶⁰, Dry fufu CIRAD/UAC-FSA⁵⁷, Dry fufu CIRAD/IITA⁵⁶, Bâton CIRAD/IITA⁵⁴, Wet Fufu IITA⁶⁰ and Boiled cassava UAC/FSA⁵⁹ data.

Furthermore, this review addressed a clear gender dimension as processing work is often dominated by women because of existing norms that often push women into monotonous and drudgery related tasks. As social impact through social and gender inclusiveness is a particular outcome aimed by RTB breeding⁷³ and part of the sustainable development goals, increasing productivity and limiting perceived drudgery in RTB processing, is therefore crucial within the empowerment of women from below. Gender transformative approaches that aim to change gender roles are important but often slow and are not always able to address the concrete working conditions, livelihoods and their context (including the existing norms) that women are largely dependent on^{74, 75} to make a living and increase their income and independence.

A special effort should be made by RTB breeders to evaluate the yields of the final product and the additional fatigue that could be brought by the introduction of new genotypes. This is particularly necessary for the more complex processed products, especially those from cassava, because of the additional need to detoxify them, reduce weight of the marketable product by removing water and the need to increase shelf live. Moreover, addressing labour requirements and associated drudgery related to processing is not enough because breeders must also consider consumer preferences as well. There is a large diversity of requirements for granulated and soaked products on one hand, and pasty and dough products such as pounded yam and matooke, on the other. All these products are obtained after complex processing steps^{27,34,35,36}.

The processing efficiency or productivity that determines the amount of labour required from processors and the related drudgery are key factors in the varietal adoption process. The processors, who are often the decision makers for the purchase of raw materials, strongly contribute to the creation of a market segment for new genotypes or to their rejection depending on the difficulty of processing and/or the food product yields obtained. The views of consumers, traders and growers have already been incorporated into the definition of breeding goals. This review highlights, however, the critical importance of processors in the final varietal adoption⁷⁶. Their perception of each genotype

provides relevant information that must be integrated upstream in the breeding pipeline to increase the chances of success of new varieties.

Acknowledgements

In memoriam of our colleague and friend Geneviève Fliedel, Ph.D., Food technologist at CIRAD who suggested and contributed to this work and for the passion and enthusiasm that characterised her work as a researcher.

The authors are grateful to the grant opportunity INV-008567 (formerly OPP1178942): Breeding RTB Products for End User Preferences (RTBfoods), to the French Agricultural Research Centre for International Development (CIRAD), Montpellier, France, by the Bill & Melinda Gates Foundation (BMGF): <https://rtbfoods.cirad.fr>.

The authors thank the various partner institutions of the RTBfoods project for allowing the sharing of additional data not contained in the diagnostic reports mentioned in the bibliographic list.

The final proofreading of the manuscripts by Clair Hershey improved greatly the quality of this manuscript.

Credit Authorship Contribution Statement

Conceptualization: AB, DD, TM, FKA, GF, LF, BT

Data curation: HC, BA, DO, DOO, DO, AA, BO, MO, SN, NO, LA

Formal Analysis: AB, DD, HC, AB, BT, BO, SN, LA

Investigation: AB, AB, BT, DOO, MO, SN, NO, LA, DD

Methodology: AB, GF, MT, LA, LF, DD

Supervision: AB, BT, TM, LA, GF, LF, DD

Writing: Original draft: AB, HC, DD

Writing reviewing and editing: AB, HC, DD, LF, BT

Project administration: DD

Editorial Policies and Ethical Considerations

The authors declare no conflict of interests. Authors have agreed to this publication. All funding has been acknowledged.

References

1. Nweke FI and Enete A, Gender Surprises in Food Production, Processing, and Marketing with Emphasis on Cassava in Africa. COSCA – Collaborative Study of Cassava in Africa. Working Paper n°19 · International Institute of Tropical Agriculture IITA, Ibadan, Nigeria (1999). https://www.google.fr/books/edition/Gender_Surprises_in_Food_Production_Proc/wYHl6osg-ToC?hl=fr&gbpv=1&dq
2. Doss C, Meinzen-Dick R, Quisumbing A and Theis S, Women in agriculture: Four myths. *Glob. Food Secur.*, **16**:69-74 (2018). <https://doi.org/10.1016/j.gfs.2017.10.001>
Nweke FI, Cassava Processing in Sub-Saharan Africa: the Implications for Expanding Cassava Production. *Outlook Agric.* **23**(3):197-205 (1994). <https://doi.org/10.1177/003072709402300307>
3. Curran S, Leigh-Anderson C, Gugerty MK, Cook J, Yorgy G and Gockel R, Gender and cropping: cassava in Sub-Saharan Africa. Evans School Policy Analysis and Research (EPAR). p. 6. <https://evans.uw.edu/policy-impact/epar/research/gender-cropping-sub-saharan-africa-cassava>
4. Teeken B, Olaosebikan O, Haleegoah J, Oladejo E, Madu T, Bello A *and al.*, Cassava trait preferences of men and women farmers in Nigeria: implications for breeding. *Econ. Bot.*, **20**, 1– 15. <https://doi.org/10.1007/s12231-018-9421-7>
5. Forsythe L, Martin AM and Posthumus H, Cassava market development: a path to women's empowerment or business as usual? *Food Chain*, **5**(1-2): 11-27(2015). <https://doi.org/10.3362/2046-1887.2015.003>
6. Forsythe L, Posthumus H and Martin A, A crop of one's own? Women's experiences of cassava commercialization in Nigeria and Malawi. *J. gen., agric. food secur.* , **1**(2): 110-128 (2016). <https://doi.org/10.22004/ag.econ.246033>
7. Nweke FI, Spencer DSC and Lynam JK, The Cassava transformation: Africa's best-kept secret. Michigan State University Press: East Lansing, MI, 273 p. (2012). <https://msupress.org/9781609172831/the-cassava-transformation/>
8. FAO, The State of Food and Agriculture 2010-2011: Women in Agriculture: Closing the Gender Gap for Development. Food and Agriculture Organization of the United Nations, Rome, Italy (2011). <https://www.fao.org/3/i2050e/i2050e.pdf>
9. Abass AB, Awoyale W, Ogundapo A, Oluwasoga O, Nwaoliwe G, Oyelekan J *and al.*, Adoption of improved cassava varieties by processors is linked to processing characteristics and products biophysical attributes, *J. Food Process. Preserv.*, **46**(3):e16350 (2022). <https://doi.org/10.1111/jfpp.16350>
10. Bishop-Sambrook C, How to do Reducing rural women's domestic workload through labour-saving technologies and practices - Gender, targeting and social inclusion. IFAD, Rome, Italy (2016). <https://www.ifad.org/documents/38714170/db859c93-9066-411a-ad40-a0204c98351c> ; <https://www.ifad.org/documents/10180/c86179dd-ad6d-4622-864a-ed590016250d>
11. Polar V and Demont M, Designer Crops of the Future Must Be Better Tailored for Women in Agriculture - Nutrition levels and taste should be considered along with traits such as higher yields. *Sci. Am.* **327**(2):Jul 25, (2022). <https://www.scientificamerican.com/article/designer-crops-of-the-future-must-be-better-tailored-for-women-in-agriculture/>
12. Weltzien E, Rattunde F, Christinck A and Ashby J, Gender and farmer preferences for varietal traits: evidence and issues for crop improvement. *Plant Breed. Rev.*, **43**:243-278. (2020). <https://doi.org/10.1002/9781119616801.ch7>
13. Adenugba AA and John P, Hazardous conditions of women in gari processing industry in Ibadan, South West, Nigeria. *J. Educ. Soc. Res.*, **4**(3):511. (2014). <https://doi.org/10.5901/jesr.2014.v4n3p511>

15. Samuel TM, Igbeka JC and Kolawole OP, A survey of improved gari frying methods. *Int. J. Food Eng.*, **6**(2):1-12. (2010). <https://doi.org/10.2202/1556-3758.1473>
16. Okwor TJ, Ozoh OB, Okonkwo IJ and Osibogun A. Occupational exposure to particulate matter from biomass smoke and its relationship to respiratory symptoms and pulmonary function among rural women involved in cassava processing in Nigeria. *Open J. Prev. Med.*, **7**: 41–54(2017). <https://doi.org/10.4236/ojpm.2017.73004>
17. Vemireddy V and Choudhary A, Labor-saving technologies designed for women can reduce their drudgery. CGIAR GENDER PLATFORM Evidence Explainer. Nairobi, Kenya: CGIAR GENDER Platform (2021). <https://hdl.handle.net/10568/114807>.
18. Bello A, Afolabi A, Olaosebikan O, Edughaen G, Dufour D, Bouniol A and al., Genetic and environmental effects on processing productivity, gari yields and eba acceptability: drudgery of women's work. *J. Sci. Food Agric.* Same Special Issue: Tropical roots, tubers and bananas: New breeding tools and methods to meet consumer preferences. Manuscript JSFA-23-0360, under review. (2023).
19. Mrunalini A and Snehalatha Ch (2010) Drudgery experiences of gender in crop production activities. *J. Agric. Sci.*, **1**(1), 49-51. <https://doi.org/10.1080/09766898.2010.11884654>
20. Bal, SK, Shivani S and Kaur H (2013). Assessment of drudgery experience of rural women while performing different farm operations. *Adv. Res. J. Soc. Sci.*, **4**(1), 68-71. http://researchjournal.co.in/upload/assignments/4_68-71.pdf
21. Isaksen J (2000). Constructing meaning despite the drudgery of repetitive work. *J. Humanist. Psychol.*, **40**(3), 84-107. <https://doi.org/10.1177/0022167800403008>
22. Luna J, Dufour D, Tran T, Pizarro M, Calle F, García Domínguez M, and al, Post-harvest physiological deterioration in several cassava genotypes over sequential harvests and effect of pruning prior to harvest. *Int. J. Food Sci. Technol.*, **56**(3):1322-1332 (2021). <https://doi.org/10.1111/ijfs.14711>
23. Ravi V, Aked J and Balagopalan C, Review on tropical root and tuber crops I. Storage methods and quality changes. *Crit Rev Food Sci Nutr*, **36**(7):661-709 (1996). <https://doi.org/10.1080/10408399609527744>
24. Kouamé AC, Kouassi KN, N'dri YD and Amani NG, Glycaemic index and load values tested in normoglycemic adults for five staple foodstuffs: pounded yam, pounded cassava-plantain, placali, attieke and maize meal stiff porridge. *Nutrients*, **7**(2):1267-1281 (2015). <https://doi.org/10.3390/nu7021267>
25. Marimo P, Akankwasa K, Khamila S, Tinyoro SE, Bouniol A, Ndagire L, and al., RTBfoods Participatory Processing Diagnosis of Matooke in Uganda. RTBfoods/CIRAD, Kampala, Uganda, 38 p. (2021). <https://doi.org/10.18167/agritrop/00615>
26. Otegbayo B, Madu T, Oroniran O, Chijioke U, Fawehinmi O, Okoye B, and al., End-user preferences for pounded yam and implications for food product profile development. *Int. J. Food Sci. Technol.*, **56**(3):1458-1472 (2021). <https://doi.org/10.1111/ijfs.14770>
27. Dufour D, Hershey C, Hamaker BR and Lorenzen J, Integrating end-user preferences into breeding programmes for roots, tubers and bananas. *Int. J. Food Sci. Technol.*, **56**(3):1071-1075 (2021). <https://doi.org/10.1111/ijfs.14911>
28. Bokanga M, Distribution of cyanogenic potential in cassava germplasm. In: International Workshop on Cassava Safety, edited By Bokanga M, Essers AJA, Poulter N, Rosling H and Tewe O, WOCAS/ISHS /ISTRIC, The Netherland, *Acta Hort.*, **375**: 117-124. (1994). https://www.actahort.org/books/375/375_9.htm
29. Sánchez T, Salcedo E, Ceballos H, Dufour D, Mafla G, Morante N and al., Screening of starch quality traits in cassava (*Manihot esculenta* Crantz). *Starch/Stärke*, **61**(1):12–19 (2009). <http://dx.doi.org/10.1002/star.200800058> ; Erratum: *Starch/Stärke*, **61**(5):310 (2009). <http://dx.doi.org/10.1002/star.200990027>

30. Chiwona-Karltun L, Brimer L, Kalenga Saka JD, Mhone AR, Mkumbira J, Johansson L *and al.*, Bitter taste in cassava roots correlates with cyanogenic glucoside levels. *J. Sci. Food Agric.*, **84**:581-90 (2004). <https://doi.org/10.1002/jsfa.1699>
31. Nambisan B, Strategies for elimination of cyanogens from cassava for reducing toxicity and improving food safety. *Food Chem. Toxicol.*, **49**(3):690-693. (2011). <https://doi.org/10.1016/j.fct.2010.10.035>
32. FAO/WHO Maximum levels for hydrocyanic acid in cassava and cassava products. Food standards programme codex committee on contaminants in foods, Seventh session, Moscow, Russian Federation, 8 – 12 april 2013. Agenda Item 10, CX/CF 13/7/10 (2013). https://www.fao.org/fao-who-codexalimentarius/sh-proxy/en/?lnk=1&url=https%253A%252F%252Fworkspace.fao.org%252Fsites%252Fcodex%252Fshared%252Fdocuments%252Farchive%252Fmeetings%252Fccccf%252Fccccf7%252Fcf07_11e.pdf
33. Montagnac JA, Davis CR. and Tanumihardjo SA, Processing techniques to reduce toxicity and Antinutrients of Cassava for Use as a Staple Food. *Compr. Rev. Food Sci. Food Saf.*, **8**:17-27 (2009). <https://doi.org/10.1111/j.1541-4337.2008.00064.x>
34. Béchoff A, Tomlins K, Fliedel G, Becerra Lopez-lavalle LA, Westby A, Hershey C *and al.*, Cassava traits and end-user preference: Relating traits to consumer liking, sensory perception, and genetics. *Crit Rev Food Sci Nutr*, **58**(4):547-567 (2018). <https://doi.org/10.1080/10408398.2016.1202888>
35. Lancaster PA, Ingram JS, Lira MY and Coursey DG, Traditional cassava-based foods: Survey of processing techniques. *Econ. Bot.*, **36**:12-45 (1982). <https://doi.org/10.1007/BF02858697>
36. Westby A, Cassava utilization, storage and small-scale processing. In: *Cassava: Biology, Production and Utilization*, edited by Hillocks RJ., Thresh JM and Bellotti AC. CABI Publ., New York. Pp. 281–300 (2002). http://ciat-library.ciat.cgiar.org/articulos_ciat/cabi_17ch14.pdf
37. Forsythe L, Tufan H, Bouniol A, Kleih U and Fliedel G, An interdisciplinary and participatory methodology to improve user acceptability of root, tuber and banana varieties. *Int. J. Food Sci. Technol.*, **56**(3):1115-1123 (2021). <https://doi.org/10.1111/ijfs.14680>
38. Forsythe L, Olaosebikan O, Teeken B, Ngoh Newilah G, Mayanja S, Nanyonjo AR, *and al.*, A case of inter/trans-disciplinarity and collaborative decision making: the co-construction of endered Food Product Profiles. *J. Sci. Food Agric.* Same Special Issue: Tropical roots, tubers and bananas: New breeding tools and methods to meet consumer preferences. Manuscript JSFA-23-0282, under review. (2023).
39. Fliedel G, Bouniol A, Kleih U, Tufan H and Forsythe L, RTBfoods Step 3: Participatory Processing Diagnosis and Quality Characteristics, RTBfoods/CIRAD, Montpellier, France, 29 p. (2018). <https://doi.org/10.18167/agritrop/00570>
40. Ndjouenkeu R, Ngoualem KF, Teeken B, Okoye B, Madu T, Olaosebikan OD, *and al.*, From cassava to gari: mapping of quality characteristics and end-user preferences in Cameroon and Nigeria. *Int. J. Food Sci. Technol.*, **56**(3):1223-1238 (2021). <https://doi.org/10.1111/ijfs.14790>
41. Chijioke U, Madu T, Okoye B, Ogunka AP, Ejechi M, Ofoeze M, *and al.*, Quality attributes of fufu in South-East Nigeria: guide for cassava breeders. *Int. J. Food Sci. Technol.*, **56**(3):1247-1257 (2021). <https://doi.org/10.1111/ijfs.14875>
42. Ferris R. S. B., Ortiz R., Vuylsteke D., Fruit quality evaluation of plantains, plantain hybrids, and cooking bananas. *Postharvest Biol. Technol.* **15**(1):73– 81(1999). [https://doi.org/10.1016/S0925-5214\(98\)00067-2](https://doi.org/10.1016/S0925-5214(98)00067-2)
43. Gibert O, Dufour D, Giraldo A, Sánchez T, Reynes M, Pain JP *and al.*, Differentiation between Cooking Bananas and Dessert Bananas. 1. Morphological and Compositional Characterization of Cultivated Colombian Musaceae (*Musa* sp.) in Relation to Consumer Preferences. *J. Agric. Food Chem.* **57**(17): 7857–7869 (2009). <https://doi.org/10.1021/jf901788x>

44. Mäder J., Rawel H. Kroh L.W., Composition of Phenolic Compounds and Glycoalkaloids α -Solanine and α -Chaconine during Commercial Potato Processing. *J. Agric. Food Chem.* **57**(14), 6292–6297 (2009). <https://doi.org/10.1021/jf901066k>
45. Pathak P. D., Mandavgane S. A., Puranik N. M., Jambhulkar S. J., Kulkarni B. D. Valorization of potato peel: a biorefinery approach, *Crit. Rev. Biotechnol.* **38**(2), 218-230, (2018). <https://doi.org/10.1080/07388551.2017.1331337>
46. Visalakshi C, Shirly Raichal Anil, Sheela MN, Vivek Hegde, Jyothi AN and Sreekumar J, Genetic variation for important processing traits and identification of potential parents for developing hybrid clones suitable for processing in sweet potato (*Ipomoea batatas* L Lam). *S. Afr. J. Bot.*, 141:255-264 (2021). <https://doi.org/10.1016/j.sajb.2021.04.005>
47. Adetan DA, Adekoya LO, Aluko OB, Theory of a mechanical method of peeling cassava tubers with knives. *Int Agrophys*, **20**(4), 269–276. (2006). <http://www.international-agrophysics.org/pdf-106611-37453?filename=Theory%20of%20a%20mechanical.pdf>
48. Bouniol A, Adinsi L, Honfozo L, Hounhouigan J, Fliedel G and Akissoe N RTBfoods Participatory Processing Diagnosis of Boiled Yam in Benin. RTBfoods/CIRAD, Cotonou, Benin, 27 p. (2021). <https://doi.org/10.18167/agritrop/00627>
49. Madu T, Ofoeze M, Onyemauwa N and Okoye B, RTBfoods Participatory Processing Diagnosis of Boiled Yam in Nigeria. RTBfoods/CIRAD, Umudike, Nigeria, 17 p. (2021). <https://doi.org/10.18167/agritrop/00619>
50. Otegbayo B, Oroniran O, Fawehinmi O and Tanimola A, RTBfoods Participatory Processing Diagnosis of Pounded Yam in Nigeria. RTBfoods/CIRAD, Iwo, Nigeria, 22 p. (2021). <https://doi.org/10.18167/agritrop/00616>
51. Tinyiro SE, Nakitto M, Kisakye S, Bamwirire D, Mirembe A, Okonya, J. *et al.*, RTBfoods Participatory Processing Diagnosis of Boiled Sweetpotato in Uganda. RTBfoods/CIRAD, Kampala, Uganda, 25p. (2021). <https://doi.org/10.18167/agritrop/00623>
52. Nakitto M, Tinyiro SE, Kisakye S, Bamwirire D, Mirembe A, Okonya J *et al.*, RTBfoods Participatory Processing Diagnosis for Boiled Potato in Uganda. RTBfoods/CIRAD, Kampala, Uganda, 21 p. (2021). <https://doi.org/10.18167/agritrop/00622>
- Ngoh Newilah G, Kendine Vepowo C and Takam Ngouno A, RTBfoods Participatory Processing Diagnosis for Boiled Plantain in Cameroon, RTBfoods/CIRAD, Yaoundé, Cameroon, 27 p. (2021). <https://doi.org/10.18167/agritrop/00626>
54. Bouniol A, Prin L, Hanna R, Fosto A and Fliedel G, Assessment of the processability of improved cassava varieties into a traditional food product (“baton” or “chikwangue”) in Cameroon. RTB Working Paper n°2012-1, CGIAR research program on Roots, Tubers and Bananas (RTB), 28 p. (2017). <http://agritrop.cirad.fr/595635/>
55. Bouniol A, Adinsi L., Hotegni F, Delpech A, Akissoé N, Fliedel G., *et al.*, Assessment of the processing ability of improved cassava varieties into a traditional food product (Gari) in Benin. Savalou. RTB Working Paper n°2012-1, CGIAR research program on Roots Tubers & Bananas (RTB), 34 p. (2020). <https://agritrop.cirad.fr/597596/>
56. Bouniol A, Ospina MA, Fosto A, Hanna R and Dufour D, Assessment of the processing ability of improved cassava varieties into a traditional food product (FuFu) in Cameroon. Bertoua, RTB Working Paper n°2012-1, CGIAR Research Program on Roots, Tubers and Bananas (RTB), 24 p. (2017). <https://agritrop.cirad.fr/597597/>
57. Bouniol A, Adinsi L, Padonou SW, Hotegni F, Gnanvossou D, Akissoé N, *et al.*, Assessment of the processing ability of improved cassava varieties into a traditional food product (Lafun) in Benin. Kétou. RTB Working Paper n°2012-1, CGIAR Research Program on Roots, Tubers and Bananas (RTB), 29 p. (RTB Working Paper, 2012-1) (2019). <https://agritrop.cirad.fr/597594/>

58. Hamba S, Nanyonjo AR, Kanaabi M and Kawuki RS, RTBfoods Participatory Processing Diagnosis of Boiled Cassava in Uganda. RTBfoods/CIRAD, Kampala, Uganda, 25 p. (2021). <https://doi.org/10.18167/agritrop/00625>
59. Bouniol A, Adinsi L, Honfozo L, Hounhouigan J and Akissoe N, RTBfoods Participatory Processing Diagnosis of Boiled Cassava in Benin. RTBfoods/CIRAD, Cotonou, Benin, 24p. (2021). <https://doi.org/10.18167/agritrop/00624>
60. Teeken B, Agbona A, Bello A, Olaosebikan O, Alamu E, Adesokan M, *et al.*, Understanding cassava varietal preferences through pairwise ranking of *gari-eba* and *fufu* prepared by local farmer-processors. *Int. J. Food Sci. Technol.*, **56**(3):1258-1277 (2021). <https://doi.org/10.1111/ijfs.14862>
- Madu T, Okoye B, Onyemauwa N and Ofoeze M, RTBfoods Participatory Processing Diagnosis of Gari/Eba in Nigeria. RTBfoods/CIRAD, Umudike, Nigeria, 21 p. (2021). <https://doi.org/10.18167/agritrop/00614>
- Madu T, Okoye B, Onyemauwa N and Ofoeze M, RTBfoods Participatory Processing Diagnosis of Fufu in Nigeria. RTBfoods/CIRAD, Umudike, Nigeria, 18 p. (2021). <https://doi.org/10.18167/agritrop/00620>
63. Diby N'Nan AS, Ebah Djedji C, Kanon L, Yapi E, Bouniol A, Bechoff A and al., Participatory processing diagnosis for attiéke in Cote d'Ivoire. Surveys on Trait Preferences, WP1 - Activity 4. Abidjan : RTBfoods Project-CIRAD, 26 p. (2021). <https://doi.org/10.18167/agritrop/00724>
64. Nweke FI, Lutete D, Dixon AGO, Ugwu BO, Ajobo O, Kalombo N and al., Cassava production and processing in the Democratic Republic of Congo. Collaborative Study on Cassava in Africa (COSCA) Working Paper N°22, 87-93 (2000). https://books.google.fr/books?id=A4VY_jli7kAC&lpg=PP1&hl=fr&pg=PP1#v=onepage&q&f=false
65. Pérez JC, Lenis JI, Calle F, Morante N, Sánchez T, Debouck D *and al.* Genetic variability of root peel thickness and its influence in extractable starch from cassava (*Manihot esculenta* Crantz) roots. *Plant Breed.*, **130**: 688-693 (2011). <https://doi.org/10.1111/j.1439-0523.2011.01873.x>
66. Kawuki RS, Ferguson M., Labuschagne MT, Herselman L, Orone J, Ralimanana an al., Variation in qualitative and quantitative traits of cassava germplasm from selected national breeding programmes in sub-Saharan Africa, *Field Crops Res.*, **122**(2), 151-156 (2011). <https://doi.org/10.1016/j.fcr.2011.03.006>.
67. Krishnakumar T, Sajeev MS, Pradeepika C, Namrata AG, Sanket JM, Jeevarathinam G *and al.*, Physical and mechanical properties of cassava (*Manihot esculenta* Crantz) cultivars: Implications for the design of mechanical peeling machines. *J. Food Process Eng.*, **45**(6), e13923 (2022). <https://doi.org/10.1111/jfpe.13923>
- Hauser S, Bakelana TZ, Bungu D and Mwangu M, Cassava root yield losses caused by root shape, rot and peeling and cut-off for processing of four cassava varieties on different soils in DR Congo. *Advances in Nutrition and Food Science*, **5**:1-10. (2020). <https://cgspace.cgiar.org/handle/10568/119338>
69. Bello A., Olaosebikan O., Osunbgbade A., Teeken B. and Bouniol A., Participatory processing diagnosis for Gari/Eba in Nigeria. Understanding the drivers of trait preferences and the development of multi-user RTB product profiles, WP1, step 3. Ibadan : RTBfoods Project-CIRAD, 34 p. (2020). <https://doi.org/10.18167/agritrop/00621>
70. Ezeocha, C.V., Ihesie, L.C. & Kanu, A.N., Comparative evaluation of toasting variables and the quality of gari produced by different women in Ikwuano LGA, Abia State, Nigeria. *J. Food Process. Preserv.*, **43**, e14060. (2019). <https://doi.org/10.1111/jfpp.14060>
71. Dahdouh L, Escobar A, Rondet E, Ricci J, Fliedel G, Adinsi L, *and al.*, Role of dewatering and roasting parameters on the quality of handmade gari. *Int. J. Food Sci. Technol.*, **56**(3):1298-1310 (2021). <https://doi.org/10.1111/ijfs.14745>
72. Escobar A, Rondet E, Dahdouh L, Ricci J, Akissoé N, Dufour D and al., Identification of robust versus critical unit operations in the elaboration of gari quality. *Int. J. Food Sci. Technol.*, **56**(3):1311-1321 (2021). <https://doi.org/10.1111/ijfs.14857>

73. Polar V, Teeken B, Mwende J, Marimo P, Tufan HA, Ashby JA and al., Building Demand-Led and Gender-Responsive Breeding Programs. *In*: Thiele G, Friedmann M, Campos H, Polar V and Bentley JW (eds). Root, Tuber and Banana Food System Innovations. Springer, Cham. (2022). https://doi.org/10.1007/978-3-030-92022-7_16
74. Adams, R. (2008). Empowerment, participation and social work; 4th edition. (4 ed.) (Practical social work series). Palgrave Macmillan.
75. Njuki J, Eissler S, Malapit H, Meinzen-Dick R, Bryan E and Quisumbing A, A Review of Evidence on Gender Equality, Women's Empowerment, and Food Systems. *In*: von Braun J, Afsana K, Fresco LO, Hassan M.H.A. (eds) Science and Innovations for Food Systems Transformation. Springer, Cham. (2023). https://doi.org/10.1007/978-3-031-15703-5_9
76. Thiele G, Dufour D, Vernier P, Mwangi ROM, Parker M, Schulte Geldermann E *and al.*, Review of varietal change in roots, tubers and bananas: consumer preferences and other drivers of adoption and implications for breeding. *Int. J. Food Sci. Technol.*, **56**(3):1076-1092. <https://doi.org/10.1111/ijfs.14684>