



## King's Research Portal

DOI:

[10.1146/annurev-physiol-042222-022728](https://doi.org/10.1146/annurev-physiol-042222-022728)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Brunello, E., & Fusi, L. (2023). Regulating Striated Muscle Contraction: Through Thick and Thin. *Annual Review of Physiology*, 86. <https://doi.org/10.1146/annurev-physiol-042222-022728>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

# Regulating Striated Muscle Contraction: Through Thick and Thin

Volume 86 of the *Annual Review of Physiology*

**Elisabetta Brunello<sup>1</sup>, Luca Fusi<sup>1,2</sup>**

<sup>1</sup>Randall Centre for Cell and Molecular Biophysics, School of Basic and Medical Biosciences and BHF Centre of Research Excellence, King's College London, London SE1 1UL, UK.

<sup>2</sup>Centre for Human and Applied Physiology, School of Basic and Medical Biosciences, King's College London, London SE1 1UL, UK.

Keywords: Muscle regulation, myosin, troponin, Length-dependent activation, cardiac muscle, skeletal muscle.

## **Abstract (149/150)**

Force generation in striated muscle is primarily controlled by structural changes in the actin-containing thin filaments triggered by an increase in intracellular calcium concentration. However, recent studies have elucidated a new class of regulatory mechanisms based on the myosin-containing thick filament that control the strength and speed of contraction by modulating the availability of the myosin motors for the interaction with actin. This review summarizes the mechanisms of thin and thick filament activation that regulate the contractility of skeletal and cardiac muscle. A novel dual-filament paradigm of muscle regulation is emerging, in which the dynamics of force generation depends on the coordinated activation of thin and thick filaments. We highlight the interfilament signaling pathways based on titin and myosin-binding protein-C that couple thin and thick filament regulatory mechanisms. This dual-filament regulation mediates the length-dependent activation of cardiac muscle that underlies the control of the cardiac output in each heartbeat.

## **1. Introduction**

The contraction of striated muscle is generated by relative sliding of actin-containing thin filaments and myosin-containing thick filaments driven by transient interactions of the myosin motors with actin coupled to the hydrolysis of ATP. Regular arrays of thin and thick filaments are organized into  $\sim 2\mu\text{m}$ -long contractile units, called sarcomeres (Fig. 1), which are arranged in series along the length of the muscle cell. The start signal for contraction is provided by the action potential that travels down transverse invaginations of the sarcolemma (T-tubules) into the interior of the muscle cell, activating the voltage-gated calcium channels, known as dihydropyridine receptors (DHPR) (1, 2). In skeletal muscle the DHPRs are mechanically linked to the calcium channels situated in the membrane of the sarcoplasmic reticulum, called ryanodine receptors (RyR). The voltage-induced activation of the DHPRs directly triggers the opening of the RyRs, causing a large increase in the

cytoplasmic free calcium concentration within 2 ms from the action potential (3), followed by the recovery of the calcium concentration before the electrical stimulation due to the  $\text{Ca}^{2+}$  reuptake in the sarcoplasmic reticulum driven by the calcium pumps. Instead, in cardiac muscle the activation of the DHPRs, that are uncoupled from the RyRs on the sarcoplasmic reticulum, allows extracellular calcium ions to flow into the myoplasm. These calcium ions bind and open the RyRs, triggering the release of the calcium ions stored in the sarcoplasmic reticulum, a mechanism known as calcium-induced calcium release (1). Due to the differences in the mechanism of calcium release from intracellular stores, the increase in free calcium concentration after a single action potential in the cardiac muscle cell is ten times smaller and slower than that in skeletal muscle (Fig. 1A) (4, 5).

Until recently the classical paradigm for the regulation of the interaction between myosin and actin in the muscle sarcomere was exclusively focused on the structural changes in the regulatory proteins of the thin filament, troponin and tropomyosin, triggered by the transient increase in intracellular calcium concentration following the electrical stimulation of muscle (6). In the classical view of thin-filament regulation of muscle contractility, troponin and tropomyosin are regarded as “gate-keepers” of the thin filament, switching off the thin filament at low calcium concentration in resting muscle and allowing myosin to bind actin when calcium ions are bound to troponin. Therefore, according to this view, the kinetics of force generation and relaxation in the force twitch triggered by the action potential are exclusively accounted for by the dynamics of the calcium transient and of the associated structural changes in the thin filament (Fig. 1C).

However, studies in the last decade shed light on a novel class of regulatory mechanisms based on the thick filament, that regulate the contractility of skeletal and heart muscle in addition to the classical thin-filament based regulation (7-9). The evidence that the myosin motors at low calcium concentrations are switched off onto the surface of the thick filament backbone in an inhibited conformation, called interacting-heads motif (10, 11), challenged the classical paradigm of muscle regulation, suggesting that the myosin motors are not immediately available for the interaction with actin following electrical stimulation. Since then, it became clear that the contraction of striated muscle requires not only the activation of the thin filament by calcium, but also the removal of myosin inhibition on the thick filament, via independent mechanisms which do not necessarily require calcium (Fig. 1C) (12, 13). Similarly, mechanical relaxation of muscle triggered by the decrease in myoplasmic calcium concentration cannot be solely explained by the inactivation of the thin filament triggered by the removal of calcium from troponin. The mechanism of muscle relaxation is still unclear, but its rate depends primarily on the kinetics of detachment of myosin motors from actin, which is likely controlled by the coordinated inactivation of thin (14) and thick filaments (15-17) and by the inter-sarcomere dynamics (17, 18). Therefore, according to the myosin-based paradigm of muscle regulation emerged from these studies, the thick filament with its regulatory proteins Myosin-Binding Protein-C (MyBP-C) and titin modulates the number of myosin motors available for the interaction with actin during contraction. This review is focused on the regulatory mechanisms in thin and thick filaments that underpin the control of the dynamics of force generation in skeletal and cardiac muscle, and on the coordination of thin and thick filament-based regulatory pathways in a novel dual-filament mechanism of regulation of muscle contraction.

## **2. Thin filament-based regulation in striated muscle.**

### **2.1 Structural changes in the thin filament induced by calcium and myosin motors.**

In muscle cells at low calcium concentration the tropomyosin strand on the thin filament is held by troponin in a position that blocks the myosin binding sites on actin, inhibiting contraction (Fig. 1C). Following the electrical stimulation of muscle calcium binding to troponin C (TnC) triggers structural changes in the troponin complex which lead to the azimuthal motion of tropomyosin away from its inhibitory position on the thin filament, partially uncovering the myosin binding sites on actin and enabling contraction. The control of the ON/OFF state of the thin filament by calcium ions is at the basis of the Steric Blocking mechanism of regulation of muscle contraction proposed nearly 50 years ago (19-22). Recent cryo-electron microscopy (Cryo-EM) studies have provided high-resolution three-dimensional (3D) reconstructions of cardiac thin filaments in the absence of calcium and at saturating (23) and submaximal (24) calcium concentrations, greatly advancing our understanding of the structural basis of thin filament regulation by calcium (25, 26).

In vitro studies of the mechanism of thin filament regulation described a third structural state of tropomyosin on the thin filament, in which myosin motors strongly bound to actin in the ATP-free state (rigor state) further shift the tropomyosin strand away from the myosin-binding sites on actin (27, 28). These studies therefore introduced the concept of three regulatory states of the thin filament, corresponding to three azimuthal positions of tropomyosin, the blocked ( $\text{Ca}^{2+}$ -free), closed ( $\text{Ca}^{2+}$ -bound) and open (Myosin-bound) states (Fig. 2). The comparison of 3D-reconstructions of isolated thin filaments in the presence and in the absence of calcium (22) and high-resolution structures of the thin filaments decorated with myosin motors in the rigor state (22, 29), together with the evidence of cooperative binding of myosin motors along the thin filament (30), provides support to that concept. More recently, cryo-electron tomograms of the sarcomere in skeletal myofibrils in the rigor state showed that in the region of filament overlap (A-band) the myosin motors attached to actin hold the thin filaments in the open state, whereas in the absence of myosin at no-overlap in the I-band the thin filaments are in the closed state (31, 32). However, until recently the role of myosin binding to actin in the activation of the thin filament during contraction was unclear.

The mechanism of thin filament regulation has been studied in demembranated skeletal (33-35) and cardiac muscle (36-38) using fluorescence polarization from bifunctional rhodamine probes on troponin-C (TnC), a technique which allow to determine the orientation of specific protein domains in situ under physiological conditions (39). A recent study using those TnC probes has elucidated the contribution of  $\text{Ca}^{2+}$  and myosin to the regulatory structural changes in the thin filament in demembranated muscle fibers from rabbit skeletal muscle (40). The results of this study showed that at near physiological temperature ( $26^{\circ}\text{C}$ ) and lattice spacing, conditions that are required to preserve the thick filament regulatory mechanisms in demembranated muscle fibers (41-43), the calcium sensitivity and the cooperativity of the regulatory structural changes in the thin filament are increased by the attachment of force generating myosin motors to actin. This is consistent with the effect of myosin on thin filament activation measured with fluorescent probes on the inhibitory subunit of troponin (TnI) in human slow-twitch fibers (44). These results suggest that during contraction of skeletal muscle the thin filaments are only partially activated by calcium, and that full activation and co-operativity of the thin filaments require myosin binding to actin, extending the concept of three regulatory state of the thin filament from in vitro studies to the myofilament lattice in physiological conditions. The myosin-dependent increase in calcium sensitivity and cooperativity of the orientation change of troponin is blunted by cooling the muscle fibers at  $11^{\circ}\text{C}$  (34), supporting the conclusion from mechanical studies that the cooperativity of thin filament activation depends not only on the attachment of myosin to actin, but also on the force generated by the myosin motors (45).

In demembrated cardiac muscle at 20°C myosin attachment to actin increases the calcium sensitivity and cooperativity of the orientation change of both the C- and N-lobe of TnC (38), although the myosin-dependent contribution to the activation of the thin filament is much smaller than that observed in skeletal muscle (40). This is consistent with the evidence from electron microscopy studies that the amplitude of the myosin-dependent motion of tropomyosin in the cardiac thin filament is smaller than that in the skeletal thin filament (46), suggesting that the mechanism of activation of the thin filament in cardiac muscle might be different from that in skeletal muscle. However, in relaxed cardiac muscle at temperatures below 27°C the thick filament is partially activated in the absence of calcium (47). Due to the coupling between thin and thick filament regulatory mechanisms (40), it is likely that the changes in the regulatory state of the thick filament induced by the low temperature might also alter the signaling pathway of thin filament activation during contraction. Therefore, further work is required to determine the contribution of calcium and myosin to the regulatory structural changes in the cardiac thin filament and the coupling between thin and thick filament activation under more physiological conditions.

## **2.2 Structural dynamics of the thin filament during muscle contraction.**

The kinetics of force generation during contraction depends on the speed of activation of thin and thick filaments following the electrical stimulation of muscle. How fast does the tropomyosin move on the thin filament during muscle activation? In vitro experiments using a pyrene label attached to tropomyosin have shown that the kinetics of the transition of the thin filament from the closed to the open state triggered by myosin motors in solution has a rate constant of  $\sim 1000 \text{ s}^{-1}$  (28, 48). The motion of tropomyosin in electrically stimulated skeletal muscle has been measured by X-ray diffraction, the only label-free technique that allows the structural dynamics of the myofilaments to be measured in situ on the millisecond time scale in the quasi-crystalline lattice of the muscle cell under physiological conditions (49). In their pioneering study Kress et al. measured the time course of the intensity of the second actin layer line (ALL2) reflection, associated with azimuthal shift in the position of tropomyosin on the thin filament, during the isometric contraction of frog skeletal muscle (14). Their results showed that at 22°C the kinetics of the structural changes in tropomyosin on the thin filament is much faster than the dynamics of force generation, whereas during relaxation after the end of the electrical stimulation it is similar to the kinetics of the force decay (14). Similar dynamic changes in tropomyosin conformation were observed by Yagi and coworkers during twitch contractions of frog skeletal muscle at 16°C, elicited by a single electrical stimulus (50). These results confirmed that the motion of tropomyosin triggered by calcium binding to troponin is the first step in the signaling pathway of muscle activation. In overstretched muscle at no-overlap the return of tropomyosin to its resting position during relaxation is faster than that at optimal filament overlap, suggesting that the myosin motors attached to actin might interfere with the recovery of the inhibitory position of tropomyosin (14). This is consistent with the evidence that forced detachment of myosin motors from the thin filament induced by a ramp shortening applied at the peak of the force twitch accelerates the recovery of the resting inhibitory position of tropomyosin (51).

The speed of thin filament activation following a rapid jump in calcium concentration produced by photolysis of caged calcium has been measured by X-ray diffraction with a time resolution of 4 ms in demembrated muscle fibers from rabbit psoas muscle at 4°C (52). In that study the time course of the intensity of the tropomyosin reflection was fitted with a double-exponential function in which the faster component, representing  $\sim 60\%$  of the motion of tropomyosin on the thin filament, was associated with the calcium-induced motion of tropomyosin and had a rate constant of  $\sim 160 \text{ s}^{-1}$ , whereas the slower component ( $\sim 40\%$ ) had a rate constant of  $\sim 18 \text{ s}^{-1}$ , similar to the rate of force

generation, and was associated with the attachment of myosin motors to actin. Moreover, the shortening of the muscle fiber at zero load applied at the plateau of contraction triggered the partial recovery of the inhibitory position of tropomyosin on the thin filament (52), suggesting a role of the myosin motors in the activation of the thin filament during contraction.

However, despite the remarkable combination of the X-ray diffraction technique with sophisticated physiological methods applied on demembrated muscle fibers, under the experimental conditions used in that study the physiological mechanisms of thin and thick filament activation in the muscle fiber are severely disrupted (41-43). A more recent X-ray diffraction study in demembrated skeletal muscle fibers conducted at 30°C investigated the role of myosin in the activation of the thin filament in situ, and it showed that during muscle shortening against a load, corresponding to ~30% of the tetanic force, the tropomyosin strand remains in the same conformation as that at the tetanus plateau, suggesting that even a small fraction of myosin motors attached to actin might hold the thin filament in the open state (53). Currently, there is no published X-ray diffraction study to our knowledge of the structural dynamics of the thin filament in electrically stimulated mammalian skeletal or cardiac muscle at near-physiological temperature. Measuring the motion of tropomyosin by X-ray diffraction in electrically paced cardiac muscle is even more challenging. Indeed, in cardiac muscle the changes in intensity of the tropomyosin reflection are expected to be smaller than those observed in skeletal muscle, firstly because the thin filament is only partially activated at submaximal calcium concentration, and secondly because the amplitude of the azimuthal motion of tropomyosin induced by calcium and myosin is smaller than that in skeletal muscle, according to EM reconstruction of thin filaments (46, 49). However, these limitations might be partially overcome in future studies exploiting the recent advancements in synchrotron technology and X-ray photon-counting detectors with higher sensitivity and higher temporal and spatial resolution (54).

Thanks to the high signal-to-noise ratio of fluorescence, the fluorescence polarization technique represents a powerful tool to measure with high specificity and high temporal resolution the structural dynamics of regulatory protein domains in the native environment of skeletal and cardiac muscle cells during contraction (55). Recently, a fluorescence polarization study using probes on TnC and on the regulatory light chain (RLC) of myosin has characterized the speed of activation of troponin and of the myosin motors with 0.12 ms time resolution in skeletal muscle fibers from rabbit psoas muscle activated by photolysis of caged calcium in near-physiological conditions (40). In those experiments calcium- and myosin-dependent structural changes in the thin filament could be kinetically separated, thanks to the difference of two orders of magnitude in their rate constants. Calcium binding to troponin triggers a fast reorientation of the troponin complex on the thin filament with a rate constant of 4,000-7,000 s<sup>-1</sup>, followed by a slower orientation change in troponin that tracks myosin binding to actin and the kinetics of force generation (~30 s<sup>-1</sup>) (40). The attachment of force-generating myosin motors to actin triggers a positive feedback loop in the activation of the thin filament, called “myosin-sensing” (Fig. 2), which dynamically couples the regulatory state of the thin filament to that of the thick filament during contraction at high-loads and allows skeletal muscle to achieve rapid and cooperative activation following electrical stimulation.

In cardiac muscle the structural dynamics of the thin filament on the timescale of the heartbeat has not been elucidated. A time-resolved fluorescence polarization study using fluorescent probes on TnC in demembrated cardiac trabeculae activated by caged calcium showed that the calcium-induced structural changes in troponin are ten times faster than force generation (56), suggesting that also in cardiac muscle the calcium-dependent activation of the thin filament does not limit the rate of force generation. This result is consistent with the evidence from a recent X-ray diffraction study that the kinetics of force generation and relaxation in electrically-paced intact cardiac muscle

is largely controlled by the dynamics of the structural changes in the thick filament (16), as discussed below.

### **3. Thick filament-based regulation in striated muscle**

#### **3.1 The conformation of the myosin motors on the thick filament in resting muscle.**

The myosin filament is a centrosymmetric structure containing two arrays of myosin motors (Fig. 1B). Each array in the half-filament consists of 49 layers of motors with an axial periodicity of ~14.5 nm. Each half-filament is associated with six titin molecules connecting the thick filament tips to the ends of the sarcomere in the Z-band. The thick-filament binding region of titin contains repeating immunoglobulin/fibronectin domains arranged in 11 x 11-domain (C-type) super-repeats or 6 x 7-domain (D-type) super-repeats, in the central and distal region of the thick filament, respectively (57, 58). MyBP-C is located in the middle third of each half thick filament with an axial periodicity of ~43 nm and it is anchored to the myosin backbone via its C-terminal domain, whereas its N-terminal region is thought to be in a dynamic equilibrium between actin-bound and myosin-bound states, modulating the regulatory state of thin and thick filaments, respectively (59-62). The position of MyBP-C defines the C-zone of the thick filament, whereas the neighboring filament regions lacking MyBP-C are referred to as proximal (P) and distal (D) zones (Fig. 1B).

Recently it has become clear that in resting muscle the myosin motors are inhibited by folding onto the surface of the thick filament backbone and therefore they are not immediately available for the interaction with actin following electrical stimulation (Fig. 1C). The structural basis of myosin inhibition has been elucidated for the first time in electron microscopy studies of relaxed thick filaments from tarantula muscle (11), showing that the two motor domains of the myosin molecule are folded onto the myosin tail in an asymmetric conformation called interacting heads motif (IHM), originally observed in two-dimensional crystals of smooth muscle myosin (10), which inhibits both the myosin ATP-ase and the interaction of myosin with actin. Early 3D-reconstructions of the C-zone of human cardiac thick filaments also show the characteristic IHM structure (63-65), and subsequent studies demonstrated that folding of myosin motors into the IHM is a highly conserved mechanism for the inhibition of myosin II (66, 67). Recent cryo-electron microscopy studies have provided high-resolution structures of the purified human  $\beta$ -cardiac myosin IHM in the relaxed state (68) and of the C-zone of cardiac thick filaments either isolated (69) or in the sarcomere of relaxed myofibrils (70) in the presence of the myosin inhibitor Mavacamten. The *in vitro* and *in situ* reconstructions of the thick filaments have elucidated distinct conformations of the folded myosin motors in the C-zone and the structural organisation of MyBP-C and of the titin C-repeats on the surface of the thick filament (69, 70). Moreover, cryo-electron tomograms of the C-zone in relaxed myofibrils (70) and in chemically-fixed cardiac muscle (71) revealed MyBP-C links between thin and thick filaments, although the regulatory role of such links in the interfilament signalling pathway of muscle regulation remains unclear.

Because the myosin ATP-ase is inhibited in the IHM, the folded state of myosin has been generally associated with the “super-relaxed (SRX)” or OFF state of myosin, a biochemical state of myosin characterized by a slow ATP turnover, identified in relaxed skeletal (72, 73) and cardiac muscle (73,

74). The equilibrium between the SRX state and the disordered-relaxed (DRX) state of myosin can be pharmacologically modulated to alter the resting muscle metabolism (75, 76). However, a recent X-ray diffraction study using 2'-deoxy-ATP (dATP) to destabilize the folded state of myosin in relaxed porcine cardiac muscle showed that the structural transition of the myosin motors caused by dATP is not equivalent to the biochemical transition from SRX to DRX state (77). Therefore, the relationship between the SRX/DRX state of myosin and the regulatory structural states of the myosin motors remains to be elucidated.

The evidence that in resting muscle the myosin motors are sequestered in the folded state onto the filament surface led to the hypothesis that the contractility of skeletal muscle is largely controlled by regulatory structural changes in the thick filament (7), in addition to the classical calcium-dependent regulation of the thin filament. Moreover, it has been hypothesized that in the heart the destabilization of the IHM by mutations in thick filament proteins which cause hypertrophic cardiomyopathy (HCM) might disrupt the thick-filament based regulation of contraction, potentially explaining the hyper-contractile phenotype observed in HCM (60, 78-80).

The in-situ conformation of the myosin motors on the thick filament in resting skeletal muscle (15, 81) and in cardiac muscle in diastole (16, 82) derived from X-ray diffraction studies is largely consistent with the IHM, indicating that under physiological conditions most of the myosin motors are inhibited by folding onto the filament surface. However, a recent X-ray diffraction study on cardiac muscle described two populations of folded motors in diastole, the folded-helically ordered motors located in the C-zone (Fig. 1C), and the folded non-helically ordered motors located outside the C-zone (Fig. 1C)(16). The folded-helical conformation of the myosin motors might be stabilized by MyBP-C, in agreement with the 3D-reconstructions of the C-zone of the thick filament (69, 70), and might account for the enrichment of the SRX state of myosin in the C-zone of the filament observed in isolated skeletal (83, 84) and cardiac myofibrils (85). However, in cardiac myofibrils isolated from mice expressing a cardiac MyBP-C with a N-terminal truncation the fraction of SRX state remains higher in the central regions (P and C) of the thick filament (85), suggesting that other regulatory factors might contribute to the stabilization of the myosin SRX in these regions.

X-ray diffraction and fluorescence polarization studies showed that the physiological resting structure of the thick filament and the folded conformation of the myosin motors are largely preserved in demembranated skeletal (41, 42) and cardiac muscle (86, 87) under relaxing conditions, but only at temperatures higher than 26°C and after recovering the physiological lattice distance typical of the intact muscle using the osmotic agent Dextran T-500. Under these conditions therefore the demembranated muscle preparations represent a suitable model to investigate the regulatory mechanisms of the thick filament. Fluorescence polarization studies using bifunctional rhodamine probes on the regulatory light chain of myosin exchanged in demembranated skeletal muscle fibers (42) and cardiac trabeculae (87) concluded that in near-physiological conditions in the absence of calcium about 2/3 of the myosin motors on the thick filament are folded in the IHM, whereas 1/3 of the motors are more perpendicular to the filament axis, consistent with the fraction of folded and isotropic motors, respectively, in diastole determined in X-ray diffraction experiments (16). These isotropic motors might be a population of "constitutively ON" (12) or "sentinel motors" (66) (Fig. 1, 2) which are immediately available for the interaction with actin when the thin filament is partially activated by calcium. Although there are currently no high-resolution structures of the D-zone of the thick filament, it has been suggested that the isotropic motors are mainly located in the D-zone (Fig. 1) (16), consistent with the enrichment of DRX myosin motors in this filament domain (83-85).

In conclusion, in resting muscle the myosin motors on the thick filament have multiple conformations that cannot be described only in terms of binary structural OFF/ON or biochemical

SRX/DRX states. These myosin conformations might be confined to specific filament domains with different protein composition, and therefore new structural approaches are necessary to characterize their spatial distribution and regulatory function in sarcomere subdomains under physiological conditions.

### **3.2 The mechanism of activation of the thick filament in skeletal muscle**

Recent X-ray diffraction studies of amphibian skeletal muscle showed that the activation of the folded myosin motors on the thick filament is directly controlled by filament stress, based on the evidence that the activation of the thick filament is prevented by imposing shortening at zero load after the start of the stimulation (12). The stress-dependent activation of the thick filament was further characterized in a fluorescence polarization study in demembranated muscle fibers showing that the passive tension transmitted to the thick filament by titin partially activates the folded myosin motors, independently of calcium (13). According to the mechano-sensing mechanism the force generated by the constitutively ON motors triggers a positive feedback loop in the activation of the thick filament (Fig. 2) (7), which might account for the high cooperativity in the activation of the folded myosin motors in demembranated muscle fibers activated at constant calcium concentrations (13). The release of folded motors from the filament surface determines an increase in the number of actin-bound motors which sensitizes the thin filament to calcium and increases the cooperativity of thin filament activation, triggering the “myosin-sensing” feedback loop in the activation of the thin filament (40) (Fig. 2). Such a coupling between mechano-sensing in the thick filament and myosin-sensing in the thin filament might account for the rapid activation of skeletal muscle following electrical stimulation.

It has been recently shown that the mechano-sensing mechanism controls the activation of the myosin motors also in the fast-twitch muscle of the mouse (88). However, there appear to be some differences in the structural dynamics of the thick filament in mammalian skeletal muscle, with respect to that in amphibian muscle (81, 89). Firstly, a small component of thick filament activation, which consisted of a slight elongation of the filament backbone signaled by the increase in the spacing of the M6 reflection ( $S_{M6}$ ), was detected during shortening at zero load after the start of the stimulation. Secondly, the analysis of the M3 reflection, associated with the axial periodicity (14.5 nm) of the myosin motors along the thick filament, revealed a new component peak indexing at a slightly longer periodicity of  $\sim 14.8$  nm, which appears in the early phase of force generation. This extra peak also appears during activation of demembranated skeletal muscle at submaximal calcium concentrations close to the threshold value for force generation (pCa 7-6.4) (90). It has been proposed that this longer periodicity could be associated with a population of myosin motors interacting with the titin C-repeats on the thick filament (41). However, this hypothesis is not consistent with the recent high-resolution reconstructions of the C-zone of the thick filament that show a single axial periodicity matching that of the myosin helix (43nm) (69, 70). An alternative hypothesis is that the myosin motors with longer periodicity might be associated with the D-type repeats of titin in the D-zone of the thick filament, and might be involved in a transient activation state of the thick filament in the early phase of force generation (90). Finally, the activation of the different zones of the thick filament during isometric contraction is sequential. The constitutively ON motors are preferentially located at the tips of the thick filament (Fig. 1) and trigger the stress-dependent activation of the myosin motors in the P-zone, followed by the activation of motors in the C-zone (88). In conclusion, filament stress is the major determinant of thick filament activation in fast-twitch mammalian skeletal muscle, but an additional pathway of myosin activation independent of stress cannot be excluded.

The time course of the structural changes in the thick filament triggered by a single action potential has been measured in EDL muscle of the mouse by time-resolved X-ray diffraction (15). In response to a single electrical stimulus the length of the thick filament backbone, signaled by the X-ray parameter  $S_{M6}$ , slightly increases triggering the partial disruption of the helical order of the myosin motors on the thick filament ( $I_{ML1}$ ) and the conformational change of the myosin motors from the folded to the perpendicular actin-attached state ( $L_{M3}$ ), with time courses similar to that of force generation (Fig. 3). The kinetics of thin filament activation triggered by the action potential has not been measured by X-ray diffraction in mammalian muscle, but according to the structural dynamics of troponin measured by fluorescence polarization in demembrated muscle fibers (40) the calcium-induced structural changes in the thin filament are expected to be transiently complete at the peak of the calcium transient,  $\sim 2$ ms after the electrical stimulus. Therefore, according to these results, the speed of force generation in the twitch is largely rate-limited by the dynamics of activation of the myosin motors on the thick filament. Mathematical models of muscle contraction that include the stress-dependent activation of the myosin motors in the thick filament simulate the time course of force generation during an isometric contraction (91), supporting the conclusion that the dynamics of force generation in skeletal muscle is largely determined by the structural dynamics of the thick filament.

The kinetics of the recovery of the resting length of the backbone and of the folded-helical conformation of the myosin motors on the thick filament are similar to the kinetics of force relaxation (Fig. 3), indicating that in the force twitch the inactivation of the thick filament is rate-limiting for the kinetics of mechanical relaxation. Muscle relaxation from the tetanus plateau occurs in two phases. In the first phase, called isometric relaxation, force declines at constant sarcomere length following myosin detachment from actin, but the myosin filament remains in the active state (15, 17). Instead, in the second phase, called chaotic relaxation due to sarcomere yielding, the thick filament partially recovers the resting structure with a kinetics slower than that of force (15). Although the mechanism of muscle relaxation is still not understood, these results suggest that thick filament inactivation alone cannot explain the kinetics of force relaxation from high loads and that the coordinated inactivation of thin and thick filaments might be required to inhibit the positive myosin- and mechano-sensing feedbacks, in thin and thick filaments respectively.

### **3.3 The mechanism of activation of the thick filament in cardiac muscle**

It has been recently proposed that the activation of the thick filament in cardiac muscle is controlled by a mechano-sensing mechanism similar to that observed in skeletal muscle (8). This conclusion is based on the evidence from X-ray diffraction experiments on cardiac trabeculae from rat hearts that at the peak of the force twitch the activation level of the thick filament is proportional to the systolic force (82). A fluorescence polarization study using RLC probes exchanged in demembrated rat trabeculae showed that the activation of the myosin motors from the folded state at constant calcium concentrations and at different levels of RLC phosphorylation depends on filament stress, providing further support to the mechanism of stress-dependent activation of the thick filament (87). The mechano-sensing transition of the myosin motors has been included in mathematical models of cardiac muscle contraction to simulate the time course of force in the cardiac twitch, as well as the length-dependent activation of cardiac muscle (92-95).

However, more recently, an X-ray diffraction study on demembrated porcine cardiac muscle proposed that the thick filament in cardiac muscle can be directly activated by calcium (96). In that study the activation of the thin filament by calcium was pharmacologically inhibited to determine

the effect of constant calcium concentrations in the absence of force on the structure of the thick filament. The results showed that the thick filament is partially switched on at constant  $[Ca^{2+}]$ , suggesting a direct role of calcium in the activation of the myosin motors (96), although the calcium binding site on the thick filament is unknown. Biochemical evidences showed that porcine cardiac myosin RLC can bind  $Ca^{2+}$ , but under physiological conditions the binding sites would be mainly occupied by  $Mg^{2+}$  in diastole and the exchange of  $Mg^{2+}$  with  $Ca^{2+}$  during contraction would be too slow to trigger the activation of the thick filament on the time scale of the heartbeat (97). An alternative hypothesis is that calcium signaling to the thick filament might be mediated by MyBP-C (98). Previous X-ray diffraction studies on intact papillary muscle isolated from the right ventricle of canine hearts reported an increase in the transfer of mass of myosin motors from the thick to the thin filament in diastole at higher frequencies of stimulation (99, 100), suggesting that additional signaling pathways might control the regulatory state of the thick filament in cardiac muscle. In conclusion, further work is required to solve the controversy about the role of calcium and filament stress in the activation of the cardiac thick filament (101).

The structural dynamics of the thick filament during contraction of cardiac muscle has been recently elucidated in time-resolved X-ray diffraction studies on electrically paced cardiac trabeculae from rat hearts (16). Both the time course of the force twitch and the kinetics of the regulatory structural changes in the thick filament, associated with the elongation of the thick filament backbone and the release of the myosin motors from the folded conformation, are ten times slower than those observed in skeletal muscle, although their amplitudes are similar (Fig. 3). The difference in the speed of activation of the thick filament might be accounted for by the different myosin isoforms in these two muscle types and by the different time course of the intracellular calcium transient, which is ten times smaller and slower in the cardiac muscle cell (4) with respect to that in skeletal muscle (3) (Fig. 3). In cardiac muscle the activation of the myosin motors starts in the D-zone and it propagates to the central regions of the thick filament, and at the peak of the force twitch the actin-bound myosin motors contributing to force generation are only confined to the C-zone of the thick filament (16), likely due to the sensitization of the thin filament to calcium in the C-zone induced by MyBP-C links (Fig. 2) (102, 103).

During relaxation the deactivation of the thick filament occurs in the zonal sequence D-C-P, following the spatial gradient of force distribution on the filament backbone, in agreement with the mechanosensing hypothesis, and it is accompanied by an increase in sarcomere inhomogeneity (16). However, in contrast with the fast reformation of the myosin helix observed in skeletal muscle, in cardiac muscle the recovery of the folded-helical conformation of the myosin motors occurs only at a later stage of mechanical relaxation (Fig. 3C,  $t > 300ms$ ), suggesting that the helical order of the thick filament in the C-zone of the cardiac thick filament requires additional inter-molecular interactions between myosin motors and MyBP-C. In conclusion, the zonal dynamics of thin and thick filament activation, and their coupling by an interfilament signaling pathway based on MyBP-C, control the strength and the dynamics of contraction in cardiac muscle.

#### **4. The role of titin in the regulation of muscle contractility**

Titin is traditionally regarded as a passive spring that resists the lengthening of the sarcomere in resting muscle, the elasticity of which is associated with its I-band domain and is tuned to the operative sarcomere length range in cardiac and skeletal muscle (104, 105). The A-band domain of titin associated with the thick filament backbone acts as a molecular ruler and template for the

assembly of the thick filament (58, 106). However, recent studies on skeletal muscle suggested that titin might have a direct role in the regulation of muscle contractility. In actively contracting skeletal muscle titin exhibits a dynamic stiffness that is about 100 times higher than that in the muscle at rest (107), which is likely due to the stiffening of the skeletal titin isoform induced by calcium (13, 108, 109), and it has been proposed that during contraction titin switches from an extensible spring to a “mechanical rectifier” that allows free shortening and resists the elongation of the sarcomere (110). The increased stiffness of titin in the active sarcomere might buffer force imbalances due to sarcomere length inhomogeneity during high-load contractions, as suggested by the evidence that the cleavage of titin *in situ* leads to sarcomere disorganization upon muscle activation (111). Fluorescence polarization experiments using probes on the myosin RLC in demembranated skeletal muscle fibers showed that the titin-based passive stress applied to the thick filament triggers the partial release of myosin motors from the folded state, suggesting a possible role of titin in the mechano-sensing transition of the thick filament (13) (Fig. 2). The activation of the thick filament by the titin-based passive force was also recently shown in X-ray diffraction experiments on relaxed demembranated skeletal muscle fibers isolated from a mouse model in which titin could be specifically cleaved (112). More recently, X-ray diffraction experiments on electrically stimulated muscle fibers of the frog, in which force generation was suppressed using the myosin inhibitor para-nitro-blebbistatin, showed that in the presence of calcium the stress transmitted by titin to the thick filament backbone alters the resting conformation of the myosin motors (110).

Therefore, in skeletal muscle, titin stress triggers the mechano-sensing transition of the thick filament, but what is its physiological role in muscle contraction? Recent time-resolved fluorescence polarization experiments in demembranated skeletal muscle fibers activated by photolysis of caged calcium showed that the kinetics of activation of the myosin motors folded on the thick filament surface can be accelerated by stretching the muscle fiber during the early phase of contraction (40). The titin stress transmitted to the thick filament therefore triggers the rapid recruitment of active myosin motors from the folded state (Fig. 2), that can attach to actin and generate a high resistive force, accounting for the fast and efficient braking response of active skeletal muscle to the external stretch (113, 114) .

In contrast with the effect observed in skeletal muscle, in cardiac muscle in diastole the passive tension generated by increasing the sarcomere length in the physiological range (1.9-2.3 $\mu\text{m}$ ) is small and does not trigger the activation of the thick filament (16, 87, 115, 116). Moreover, the stiffness of the cardiac titin is already high in diastole and is not altered in the presence of calcium (109), suggesting that the role of titin in buffering force imbalances between sarcomeres might be important in systole, as well as in diastole. However, the regulatory role of titin in the control of the contractility of cardiac muscle is poorly understood, limiting our understanding of the mechanisms by which titin mutations can cause cardiac diseases (117-120).

## **5. Myofilament length-dependent activation**

The contractility of cardiac muscle is largely controlled by an auto-regulatory mechanism, called the “Frank-Starling Law of the Heart”, according to which the ventricular stroke volume, i.e. the amount of blood pumped in each heartbeat, increases in response to an increase in the ventricular volume in diastole (121, 122). The cellular mechanism underlying this phenomenon consists of a stretch-induced sensitization of the myofilaments to calcium, called “length-dependent activation”

(LDA)(121, 123), which causes an increase in the strength of contraction in response to the stretch of the cardiac muscle cell in diastole (16). LDA increases the efficiency of cardiac muscle contraction (124) and its disruption is associated with decreased contractile performance of the heart in human hypertrophic cardiomyopathies (125) and in patients with ischemic heart failure (126).

Despite decades of research, the length-sensing mechanism that modulates the calcium sensitivity of the myofilaments in response to changes in sarcomere length is still unknown. It has been recently hypothesized that the stretch-induced potentiation of cardiac muscle contractility could be explained by an increase in diastole in the number of myosin motors available for the interaction with actin at longer sarcomere lengths (116). However, recent studies conducted by different laboratories suggest that this is not the case. X-ray diffraction experiments on electrically paced cardiac trabeculae isolated from rat hearts showed that increasing the sarcomere length in the range 1.95-2.40  $\mu\text{m}$  in diastole does not alter the relaxed structure of the thick filament (16, 115, 116). This result is in agreement with the conclusion from fluorescent polarization experiments on demembranated rat cardiac trabeculae showing that in the absence of calcium, or at calcium concentrations close to the diastolic value ( $\sim 0.1\mu\text{M}$ ), the folded myosin motors are insensitive to the passive force applied by titin to the thick filament (87). In contrast with these results, a recent X-ray diffraction study on demembranated porcine myocardium concluded that the stretch in the absence of calcium triggers a partial release of myosin motors from the folded state (127).

In conclusion, the majority of studies indicate that LDA is not mediated by stretch-induced changes in the structure of the thick filament in diastole, although these results remain still controversial. Stretch of the cardiac muscle cell at calcium concentrations in the systolic range (0.2-1 $\mu\text{M}$ ), which partially activate the thin filament, triggers the release of myosin motors from the folded state contributing to the stretch-induced potentiation of cardiac contractility (87). Conversely, shortening of active cardiac muscle might induce deactivation of the myosin motors on the thick filament accelerating muscle relaxation (128). All together these results suggest that LDA in cardiac muscle might be explained by a dual-filament mechanism which couples length-dependent structural changes in thin and thick filaments in the presence of calcium (87). The molecular mechanism of sarcomere-length sensing remains unclear, but it might involve interfilament communication pathways based on interactions between thin filament proteins and MyBP-C (87) or titin (121, 122).

## 6. Conclusions

The generation of force in striated muscle is controlled by a dual-filament regulatory mechanism that positively couples the calcium dependent structural changes in the thin filament with the stress-dependent activation of the thick filament. In skeletal muscle the activation of thin and thick filament is coordinated by two positive feedback mechanisms, mechano-sensing in the thick filament and myosin-sensing in the thin filament, that account for the rapid and cooperative activation of muscle following electrical stimulation. An additional pathway based on titin stress might accelerate the kinetics of the mechano-sensing transition of the thick filament in response to a rapid increase in the external load, allowing skeletal muscle to act as an efficient brake. A dual-filament regulation mechanism also operates in the heart, although the coupling between thin and thick filament regulatory mechanisms has not been fully elucidated. The dynamics of contraction in cardiac muscle is controlled by the zonal dynamics of thin and thick filament domains that are

coordinated by an interfilament signaling pathway based on MyBP-C. The coordinated activation of thin and thick filaments in this dual-filament model might underlie the length-dependent activation of cardiac muscle as the basis of the Frank-Starling Law of the heart.


### **Future issues**

- What is the structural basis of mechano-sensing in the thick filament?
- What is the mechanism of regulation of the myosin motors in the D-zone of the thick filament?
- Is the thick filament directly activated by  $\text{Ca}^{2+}$ ?
- What is the mechanism of muscle relaxation?
- What is the role of titin and MyBP-C in the regulation of cardiac contractility?
- What is the length sensor at the basis of LDA in cardiac muscle?

### **Acknowledgments**

We are grateful to M. Irving and D. Warshaw for helpful comments on the manuscript. EB was funded by a British Heart Foundation (BHF) Intermediate Basic Science Research Fellowship (FS/17/3/32604). LF was funded by a Sir Henry Dale Fellowship awarded by the Wellcome Trust and the Royal Society (210464/Z/18/Z).


## Figures




concentration ( $[Ca^{2+}]_i$ ). In cardiac muscle the voltage-gated calcium channels are uncoupled from the RyRs, so that the extracellular calcium entering the cell triggers the calcium release from the sarcoplasmic reticulum (calcium induced-calcium release). The time courses of the intracellular calcium concentration and of the force twitch are shown in the lower panels.

B. Schematic of the sarcomere, the structural unit of striated muscle. The thin filaments are anchored to the Z-bands and contain actin (white circles), tropomyosin (brown lines) and the troponin complex (light green). Thick filaments (purple) are aligned in the center of the sarcomere, the M-band, and contain two antiparallel arrays of myosin motors. Each half-filament is divided into a proximal or P-zone, a Myosin-binding Protein-C (MyBP-C)-containing C-zone and a distal or D-zone. In the C-zone MyBP-C molecules (orange) form links with actin filaments. Titin filaments span the half-sarcomere length and run along the thick filament surface; I-band titin, black; A-band titin C-type repeats, blue, and D-type repeats, light blue.

C. Snapshots of the structure of the myofilaments at low (top panels) and high (bottom panels)  $[Ca^{2+}]_i$  in the C-zone (left) and D-zone (right). At low  $[Ca^{2+}]_i$ , tropomyosin (brown lines) blocks the myosin binding sites on the actin filament (black). In the C-zone of the thick filament (top left), most of the myosin motors are folded and helically ordered on the thick filament surface (magenta), whereas in the D-zone, most of the myosin motors are disordered (top right, grey). Calcium binding to troponin (bottom right, yellow circles) triggers the partial activation of the thin filament. In the D-zone the disordered myosin motors are immediately available to bind actin, whereas in the C-zone they need to be released from the folded conformation before attaching to actin.



**Figure 2.** Schematic of dual-filament regulation in striated muscle. The contraction of striated muscle is controlled by two positive feedback loops modulating the activation of the thick (*mechano-sensing*) and thin (*myosin-sensing*) filaments, respectively. In resting muscle, the thin filament is blocked, as the myosin binding sites on actin are covered by tropomyosin, and most of the myosin motors on the thick filament are unavailable to interact with actin as they are folded and helically ordered on the thick filament surface. Few myosin motors are disordered. Myosin binding protein C, MyBP-C, stabilizes the folded conformation of the myosin motors in skeletal muscle at rest or in cardiac muscle in diastole. MyBP-C sensitizes the thin filament to calcium in the C-zone. Calcium binding to TnC triggers the movement of tropomyosin on the actin filament partially uncovering the myosin binding sites on actin (thin filament closed state). Myosin binding to actin induces a further movement of tropomyosin triggering the transition of the thin filament to the open state. Filament stress triggers the release of folded motors from the filament surface (mechano-sensing), increasing the pool of available motors (grey) and consequently the number of actin-bound motors. In skeletal muscle, titin stress accelerates the release of folded motors during stretch of active muscle.



**Figure 3.** Structural dynamics of the thick filament triggered by a single action potential in skeletal and cardiac muscle. A. Calcium transient (black thin line; from Baylor & Hollingworth, 2003) and force response (black thick line; from Hill et al., 2021) in EDL skeletal muscle from the mouse; Calcium transient (magenta thin line; from Janssen et al., 2002) and force response (magenta thick line; from Brunello et al., 2020) in rat ventricular cardiac trabeculae. Horizontal dashed lines indicate values at rest (black) or in diastole (magenta), in skeletal and cardiac muscle, respectively. Vertical dashed lines, peak force of the skeletal (black) and heart (magenta) muscle twitch. The vertical green line marks the electrical stimulus. B. Changes in the periodicity of the thick filament backbone, reported by the spacing of the M6 reflection ( $S_{M6}$ ). C. Changes in the intensity of the ML1 reflection ( $I_{ML1}$ ), associated with the folded helical conformation of the myosin motors on the filament surface (inset, magenta). D. Changes in fractional intensity of the low-angle peak of the M3 reflection ( $L_{M3}$ ), associated with the conformation of myosin motors, moving from a folded (inset, magenta) to a more perpendicular actin-bound conformation (inset, dark green). Black symbols, EDL skeletal muscle, data from Hill et al. (2021). Magenta symbols, rat cardiac trabeculae electrically paced at 1Hz; data from Brunello et al., 2020. Spatial

calibration in B for both skeletal and heart muscle as described in Brunello et al., 2020.  
Temperature, 26-28°C.

## References

1. Bers DM. 2000. Calcium fluxes involved in control of cardiac myocyte contraction. *Circ Res* 87: 275-81.
2. Lamb GD. 2000. Excitation-contraction coupling in skeletal muscle: comparisons with cardiac muscle. *Clin Exp Pharmacol Physiol* 27: 216-24. doi:10.1046/j.1440-1681.2000.03224.x
3. Baylor SM, Hollingworth S. 2003. Sarcoplasmic reticulum calcium release compared in slow-twitch and fast-twitch fibres of mouse muscle. *J Physiol* 551: 125-38. doi:10.1113/jphysiol.2003.041608
4. Janssen PM, Stull LB, Marban E. 2002. Myofilament properties comprise the rate-limiting step for cardiac relaxation at body temperature in the rat. *Am J Physiol Heart Circ Physiol* 282: H499-507. doi:10.1152/ajpheart.00595.2001
5. Backx PH, Ter Keurs HE. 1993. Fluorescent properties of rat cardiac trabeculae microinjected with fura-2 salt. *Am J Physiol* 264: H1098-110. doi:10.1152/ajpheart.1993.264.4.H1098
6. Gordon AM, Homsher E, Regnier M. 2000. Regulation of contraction in striated muscle. *Physiological Reviews* 80: 853-924.
7. Irving M. 2017. Regulation of Contraction by the Thick Filaments in Skeletal Muscle. *Biophys J* 113: 2579-94. doi:10.1016/j.bpj.2017.09.037
8. Piazzesi G, Caremani M, Linari M, Reconditi M, Lombardi V. 2018. Thick Filament Mechano-Sensing in Skeletal and Cardiac Muscles: A Common Mechanism Able to Adapt the Energetic Cost of the Contraction to the Task. *Front Physiol* 9: 736. doi:10.3389/fphys.2018.00736
9. Marcucci L. 2023. Muscle Mechanics and Thick Filament Activation: An Emerging Two-Way Interaction for the Vertebrate Striated Muscle Fine Regulation. *Int J Mol Sci* 24. doi:10.3390/ijms24076265
10. Wendt T, Taylor D, Trybus KM, Taylor K. 2001. Three-dimensional image reconstruction of dephosphorylated smooth muscle heavy meromyosin reveals asymmetry in the interaction between myosin heads and placement of subfragment 2. *Proceedings of the National Academy of Sciences of the United States of America* 98: 4361-6. doi:10.1073/pnas.071051098
11. Woodhead JL, Zhao FQ, Craig R, Egelman EH, Alamo L, Padron R. 2005. Atomic model of a myosin filament in the relaxed state. *Nature* 436: 1195-9. doi:10.1038/nature03920
12. Linari M, Brunello E, Reconditi M, Fusi L, Caremani M, et al. 2015. Force generation by skeletal muscle is controlled by mechanosensing in myosin filaments. *Nature* 528: 276-9. doi:10.1038/nature15727
13. Fusi L, Brunello E, Yan Z, Irving M. 2016. Thick filament mechano-sensing is a calcium-independent regulatory mechanism in skeletal muscle. *Nature Communications* 7: 13281. doi:10.1038/ncomms13281
14. Kress M, Huxley HE, Faruqi AR, Hendrix J. 1986. Structural changes during activation of frog muscle studied by time-resolved X-ray diffraction. *J Mol Biol* 188: 325-42.
15. Hill C, Brunello E, Fusi L, Ovejero JG, Irving M. 2021. Myosin-based regulation of twitch and tetanic contractions in mammalian skeletal muscle. *Elife* 10. doi:10.7554/eLife.68211
16. Brunello E, Fusi L, Ghisleni A, Park-Holohan SJ, Ovejero JG, et al. 2020. Myosin filament-based regulation of the dynamics of contraction in heart muscle. *Proc Natl Acad Sci U S A* 117: 8177-86. doi:10.1073/pnas.1920632117
17. Brunello E, Fusi L, Reconditi M, Linari M, Bianco P, et al. 2009. Structural changes in myosin motors and filaments during relaxation of skeletal muscle. *Journal of Physiology* 587: 4509-21. doi:DOI 10.1113/jphysiol.2009.176222
18. Poggesi C, Tesi C, Stehle R. 2005. Sarcomeric determinants of striated muscle relaxation kinetics. *Pflugers Arch* 449: 505-17. doi:10.1007/s00424-004-1363-5

19. Vibert PJ, Haselgrove JC, Lowy J, Poulsen FR. 1972. Structural changes in actin-containing filaments of muscle. *J Mol Biol* 71: 757-67. doi:10.1016/s0022-2836(72)80036-6
20. Parry DA, Squire JM. 1973. Structural role of tropomyosin in muscle regulation: analysis of the x-ray diffraction patterns from relaxed and contracting muscles. *J Mol Biol* 75: 33-55. doi:10.1016/0022-2836(73)90527-5
21. Lehman W. 2016. Thin Filament Structure and the Steric Blocking Model. *Compr Physiol* 6: 1043-69. doi:10.1002/cphy.c150030
22. Poole KJ, Lorenz M, Evans G, Rosenbaum G, Pirani A, et al. 2006. A comparison of muscle thin filament models obtained from electron microscopy reconstructions and low-angle X-ray fibre diagrams from non-overlap muscle. *J Struct Biol* 155: 273-84. doi:10.1016/j.jsb.2006.02.020
23. Yamada Y, Namba K, Fujii T. 2020. Cardiac muscle thin filament structures reveal calcium regulatory mechanism. *Nat Commun* 11: 153. doi:10.1038/s41467-019-14008-1
24. Risi CM, Pepper I, Belknap B, Landim-Vieira M, White HD, et al. 2021. The structure of the native cardiac thin filament at systolic Ca(2+) levels. *Proc Natl Acad Sci U S A* 118. doi:10.1073/pnas.2024288118
25. Tobacman LS. 2021. Troponin Revealed: Uncovering the Structure of the Thin Filament On-Off Switch in Striated Muscle. *Biophys J* 120: 1-9. doi:10.1016/j.bpj.2020.11.014
26. Lehman W, Pavadai E, Rynkiewicz MJ. 2021. C-terminal troponin-I residues trap tropomyosin in the muscle thin filament blocked-state. *Biochem Biophys Res Commun* 551: 27-32. doi:10.1016/j.bbrc.2021.03.010
27. McKillop DF, Geeves MA. 1993. Regulation of the interaction between actin and myosin subfragment 1: evidence for three states of the thin filament. *Biophys J* 65: 693-701. doi:10.1016/S0006-3495(93)81110-X
28. Geeves MA. 2012. Thin Filament Regulation. In *Comprehensive Biophysics*, ed. EH Egelman, pp. 251-67: Elsevier
29. von der Ecken J, Heissler SM, Pathan-Chhatbar S, Manstein DJ, Raunser S. 2016. Cryo-EM structure of a human cytoplasmic actomyosin complex at near-atomic resolution. *Nature* 534: 724-8. doi:10.1038/nature18295
30. Desai R, Geeves MA, Kad NM. 2015. Using fluorescent myosin to directly visualize cooperative activation of thin filaments. *J Biol Chem* 290: 1915-25. doi:10.1074/jbc.M114.609743
31. Wang Z, Grange M, Wagner T, Kho AL, Gautel M, Raunser S. 2021. The molecular basis for sarcomere organization in vertebrate skeletal muscle. *Cell* 184: 2135-50 e13. doi:10.1016/j.cell.2021.02.047
32. Wang Z, Raunser S. 2023. Structural Biochemistry of Muscle Contraction. *Annu Rev Biochem* 92: 411-33. doi:10.1146/annurev-biochem-052521-042909
33. Ferguson RE, Sun YB, Mercier P, Brack AS, Sykes BD, et al. 2003. In situ orientations of protein domains: troponin C in skeletal muscle fibers. *Mol Cell* 11: 865-74.
34. Sun YB, Brandmeier B, Irving M. 2006. Structural changes in troponin in response to Ca<sup>2+</sup> and myosin binding to thin filaments during activation of skeletal muscle. *Proceedings of the National Academy of Sciences of the United States of America* 103: 17771-6. doi:10.1073/pnas.0605430103
35. Knowles AC, Irving M, Sun YB. 2012. Conformation of the Troponin Core Complex in the Thin Filaments of Skeletal Muscle during Relaxation and Active Contraction. *J Mol Biol* 421: 125-37. doi:10.1016/j.jmb.2012.05.005
36. Sevrieva I, Knowles AC, Kampaourakis T, Sun YB. 2014. Regulatory domain of troponin moves dynamically during activation of cardiac muscle. *J Mol Cell Cardiol* 75: 181-7. doi:10.1016/j.yjmcc.2014.07.015
37. Sun YB, Lou F, Irving M. 2009. Calcium- and myosin-dependent changes in troponin structure during activation of heart muscle. *J Physiol* 587: 155-63. doi:10.1113/jphysiol.2008.164707

38. Zhang X, Kampourakis T, Yan Z, Sevrieva I, Irving M, Sun YB. 2017. Distinct contributions of the thin and thick filaments to length-dependent activation in heart muscle. *Elife* 6. doi:10.7554/eLife.24081
39. Corrie JE, Brandmeier BD, Ferguson RE, Trentham DR, Kendrick-Jones J, et al. 1999. Dynamic measurement of myosin light-chain-domain tilt and twist in muscle contraction. *Nature* 400: 425-30. doi:10.1038/22704
40. Brunello E, Marcucci L, Irving M, Fusi L. 2023. Activation of skeletal muscle is controlled by a dual-filament mechano-sensing mechanism. *Proc Natl Acad Sci U S A* 120: e2302837120. doi:10.1073/pnas.2302837120
41. Caremani M, Fusi L, Linari M, Reconditi M, Piazzesi G, et al. 2021. Dependence of thick filament structure in relaxed mammalian skeletal muscle on temperature and interfilament spacing. *J Gen Physiol* 153. doi:10.1085/jgp.202012713
42. Fusi L, Huang Z, Irving M. 2015. The Conformation of Myosin Heads in Relaxed Skeletal Muscle: Implications for Myosin-Based Regulation. *Biophysical Journal* 109: 783-92. doi:10.1016/j.bpj.2015.06.038
43. Caremani M, Brunello E, Linari M, Fusi L, Irving TC, et al. 2019. Low temperature traps myosin motors of mammalian muscle in a refractory state that prevents activation. *J Gen Physiol* 151: 1272-86. doi:10.1085/jgp.201912424
44. Lopez-Davila AJ, Chalovich JM, Zittrich S, Piep B, Matinmehr F, et al. 2020. Cycling Cross-Bridges Contribute to Thin Filament Activation in Human Slow-Twitch Fibers. *Front Physiol* 11: 144. doi:10.3389/fphys.2020.00144
45. Caremani M, Marcello M, Morotti I, Pertici I, Squarci C, et al. 2022. The force of the myosin motor sets cooperativity in thin filament activation of skeletal muscles. *Commun Biol* 5: 1266. doi:10.1038/s42003-022-04184-0
46. Risi C, Eisner J, Belknap B, Heeley DH, White HD, et al. 2017. Ca(2+)-induced movement of tropomyosin on native cardiac thin filaments revealed by cryoelectron microscopy. *Proc Natl Acad Sci U S A* 114: 6782-87. doi:10.1073/pnas.1700868114
47. Ovejero JG, Fusi L, Park-Holohan SJ, Ghisleni A, Narayanan T, et al. 2022. Cooling intact and demembranated trabeculae from rat heart releases myosin motors from their inhibited conformation. *J Gen Physiol* 154. doi:10.1085/jgp.202113029
48. Geeves MA, Lehrer SS. 1994. Dynamics of the muscle thin filament regulatory switch: the size of the cooperative unit. *Biophys J* 67: 273-82. doi:10.1016/S0006-3495(94)80478-3
49. Ma W, Irving TC. 2022. Small Angle X-ray Diffraction as a Tool for Structural Characterization of Muscle Disease. *Int J Mol Sci* 23. doi:10.3390/ijms23063052
50. Matsuo T, Yagi N. 2008. Structural changes in the muscle thin filament during contractions caused by single and double electrical pulses. *J Mol Biol* 383: 1019-36. doi:10.1016/j.jmb.2008.09.007
51. Iwamoto H, Suzuki T, Fujisawa T. 2000. Time-resolved two-dimensional X-ray diffraction study of the effect of shortening on activation of contracting skeletal muscle. *Pflugers Arch* 439: 646-9. doi:10.1007/s004249900224
52. Tamura T, Wakayama J, Inoue K, Yagi N, Iwamoto H. 2009. Dynamics of thin-filament activation in rabbit skeletal muscle fibers examined by time-resolved x-ray diffraction. *Biophysical Journal* 96: 1045-55. doi:10.1016/j.bpj.2008.09.022
53. Bershtsky SY, Koubassova NA, Ferenczi MA, Kopylova GV, Narayanan T, Tsaturyan AK. 2017. The Closed State of the Thin Filament Is Not Occupied in Fully Activated Skeletal Muscle. *Biophys J* 112: 1455-61. doi:10.1016/j.bpj.2017.02.017
54. Narayanan T, Sztucki M, Van Vaerenbergh P, Leonardon J, Gorini J, et al. 2018. A multipurpose instrument for time-resolved ultra-small-angle and coherent X-ray scattering. *J Appl Crystallogr* 51: 1511-24. doi:10.1107/S1600576718012748

55. Fusi L, Brunello E, Sevrieva IR, Sun YB, Irving M. 2014. Structural dynamics of troponin during activation of skeletal muscle. *Proceedings of the National Academy of Sciences of the United States of America* 111: 4626-31. doi:10.1073/pnas.1321868111
56. Bell MG, Lankford EB, Gonye GE, Ellis-Davies GC, Martyn DA, et al. 2006. Kinetics of cardiac thin-filament activation probed by fluorescence polarization of rhodamine-labeled troponin C in skinned guinea pig trabeculae. *Biophys J* 90: 531-43. doi:10.1529/biophysj.105.072769
57. Tonino P, Kiss B, Gohlke J, Smith JE, 3rd, Granzier H. 2019. Fine mapping titin's C-zone: Matching cardiac myosin-binding protein C stripes with titin's super-repeats. *J Mol Cell Cardiol* 133: 47-56. doi:10.1016/j.yjmcc.2019.05.026
58. Bennett P, Rees M, Gautel M. 2020. The Axial Alignment of Titin on the Muscle Thick Filament Supports Its Role as a Molecular Ruler. *J Mol Biol* 432: 4815-29. doi:10.1016/j.jmb.2020.06.025
59. Harris SP. 2021. Making waves: A proposed new role for myosin-binding protein C in regulating oscillatory contractions in vertebrate striated muscle. *J Gen Physiol* 153. doi:10.1085/jgp.202012729
60. Nag S, Trivedi DV, Sarkar SS, Adhikari AS, Sunitha MS, et al. 2017. The myosin mesa and the basis of hypercontractility caused by hypertrophic cardiomyopathy mutations. *Nat Struct Mol Biol*. doi:10.1038/nsmb.3408
61. Ponnam S, Sevrieva I, Sun YB, Irving M, Kampaourakis T. 2019. Site-specific phosphorylation of myosin binding protein-C coordinates thin and thick filament activation in cardiac muscle. *Proc Natl Acad Sci U S A* 116: 15485-94. doi:10.1073/pnas.1903033116
62. Previs MJ, Beck Previs S, Gulick J, Robbins J, Warshaw DM. 2012. Molecular mechanics of cardiac myosin-binding protein C in native thick filaments. *Science* 337: 1215-8. doi:10.1126/science.1223602
63. Zoghbi ME, Woodhead JL, Moss RL, Craig R. 2008. Three-dimensional structure of vertebrate cardiac muscle myosin filaments. *Proceedings of the National Academy of Sciences of the United States of America* 105: 2386-90. doi:10.1073/pnas.0708912105
64. Al-Khayat HA, Kensler RW, Squire JM, Marston SB, Morris EP. 2013. Atomic model of the human cardiac muscle myosin filament. *Proceedings of the National Academy of Sciences of the United States of America* 110: 318-23. doi:10.1073/pnas.1212708110
65. Alamo L, Ware JS, Pinto A, Gillilan RE, Seidman JG, et al. 2017. Effects of myosin variants on interacting-heads motif explain distinct hypertrophic and dilated cardiomyopathy phenotypes. *Elife* 6. doi:10.7554/eLife.24634
66. Craig R, Padron R. 2022. Structural basis of the super- and hyper-relaxed states of myosin II. *J Gen Physiol* 154. doi:10.1085/jgp.202113012
67. Lee KH, Sulbaran G, Yang S, Mun JY, Alamo L, et al. 2018. Interacting-heads motif has been conserved as a mechanism of myosin II inhibition since before the origin of animals. *Proc Natl Acad Sci U S A* 115: E1991-E2000. doi:10.1073/pnas.1715247115
68. Grinzato A, Auguin D, Kikuti C, Nandwani N, Moussaoui D, et al. 2023. Cryo-EM structure of the folded-back state of human beta-cardiac myosin. *Nat Commun* 14: 3166. doi:10.1038/s41467-023-38698-w
69. Dutta D, Nguyen V, Campbell KS, Padron R, Craig R. 2023. Cryo-EM structure of the human cardiac myosin filament. *bioRxiv*. doi:10.1101/2023.04.11.536274
70. Tamborrini D, Wang Z, Wagner T, Tacke S, Stabrin M, et al. 2023. In situ structures from relaxed cardiac myofibrils reveal the organization of the muscle thick filament. *bioRxiv*: 2023.04.11.536387. doi:10.1101/2023.04.11.536387
71. Huang X, Torre I, Chiappi M, Yin Z, Vydyanath A, et al. 2023. Cryo-electron tomography of intact cardiac muscle reveals myosin binding protein-C linking myosin and actin filaments. *J Muscle Res Cell Motil*. doi:10.1007/s10974-023-09647-3
72. Stewart MA, Franks-Skiba K, Chen S, Cooke R. 2010. Myosin ATP turnover rate is a mechanism involved in thermogenesis in resting skeletal muscle fibers. *Proceedings of the*

- National Academy of Sciences of the United States of America* 107: 430-5.  
doi:10.1073/pnas.0909468107
73. Walklate J, Kao K, Regnier M, Geeves MA. 2022. Exploring the super-relaxed state of myosin in myofibrils from fast-twitch, slow-twitch, and cardiac muscle. *J Biol Chem* 298: 101640. doi:10.1016/j.jbc.2022.101640
  74. Hooijman P, Stewart MA, Cooke R. 2011. A new state of cardiac myosin with very slow ATP turnover: a potential cardioprotective mechanism in the heart. *Biophys J* 100: 1969-76. doi:10.1016/j.bpj.2011.02.061
  75. Cooke R. 2011. The role of the myosin ATPase activity in adaptive thermogenesis by skeletal muscle. *Biophys Rev* 3: 33-45. doi:10.1007/s12551-011-0044-9
  76. Nogara L, Naber N, Pate E, Canton M, Reggiani C, Cooke R. 2016. Piperine's mitigation of obesity and diabetes can be explained by its up-regulation of the metabolic rate of resting muscle. *Proc Natl Acad Sci U S A* 113: 13009-14. doi:10.1073/pnas.1607536113
  77. Ma W, McMillen TS, Childers MC, Gong H, Regnier M, Irving T. 2023. Structural OFF/ON transitions of myosin in relaxed porcine myocardium predict calcium-activated force. *Proc Natl Acad Sci U S A* 120: e2207615120. doi:10.1073/pnas.2207615120
  78. Alamo L, Pinto A, Sulbaran G, Mavarez J, Padron R. 2017. Lessons from a tarantula: new insights into myosin interacting-heads motif evolution and its implications on disease. *Biophys Rev*. doi:10.1007/s12551-017-0292-4
  79. Trivedi DV, Adhikari AS, Sarkar SS, Ruppel KM, Spudich JA. 2017. Hypertrophic cardiomyopathy and the myosin mesa: viewing an old disease in a new light. *Biophys Rev*. doi:10.1007/s12551-017-0274-6
  80. Toepfer CN, Wakimoto H, Garfinkel AC, McDonough B, Liao D, et al. 2019. Hypertrophic cardiomyopathy mutations in MYBPC3 dysregulate myosin. *Sci Transl Med* 11. doi:10.1126/scitranslmed.aat1199
  81. Reconditi M, Brunello E, Linari M, Bianco P, Narayanan T, et al. 2011. Motion of myosin head domains during activation and force development in skeletal muscle. *Proceedings of the National Academy of Sciences of the United States of America* 108: 7236-40. doi:10.1073/pnas.1018330108
  82. Reconditi M, Caremani M, Pinzauti F, Powers JD, Narayanan T, et al. 2017. Myosin filament activation in the heart is tuned to the mechanical task. *Proc Natl Acad Sci U S A* 114: 3240-45. doi:10.1073/pnas.1619484114
  83. Nelson SR, Li A, Beck-Previs S, Kennedy GG, Warshaw DM. 2020. Imaging ATP Consumption in Resting Skeletal Muscle: One Molecule at a Time. *Biophys J* 119: 1050-55. doi:10.1016/j.bpj.2020.07.036
  84. Pilagov M, Heling L, Walklate J, Geeves MA, Kad NM. 2023. Single-molecule imaging reveals how mavacamten and PKA modulate ATP turnover in skeletal muscle myofibrils. *J Gen Physiol* 155. doi:10.1085/jgp.202213087
  85. Nelson S, Beck-Previs S, Sadayappan S, Tong C, Warshaw DM. 2023. Myosin-binding protein C stabilizes, but is not the sole determinant of SRX myosin in cardiac muscle. *J Gen Physiol* 155. doi:10.1085/jgp.202213276
  86. Ovejero JG, Fusi L, Park-Holohan S, Ghisleni A, Narayanan T, et al. 2019. The OFF-to-ON transition of thick filaments in isolated trabeculae from rat heart induced by cooling. *Biophysical Journal* 116: 263A. doi:10.1016/j.bpj.2018.11.1429
  87. Park-Holohan SJ, Brunello E, Kampourakis T, Rees M, Irving M, Fusi L. 2021. Stress-dependent activation of myosin in the heart requires thin filament activation and thick filament mechanosensing. *Proc Natl Acad Sci U S A* 118. doi:10.1073/pnas.2023706118
  88. Hill C, Brunello E, Fusi L, Ovejero JG, Irving M. 2022. Activation of the myosin motors in fast-twitch muscle of the mouse is controlled by mechano-sensing in the myosin filaments. *J Physiol* 600: 3983-4000. doi:10.1113/JP283048

89. Brunello E, Bianco P, Piazzesi G, Linari M, Reconditi M, et al. 2006. Structural changes in the myosin filament and cross-bridges during active force development in single intact frog muscle fibres: stiffness and X-ray diffraction measurements. *J Physiol* 577: 971-84. doi:10.1113/jphysiol.2006.115394
90. Caremani M, Fusi L, Reconditi M, Piazzesi G, Narayanan T, et al. 2023. Dependence of myosin filament structure on intracellular calcium concentration in skeletal muscle *Under review*.
91. Marcucci L, Reggiani C. 2016. Mechanosensing in Myosin Filament Solves a 60 Years Old Conflict in Skeletal Muscle Modeling between High Power Output and Slow Rise in Tension. *Front Physiol* 7: 427. doi:10.3389/fphys.2016.00427
92. Marcucci L, Washio T, Yanagida T. 2017. Titin-mediated thick filament activation, through a mechanosensing mechanism, introduces sarcomere-length dependencies in mathematical models of rat trabecula and whole ventricle. *Sci Rep* 7: 5546. doi:10.1038/s41598-017-05999-2
93. Campbell KS, Janssen PML, Campbell SG. 2018. Force-Dependent Recruitment from the Myosin Off State Contributes to Length-Dependent Activation. *Biophys J* 115: 543-53. doi:10.1016/j.bpj.2018.07.006
94. Mijailovich SM, Prodanovic M, Poggesi C, Geeves MA, Regnier M. 2021. Multiscale modeling of twitch contractions in cardiac trabeculae. *J Gen Physiol* 153. doi:10.1085/jgp.202012604
95. Marcucci L, Washio T, Yanagida T. 2019. Proposed mechanism for the length dependence of the force developed in maximally activated muscles. *Sci Rep* 9: 1317. doi:10.1038/s41598-018-36706-4
96. Ma W, Nag S, Gong H, Qi L, Irving TC. 2022. Cardiac myosin filaments are directly regulated by calcium. *J Gen Physiol* 154. doi:10.1085/jgp.202213213
97. Holroyde MJ, Potter JD, Solaro RJ. 1979. The calcium binding properties of phosphorylated and unphosphorylated cardiac and skeletal myosins. *J Biol Chem* 254: 6478-82.
98. Previs MJ, Mun JY, Michalek AJ, Previs SB, Gulick J, et al. 2016. Phosphorylation and calcium antagonistically tune myosin-binding protein C's structure and function. *Proc Natl Acad Sci U S A* 113: 3239-44. doi:10.1073/pnas.1522236113
99. Matsubara I, Yagi N, Endoh M. 1979. Movement of myosin heads during a heart beat. *Nature* 278: 474-6. doi:10.1038/278474a0
100. Matsubara I, Yagi N, Endoh M. 1980. The states of myosin heads in heart muscle during systolic and diastolic phases. *Eur Heart J Suppl A*: 17-20.
101. Previs MJ. 2023. Calcium activation through thick and thin? *J Gen Physiol* 155. doi:10.1085/jgp.202213265
102. Mun JY, Previs MJ, Yu HY, Gulick J, Tobacman LS, et al. 2014. Myosin-binding protein C displaces tropomyosin to activate cardiac thin filaments and governs their speed by an independent mechanism. *Proceedings of the National Academy of Sciences of the United States of America* 111: 2170-5. doi:10.1073/pnas.1316001111
103. Kampourakis T, Yan Z, Gautel M, Sun YB, Irving M. 2014. Myosin binding protein-C activates thin filaments and inhibits thick filaments in heart muscle cells. *Proceedings of the National Academy of Sciences of the United States of America* 111: 18763-8. doi:10.1073/pnas.1413922112
104. Tskhovrebova L, Trinick J. 2003. Titin: properties and family relationships. *Nat Rev Mol Cell Biol* 4: 679-89. doi:10.1038/nrm1198
105. Linke WA. 2018. Titin Gene and Protein Functions in Passive and Active Muscle. *Annu Rev Physiol* 80: 389-411. doi:10.1146/annurev-physiol-021317-121234
106. Tonino P, Kiss B, Strom J, Methawasin M, Smith JE, 3rd, et al. 2017. The giant protein titin regulates the length of the striated muscle thick filament. *Nat Commun* 8: 1041. doi:10.1038/s41467-017-01144-9

107. Powers JD, Bianco P, Pertici I, Reconditi M, Lombardi V, Piazzesi G. 2019. Contracting striated muscle has a dynamic I-band spring with an undamped stiffness one hundred times larger than the passive stiffness. *Journal of Physiology*.
108. Labeit D, Watanabe K, Witt C, Fujita H, Wu Y, et al. 2003. Calcium-dependent molecular spring elements in the giant protein titin. *Proceedings of the National Academy of Sciences of the United States of America* 100: 13716-21. doi:10.1073/pnas.2235652100
109. Cornachione AS, Leite F, Bagni MA, Rassier DE. 2016. The increase in non-cross-bridge forces after stretch of activated striated muscle is related to titin isoforms. *Am J Physiol Cell Physiol* 310: C19-26. doi:10.1152/ajpcell.00156.2015
110. Squarci C, Bianco P, Reconditi M, Pertici I, Caremani M, et al. 2023. Titin activates myosin filaments in skeletal muscle by switching from an extensible spring to a mechanical rectifier. *Proc Natl Acad Sci U S A* 120: e2219346120. doi:10.1073/pnas.2219346120
111. Li Y, Hessel AL, Unger A, Ing D, Recker J, et al. 2020. Graded titin cleavage progressively reduces tension and uncovers the source of A-band stability in contracting muscle. *Elife* 9. doi:10.7554/eLife.64107
112. Hessel AL, Ma W, Mazara N, Rice PE, Nissen D, et al. 2022. Titin force in muscle cells alters lattice order, thick and thin filament protein formation. *Proc Natl Acad Sci U S A* 119: e2209441119. doi:10.1073/pnas.2209441119
113. Brunello E, Reconditi M, Elangovan R, Linari M, Sun YB, et al. 2007. Skeletal muscle resists stretch by rapid binding of the second motor domain of myosin to actin. *Proc Natl Acad Sci U S A* 104: 20114-9. doi:10.1073/pnas.0707626104
114. Fusi L, Reconditi M, Linari M, Brunello E, Elangovan R, et al. 2010. The mechanism of the resistance to stretch of isometrically contracting single muscle fibres. *Journal of Physiology-London* 588: 495-510. doi:DOI 10.1113/jphysiol.2009.178137
115. Caremani M, Pinzauti F, Powers JD, Governali S, Narayanan T, et al. 2018. Inotropic interventions do not change the resting state of myosin motors during cardiac diastole. *J Gen Physiol*. doi:10.1085/jgp.201812196
116. Ait-Mou Y, Hsu K, Farman GP, Kumar M, Greaser ML, et al. 2016. Titin strain contributes to the Frank-Starling law of the heart by structural rearrangements of both thin- and thick-filament proteins. *Proc Natl Acad Sci U S A*. doi:10.1073/pnas.1516732113
117. Kellermayer D, Smith JE, 3rd, Granzier H. 2019. Titin mutations and muscle disease. *Pflugers Arch* 471: 673-82. doi:10.1007/s00424-019-02272-5
118. Loescher CM, Hobbach AJ, Linke WA. 2022. Titin (TTN): from molecule to modifications, mechanics, and medical significance. *Cardiovasc Res* 118: 2903-18. doi:10.1093/cvr/cvab328
119. Herman DS, Lam L, Taylor MR, Wang L, Teekakirikul P, et al. 2012. Truncations of titin causing dilated cardiomyopathy. *N Engl J Med* 366: 619-28. doi:10.1056/NEJMoa1110186
120. Rees M, Nikoopour R, Fukuzawa A, Kho AL, Fernandez-Garcia MA, et al. 2021. Making sense of missense variants in TTN-related congenital myopathies. *Acta Neuropathol* 141: 431-53. doi:10.1007/s00401-020-02257-0
121. de Tombe PP, Mateja RD, Tachampa K, Mou YA, Farman GP, Irving TC. 2010. Myofilament length dependent activation. *Journal of Molecular and Cellular Cardiology* 48: 851-58. doi:DOI 10.1016/j.yjmcc.2009.12.017
122. Sequeira V, van der Velden J. 2017. The Frank-Starling Law: a jigsaw of titin proportions. *Biophys Rev* 9: 259-67. doi:10.1007/s12551-017-0272-8
123. Allen DG, Kentish JC. 1985. The Cellular Basis of the Length Tension Relation in Cardiac-Muscle. *Journal of Molecular and Cellular Cardiology* 17: 821-40. doi:Doi 10.1016/S0022-2828(85)80097-3
124. Han JC, Taberner AJ, Loisel DS, Tran K. 2022. Cardiac efficiency and Starling's Law of the Heart. *J Physiol* 600: 4265-85. doi:10.1113/JP283632

125. Sequeira V, Wijnker PJ, Nijenkamp LL, Kuster DW, Najafi A, et al. 2013. Perturbed length-dependent activation in human hypertrophic cardiomyopathy with missense sarcomeric gene mutations. *Circ Res* 112: 1491-505. doi:10.1161/CIRCRESAHA.111.300436
126. Tanner BCW, Awinda PO, Agonias KB, Attili S, Blair CA, et al. 2023. Sarcomere length affects Ca<sup>2+</sup> sensitivity of contraction in ischemic but not non-ischemic myocardium. *J Gen Physiol* 155. doi:10.1085/jgp.202213200
127. Ma W, Henze M, Anderson RL, Gong H, Wong FL, et al. 2021. The Super-Relaxed State and Length Dependent Activation in Porcine Myocardium. *Circ Res* 129: 617-30. doi:10.1161/CIRCRESAHA.120.318647
128. Kampourakis T, Irving M. 2021. The regulatory light chain mediates inactivation of myosin motors during active shortening of cardiac muscle. *Nat Commun* 12: 5272. doi:10.1038/s41467-021-25601-8