

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.



ISSN: 0015-5659

e-ISSN: 1644-3284

Oxidative stress mediates hippocampal neuronal apoptosis through ROS/JNK/P53 pathway in rats with PTSD triggered by high-voltage electrical burn

Authors: Ying Lv, Xuegang Zhao, Rui Zhang, Zhaopeng He, Yanfeng Xu, Lihong Tu, Lei Jiang, Shunjiang Xu, Qingfu Zhang

DOI: 10.5603/fm.95727

Article type: Original article

Submitted: 2023-05-28

Accepted: 2023-08-21

Published online: 2023-09-07

This article has been peer reviewed and published immediately upon acceptance. It is an open access article, which means that it can be downloaded, printed, and distributed freely, provided the work is properly cited.

Articles in "Folia Morphologica" are listed in PubMed.

Oxidative stress mediates hippocampal neuronal apoptosis through ROS/JNK/P53 pathway in rats with PTSD triggered by high-voltage electrical burn

Ying Lv et al., Oxidative stress mediates hippocampal neuronal apoptosis through ROS/JNK/P53 pathway

Ying Lv^{1,2}, Xuegang Zhao³, Rui Zhang³, Zhaopeng He¹, Yanfeng Xu¹, Lihong Tu¹, Lei Jiang³, Shunjiang Xu³, Qingfu Zhang¹

¹Department of Burns and Plastic Surgery, the First Hospital of Hebei Medical University, Shijiazhuang, China

²Department of Cardiac Surgery, the First Hospital of Hebei Medical University, Shijiazhuang, China

³Hebei Key Laboratory of Brain Science and Psychiatric-Psychologic Disease, Shijiazhuang, China

Address for correspondence: Qingfu Zhang and Shunjiang Xu, The First Hospital of Hebei Medical University, No. 89, Donggang Road, Shijiazhuang, 050031, Hebei Province, China, tel/fax: 0311-87156771, e-mail: zhangqfdoctor2@163.com; xushunjiang@hebmh.edu.cn

ABSTRACT

Background: The pathogenesis of post-traumatic stress disorder (PTSD) triggered by high-voltage electrical burn (HVEB) remains unclear and the oxidative stress plays a role in this process. The purpose of this study is to investigate the underlying mechanism of oxidative stress mediates hippocampal neuronal apoptosis in rats with PTSD triggered by HVEB.

Materials and methods: The PTSD rat model was developed by stimulating with high voltage electricity and screened using behavioral performance including Morris water maze (MWM), elevated plus-maze (EPM) and open-field test (OFT). The

reactive oxygen species (ROS) generation was measured by DHE fluorescence staining or flow cytometry. Western blotting assay was used to detect the proteins of p-JNK, JNK, P53, PUMA, Bcl-2 and Bax in hippocampal tissue or HT22 cells treated with electrical stimulation.

Results: The serum MDA and 8-OHdG levels were increased ($P < 0.001$), while the activities of SOD and CAT were decreased ($P < 0.001$) significantly in patients with HVEB. Behavioral test results showed that high-voltage electric stimulation induced the PTSD-like symptoms and the ROS-JNK-P53 pathway was involved in the neuronal apoptosis in rats with PTSD induced by HVEB. In vitro experiments further confirmed the electrical stimulation induced neuronal apoptosis through ROS/JNK/P53 signaling pathway and the antioxidant NAC could rescued the ROS generation, activation of JNK/P53 proteins and improved the cell apoptosis rate in HT22 cells. Finally, the JNK inhibitor SP600125 could significantly inhibited the percentage of HT22 cell apoptosis induced by electrical stimulation ($P < 0.001$).

Conclusions: These results indicated that oxidative stress mediates hippocampal neuronal apoptosis through ROS/JNK/P53 pathway in rats with PTSD triggered by HVEB.

Key words: high voltage electric burn, P53 protein, oxidative stress, brain injury, PTSD

INTRODUCTION

High-voltage electrical burn (HVEB) is a rare yet destructive class of injury that has been considered as a serious public health issue worldwide [1, 2, 3, 4]. The high-voltage alternating current causes a series of damages such as muscle tonic contraction and intracellular ion disturbances through thermal and chemical effects, resulting in progressive damage to the body [5, 6, 7]. Post-traumatic stress disorder (PTSD), a mental health disorder, is triggered by a frightening and/or traumatic event — whether experienced or witnessed. The clinical symptoms of PTSD may include

recurrence of traumatic events, avoidance of traumatic cues, continuous hypervigilance, selective forgetting, emotional numbness, nightmares and severe anxiety [8, 9, 10]. It is a state of post-traumatic psychological imbalance, which can easily cause intense psychological pain and severe damage to social functions of patients, and is the main manifestation of stress disorder [11, 12]. As the HVEB is a terrifying life event, researches have shown that 41 % of electrical burn patients have the post-traumatic stress response, and 7.6 % of them have been diagnosed as PTSD [13, 14]. To our knowledge, there is no accurate statistical data on the incidence of electrical burn patients complicated by mental disorders in China and even in the world. At the present stage, researches on the pathogenesis of PTSD triggered by high-voltage electric burn are still in its infancy, and the exact molecular mechanism remains unclear.

Oxidative stress is a status of imbalance between cellular generation of reactive oxygen species (ROS) and the antioxidant defense systems, causing to cellular injury. Researches have suggested the link between oxidative stress and high-voltage electric burn injury in patients or animal models [15, 16, 17, 18]. Severe life stress events, such as HVEB, could stimulate the increase of sympathetic nerve-adrenergic medulla responsiveness, leading to the dysfunction of the hypothalamic-pituitary-adrenal axis (HPA axis) of the nervous system and contribute to the pathogenesis of PTSD [19, 20, 21]. The main reason for oxidative stress triggered by high-voltage electric burns is that the electrothermal effects can directly cause the cell damage of multiple organs. The ROS generation, 8-oxo-deoxyguanosine (8-OHdG) and the levels of end product of lipid peroxides (malondialdehyde, MDA) are most commonly used markers for oxidative stress studies. Meanwhile, the activities of SOD and CAT represent the main antioxidant enzymes in the body. The p53 protein is a key mediator of the DNA damage response and tumor suppression and plays an important role in the cellular response to oxidative stress [22, 23, 24]. Studies have found that the oxidative stress can activate P53 protein through ROS-JNK pathway [25]. P53 is involved in apoptosis and is also related to cell death pathways such as programmed cell necrosis

and autophagy [26, 27]. When the body is in a continuous and high-level oxidative stress state, the P53 protein on the one hand triggers the overexpression of P53-inducible gene 3 (PIG3) and proline oxidase (proline oxidase) to reduce the mitochondrial membrane Potential, aggravate oxidative stress damage, leading to DNA damage and apoptosis [28]. On the other hand, it promotes apoptosis by inducing pro-apoptotic proteins (PUMA and Bax) [29]. Imaging studies suggest that the continuous reduction of hippocampal volume in PTSD patients might be related to the activation of P53 protein by high levels of oxidative stress to trigger neuronal apoptosis [30, 31]. However, the role of oxidative stress and its underlying mechanism in the pathogenesis of PTSD triggered by High-voltage electrical burn need to be further investigated.

In this study, the PTSD rat model triggered by HVEB and the HT22 cell model stimulating with electrical current were used to detect the oxidative stress biomarkers and the proteins related to the ROS/JNK/P53 pathway and the neuronal apoptosis. The research will provide an experimental basis for further elucidating the mechanism of high-voltage electrical burn-induced PTSD and screening the potential intervention targets in clinic.

MATERIALS AND METHODS

Participants and serum samples

This study included 30 patients with HVEB and 30 subjects of healthy control. The patients with HVEB were recruited from the department of burns and plastic surgery, the First Hospital of Hebei Medical University. The informed consent was obtained from all participants in accordance with the protocol approved by the Research Ethics Committee of the First Hospital of Hebei Medical University. The demographic characteristics of the participants in the study were shown in Table 1. A fasting blood sample (3 mL) was collected from each subject, and placed in a test tube (BD Biosciences, NJ, USA), and placed vertically for 30 min at room temperature. Blood samples were centrifuged at 3000 g for 10 minutes at room

temperature and then at 16,000 g for 10 minutes at 4°C. The serum was aliquoted and stored at -80°C until use. Avoid repeated freeze/thaw cycles. The neuropsychological symptoms of patients with high-voltage burn were evaluated during the acute phase in Table 2.

Animals

Fifty healthy male SD rats, weighing 200-250 g, were purchased from the Experimental Animal Center of Hebei Medical University. The experimental animals were kept in a clean-grade animal room with a temperature of (24±3) °C, a relative humidity of (40±5) %, and natural light. The experimental protocol was reviewed and approved by the Experimental Animal Ethics Committee of the First Hospital of Hebei Medical University. The SD rats were randomly divided into control group (without electric stimulation), Single-Prolonged Stress (SPS) group and PTSD with high voltage electric burn (HVEB) group. Ten rats in the control group were selected as the sham shock, ten rats in the SPS group were selected as positive control group, and the others were given high-voltage electrical stimulation as the HVEB group to establish a high-voltage electrical PTSD model. After the high-voltage electrical stimulation, the rats in the three groups were subjected to a 5-day behavioral test using the Morris water maze automatic recorder. Behavioral tests (water maze test, open field test and elevated plus-maze test) were performed after two weeks of high-voltage electric stimulation.

Preparation of high-voltage electric burn triggered PTSD animal model

The laboratory humidity was at 35~45 % RH and the temperature was at 24~26°C. The rats were supine on the insulating table after they were anesthetized with 10 ml/kg of 3 % pentobarbital via intraperitoneal (ip) injection. Their limbs were fixed, and send it between the high-voltage output circuits. Fix the current inlet electrode plate on the rat's head and the outlet electrode plate in the hair removal area of the left lower extremity, after confirming that the high-voltage line is connected

correctly, turn on the 220 V power supply, turn on the voltage regulator, and turn the voltage regulator button. When the voltage rises to 2 kV, the switch is turned on and the current is actually passed (1.92 ± 0.24) A. Power on for 3 seconds, immediately disconnect the switch and release the electrode plate. Check the vital signs of the rat, and immediately perform cardiac compression resuscitation for respiratory and cardiac arrest. Immediately after the electric shock, the rats showed muscle spasms all over the body, and the skin of the current outlet and inlet was browned, and some of them became black and carbonized. The model was successfully replicated. The control group was subjected to a sham electric experiment. Before the experiment, the switch was turned off to ensure that no current passed through. The rest of the process was replicated with the animal model of high-voltage electric burn. The rats received high-voltage electric stimulation were treated with low-dose meloxicam (1mg/kg, subcutaneous injection, Shandong, China) as the analgetic once daily for 3 days. After two weeks of high-voltage electric stimulation, the behavioral tests of the three groups of rats were performed to screen the rats with PTSD, including Morris water maze test, open field test and elevated plus-maze apparatus.

PTSD model of Single-Extended Stress (SPS)

The protocol of SPS model used in this study was described previously [32]. Rats were restrained in clear plexiglass restraint tubes for 2 hours, and then immediately swam in clear acrylic cylinders (24 cm in diameter, 50 cm in height) in 24 °C water for 20 minutes. Then rats were placed back to recovery for 15 minutes, and exposed to diethyl ether until they became unconsciousness, all under bright white light conditions.

Behavioral Test

Survival of rats after high-voltage electrical stimulation: Except for 10 rats in control group and 10 rats in SPS group as positive control, 11 rats of the remaining 30 rats in HVEB group had died, and the mortality rate was 36 %. After two week of

high-voltage electric stimulation, the behavioral tests for the three groups of rats were performed, including Morris water maze test, open field test and elevated plus-maze test. Each group selected 6 animals for the subsequential experiment.

Morris Water Maze (MWM) Test

The rats in the control group, SPS group and the HVEB group were randomly selected, and the experiment included a concealed platform test. In the concealed platform test, the rats were placed into the water from one of the four quadrants facing the pool wall. Each time the experimental rats swim for 60 seconds to find the hidden platform. Record the swimming path and the time to find the hidden platform in the pool, which is the escape latency (EL). If the rat cannot find the platform within 60 seconds, guide it to the stage and stand for 10 seconds. Analysis software was used to analyze percentage of time spent in correct quadrant of MWM and latency to finding a hidden platform of rats. Test the spatial learning and memory abilities of rats.

Open Field (OF) Test

Randomly selected the rats in the control group, SPS group and HVEB group, and recorded the distance and time of central movement of each rat in the central area of the open field for 20 min. This experiment is used to assess the animal's alertness level, anxiety state and ability to adapt to the environment.

Elevated Plus-maze (EPM) Test

An elevated plus labyrinth device, located in a dark test room, was used to measure anxiety. The device had a maze of two open arms (50 cm × 10 cm) and two closed arms (50 cm × 10 cm × 40 cm) connected by a central square (10 cm × 10 cm) and elevated 50 cm above the floor centimeter. The maze was made of opaque black plexiglass, and a camera was installed above the maze to transmit video to a computer monitor. At the beginning of the experimental study, the rats were placed in the central area facing the open arms, and the number of open/closed arm entries (OA and CA, respectively) and open/closed arm time (OT/CT) were recorded for each rat.

Cell culture

HT-22 cells were obtained from Life Technologies (Waltham, MA, USA) and cultured in complete Dulbecco's Modification of Eagle's Medium (DMEM, Gibco, Invitrogen GmbH, Karlsruhe, Germany) supplemented with 10 % fetal bovine serum, 1 % penicillin and streptomycin at 37 °C, 5 % CO₂, and 95 % saturated humidity. When the cells reach the logarithmic growth phase, they are subcultured in proportion.

Establishment of electrical stimulation cell model

The HT22 cells were seeded onto 24-wellplates (the number of cells per well is 5×10^4) and divided into control group and electrical stimulation group. The control group was added with normal culture medium, and the electrical stimulation group was given different voltage stimulation conditions (0, 25, 50, 75, and 100 V) for 5 s with the 2×5 mm carbon graphite electrodes. Cell Counting Kit-8 (CCK-8, Dojindo Laboratories, Japan) measured the cell viability as instructed. Choose the optimal electrical voltage to prepare the electrical stimulation cell model.

Treatment of HT22 cells with N-acetylcysteine (NAC)

NAC (N-acetyl-L-cysteine) is a commonly used antioxidant. The HT22 cells were divided into control group and experimental group. Normal culture medium (without NAC) was added to the cells of control group, and the different concentrations of NAC (0.25, 0.5, 1, 2, and 4 mmol/L) were added to the cells of experimental group, and the cells were incubated for 24 h. Then the cell activity was determined by CCK-8 assay.

Cell grouping

The HT22 cells were seeded in a 96-well cell culture plate (the number of cells was 1×10^5 per well). The HT22 cells were divided into four groups, including control group, ES group (50V), NAC treatment group (1mmol/L) and ES+NAC group. Each

group has at least 3 replicate wells.

Detection of oxidative stress biomarkers

The levels of oxidative stress biomarkers, including Superoxide dismutase (SOD), Malondialdehyde (MDA), 8-oxo-deoxyguanosine (8-OHdG) and Catalase (CAT) in serum of patients or animals were detected using an enzyme-linked immunosorbent assay (ELISA) assay kits and Colorimetric Assay Kits (Elabscience Wuhan, China).

Dihydroethidium (DHE) staining

The hippocampal tissue sections were stained with Dihydroethidium (DHE, ThermoFisher, MA, USA) to detect the ROS generation in the hippocampus from rats.

Determination of the apoptotic rate of HT22 cells

Annexin V-FITC and propidium iodide (PI) (Biyuntian, Shanghai, China) fluorescence were used to analyze the percentage of apoptosis. HT22 cells were seeded in a six-well plate at a ratio of 2×10^5 cells/well, and processed as described above. The cells were washed twice with cold phosphate buffered saline (PBS), centrifuged at 4°C for 5 minutes, discarded the supernatant, added 100ul 1× Binding Buffer, and blown to a single cell suspension. Add 5 µl Annexin V-FITC and 5 µl PI Staining Solution to each tube, incubate at 25°C for 10 min in the dark, add 400 µl 1×Binding Buffer, mix and stain the sample within 1h, finally the cells were detected using flow cytometry (Kraemer Blvd, Brea, CA, USA).

Determination of ROS generation

The levels of intracellular ROS were detected by Oxidation-sensitive fluorogenic probe, 2',7'-Dichlorofluorescein Diacetate (DCFH-DA) (Biyuntian Biotechnology Company, Shanxi, China). Collect the processed cells by conventional centrifugation and discard the supernatant, add PBS to resuspend and wash 3 times, mix with DCFH-DA (10 µmol/L) solution, incubate at 37 °C for 20 min, invert and mix every

3-5 min, centrifuge remove the supernatant, resuspend in PBS, and repeat 3 times. The non-specific esterase of DCFH-DA cells removes the acetyl group, and the ROS is further oxidized to produce the fluorescent compound 2,7-dichlorofluorescein (DCF). The fluorescence signal intensity of DCF was measured by the flow cytometer, and the fluorescence intensity was measured at the excitation wavelength of 488 nm and the emission wavelength of 525 nm. The pure DCFH-DA solution was used for zero adjustment, and the value of the blank treatment group was 100 %. The above experimental steps were repeated 3 times.

Western blot analysis

Take the hippocampal tissue specimen of the experimental animal, put it in the cold lysis buffer (Bebo, Shanghai, China), homogenized at low temperature, centrifuged and take the supernatant. The expression of p-JNK, JNK, P53, PUMA, Bcl-2 and Bax proteins in the hippocampal tissue of rats were detected by Western blot assay 40 days after high-voltage stimulation. Take HT22 cells, inoculate them in a 6-well plate, culture for 24 h until the cells grow to 85 % confluence, divide the cells into control group, electrical stimulation (ES) group, NAC group, ES+NAC group. After 24 h of treatment, the cells are lysed and centrifuged. Then take the supernatant for subsequent experiments. Both the animal hippocampal tissue and HT22 cells were used to determine the protein content by the BCA (Solebold, Shanghai, China) method. After protein denaturation, 12 % polyacrylamide gel (SDS-PAGE) (Bio-rad, Shanghai, China) was used for electrophoresis to separate the protein in the gel to the nitrocellulose PVDF membrane. Block with the bovine protein BSA at room temperature for 1 h, and added the corresponding primary antibodies JNK (9252; Cell Signaling Technology, Beverly, MA, USA), p-JNK (Thr183/Tyr185) (4668; Cell Signaling Technologies, Beverly, MA, USA), P53 (ab202026, Abram, Cambridge, UK), PUMA (ab9643, Abram, Cambridge, UK), Bax (ab3250, Abram, Cambridge, UK) and Bcl-2 (ab59348, Abram, Cambridge, UK) specific antibodies (1:500) and β -actin (10004156, Proteintech, Wuhan, China)

(1:1000), incubate overnight at 4°C, wash the membrane with TBST after 3 times, add horseradish peroxidase HRP-labeled goat anti-rabbit secondary antibody (A23920, Abbkine, Beijing, China) (1:1000), incubate for 1 h at room temperature, wash the membrane 3 times with TBST, and develop the color by ECL luminescence. After washing the membrane, it will be developed. And observe the results after quantification. β -actin is used as an internal reference, and finally the target band is analyzed by scanning density, and the ratio of the gray value of each target protein/ β -actin gray value is used to reflect the relative protein expression level of each group. Repeat 3 times to get the average value. Quantity One gel analysis system analyzes the protein content of protein bands.

Statistical analysis

All data were presented as mean \pm SD, and statistical analysis were performed using SPSS software version 16.0 (SPSS, Inc., Chicago, IL, USA). The two-tailed Student's t-test was used for two-group comparisons□the ANOVA analysis was followed Dunnett's or Tukey's test for three parameter groups. A *P* value of < 0.05 was considered to be statistically significant. All experiments were independently repeated at least three times.

Ethics Approval and Consent to Participate

This study was conducted in accordance with the Guidelines for the Care and Use of Laboratory Animals of Hebei Medical University, and the experimental protocols used in this research were approved the First Hospital of Hebei Medical University.

RESULTS

The oxidative stress is involved in the process of high-voltage electric burn

To understand whether the oxidative stress take a part in the process of high-voltage electric burn, the oxidative stress biomarkers in serum including the content of malondialdehyde (MDA), the activities of superoxide dismutase (SOD), 8-oxo-deoxyguanosine (8-OHdG) level and Catalase (CAT) activity were detected by

Enzyme-linked immunoassay (ELISA) assay and Colorimetric Assay Kits. The results showed that the serum MDA content (Figure 1A) and the 8-OHdG levels (Figure 1C) were significantly increased ($P < 0.001$), while the SOD (Figure 1B) and CAT (Figure 1D) activity were significantly decreased ($P < 0.001$) in HVEB patient group compared with the control group. These results indicated that the oxidative stress injury is involved in the process of high-voltage electric burn in patients. The patients with high-voltage burn usually have different neuropsychological symptoms during acute phase, including anxiety (58 %), sleep difficulties (60 %), depression (49 %), and memory/attention difficulties (62 %) and so on.

Changes of behavioral tests in rats with high-voltage electric burn

To develop a PTSD animal model triggered by high-voltage electric burn, the rats were treated with high-voltage electric stimulation and the behavioral tests were performed to screen the rats with PTSD, including Morris water maze test, open field test and elevated plus-maze apparatus (Figure 2A). The single-prolonged stress (SPS) model was used as the positive control of the PTSD. The representative images of swimming trajectory to find the platform for rats of each group in MWM test were shown in (Figure 2B). The results of MWM test showed that the time spent in the target quadrant was decreased, the distance traveled to platform and the escape latency were increased in rats of SPS and PTSD groups compared with that of control group ($P < 0.05$) (Figure 2C-F). In the open field test, the distance in central movement, as well as the time in central region of rats in SPS and PTSD groups were reduced significantly compared with that of control group ($P < 0.001$) (Figure 2G). The results of time spent in open arms and the number of entry in open arms in elevated plus-maze (EPM) test also showed the similar trend for rats in SPS and PTSD groups (Figure 2H). These results suggested that high-voltage electric stimulation could induced the PTSD-like symptoms, such as the reduced ability of spatial learning and memory, alertness and anxiety, conditioned fear and sensitized fear and decreased environmental adaptability.

ROS-JNK-P53 pathway is involved in the neuronal apoptosis in rats with PTSD induced by HVEB

To investigate the mechanisms of oxidative stress in the pathogenesis of PTSD triggered by High-voltage electrical burn, we tested the oxidative stress biomarkers, apoptosis related proteins and its upstream signaling molecular in the hippocampus of rat model with PTSD induced by HVEB. The results of ELISA showed that the MDA contents were increased ($P < 0.05$) (Figure 3A) and the SOD activity was decreased ($P < 0.001$) (Figure 3B) in the serum of rats with PTSD compared with the control group. DHE staining results showed that the ROS generation in the hippocampal CA3 area of rats with PTSD was increased compared with the control group (Figure 3C). The expression levels of p-JNK, P53, PUMA and Bax proteins were higher, but the Bcl-2 protein was lower in the hippocampus of rats with PTSD than that of control group ($P < 0.05$) (Figure 3D). These results suggested that oxidative stress mediated hippocampal neuronal apoptosis may through ROS-JNK-P53 pathway in rats with PTSD induced by high-voltage electrical burns.

Oxidative stress participated in the neuronal apoptosis induced by electrical stimulation

To verify if the oxidative stress participated in the neuronal apoptosis induced by electrical stimulation, we developed an in vitro electrical stimulation (ES) cell model using HT-22 cell lines. The HT22 cells were treated with different voltage of electrical stimulation (ES) (0 to 100 V) for 5 s and continuously cultured for 24 h, CCK-8 reagent was used to determine the cell viability for screening the stimulation conditions preliminarily. The results showed that after 24 hours of treatment of HT22 cells with different voltage of electrical stimulation for 5s, the cell viability was gradually decline and caused significant decrease at 75 v and 100 v compared to control group, and the cell survival rate was significantly reduced in a voltage-dependent manner ($P < 0.001$) (Figure 4A). So we selected the 50 V voltage as the best stimulation

condition to prepare the ES cell model. In order to explore whether the antioxidant N-Acetyl-L-cysteine (NAC) has a protective effect on the injury induced by electrical stimulation in HT22 cells [33], different concentrations of NAC were used to incubate the HT22 cells for 24 h, and the cell survival rate was detected by CCK-8 assay. The results showed that the cell viability was decreased significantly at the concentration of 2 mmol/L NAC ($P < 0.001$) (Figure 4B). Finally, we selected the concentration of 1 mmol/L NAC for the subsequent experiments. Next, the HT22 cells were stimulated with 50 V voltage electrical current for 5 s and continuously incubated with or without 1mmol/L NAC for 24 h. We found that the viability of HT22 cells were decreased ($P < 0.001$) (Figure 4C-F), but the ROS generation ($P < 0.001$) (Figure 4D-G) and the cell apoptosis rate ($P < 0.001$) (Fig. 4E, H) were increased significantly by the electrical stimulation, and the antioxidant NAC could rescued the ROS generation and improved the cell apoptosis rate in HT22 cells. These results suggested that oxidative stress participated in the neuronal apoptosis induced by electrical stimulation in vitro.

The electrical stimulation induced neuronal apoptosis through ROS/JNK/P53 signaling pathway

The results of in vivo experiment have indicated that oxidative stress mediated hippocampal neuronal apoptosis may through ROS-JNK-P53 pathway in rats with PTSD induced by high-voltage electrical burns. To validate this conclusion, we used western blot assay to detect the expression of the JNK/P53 signaling molecular and the apoptosis related proteins in electrical stimulation-induced HT22 cells. The results showed that the expression of p-JNK, P53, PUMA, and Bax proteins were increased significantly ($P < 0.01$) in HT22 cells treated with electrical stimulation compared with the control group, and the antioxidant NAC could block these changes in the JNK/P53 signaling proteins and cell apoptosis related proteins (Figure 5A-B). The above results indicate that JNK and P53 are involved in the apoptosis induced by electrical stimulation, and NAC reversed the activation of JNK and P53, and inhibited

the up-regulation of PUMA, Bax and the down-regulation of Bcl-2. In addition, the JNK inhibitor SP600125 could significantly inhibited the percentage of HT22 cell apoptosis induced by electrical stimulation ($P < 0.001$) (Figure 5C-D). These results suggested that the ROS-JNK-P53 pathway plays a crucial role in the neuronal apoptosis induced by electrical stimulation.

DISCUSSION

Post-traumatic stress disorder (PTSD) caused by high-voltage electrical burn has a high disability and fatality rate in clinic [34, 35], its pathogenesis is still unknown, and there has been a lack of effective clinical treatment measures. Oxidative stress induced by burns causes cell apoptosis is a research hotspot in the field of burn medicine. Current studies have found that PTSD caused by high-voltage electric burn is closely related to oxidative stress [36]. This study reported that oxidative stress mediates hippocampal neuronal apoptosis through ROS/JNK/P53 pathway in PTSD rats induced by High-voltage electrical burn, which will provide a experimental basis for the prevention and treatment of PTSD induced by high-voltage electrical burn.

Oxidative stress triggered by burns increases the production of oxygen free radicals in the body and/or decreases the body's antioxidant capacity, which causes lipid peroxidation, protein oxidation, DNA damage and so on, and result in multiple organ dysfunction or even failure in burn patients, eventually leads to the death of burn patients [37, 38, 39]. But how the oxidative stress participate the pathogenesis of PTSD caused by high-voltage electric burn and it underlying mechanisms is still not unclear. In this study, we confirmed that the oxidative stress injury is involved in the disease process of high-voltage electric burn patients. At the same time, we developed a PTSD rat model triggered high-voltage electric burn. The results of water maze, open field tests and elevated plus-maze apparatus tests showed that the high-voltage electric burn PTSD model rats exhibited PTSD-like symptoms, such as the reduced ability of spatial learning and memory, increased alertness and anxiety, and decreased environmental adaptability. We found that the levels of MDA contents in the PTSD

model rats were increased significantly after high-voltage electric burns, and the SOD activities was decreased significantly. The related proteins in the ROS/JNK/P53 pathway in the hippocampus were confirmed that oxidative stress induced by high-voltage electric burn mediated the upregulation of p-JNK, P53, PUMA proteins and the downregulation of Bcl-2 protein. Previous studies have found that PUMA is a master regulator of Bax activation and neuronal apoptosis induced by oxidative stress. In this study, the expression of Bax protein has a very obvious change, which is considered to have a certain relationship with PUMA's dominant role in regulating Bax activation [40]. The above experiments confirmed that oxidative stress is closely related to the onset of PTSD in high-voltage electric burn to a certain extent.

Previous studies have found that the production of ROS plays a critical role in inducing cell apoptosis[41, 42]. ROS can induce DNA damage, mitochondrial permeability and activation of apoptosis signals, leading to cell death. In this experiment, we used electrical stimulation induced HT22 cells to prepare oxidative damage models. The study showed that the electrical stimulation significantly increased the production of ROS in the oxidative stressed HT22 cell model. As an antioxidant, NAC can effectively protect HT22 cells from apoptosis induced by ROS. The reason for the anti-ROS activity of NAC is to scavenge free radicals through the redox potential of thiols or increase the level of glutathione in cells [43]. Therefore, protecting hippocampal neurons from apoptosis induced by oxidative stress may have a very critical therapeutic value in the treatment of PTSD.

When the cells received ischemia and hypoxia stimulation persistently, the JNK signaling pathway is activated and participates in cell apoptosis. JNK is one of the members of the mitogen-activated protein kinase (MAPK) pathway. Phosphorylated JNK can induce cell apoptosis, which is very important in the exogenous and endogenous apoptotic pathways [44]. This experiment showed that high-voltage electrical stimulation induced an increase in p-JNK in the hippocampus of PTSD rats through oxidative stress, which is consistent with the increase in ROS levels. At the same time, cell experiments proved that the electrical stimulation significantly

induced the death of HT22 cells in a voltage-dependent manner, which is consistent with previous studies. This is consistent with previous research. The electrical stimulation can induce JNK phosphorylation (p-JNK), while the JNK inhibitor SP600125 can inhibit cell apoptosis, and the NAC can inhibit JNK activation [45]. The above studies have shown that the increase in p-JNK is inseparable from the increase in ROS generation. JNK is a key mediator of inducing cell apoptosis. In addition, studies have reported that the JNK signaling pathway can regulate the P53 signaling pathway and induce cell apoptosis. As a "genome guardian", P53 protein has anti-proliferation and pro-apoptosis functions. It is an important transcription factor and tumor suppressor in the body. In different types of cell stress, P53 undergoes different post-translational modifications and transactivation target genes play a critical role in different cellular pathways, such as apoptosis, autophagy and DNA repair [46]. Current studies have shown that when the body is in a state of sustained high levels of oxidative stress, DNA damage induced by ROS can activate P53 [47] and P53 can activate pro-apoptotic genes Bax, PUMA, etc. through the endogenous (mitochondrial) pathway [48, 49]. or by inhibiting the anti-apoptotic gene Bcl-2 to promote endogenous cell apoptosis [50, 51]. P53 can also directly activate Bax in the cytoplasm through non-transcriptional pathways, and induce endogenous cell apoptosis.

As we all know, when the body encounters a serious life event, it induces the body to produce a high level of oxidative stress, enter the resistance phase and exhaustion phase, and develop into PTSD [52, 53]. The continuous reduction of hippocampal volume in patients with PTSD may be related to the activation of P53 protein induced by high levels of oxidative stress and trigger neuronal apoptosis. However, it is still unclear how hippocampal neurons are damaged by high-voltage electric burn. In this study, we detected the expression of p-JNK, P53 and other related apoptotic proteins in the hippocampus of PTSD model rats through western blot assay, and the results proved that high-voltage electric burn activate the ROS/JNK/P53 pathway. By activating the P53 protein, it triggers the expression of the

pro-apoptotic protein PUMA, down-regulates Bcl-2, and promotes neuronal apoptosis. In addition, the in vitro experiment of this study also proved that electrical stimulation induced an increase in intracellular ROS production in HT22 cells, and increased the expression of p-JNK and P53 proteins, and caused apoptosis through the endogenous (mitochondrial) pathway, which was manifested in the decreased expression of anti-apoptotic molecule Bcl-2 protein and the increased expression of Bax and PUMA protein. The antioxidant NAC can prevent electrical stimulation induced cell death through JNK inactivation, and inhibit the accumulation of P53 by inhibiting the production of ROS. These results indicated that the development of PTSD caused by high-voltage electrical burn and oxidative stress in the cells model, the ROS-JNK-P53 pathway was involved in the regulation of cell apoptosis.

CONCLUSIONS

In summary, this study concluded that oxidative stress mediates hippocampal neuronal apoptosis through ROS/JNK/P53 pathway in rats with PTSD triggered by high-voltage electrical burn, which will provide an experimental basis for further exploring the pathogenesis of PTSD caused by high-voltage electric burn and screening the potential therapeutic targets.

Acknowledgments

This study was supported by the National Natural Science Foundation of China (81871560), and Key Project of Natural Science Foundation of Hebei Province of China (H2016206575).

Conflict of interest: None declared

REFERENCES

1. Kasana RA, Baba PU, Wani AH. Pattern of high voltage electrical injuries in the Kashmir valley: a 10-year single centre experience. *Ann Burns Fire Disasters*. 2016;29(4): 259-263. PMID:

- 28289358; PMID: PMC5347316.
2. Nielsen KJ, Carstensen O, Kærgaard A, et al. Neurological symptoms and disorders following electrical injury: A register-based matched cohort study. *PLoS One*. 2022;17(3): e0264857. doi: 10.1371/journal.pone.0264857. PMID: 35235596; PMID: PMC8890633.
 3. DePamphilis MA, Cauley RP, Sadeq F, et al. Reconstruction of the Upper Extremity High-Voltage Electrical Injury: A Pediatric Burn Hospital's 13-Year Experience. *J Burn Care Res*. 2022;43(3):696-703. doi: 10.1093/jbcr/irab177. PMID: 34534315.
 4. Shih JG, Shahrokhi S, Jeschke MG. Review of adult electrical burn injury outcomes worldwide: an analysis of low-voltage vs high-voltage electrical injury. *J Burn Care Res*. 2017;38(1):e293-298. doi: 10.1097/BCR.0000000000000373. PMID: 27359191; PMID: PMC5179293.
 5. Keshavarzi A, Rahgozar Z, Mortazavi M, et al. Epidemiology, Geographical Distribution, and Outcome Analysis of Patients with Electrical Burns Referred to Shiraz Burn Center, Shiraz, Iran during 2008-2019. *World J Plast Surg*. 2022;11(2):102-109. doi: 10.52547/wjps.11.2.102. PMID: 36117901; PMID: PMC9446130.
 6. Gentges J, Schieche C. Electrical injuries in the emergency department: an evidence-based review. *Emerg Med Pract*. 2018;20(11):1-20. PMID: 30358379.
 7. Salehi SH, Sadat Azad Y, Bagheri T, et al. Epidemiology of Occupational Electrical Injuries. *J Burn Care Res*. 2022;43(2):399-402. doi: 10.1093/jbcr/irab171. PMID: 34562012.
 8. Maercker A, Cloitre M, Bachem R, et al. Complex post-traumatic stress disorder. *Lancet*. 2022;400(10345):60-72. doi: 10.1016/S0140-6736(22)00821-2. PMID: 35780794.
 9. Barbano AC, van der Mei WF, Bryant RA, et al. Clinical implications of the proposed ICD-11 PTSD diagnostic criteria. *Psychol Med*. 2019;49(3):483-490.
 10. Merians AN, Spiller T, Harpaz-Rotem I, et al. Post-traumatic Stress Disorder. *Med Clin North Am*. 2023;107(1):85-99. doi: 10.1016/j.mcna.2022.04.003. Epub 2022 Oct 28. PMID: 36402502.
 11. Ressler KJ, Berretta S, Bolshakov VY, et al. Post-traumatic stress disorder: clinical and translational neuroscience from cells to circuits. *Nat Rev Neurol*. 2022;18(5):273-288. doi: 10.1038/s41582-022-00635-8. Epub 2022 Mar 29. PMID: 35352034; PMID: PMC9682920.
 12. Schrader C, Ross A. A review of PTSD and current treatment strategies. *Mo Med*. 2021;118(6): 546-551. PMID: 34924624; PMID: PMC8672952.
 13. Tamayo-Gómez A, Velásquez-Suárez J, Páramo-Duque L, et al. Epidemiology and factors associated with acute stress disorder in burned patients: a case-control study. *Burns*. 2022;48(4):995-1003. doi: 10.1016/j.burns.2021.09.024. Epub 2021 Oct 4. PMID: 34696951.
 14. Piotrowski A, Fillet AM, Perez P, et al. Outcome of occupational electrical injuries among French electric company workers: a retrospective report of 311 cases, 1996-2005. *Burns*. 2014;40(3):480-488.
 15. Radulovic N, Mason SA, Rehou S, et al. Acute and long-term clinical, neuropsychological and return-to-work sequelae following electrical injury: a retrospective cohort study. *BMJ Open*. 2019;9(5): e025990.
 16. Gürünlüoğlu K, Demircan M, Taşçi A, et al. Effects of high-voltage electrical burns and other burns on levels of serum oxidative stress and telomerase in children. *Burns*. 2018;44(8): 2034-2041. doi: 10.1016/j.burns.2013.08.008. Epub 2013 Sep 10. PMID: 24028742.
 17. Mobayen M, Sadeghi M. Prevalence and Related Factors of Electrical Burns in Patients Referred to Iranian Medical Centers: A Systematic Review and Meta-Analysis. *World J Plast Surg*. 2022;

- 11(1):3-11. doi: 10.52547/wjps.11.1.3. PMID: 35592228; PMCID: PMC9018033.
18. Gentges J, Schieche C. Electrical injuries in the emergency department: an evidence-based review. *Emerg Med Pract.* 2018;20(11):1-20. PMID: 30358379.
 19. Shahid F, Gondal MF, Us Sabah N, et al. Psychosocial Impact of Electrical Burn in Children: A Follow-Up Study Conducted at a Tertiary Care Hospital. *Cureus.* 2022;14(12):e32816. doi: 10.7759/cureus.32816. PMID: 36694523; PMCID: PMC9862975.
 20. Ritov G, Boltyansky B, Richter-Levin G. A novel approach to PTSD modeling in rats reveals alternating patterns of limbic activity in different types of stress reaction. *Mol Psychiatry.* 2016;21(5):630-641. doi: 10.1038/mp.2015.169. Epub 2015 Nov 10. PMID: 26552592; PMCID: PMC5414084.
 21. Souza RR, Noble LJ, McIntyre CK. Using the tingle prolonged stress model to examine the pathophysiology of PTSD. *Front Pharmacol.* 2017;8:615. doi: 10.3389/fphar.2017.00615. PMID: 28955225; PMCID: PMC5600994.
 22. Seetharaman S, Fleshner M, Park CR, et al. Influence of daily social stimulation on behavioral and physiological outcomes in an animal model of PTSD. *Brain Behav.* 2016; 6(5):e00458. doi: 10.1002/brb3.458. PMID: 27110436; PMCID: PMC4834360.
 23. Simabuco FM, Morale MG, Pavan ICB, et al. P53 and metabolism: from mechanism to therapeutics. *Oncotarget.* 2018;9(34): 23780-23823. doi: 10.18632/oncotarget.25267.
 24. Jiso A, Demuth P, Bachowsky M, et al. Natural merosoterpenes activate the DNA damage response via DNA strand break formation and trigger apoptotic cell death in p53-wild-type and mutant colorectal cancer. *Cancers (Basel).* 2021;13(13):3282. doi: 10.3390/cancers13133282. PMID: 34209047; PMCID: PMC8268692.
 25. Wang P, Zhang SD, Jiao J, et al. ROS-mediated p53 activation by juglone enhances apoptosis and autophagy in vivo and in vitro. *Toxicol Appl Pharmacol.* 2019;379:114647. doi: 10.1016/j.taap.2019.114647. Epub 2019 Jul 5. PMID: 31283929.
 26. Di J, Tang J, Qian H, et al. P53 upregulates PLC ϵ -IP3-Ca²⁺ pathway and inhibits autophagy through its target gene rap2B. *Oncotarget.* 2017;8(39):64657-64669. doi: 10.18632/oncotarget.18112. PMID: 29029384; PMCID: PMC5630284.
 27. Jiang L, Kon N, Li T, et al. Ferroptosis as a p53-mediated activity during tumour suppression. *Nature.* 2015;520(7545): 57-62. doi: 10.1038/nature14344. Epub 2015 Mar 18. PMID: 25799988; PMCID: PMC4455927.
 28. Tamura RE, Hunger A, Fernandes DC, et al. Induction of oxidants distinguishes susceptibility of prostate carcinoma cell lines to p53 gene transfer mediated by an improved adenoviral vector. *Hum Gene Ther.* 2017;28(8): 639-653. doi: 10.1089/hum.2016.139. Epub 2017 Feb 6. PMID: 28181816.
 29. Garufi A, Pistrutto G, Baldari S, et al. P53-dependent PUMA to dram antagonistic interplay as a key molecular switch in cell-fate decision in normal/high glucose conditions. *J Exp Clin Cancer Res.* 2017;36(1): 126. doi: 10.1186/s13046-017-0596-z. PMID: 28893313; PMCID: PMC5594515.
 30. Logue MW, van Rooij SJH, Dennis EL, et al. Smaller hippocampal volume in posttraumatic stress disorder: A multisite enigma-pgc study: subcortical volumetry results from posttraumatic stress disorder consortia. *Biol Psychiatry.* 2018;83(3): 244-253. doi: 10.1016/j.biopsych.2017.09.006. PMID: 29217296; PMCID: PMC5951719.
 31. Hasin D, Hatzenbuehler ML, Keyes K, et al. Substance use disorders: diagnostic and statistical

- manual of mental disorders, fourth edition (DSM-IV) and international classification of diseases, tenth edition (ICD-10). *Addiction*. 2006;101: 59-75. doi: 10.1111/j.1360-0443.2006.01584.x. PMID: 16930162.
32. Liberzon I, Krstov M, Young EA. Stress-restress: Effects on ACTH and fast feedback *Psychoneuroendocrinology*. 1997;22(6): 443-453. doi: 10.1016/s0306-4530(97)00044-9. PMID: 9364622.
 33. Liu X, Wang L, Cai J, et al. N-acetylcysteine alleviates H₂O₂-induced damage via regulating the redox status of intracellular antioxidants in H9c2 cells. *Int J Mol Med*. 2019;43(1): 199-208. doi: 10.3892/ijmm.2018.3962.. PMID: 30387809; PMCID: PMC6257848.
 34. Hyland P, Vallières F, Cloitre M, et al. Trauma, PTSD, and complex PTSD in the republic of Ireland: prevalence, service use, comorbidity, and risk factors. *Soc Psychiatry Psychiatr Epidemiol*. 2021;56(4): 649-658. doi: 10.1007/s00127-020-01912-x. PMID: 32632598.
 35. Daskal Y, Beicker A, Dudkiewicz M, et al. High voltage electric injury: mechanism of injury, clinical features and initial evaluation. *Harefuah*. 2019;158(1):65-69. PMID: 30663297.
 36. Clausen AN, Francisco AJ, Thelen J, et al. PTSD and cognitive symptoms relate to inhibition-related prefrontal activation and functional connectivity. *Depress Anxiety*. 2017;34(5): 427-436. doi: 10.1002/da.22613. PMID: 28370684; PMCID: PMC5408317.
 37. Rosenberg M, Mehta N, Rosenberg L, et al. Immediate and long-term psychological problems for survivors of severe pediatric electrical injury. *Burns*. 2015;41(5):1823-1830. doi: 10.1016/j.burns.2015.06.006. PMID: 26182828.
 38. Imbriani P, Martella G, Bonsi P, Pisani A. Oxidative stress and synaptic dysfunction in rodent models of Parkinson's disease. *Neurobiol Dis*. 2022;173:105851. doi: 10.1016/j.nbd.2022.105851. PMID: 36007757.
 39. Boeke EA, Moscarello JM, LeDoux JE, et al. Active avoidance: neural mechanisms and attenuation of pavlovian conditioned responding. *J Neurosci*. 2017;37(18): 4808-4818. doi: 10.1523/JNEUROSCI.3261-16.2017. PMID: 28408411; PMCID: PMC5426570.
 40. Steckley D, Karajgikar M, Dale LB, et al. Puma is a dominant regulator of oxidative stress induced Bax activation and neuronal apoptosis. *J Neurosci*. 2007;27(47): 12989-12999. doi: 10.1523/JNEUROSCI.3400-07.2007. PMID: 18032672; PMCID: PMC6673275.
 41. Jeong CH, Joo SH. Downregulation of reactive oxygen species in apoptosis. *J Cancer Prev*. 2016;21(1):13-20. doi: 10.15430/JCP.2016.21.1.13. PMID: 27051644; PMCID: PMC4819661.
 42. Suroto H, Asriel A, De Vega B, et al. Early and late apoptosis protein expression (Bcl-2, BAX and p53) in traumatic brachial plexus injury. *J Musculoskelet Neuronal Interact*. 2021;21(4):528-532. PMID: 34854392; PMCID: PMC8672401.
 43. Roy K, Wu Y, Meitzler JL, et al. NADPH oxidases and cancer. *Clin Sci (Lond)*. 2015;128(12):863-875. doi: 10.1042/CS20140542. PMID: 25818486.
 44. Yang H, Xie Y, Yang D, et al. Oxidative stress-induced apoptosis in granulosa cells involves JNK, p53 and puma. *Oncotarget*. 2017;8(15): 25310-25322. doi: 10.18632/oncotarget.15813. PMID: 28445976; PMCID: PMC5421932.
 45. Suha T, Asli M, Aynur S, et al. Effects of N-acetylcysteine and ethyl pyruvate on ischemia-reperfusion injury in experimental electrical burn model. *Am J Emerg Med*. 2016;34(7): 1217-1224. doi:10.1016/j.ajem.2016.03.032. PMID: 27085456.

46. Beyfuss K, Hood DA. A systematic review of p53 regulation of oxidative stress in skeletal muscle. *Redox Rep.* 2018;23(1):100-117. doi: 10.1080/13510002.2017.1416773. PMID: 29298131; PMCID: PMC6748683.
47. Shi T, van Soest DMK, Polderman PE, et al. DNA damage and oxidant stress activate p53 through differential upstream signaling pathways. *Free Radic Biol Med.* 2021;172:298-311. doi: 10.1016/j.freeradbiomed.2021.06.013. PMID: 34144191.
48. Lu Z, Miao Y, Muhammad I, et al. Colistin-induced autophagy and apoptosis involves the JNK-Bcl2-Bax signaling pathway and JNK-p53-ROS positive feedback loop in PC-12 cells. *Chem Biol Interact.* 2017;277:62-73. doi: 10.1016/j.cbi.2017.08.011. PMID: 28842171.
49. Wei R, Zhang R, Li H, et al. MiR-29 targets PUMA to suppress oxygen and glucose deprivation/r-epfusion (OGD/R)-induced cell death in hippocampal neurons. *Curr Neurovasc Res.* 2018;15(1): 47-54. doi: 10.2174/1567202615666180403170902. PMID: 29623838.
50. Kinarivala N, Patel R, Boustany RM, et al. Discovery of aromatic carbamates that confer neuroprotective activity by enhancing autophagy and inducing the anti-apoptotic protein B-cell lymphoma 2 (Bcl-2). *J Med Chem.* 2017;60(23):9739-9756. doi: 10.1021/acs.jmedchem.7b01199. PMID: 29110485; PMCID: PMC5995798.
51. La Colla A, Vasconsuelo A, Milanese L, et al. 17 β -estradiol protects skeletal myoblasts from apoptosis through p53, Bcl-2, and FoxO families. *J Cell Biochem.* 2017;118(1):104-115. doi: 10.1002/jcb.25616. PMID: 27249370.
52. Sun Y, Qu Y, Zhu J. The relationship between inflammation and post-traumatic stress disorder. *Front Psychiatry.* 2021;12:707543. doi: 10.3389/fpsy.2021.707543. PMID: 34456764; PMCID: PMC8385235.
53. Peruzzolo TL, Pinto JV, Roza TH, et al. Inflammatory and oxidative stress markers in post-traumatic stress disorder: a systematic review and meta-analysis. *Mol Psychiatry.* 2022;27(8):3150-3163. doi: 10.1038/s41380-022-01564-0. Epub 2022 Apr 27. PMID: 35477973.

Table 1. Demographic characteristics of the samples

Demographics	Control (n = 30)	HVEB (n = 30)	p-Value
Gender			
Male	27 (90 %)	25 (83.3 %)	
Female	3 (10 %)	5 (16.7 %)	0.706
Age (years)	35.90 \pm 11.96	36.90 \pm 9.42	0.72
BMI	23.39 \pm 2.50	23.35 \pm 2.34	0.948
TBSA %		11.50 (6.75,	
MDA (ng/mL)	1206.58 \pm 501.94	2362.02 \pm	< 0.001
SOD (U/mL)	261.49 \pm 57.36	151.01 \pm 53.97*	< 0.001
8-OHdG (ng/mL)	35.95 \pm 11.02	87.30 \pm 22.02*	< 0.001
CAT (U/mL)	475.34 \pm 132.16	216.45 \pm 82.95*	< 0.001

Note: HVEB, high-voltage electrical burn; BMI, body mass index; TBSA, total body surface area; MDA, malondialdehyde; SOD, superoxide dismutase; 8-OHdG, 8-oxo-deoxyguanosine; CAT, Catalase. *** $P < 0.001$ vs the control group. Data are shown as mean \pm standard deviation.

Table 2. Neuropsychological symptoms of patients with the high-voltage electrical burns during the acute phase

Neuropsychological	incidence % (n = 30)
Anxiety	58 %
Sleep difficulties	60 %
Depressed mood	49 %
Memory/concentration difficulties	62 %
Avoidance	29 %
Hypervigilance	15 %
Hyperarousal Nightmares	31 %
Social withdrawal	18 %
Suicidal ideations	5 %
Dizziness	12 %
Headaches	17 %
Phantom limb pain	11 %
Flashbacks	20 %
Avoidance Hypervigilance Flashbacks	25 %
	21 %

Figure 1. Alterations of oxidative stress biomarkers in serum of patients with the high-voltage electrical burns. The levels of MDA (A), SOD activity (B), 8-OHdG (C) and CAT activity (D) were detected by ELISA assay and Colorimetric Assay Kit . *** $P < 0.001$ vs the control group (n = 30).

Figure 2. Changes of behavioral performance tests in PTSD rat model induced by high-voltage electric burn. (A) Timeline of experimental procedures. (B) Representative images of swimming trajectory to find the platform in Morris water maze (MWM) test for rats in the indicated groups. (C) Time spent in target quadrant without the platform in MWM test. (D) Distance traveled to platform in MWM test. (E) Time spent in the target quadrant in MWM test. (F) Escape latency to finding a

hidden platform during five days of navigation training in MWM test. (G) Distance and time of central movement in the open-field test (OFT). (H) Time spent in open arms and number of entry in open arms in elevated plus-maze (EPM) test. Data were presented as mean \pm S.D (n = 6). * P < 0.05, ** P < 0.01, *** P < 0.001 vs the control group.

Figure 3. Oxidative stress mediated hippocampal neuronal apoptosis through ROS-JNK-P53 pathway in rats with PTSD induced by high-voltage electrical burns. (A-C) The serum levels of MDA content (A), and SOD activity (B) were detected by ELISA assay, and DHE fluorescence staining (C) was applied to detect ROS generation in the hippocampal CA3 region of rats in PTSD or control group. Scale bars = 100 μ m. (D) The expression of p-JNK, JNK, P53, PUMA, Bcl-2 and Bax protein were detected by Western blot analysis in the hippocampus of PTSD or control rats. The β -actin was used as a loading control. * P < 0.05, ** P < 0.01. *** P < 0.001 vs the control group (n = 6).

Figure 4. Oxidative stress participated in the neuronal apoptosis induced by electrical stimulation in HT22 cells. (A, B) HT22 cells were treated with different voltage (0, 25, 50, 75 and 100 V) of electrical stimulation (ES) for 5 s and continuously cultured for 24 h (A) or incubated with different concentrations (0, 0.25, 0.5, 1, 2 and 4 mmol/L) of N-Acetyl-L-cysteine (NAC) for 24 h (B), and then the cell viability was assessed using a CCK-8 assay. ** P < 0.01, *** P < 0.001 vs the control group. (C, F) The effects of electrical stimulation (50 V for 5 s) on the morphology (C) and the cell viability (F) of HT22 cells in the absence or presence of NAC (1 mmol/L). Scale bar = 200 μ m. (D, G) The protective effects of NAC against electrical stimulation-induced ROS generation were measured by flow cytometry. HT22 cells were treated with electrical stimulation in the absence and presence of NAC, and then were loaded with DCFH-DA to detect the generation of intracellular ROS. (E, H) The protective effects of NAC against electrical stimulation-induced cell apoptosis were measured by flow cytometry. Apoptotic HT22 cells were identified using Annexin V-FITC/PI dual staining. Data were shown as the mean \pm SD (n = 3). *** P < 0.001 vs the control group; ### P < 0.001 vs the ES group.

Figure 5. The JNK/P53 signaling pathway was involved in the electrical stimulation-induced apoptosis in HT22 cells. (A, B) HT22 cells were treated with electrical stimulation (50 V for 5 s) and co-treated with NAC (1mmol/L) for 24 h. Western blot assay was adopted for p-JNK, JNK, P53, PUMA, Bcl-2 and Bax protein examination. (C, D) The effects of SP600125 (JNK inhibitor) on the electrical stimulation-induced

cell apoptosis were measured by flow cytometry. HT22 cells were cultured in DMEM containing 10 % FBS, and were pretreated with 10 μ l SP600125 (JNK inhibitor) for 30 min. Then, cells were subjected to electrical stimulation for 5 s and continuously cultured for 24 h. Apoptotic cells were identified by FACS using Annexin V-FITC/PI dual-staining. Representative dot-plot diagrams of Annexin V-FITC/PI flow cytometry (C). The percentage of apoptotic cells were compared with the total cells (D). Data were shown as the mean \pm SD (n = 3). * $p < 0.05$, ** $P < 0.01$ and *** $P < 0.001$ vs the control group; # $P < 0.05$, ## $P < 0.01$ and ### $P < 0.001$ vs the ES group.









