

This is a provisional PDF only. Copyedited and fully formatted version will be made available soon.



P O L I S H G Y N E C O L O G Y

GINEKOLOGIA POLSKA

ORGAN POLSKIEGO TOWARZYSTWA GINEKOLOGICZNEGO
THE OFFICIAL JOURNAL OF THE POLISH GYNECOLOGICAL SOCIETY

ISSN: 0017-0011

e-ISSN: 2543-6767

Vulvar Merkel cell carcinoma combined with squamous cell carcinoma of the vulva

Authors: Marlena Cwynar, Ewa Chmielik, Grzegorz Cwynar, Piotr Ptak, Karolina Kowalczyk

DOI: 10.5603/gpl.96927

Article type: Clinical vignette

Submitted: 2023-08-11

Accepted: 2023-09-19

Published online: 2023-09-23

This article has been peer reviewed and published immediately upon acceptance. It is an open access article, which means that it can be downloaded, printed, and distributed freely, provided the work is properly cited.

Articles in "Ginekologia Polska" are listed in PubMed.

CLINICAL VIGNETTE

Vulvar Merkel cell carcinoma combined with squamous cell carcinoma of the vulva

Marlena Cwynar^{1,2}, Ewa Chmielik³, Grzegorz Cwynar², Piotr Ptak^{1,2}, Karolina Kowalczyk¹

¹*Department of Endocrinological Gynecology, Faculty of Medicine in Katowice, Medical University of Silesia, Poland*

²*Department of Gynecology and Gynecological Oncology, Hospital in Czeladz,, Poland*

³*Tumor Pathology Department, Maria Sklodowska-Curie National Research Institute of Oncology, Gliwice, Poland*

Corresponding author:

Karolina Kowalczyk

Department of Gynecological Endocrinology, School of Medicine in Katowice, Medical University of Silesia, Katowice, Poland

e-mail: karolina.kowalczyk74@gmail.com

We present a case of a 67-year-old woman who was admitted to the department of gynecological oncology with a complaint of persistent and mild pain in the area of the left labia minora. The patient has been treated with antibiotics and anti-inflammatory drugs with no improvement. She also had a dermatological consultation; however, it did not reveal dermatological causes of the patient's complaints nor any other abnormal skin lesions. A suggestion of a persistent Bartholin's gland abscess has been made. Hence, she has been referred to the surgical removal of the lesion. The general condition of the patient was stable. She reported no additional gynecologic, gastrointestinal symptoms, rapid weight loss or any other complaints.

Physical examination revealed a palpable nodular lesion about 3 cm large in the area of the left labia minora. A mild bloody discharge in the vagina was also detected. Inguinal as well as other palpable lymph nodes were not enlarged. There were no significant abnormalities on the skin of the vulva or in the transvaginal ultrasound. The

abdominal ultrasound has shown no abnormalities. The patient was qualified for surgical removal of the lesion.

The oval, solid, smooth-surface tumor approximately 4 cm large has been removed during the surgery. The recovery period was uneventful, and the patient was discharged after two days.

A definitive histopathological diagnosis revealed neuroendocrine neoplasm (NEN) combined with squamous cell carcinoma (SCC). Immunohistochemically, cells of NEN expressed characteristic immunophenotype: synaptophysin(+), CK7(+), CK20(+) CD56(+) whereas the squamous part was p63 (+) (Fig. 1).

The microscopic morphology of NEN corresponded to Merkel cell carcinoma (MCC). Based on immunohistochemistry, primary MCC combined with SCC (MCC/SCC) of the vulva was signed. Positive surgical margins were detected. The patient has been referred to adjuvant radio-chemotherapy. Unfortunately, the patient did not return for follow-up.

NENs are rare tumors arising mainly in pancreas, lungs, and gastrointestinal tract [1]. They are rarely encountered in the gynecologic tract and constitute < 2% of female reproductive tumors. NEN of the vulva accounts for < 1 % of cases. Commonly, they are admixed with other histological types, most frequently SCC [2].

Currently, WHO has divided NENs to well-differentiated neuroendocrine tumours and poorly differentiated neuroendocrine carcinomas (NECs) [3]. NECs subdivide to small-cell NECs, large-cell NECs and carcinomas admixed with NEC. This WHO classification is uniformal for NENs at majority locations, including female genital tract. However, current division has not comprised MCC [3].

MCC is an aggressive, primary cutaneous neuroendocrine carcinoma, characterized by coexpression of neuroendocrine markers and CK20 (a discriminant from other types of visceral NENs) [5]. Most commonly MCC develops in elderly patients in the sun-exposed skin areas, where its occurrence is usually related to Merkel cell polyomavirus (MCV) [3, 5]. MCCs, with or without SCC component, are extremely rare in the gynecological area [1, 5].

Vulvar MCC coexistence with SCC might indicate the role of HPV in the pathogenesis of this neoplasm. The previous study determined both malignant components of the vulva incorporated the same high-risk HPV-DNA in their genome, without MCV genome. The lack of the TP53 mutation, which is related to ultraviolet light damage, was also detected. These findings support the suspicion that a vulvar MCC is HPV-related, whereas in the other origin it is more akin to be MCV-related [5].

The diagnostic process requires the assays for a possible primary lesion, as well as secondary metastases. In the presented case, based on dermatological consultation, imaging and clinical examinations, no primary lesion in the other location has been diagnosed.

There are no specific guidelines regarding treatment. In clinical practices surgical excision plays a pivotal role in early-stage cases followed by adjuvant radiation or chemoradiation. Some patients might benefit from immunotherapy as well [1, 2].

We want to draw attention to the possibility of MCC (with or without SCC component) of the vulva, the importance of proper approach to the differential diagnosis and further follow-up of the patient in order to diagnose secondary metastasis. Moreover, we want to highlight the value of prophylactic vaccination against oncogenic HPV, not only to prevent anogenital SCC, but also HPV-associated vulvar MCC.

Ethics statement

Written informed consent for publication was obtained from the patient.

Author contributions

The authors confirm sole responsibility for the following: Author 1: Marlena Cwynar: study conception and design; Author 2: Ewa Chmielik: analysis and interpretation of results; Author 3: Grzegorz Cwynar: data collection; Author 4: Piotr Ptak: data collection; Author 5: Karolina Kowalczyk: manuscript preparation.

Acknowledgments

None.

Conflict of interest

There are no conflicts of interest.

References

1. Georgescu TA, Bohiltea RE, Munteanu O, et al. Emerging Therapeutic Concepts and Latest Diagnostic Advancements Regarding Neuroendocrine Tumors of the Gynecologic Tract. *Medicina (Kaunas)*. 2021; 57(12), doi: [10.3390/medicina57121338](https://doi.org/10.3390/medicina57121338), indexed in Pubmed: [34946283](https://pubmed.ncbi.nlm.nih.gov/34946283/).
2. Virarkar M, Vulasala SS, Morani AC, et al. Neuroendocrine Neoplasms of the Gynecologic Tract. *Cancers (Basel)*. 2022; 14(7), doi: [10.3390/cancers14071835](https://doi.org/10.3390/cancers14071835), indexed in Pubmed: [35406607](https://pubmed.ncbi.nlm.nih.gov/35406607/).
3. Neuroendocrine neoplasia in WHO Classification of Tumours Editorial Board. Female genital tumours. International Agency for Research on Cancer, Lyon (France) 2020.
4. Heath M, Jaimes N, Lemos B, et al. Clinical characteristics of Merkel cell carcinoma at diagnosis in 195 patients: the AEIOU features. *J Am Acad Dermatol*. 2008; 58(3): 375–381, doi: [10.1016/j.jaad.2007.11.020](https://doi.org/10.1016/j.jaad.2007.11.020), indexed in Pubmed: [18280333](https://pubmed.ncbi.nlm.nih.gov/18280333/).
5. Lingamaneni P, Vohra I, Upadhyay S, et al. 1402 Metastatic Merkel Cell Carcinoma With Isolated Pancreatic Metastases. *American Journal of Gastroenterology*. 2019; 114(1): S777–S777, doi: [10.14309/01.ajg.0000595136.60337.25](https://doi.org/10.14309/01.ajg.0000595136.60337.25).

Figure 1. Vulvar Merkel cell carcinoma combined with squamous cell carcinoma. **A.** Tumor histopathology with neural invasion, hematoxylin & eosin stain, 29.56x; **B.** Neuroendocrine differentiation — diffuse synatophysin expression, immunohistochemistry stain (IS), 30,24x; **C.** Cytokeratin 20, IS, 45,79x; **D.** p63 expression in a focus of squamous cell carcinoma, IS, 26,33x

