

9-12-2023

Randomized Controlled Trial of Mechanical Thrombectomy vs Catheter-Directed Thrombolysis for Acute Hemodynamically Stable Pulmonary Embolism: Rationale and Design of the PEERLESS Study

Carin F. Gonsalves

C Michael Gibson

Stefan Stortecky

Roger A Alvarez

Daren M Beam

See next page for additional authors

Follow this and additional works at: <https://jdc.jefferson.edu/radiologyfp>



Part of the [Radiology Commons](#)

[Let us know how access to this document benefits you](#)

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Department of Radiology Faculty Papers by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Carin F. Gonsalves, C Michael Gibson, Stefan Stortecky, Roger A Alvarez, Daren M Beam, James M Horowitz, Mitchell J Silver, Catalin Toma, John H Rundback, Stuart P Rosenberg, Craig D Markovitz, Thomas Tu, and Wissam A Jaber



Randomized controlled trial of mechanical thrombectomy vs catheter-directed thrombolysis for acute hemodynamically stable pulmonary embolism: Rationale and design of the PEERLESS study

Carin F. Gonsalves, MD, FSIR^a, C. Michael Gibson, MD^b, Stefan Stortecky, MD, MPH, FESC^c, Roger A. Alvarez, DO, MPH^d, Daren M. Beam, MD^e, James M. Horowitz, MD^f, Mitchell J. Silver, DO, FACC, FSVM, RPVI^g, Catalin Toma, MD^h, John H. Rundback, MD, FAHA, FSVM, FSIRⁱ, Stuart P. Rosenberg, MS^j, Craig D. Markovitz, PhD^j, Thomas Tu, MD^j, and Wissam A. Jaber, MD^k *Philadelphia, PA; Boston, MA; Weston, FL; Indianapolis, IN; New York, NY; Columbus, OH; Pittsburgh, PA; Teaneck, NJ; Irvine, CA; Atlanta, GA*

Abstract

Background The identification of hemodynamically stable pulmonary embolism (PE) patients who may benefit from advanced treatment beyond anticoagulation is unclear. However, when intervention is deemed necessary by the PE patient's care team, data to select the most advantageous interventional treatment option are lacking. Limiting factors include major bleeding risks with systemic and locally delivered thrombolytics and the overall lack of randomized controlled trial (RCT) data for interventional treatment strategies. Considering the expansion of the pulmonary embolism response team (PERT) model, corresponding rise in interventional treatment, and number of thrombolytic and nonthrombolytic catheter-directed devices coming to market, robust evidence is needed to identify the safest and most effective interventional option for patients.

Methods The PEERLESS study (ClinicalTrials.gov identifier: NCT05111613) is a currently enrolling multinational RCT comparing large-bore mechanical thrombectomy (MT) with the FlowTrier System (Inari Medical, Irvine, CA) vs catheter-directed thrombolysis (CDT). A total of 550 hemodynamically stable PE patients with right ventricular (RV) dysfunction and additional clinical risk factors will undergo 1:1 randomization. Up to 150 additional patients with absolute thrombolytic contraindications may be enrolled into a nonrandomized MT cohort for separate analysis. The primary end point will be assessed at hospital discharge or 7 days post procedure, whichever is sooner, and is a composite of the following clinical outcomes constructed as a hierarchical win ratio: (1) all-cause mortality, (2) intracranial hemorrhage, (3) major bleeding, (4) clinical deterioration and/or escalation to bailout, and (5) intensive care unit admission and length of stay. The first 4 components of the win ratio will be adjudicated by a Clinical Events Committee, and all components will be assessed individually as secondary end points. Other key secondary end points include all-cause mortality and readmission within 30 days of procedure and device- and drug-related serious adverse events through the 30-day visit.

Implications PEERLESS is the first RCT to compare 2 different interventional treatment strategies for hemodynamically stable PE and results will inform strategy selection after the physician or PERT determines advanced therapy is warranted. (Am Heart J 2023;266:128–137.)

Background

Pulmonary embolism (PE) is the third leading cause of cardiovascular death. Over the last 2 decades, the steady rise in incidence and stagnant 10% overall 30-day mortality rate underscore the unmet need for more effective PE treatment with advanced therapy.¹⁻³ However, clinical

From the ^aThomas Jefferson University Hospital, Philadelphia, PA, ^bBeth Israel Deaconess Medical Center, Boston, MA, ^cDepartment of Cardiology, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland, ^dCleveland Clinic Florida, Weston, FL, ^eIndiana University Health University Hospital, Indianapolis, IN, ^fNYU Langone Health, New York, NY, ^gOhioHealth Heart and Vascular Physicians, Columbus, OH, ^hUniversity of Pittsburgh Medical Center, Pittsburgh, PA, ⁱAdvanced Interventional & Vascular Services, LLP, Teaneck, NJ, ^jInari Medical, Irvine, CA, ^kEmory University Hospital, Atlanta, GA

Submitted February 23, 2023; accepted September 6, 2023

Reprint requests: Stefan Stortecky, MD, MPH, FESC, Associate Professor of Medicine, Department of Cardiology, Bern University Hospital, Inselspital, University of Bern, 3010 Bern, Switzerland.

E-mail address: stortecky@gmx.at.

0002-8703

© 2023 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

<https://doi.org/10.1016/j.ahj.2023.09.002>

evidence generation for advanced therapies has lagged behind the pace of innovation. There is consensus that high-risk PE, which is defined by hemodynamic instability, necessitates rapid reperfusion treatment due to an early mortality rate exceeding 30%.⁴⁻⁶ Intermediate-risk PE however, is characterized by apparent hemodynamic stability, often in the presence of right ventricle (RV) dysfunction, and is associated with an early all-cause mortality rate that ranges broadly from 3–15%.^{7,8} In-hospital PE-related mortality ranges from 2% to 6%.^{6,9,10} Although patients present without overt hemodynamic instability, intermediate-risk PE encompasses a heterogeneous spectrum of disease severity, and normotensive cardiogenic shock has been identified in 30–40% of patients upon invasive measurement.^{11,12} Current guidelines recommend anticoagulation (AC) therapy as front-line treatment for intermediate-risk PE patients and if deterioration occurs, reperfusion treatment is recommended.^{5,13}

In the absence of hemodynamic deterioration, the decision to use reperfusion treatment for intermediate-risk PE patients is more ambiguous and controversial in clinical practice. The PEITHO (Pulmonary Embolism Thrombolysis) study showed early reperfusion with systemic thrombolysis reduced the likelihood of early death or hemodynamic decompensation compared with AC alone.¹⁴ In PEITHO, the treatment benefit was primarily driven by prevention of hemodynamic decompensation, while a meta-analysis showed an additional association with decreased likelihood of PE-related mortality.¹⁵ These benefits, however, are tempered by the relatively high frequency of major bleeding events with thrombolytics.^{5,14,15} Difficulties in clearly identifying hemodynamically stable PE patients who could benefit from early advanced therapy, coupled with demonstrated challenges in identifying those at higher risk of bleeding with thrombolytics, has prohibited front-line use of systemic thrombolysis in these patients.^{13,16,17}

The recently published European Society of Cardiology (ESC) clinical consensus statement states reperfusion treatment can be considered when there is no improvement in vital signs after 24 to 48 hours of therapeutic AC or when there are signs of worsening which are expected to precede deterioration.¹³ Precisely when to intervene based on treatment failure or early warning signs remains a primary research question. Without clear data, in clinical practice, a multidisciplinary Pulmonary Embolism Response Team (PERT) would convene to make these decisions. PERT activations commonly occur in hemodynamically stable PE cases with additional risk indicators and are associated with higher utilization of catheter-directed interventions.^{8,18}

Percutaneous catheter-directed interventions, including mechanical thrombectomy (MT) and catheter-directed thrombolysis (CDT), remain an alternative to reperfusion with systemic thrombolysis.⁵ Catheter-directed interventions have been studied in hemodynamically

stable PE patients, with the aim of providing clinical benefits similar to those observed with systemic thrombolytics while reducing the risk of major bleeding.^{6,11,19,20} These interventional therapies have been shown in single-arm studies to rapidly relieve RV strain, with low reported rates of acute clinical deterioration, hemodynamic worsening, or death.²¹⁻²⁵ As a result of these observations, there has been rapid adoption of interventional treatment in clinical practice.^{26,27}

Study rationale

Although the uptake of catheter-based treatment has been accompanied by observational clinical evidence, there is an overall lack of randomized controlled trial (RCT) data for interventional strategies. This is particularly true for trials with clinical outcome-based measures of treatment effectiveness. While foundational studies of both MT and CDT have separately reported positive outcomes in patients with intermediate-risk PE, there are no prospective studies directly comparing these different strategies in randomized trials.²⁸⁻³⁰ As a result, it is currently unknown whether different interventional treatment strategies provide similar clinical benefits to patients. As multiple new thrombolytic and nonthrombolytic catheter-directed devices come to market, it is important to understand if the method of thrombus removal impacts outcomes following treatment. Large-bore MT may provide added clinical benefit due to the immediacy of thrombus extraction and RV strain relief,^{31,32} but it has also been suggested that the ability of CDT to address thrombus in the distal pulmonary vasculature could be linked to improved outcomes.³³ As clinical outcome-based data become available for devices with varying mechanisms of action, the interpretation and generalizability of results could be complicated by study differences and the lack of understanding regarding comparability of interventional strategies. Therefore, it is necessary to create a RCT framework to evaluate the safety and effectiveness of currently available interventional strategies.²⁹


The PEERLESS Study (ClinicalTrials.gov Identifier: NCT05111613) is a currently enrolling prospective, multicenter, international RCT with the primary objective of comparing the clinical outcomes of patients following large-bore MT vs CDT for the treatment of acute hemodynamically stable PE, with the goal of informing treatment decisions for patients who the physician or PERT feels are likely to benefit from catheter-based interventional management.

Methods

Patient population

The PEERLESS study will enroll 550 patients who will be randomized 1:1 to receive treatment with large-bore MT, using the FlowTrier System (Inari Medical, Irvine,

Figure 1



Enrollment flowchart and follow-up assessments. CDT, catheter-directed thrombolysis; MT, mechanical thrombectomy; sPAP, systolic pulmonary artery pressure.

CA), or CDT, using any commercially available system, at up to 60 study sites located in the US and Europe. The study enrollment flowchart is shown in Figure 1. A separate cohort of up to 150 patients who meet eligibility criteria but have absolute contraindications to thrombolytic agents (Supplementary Table 1) may be enrolled into a nonrandomized MT cohort for independent analysis if their planned primary treatment includes the FlowTrier System.

Patients ≥ 18 years of age with hemodynamically stable PE and evidence of proximal filling defect in at least 1 main or lobar pulmonary artery (PA) on echocardiogram or imaging (computed tomographic pulmonary angiogram [CTPA] or pulmonary angiogram) will be included. Additionally, symptom onset must have occurred within 14 days of confirmed PE diagnosis, and intervention must be planned to begin within 72 hours following diagnosis or arrival at the treating hospital if transferring from another hospital, whichever was later. Patients will be excluded if they are unable to receive AC with heparin, enoxaparin, or another parenteral antithrombin, or if they have a right heart clot in transit, life expectancy < 30 days, or current diagnosis or documented history of chronic thromboembolic pulmonary hypertension (CTEPH) or chronic thromboembolic disease (CTED). A complete list of study inclusion and exclusion criteria can be found in Table 1.

Randomization will be stratified by bleeding risk as measured by VTE-BLEED score ≥ 2 or < 2 (Table 2), which will occur automatically in the Electronic Data Capture system upon data entry. Following an acute VTE (venous thromboembolism) event, a VTE-BLEED score ≥ 2 or < 2 identifies patients with high or low risk of bleeding, respectively, during stable AC.³⁴ The VTE-BLEED score was not designed to predict early bleeding due to interventional treatment for VTE and therefore, stratification should not be interpreted as predictive of outcomes. The

VTE-BLEED score was selected because it is an extensively validated disease-specific score with population-based distribution data.^{35,36} The proportion of the randomized population with an elevated VTE-BLEED score will be tracked and reported. All patients who give informed consent and fulfill baseline inclusion/exclusion criteria will be randomized. The point of enrollment is when the patient meets all eligibility criteria and the primary therapeutic catheter enters the body. As shown in Table 1, patients will be excluded postrandomization if their systolic pulmonary artery pressure (sPAP) is ≥ 70 mmHg on invasive measurement at the beginning of the index procedure, prior to enrollment and insertion of the primary therapeutic catheter. A sPAP ≥ 70 mmHg suggests chronic or acute-on-chronic thrombus, and patients with this degree of pre-existing elevation in sPAP are fundamentally different from the acute PE patient population intended to be studied in PEERLESS. Patients affected by this prespecified postrandomization exclusion criterion will not be included in the analysis population. However, screen failures or postrandomization exclusions will be tracked along with the reason for exclusion.

In the original study protocol, enrolled patients were required to meet the classification of intermediate-high-risk PE determined per the 2019 ESC Guidelines⁵ fulfilling all of the following: (1) clinical signs and symptoms consistent with acute PE or risk stratification of Pulmonary Embolism Severity Index (PESI) class III to V or simplified PESI (sPESI) score ≥ 1 ; (2) hemodynamic stability; (3) RV dysfunction on echocardiogram or CTPA; and (4) elevated cardiac troponin levels. After the enrollment phase had started, concerns were raised by Investigators regarding the inability to enroll patients selected for intervention during the normal course of practice-based treatment due to the criteria requiring elevated troponin levels. Reported concerns noted the observed transient

Table 1. Full inclusion and exclusion eligibility criteria

Inclusion

1. Age ≥ 18 years
2. Echocardiographic, CTPA, or pulmonary angiographic evidence of any proximal filling defect in at least 1 main or lobar pulmonary artery
3. ALL of the following characteristics:
 - a. Clinical signs and symptoms consistent with acute PE or PESI class III-V or sPESI score ≥ 1 , AND
 - b. Hemodynamically stable, AND
 - c. RV dysfunction on echocardiography or CT, AND
 - d. Any 1 or more of the following at the time of diagnosis:
 - Elevated cardiac troponin levels
 - History of heart failure
 - History of chronic lung disease
 - Heart rate ≥ 110 beats per minute
 - SBP < 100 mmHg
 - Respiratory rate ≥ 30 breaths per minute
 - Oxygen saturation $< 90\%$
 - Syncope related to PE
 - Elevated lactate
4. Intervention planned to begin within 72 hours of the later of either:
 - a. Confirmed PE diagnosis, OR
 - b. If transferring from another hospital, arrival at the treating hospital
5. Symptom onset within 14 days of confirmed PE diagnosis

Exclusion

1. Unable to receive AC with heparin, enoxaparin, or other parenteral antithrombin
2. Index presentations with hemodynamic instability that are part of the high-risk PE definition in the ESC 2019 Guidelines, including ANY of the following:
 - a. Cardiac arrest, OR
 - b. SBP < 90 mmHg or vasopressors required to achieve a SBP ≥ 90 mmHg despite adequate filling status, AND end-organ hypoperfusion, OR
 - c. SBP < 90 mmHg or SBP drop ≥ 40 mmHg, lasting longer than 15 minutes and not caused by new-onset arrhythmia, hypovolemia, or sepsis
3. Known sensitivity to radiographic contrast agents that, as determined by the Investigator, cannot be adequately pretreated
4. Imaging evidence or other evidence that suggests, in the opinion of the Investigator, the patient is not appropriate for catheter-based intervention
5. Patient has right heart clot in transit identified at baseline screening
6. Life expectancy < 30 days as determined by the Investigator
7. Current participation in another drug or device study that, in the opinion of the Investigator, would interfere with participation in this study
8. Current or history of CTEPH or CTED diagnosis, per ESC 2019 Guidelines
9. Invasive sPAP ≥ 70 mmHg prior to the primary therapeutic catheter entering the body
10. Administration of bolus or drip/infusion thrombolytic therapy or mechanical thrombectomy for the index PE event within 48 hours prior to enrollment
11. Ventricular arrhythmias refractory to treatment at the time of enrollment
12. Known to have heparin-induced thrombocytopenia
13. Patient has any condition for which, in the opinion of the Investigator, participation would not be in the best interest of the patient. This includes a contraindication to use of the FlowTrier System or CDT System per local approved labeling
14. Patient has previously completed or withdrawn from this study
15. Patient unwilling or unable to conduct the follow up visits per protocol

AC, anticoagulation; CDT, catheter-directed thrombolysis; CT, computed tomography; CTED, chronic thromboembolic disease; CTEPH, chronic thromboembolic pulmonary hypertension; CTPA, computed tomographic pulmonary angiography; ESC, European Society of Cardiology; PE, pulmonary embolism; PESI, pulmonary embolism severity index; sPESI, simplified PESI; RV, right ventricle; SBP, systolic blood pressure; sPAP, systolic pulmonary artery pressure.

pattern of troponin increases. Consequently, the PEERLESS Steering Committee expanded the inclusion criteria to include elevated cardiac troponin levels as 1 component of a broader clinical risk factor profile identified at the time of diagnosis, as shown in [Table 1](#).

Catheter-directed thrombolysis in PEERLESS

Patients randomized to the CDT arm in PEERLESS may be treated with any commercially available CDT system cleared for use per local regulations at the discretion of the treating physician; this includes con-

Table 2. VTE-BLEED algorithm

Parameter	Points
Active cancer	2
Male patient with uncontrolled hypertension (≥ 140 mmHg)	1
Anemia	1.5
History of bleeding	1.5
Renal dysfunction (creatinine clearance 30-60 mL/min)	1.5
Age ≥ 60 y	1.5
Total	9 points

ventional (eg, Cragg-McNamara Micro Therapeutics Infusion Catheter, Medtronic, Dublin, Ireland; and Uni-Fuse Infusion Catheters, AngioDynamics, Latham, NY) and ultrasound-assisted CDT systems (EKOS Endovascular System, Boston Scientific, Marlborough, MA). Treatment with CDT in the PEERLESS study is supported by evidence from several clinical studies of both conventional and ultrasound-assisted CDT systems that included hemodynamically stable PE patients. Two studies of ultrasound-assisted CDT without comparator treatment arms reported a statistically significant decrease in RV/left ventricle (LV) ratio at 48 hours after initiation of procedure; major bleeding events occurred in 10% and 4% of patients and 30-day mortality rates were 2.7% and 1%, respectively.^{21,22} The SUNSET sPE RCT, assessed the additive benefit of ultrasound-assisted CDT over conventional CDT and showed no difference in PA thrombus reduction between the 2 therapies.³⁷ Additionally, 2 small RCTs, 1 with ultrasound-assisted CDT and 1 with conventional CDT, suggested surrogate outcomes are improved with CDT vs AC alone in patients with hemodynamically stable PE. The ULTIMA (Ultrasound Accelerated Thrombolysis of Pulmonary Embolism) study (N=59) reported greater mean reduction in RV/LV ratio at 24 hours post procedure with ultrasound-assisted CDT plus AC vs AC alone for intermediate-risk PE (0.30 ± 0.20 vs 0.03 ± 0.16 , respectively; $P < .001$). In ULTIMA, there were no major bleeding events and only 1 patient (AC arm) death (1.7%) occurred through 90 days.¹⁹ The more recently published CANARY (Catheter-Directed Thrombolysis vs Anticoagulation in Patients with Acute Intermediate-High-Risk Pulmonary Embolism) study (N = 94) of conventional CDT plus AC vs AC alone for intermediate-high-risk PE suggested improvement of RV/LV ratio among patients treated with CDT. However, this study was stopped early due to the COVID-19 pandemic and, therefore, statistical significance of the primary end point was not reached. In CANARY, 1 (2.1%) nonfatal major bleeding event occurred in the CDT arm.²⁰

Mechanical thrombectomy in PEERLESS

In contrast to the various CDT systems permitted in the PEERLESS study, patients randomized to the MT arm will be treated with the FlowTrieve System. The FlowTrieve

System is a MT device that extracts PE thrombus by large-bore syringe-based aspiration (16-24F catheter) and/or mechanical (nitinol mesh disks) modes, without employing thrombolytics. There are other smaller-bore MT devices intended to be used without a thrombolytic agent for the treatment of PE. However, given the paucity of data for their different modes of action, particularly for device safety, they are not included in this study.

Evidence supporting study treatment with the FlowTrieve System includes results from 2 previously conducted single-arm sponsored studies, FLARE and FLASH,^{23-25,38} in addition to independently conducted studies.³⁹⁻⁴¹ FLARE (FlowTrieve Pulmonary Embolectomy Clinical Study) was a prospective and multicenter investigational device exemption trial, which led to US FDA clearance of the FlowTrieve System for PE treatment in 2018.²⁵ Results were confirmed by the large, prospective, and multicenter FLASH (FlowTrieve All-Comer Registry for Patient Safety and Hemodynamics) registry in patients with intermediate- or high-risk PE. Available outcomes through the 30-day visit were recently published for the full US cohort (n = 800).²⁴ The primary end point, a major adverse event composite of device-related mortality and major bleeding within 48 hours and intraprocedural device- or procedure-related adverse events, was observed in 1.8% of patients. There were no intracranial hemorrhages or device-related deaths, and all-cause mortality was 0.8% at the 30-day visit. Immediately following the procedure, mean pulmonary artery pressure decreased by 7.6 mmHg on average. Echocardiographic assessments at the 48-hour visit showed an improvement in the baseline mean RV/LV ratio from 1.23 to 0.98, in addition to improvements in RV systolic pressure and RV function.²⁴

Interventional procedure

In this study, interventional treatment with either MT or CDT will be conducted per standard local practice and in a manner consistent with the device manufacturer's Instructions for Use. There is no specific guidance on when to terminate the procedure. The goal of PEERLESS is to compare treatment outcomes of patients treated in the standard course of clinical practice. Invasive PA pressure measurement is required prior to insertion of the primary therapeutic catheter. For patients treated with CDT, PA pressure measurement is requested, but not required per protocol, at both ≥ 10 minutes after beginning infusion through the primary therapeutic catheter and 6 ± 2 hours post procedure. For patients treated with MT, PA pressure measurement is requested, but not required per protocol, after the FlowTrieve System is removed for the last time. Estimated procedural blood loss is recorded.

The use of pre-, intra-, and postprocedural AC is guided by investigator discretion, local standard of care, and in accordance with the Instructions for Use for the assigned therapy. The dose and duration of thrombolytic agents

used for patients assigned to CDT is guided in the same manner, and no standardized thrombolytic dosing protocol is enforced. A partial thromboplastin time of 40–60 seconds is suggested to avoid bleeding during CDT administration.²¹ The rationale for the decision to not require a standardized thrombolytic dosing protocol in patients randomized to CDT is multifaceted: (1) the ultimate goal of PEERLESS is to provide data reflective of current practice patterns, which include variable thrombolytic dosing regimens during CDT; (2) as described in ESC treatment guidelines,^{5,13} the optimal thrombolytic dosing strategy for CDT is not settled and therefore institutions generally establish their own protocols; and (3) excluding sites unwilling to participate based on enforcement of a strictly controlled thrombolytic regimen could introduce bias. It is important to note, however, that the initial thrombolytic dosing strategy will be recorded, along with any dose adjustments throughout infusion, including the extension of infusion duration. Therefore, this information will be available for analysis when data become available.

Follow-up and outcomes

Follow-up assessments

Patients will have postprocedure follow-up evaluations at 24 hours (± 8 hours), hospital discharge, and 30 days ($+15$ days). The full schedule of assessments is shown in Supplementary Table 2.

Primary end point

The primary study end point is a composite end point constructed as a win ratio,⁴² which is a hierarchy of the following clinical outcomes assessed at hospital discharge or 7 days post procedure, whichever is sooner: (1) all-cause mortality, (2) intracranial hemorrhage, (3) major bleeding per the International Society on Thrombosis and Haemostasis (ISTH) definition,⁴³ (4) clinical deterioration and/or escalation to a bailout therapy, and (5) intensive care unit (ICU) admission and ICU length of stay (LOS) during the index hospitalization and following the index procedure. The first 4 components of the win ratio will be adjudicated by a Clinical Events Committee. The definitions for events included in the primary end point can be found in Table 3.

The win ratio defined by Pocock et al⁴² will be used to evaluate the primary end point and summarize the treatment difference between the randomized MT and CDT study arms. Each randomized patient treated with MT will be compared to each patient treated with CDT to determine a “winner.” Within each pair, the 5 end point components of the win ratio are compared sequentially, in the order of clinical outcome priority, until a “winner” is identified. Pairwise MT vs CDT patient comparisons without a “winner” (ie, “tie”) are not included in the win ratio calculation (eg, neither patient experiences 1 of the 5 outcomes). After all patient pairs have been compared,

the win ratio is calculated by dividing the total number of MT winners by the total number of CDT winners in the study.

In PEERLESS, an unmatched win ratio approach is utilized, in that each randomized MT patient will be compared to each CDT patient. To our best knowledge, this is the first PE RCT to employ a win ratio approach. Experience based on other cardiovascular win ratio RCTs indicates an unmatched approach is favored if a validated patient matching process is challenging to predefine,^{42,44} as is the case in hemodynamically stable PE patients. An unmatched win ratio approach was used in the COAPT trial (heart failure and functional mitral regurgitation) and the PARTNER B trial (severe symptomatic aortic stenosis), among others.^{42,44} Treatment effect determined by win ratio analysis is considered transparent and high quality.⁴² The primary advantage of using a win ratio is the prioritization of the most crucial end points. In contrast to conventional composite end points, all components are not treated equally in a win ratio composite end point design; instead, the components are prioritized by clinical importance.

Secondary end points

Another composite hierarchic win ratio consisting of only the first 4 components of the primary end point will be evaluated as a secondary end point to demonstrate comparative treatment effect without considering ICU admission or ICU LOS. Other secondary end points include the individual assessment of each of the 5 components of the primary end point, in addition to clinically relevant nonmajor (CRNM) and minor bleeding events at hospital discharge or 7 days post procedure if sooner. CRNM bleeding is defined as any sign or symptom of hemorrhage that does not fit the criteria for the ISTH definition of major bleeding but does meet at least one of the following criteria: (1) requiring medical intervention by a healthcare professional, (2) leading to hospitalization or increased level of care, or (3) prompting a face-to-face evaluation. Minor bleeding is defined as any bleeding not classified as major or CRNM bleeding.

Additionally, the change in RV/LV ratio and modified Medical Research Council (mMRC) Dyspnea scores from baseline to 24-hour visit and 30-day visit will be evaluated. The mMRC Dyspnea Scale was used in the FLASH registry and is a patient-reported assessment tool with scores from 0 to 4, with higher scores representing worse dyspnea.^{24,45} All-cause mortality and all-cause and PE-related readmission will be assessed within 30 days after index procedure. Device- and drug-related serious adverse events will be captured through the 30-day visit. The total hospital and post-index-procedure LOS will be captured through a maximum of 30 days. Other assessments at the 30-day visit include disease-specific and general health-related quality of life, assessed by PEmb-QoL and EQ-5D-5L, respectively.

Table 3. Hierarchical win ratio primary end point

Primary end point	Hierarchical clinical outcome events and definitions
Win Ratio	<ol style="list-style-type: none"> 1. All-cause mortality 2. Intracranial hemorrhage: <ul style="list-style-type: none"> • Any bleeding involving the brain parenchyma, ventricular system, or subarachnoid, subdural, or epidural regions as identified by CT scan or MRI regardless of symptoms 3. Major bleeding according to the ISTH definition in nonsurgical subjects: <ul style="list-style-type: none"> • Fatal bleeding, AND/OR • Symptomatic bleeding in a critical area or organ, such as intracranial, intraspinal, intraocular, retroperitoneal, intraarticular or pericardial, or intramuscular with compartment syndrome, AND/OR • Bleeding causing a fall in hemoglobin level of ≥ 2 g/dL (1.24 mmol/L) or leading to transfusion of ≥ 2 units of whole blood or red cells 4. A. Clinical deterioration AND/OR B. escalation to a bailout therapy <p>4A. Clinical deterioration: defined by documented objective hemodynamic or respiratory worsening that is not present at the time of enrollment, including 1 or more of the following:</p> <ul style="list-style-type: none"> • Hypotension with SBP < 90 mmHg lasting at least 30 minutes, unresponsive to fluid resuscitation, and requiring the addition of, or increased dose of, vasopressors • Fall in SBP by ≥ 40 mmHg lasting at least 30 minutes and accompanied end-organ hypoperfusion • Cardiac arrest requiring cardiopulmonary resuscitation • Bradycardia lasting longer than 10 minutes, accompanied by hypotension and requiring pharmacologic intervention or insertion of a pacemaker • Ventricular tachycardia or fibrillation requiring pharmacologic intervention or defibrillation • Requirement for an increase in fraction of inspired oxygen requirements ≥ 0.20, lasting longer than 30 minutes • Need for intubation in a previously nonintubated patient, or unplanned use of ECMO <p>4B. Escalation to a bailout therapy: defined by surgical thrombectomy; unplanned use of additional mechanical, pharmacomechanical, or pharmacologic catheter-based therapies, or systemic thrombolytics; or changing from the assigned treatment strategy after initial treatment strategy was assigned:</p> <ul style="list-style-type: none"> • If CDT was assigned and emergent or clinically driven systemic thrombolytic administration was required after CDT was initiated, this would be considered a bailout. If length of administration is simply extended and is not emergent or clinically driven, this would not qualify as a bailout • If MT was assigned, low-dose catheter-directed adjunctive thrombolytic therapy (< 10 mg tPA) that is administered intraprocedurally or postprocedurally will be strongly discouraged but not considered a bailout <ol style="list-style-type: none"> 5. ICU admission and ICU LOS during the index hospitalization and following the index procedure: <ul style="list-style-type: none"> • ICU admission and ICU LOS are characterized hierarchically as follows: 1) no ICU admission, 2) ICU admission lasting between 0–24 hours, 3) ICU admission lasting > 24 hours

CDT, catheter-directed thrombolysis; CT, computed tomography; ECMO, extracorporeal membrane oxygenation; ICU, intensive care unit; ISTH, International Society on Thrombosis and Haemostasis; LOS, length of stay; MRI, magnetic resonance imaging; MT, mechanical thrombectomy; SBP, systolic blood pressure; tPA, tissue plasminogen activator.

Statistical analysis

Sample size calculation

Assuming 80% power with one-sided alpha of 2.5%, the win ratio methodology was applied to the primary end point, consisting of the 5 components described previously, to determine the required sample size. Under these assumptions, the required sample size was determined to be 432 patients, with 216 randomized to each arm. Considering follow-up attrition and the planned assessment of secondary end points, the study aims to enroll a total of 550 patients in the randomized cohort.

End point analyses

The hierarchic win ratio composite end point will be evaluated using a modified generalized Wilcoxon test (F-S test) proposed by Finkelstein and Schoenfeld⁴⁶ to exam-

ine the performance differences between the 2 randomized treatment arms. The 95% confidence interval (CI) of the win ratio estimate will be derived via bootstrap method.

Outcomes for each of the 5 individual components of the primary end point, incidence of CRNM and minor bleeding events at the sooner of discharge or 7 days post procedure, all-cause mortality and all-cause and PE-related readmission within 30 days of index procedure, device- and drug-related serious adverse events through 30-day visit, and mMRC dyspnea score at 24-hour and 30-day visits will be compared between treatment arms using *P* values derived using Fisher or Fisher exact tests as appropriate. The change in RV/LV ratio from baseline to 24-hour visit and overall PEMB-QoL and EQ-5D-5L scores

at the 30-day visit will be assessed with *P* values derived using the Wilcoxon rank sum test. *P* values $\leq .05$ will be considered statistically significant. Data from the nonrandomized cohort of patients with absolute contraindications to thrombolytics will not be used for any protocol-defined hypothesis testing nor used to calculate any primary or secondary end points.

Funding and ethical considerations

This study is being sponsored by Inari Medical. The initial draft of this manuscript was developed with writing assistance provided by the study sponsor. All authors are members of the PEERLESS Steering Committee and contributed significantly to the overall design and study protocol, which served as the basis for this publication.

Investigators are responsible for obtaining written or, if applicable, electronic Informed Consent for each study patient in accordance with pertinent regulations. This study is approved by an appropriate institutional review board or ethics committee at each site. Boston Clinical Research Institute is serving as the Clinical Events Committee in this study, which will be utilized for the purposes of adjudicating safety-related primary and secondary end points. Site-reported safety and outcome data will be provided to the committee for review and adjudicated for all patients enrolled in the study.

Discussion

Expected impact of PEERLESS

PEERLESS is the first RCT comparing 2 different interventional strategies for the treatment of hemodynamically stable PE. This study will address the comparability of clinical outcomes following MT with the FlowTrier System vs CDT conducted per local standard of care. The PEERLESS study design is intended to generate important data to support decision-making after the physician or PERT determines intervention is warranted. The adoption of interventional treatment in real-world clinical practice is increasing based on single-arm studies demonstrating reduced short- and long-term mortality and morbidity in intermediate-risk PE patients. With the recent and ongoing emergence of multiple catheter-based systems for interventional PE management, there is a fundamental and topical imperative to understand treatment risks and value between thrombolytic and nonthrombolytic strategies.

Remaining uncertainties

PEERLESS does not include an anticoagulation treatment arm, the guideline recommended treatment for hemodynamically stable PE, and this is a significant limitation of the study. This study will not determine if interventional treatment is superior to conservative treatment with AC for these patients and does not include a specified analysis to identify patients most likely to benefit

from interventional treatment. Other RCTs either in development or underway will study patients with hemodynamically stable PE to assess outcomes for intervention compared to conservative therapy alone, including HI-PEITHO, PE-TRACT, STORM-PE, and the forthcoming PEERLESS II study.^{47–49} PEERLESS is designed to provide data supporting treatment strategy selection after the decision to intervene is made. In the absence of hemodynamic worsening, the individualized decision to use catheter-directed intervention is complex and currently made without standardized algorithmic guidance.

The results from PEERLESS will provide insight on the comparability of cardinal clinical outcomes with MT vs CDT through 30 days post intervention. However, it is important to recognize that long-term outcomes following intervention are an increasingly important part of the discussion concerning interventional therapies and are not being addressed in this study. Additionally, the rate of long-term sequelae, such as CTEPH and CTED, is a key treatment outcome and will need to be investigated in follow-on studies.

Current enrollment status

As of April 30, 2023, 59 study sites have been activated with a total of 370 patients enrolled, 298 in the randomized arms and 72 in the nonrandomized MT cohort.

Disclosures

C.F. Gonsalves reports receiving consulting fees from Inari Medical. C.M. Gibson reports receiving research grant support from Angel Medical Corporation, Bayer Corp, CSL Behring, Janssen Pharmaceuticals, Johnson & Johnson Corporation, and Portola Pharmaceuticals; and has received consulting monies from Amarin Pharma, Amgen, Arena Pharmaceuticals, Bayer Corporation, Boehringer Ingelheim, Boston Clinical Research Institute, Cardiovascular Research Foundation, Chiesi, CSL Behring, Eli Lilly, Gilead Sciences, Inc, Inari Medical, Janssen Pharmaceuticals, Johnson & Johnson Corporation, The Medicines Company, Merck & Co, Inc, Novo Nordisk, Pfizer, Pharma Mar, Portola Pharmaceuticals, Sanofi, Somahlution, St Francis Hospital, Verson Corporation, and Web MD. S. Stortecky reports receiving research grants to the institution from Edwards Lifesciences, Medtronic, Abbott Vascular, and Boston Scientific; consulting fees from Boston Scientific and Teleflex, as well as honoraria for lectures from Boston Scientific. R.A. Alvarez reports receiving consulting fees from United Therapeutics Corporation. D.M. Beam reports receiving consulting fees from Inari Medical and Janssen Pharmaceuticals. J.M. Horowitz reports receiving consulting fees from Inari Medical and Penumbra. M.J. Silver reports receiving consulting fees from Medtronic, Boston Scientific, WL Gore and Associates, Cook Medical, Contego Medical, and Inari Medical, as well as speak-

ers' bureau for Bristol Myers Squibb, Astra Zeneca, and Pfizer. C. Toma reports receiving consulting fees from Medtronic, Neptune Medical and Philips. J.H. Rundback reports being a consultant or advisory board member for Inari Medical, Philips, AngioDynamics, Boston Scientific, Abbott, Medtronic, Truvic, and Aveera Medical; has stocks/options in Protexa, Aveera, and Kalgaro Medical; is a board member of Protexa Medical and the CLI Global Society; is the site or national PI and receives institutional research support from Abbott, 4S, Terumo, Reflow Medical and AngioDynamics. S.P. Rosenberg, C.D. Markovitz, and T. Tu are employees of Inari Medical and own stock and/or have stock options in the company. W.A. Jaber reports receiving consulting fees from Medtronic and Inari Medical.

Acknowledgments

Authors acknowledge receiving writing and editorial assistance from Kelly Koch, PharmD and Jessica Parsons, PhD.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:[10.1016/j.ahj.2023.09.002](https://doi.org/10.1016/j.ahj.2023.09.002).

References

- Barco S, Valerio L, Gallo A, et al. Global reporting of pulmonary embolism-related deaths in the World Health Organization mortality database: vital registration data from 123 countries. *Res Pract Thromb Haemost* 2021;5:e12520.
- Kucher N, Rossi E, De Rosa M, Goldhaber SZ. Massive pulmonary embolism. *Circulation* 2006;113:577–82.
- R Lookstein, PERT Consortium® Quality Database Update, Presented at the 2022 PERT Consortium® Annual Meeting, September 29 - October 1, 2022, Virtual Format.
- Chopard R, Behr J, Vidoni C, et al. An update on the management of acute high-risk pulmonary embolism. *J Clin Med* 2022;11:4807.
- Konstantinides SV, Meyer G, Becattini C, et al. 2019 ESC Guidelines for the diagnosis and management of acute pulmonary embolism developed in collaboration with the European Respiratory Society (ERS). *Eur Heart J* 2020;41:543–603.
- Secemsky E, Chang Y, Jain CC, et al. Contemporary management and outcomes of patients with massive and submassive pulmonary embolism. *Am J Med* 2018;131:1506–14 e0.
- Ismayl M, Machanahalli Balakrishna A, Aboeata A, et al. Meta-analysis comparing catheter-directed thrombolysis versus systemic anticoagulation alone for submassive pulmonary embolism. *Am J Cardiol* 2022;178:154–62.
- Schultz J, Giordano N, Zheng H, et al. EXPRESS: a multidisciplinary pulmonary embolism response team (PERT) - experience from a national multicenter consortium. *Pulm Circ* 2019;9:2045894018824563.
- Becattini C, Casazza F, Forgiione C, et al. Acute pulmonary embolism: external validation of an integrated risk stratification model. *Chest* 2013;144:1539–45.
- Vanni S, Nazerian P, Pepe G, et al. Comparison of two prognostic models for acute pulmonary embolism: clinical vs. right ventricular dysfunction-guided approach. *J Thromb Haemost* 2011;9:1916–23.
- Khandhar SJ, Mehta M, Cilia L, et al. Invasive hemodynamic assessment of patients with submassive pulmonary embolism. *Catheter Cardiovasc Interv* 2020;95:13–18.
- Bangalore S, Horowitz JM, Beam D, et al. Prevalence and predictors of cardiogenic shock in intermediate-risk pulmonary embolism: insights from the FLASH registry. *JACC Cardiovasc Interv* 2023;16:958–72.
- Pruszczyk P, Klok FA, Kucher N, et al. Percutaneous treatment options for acute pulmonary embolism: a clinical consensus statement by the ESC Working Group on Pulmonary Circulation and Right Ventricular Function and the European Association of Percutaneous Cardiovascular Interventions. *EuroIntervention* 2022;18:e623–38.
- Meyer G, Vicaut E, Danays T, et al. Fibrinolysis for patients with intermediate-risk pulmonary embolism. *N Engl J Med* 2014;370:1402–11.
- Marti C, John G, Konstantinides S, et al. Systemic thrombolytic therapy for acute pulmonary embolism: a systematic review and meta-analysis. *Eur Heart J* 2015;36:605–14.
- Chatterjee S, Weinberg I, Yeh RW, et al. Risk factors for intracranial haemorrhage in patients with pulmonary embolism treated with thrombolytic therapy development of the PE-CH score. *Thromb Haemost* 2017;117:246–51.
- Rali PM, Criner GJ. Submassive pulmonary embolism. *Am J Respir Crit Care Med* 2018;198:588–98.
- Fleitas Sosa D, Lehr AL, Zhao H, et al. Impact of pulmonary embolism response teams on acute pulmonary embolism: a systematic review and meta-analysis. *Eur Respir Rev* 2022;31:220023.
- Kucher N, Boekstegers P, Muller OJ, et al. Randomized, controlled trial of ultrasound-assisted catheter-directed thrombolysis for acute intermediate-risk pulmonary embolism. *Circulation* 2014;129:479–86.
- Sadeghipour P, Jenab Y, Moosavi J, et al. Catheter-directed thrombolysis vs anticoagulation in patients with acute intermediate-high-risk pulmonary embolism: the CANARY randomized clinical trial. *JAMA Cardiol* 2022;7:1189–97.
- Piazza G, Hohlfelder B, Jaff MR, et al. A prospective, single-arm, multicenter trial of ultrasound-facilitated, catheter-directed, low-dose fibrinolysis for acute massive and submassive pulmonary embolism: the SEATTLE II study. *JACC Cardiovasc Interv* 2015;8:1382–92.
- Tapson VF, Sterling K, Jones N, et al. A randomized trial of the optimum duration of acoustic pulse thrombolysis procedure in acute intermediate-risk pulmonary embolism: the OPTALYSE PE trial. *JACC Cardiovasc Interv* 2018;11:1401–10.
- Toma C, Bunte MC, Cho KH, et al. Percutaneous mechanical thrombectomy in a real-world pulmonary embolism population: interim results of the FLASH registry. *Catheter Cardiovasc Interv* 2022;99:1345–55.
- Toma C, Jaber WA, Weinberg MD, et al. Acute outcomes for the full US cohort of the FLASH mechanical thrombectomy registry in pulmonary embolism. *EuroIntervention* 2023;18:1201–12.
- Tu T, Toma C, Tapson VF, et al. A prospective, single-arm,

- multicenter trial of catheter-directed mechanical thrombectomy for intermediate-risk acute pulmonary embolism: the FLARE study. *JACC Cardiovasc Interv* 2019;12:859–69.
26. Gayou EL, Makary MS, Hughes DR, et al. Nationwide trends in use of catheter-directed therapy for treatment of pulmonary embolism in medicare beneficiaries from 2004 to 2016. *J Vasc Interv Radiol* 2019;30:801–6.
 27. Raghupathy S, Barigheid AP, Doorogen R, et al. Prevalence, trends, and outcomes of pulmonary embolism treated with mechanical and surgical thrombectomy from a nationwide inpatient sample. *Clin Pract* 2022;12:204–14.
 28. Feroze R, Arora S, Tashtish N, et al. Comparison of large-bore thrombectomy with catheter-directed thrombolysis for the treatment of pulmonary embolism. *J Soc Cardiovasc Angiogr Interv* 2023;2:100453.
 29. Giri J, Sista AK, Weinberg I, et al. Interventional therapies for acute pulmonary embolism: current status and principles for the development of novel evidence: a scientific statement from the American Heart Association. *Circulation* 2019;140:e774–801.
 30. Inci EK, Khandhar S, Toma C, et al. Mechanical thrombectomy versus catheter directed thrombolysis in patients with pulmonary embolism: a multicenter experience. *Catheter Cardiovasc Interv* 2023;101:140–6.
 31. Alonso Martinez JL, Annicchero Sanchez FJ, Urbieto Echezarreta MA, et al. Central versus peripheral pulmonary embolism: analysis of the impact on the physiological parameters and long-term survival. *N Am J Med Sci* 2016;8:134–42.
 32. Meinel FG, Nance Jr JW, Schoepf UJ, et al. Predictive value of computed tomography in acute pulmonary embolism: systematic review and meta-analysis. *Am J Med* 2015;128:747–59 e2.
 33. Singh M, Shafi I, Rali P, et al. Contemporary catheter-based treatment options for management of acute pulmonary embolism. *Curr Treat Options Cardiovasc Med* 2021;23:44.
 34. Klok FA, Hoesel V, Clemens A, et al. Prediction of bleeding events in patients with venous thromboembolism on stable anticoagulation treatment. *Eur Respir J* 2016;48:1369–76.
 35. Klok FA, Barco S, Konstantinides SV. External validation of the VTE-BLEED score for predicting major bleeding in stable anticoagulated patients with venous thromboembolism. *Thromb Haemost* 2017;117:1164–70.
 36. Nishimoto Y, Yamashita Y, Morimoto T, et al. Validation of the VTE-BLEED score's long-term performance for major bleeding in patients with venous thromboembolisms: From the COMMAND VTE registry. *J Thromb Haemost* 2020;18:624–32.
 37. Avgerinos ED, Jaber W, Lacomis J, et al. Randomized trial comparing standard versus ultrasound-assisted thrombolysis for submassive pulmonary embolism: the SUNSET sPE trial. *JACC Cardiovasc Interv* 2021;14:1364–73.
 38. Jaber WA, Kabrhel C, Rosenfield K, et al. Percutaneous thrombectomy in emergency department patients with pulmonary embolism: the FLARE ED sub-study. *J Emerg Med* 2020;58:175–82.
 39. Buckley JR, Wible BC. In-hospital mortality and related outcomes for elevated risk acute pulmonary embolism treated with mechanical thrombectomy versus routine care. *J Intensive Care Med* 2022;37:877–82.
 40. Toma C, Khandhar S, Zalewski AM, et al. Percutaneous thrombectomy in patients with massive and very high-risk submassive acute pulmonary embolism. *Catheter Cardiovasc Interv* 2020;96:1465–70.
 41. Wible BC, Buckley JR, Cho KH, et al. Safety and efficacy of acute pulmonary embolism treated via large-bore aspiration mechanical thrombectomy using the Inari FlowTrieve device. *J Vasc Interv Radiol* 2019;30:1370–5.
 42. Pocock SJ, Ariti CA, Collier TJ, Wang D. The win ratio: a new approach to the analysis of composite endpoints in clinical trials based on clinical priorities. *Eur Heart J* 2012;33:176–82.
 43. Schulman S, Kearon C Subcommittee on Control of Anticoagulation of the Scientific and Standardization Committee of the International Society on Thrombosis and Haemostasis. Definition of major bleeding in clinical investigations of antithrombotic medicinal products in non-surgical patients. *J Thromb Haemost* 2005;3:692–4.
 44. Redfors B, Gregson J, Crowley A, et al. The win ratio approach for composite endpoints: practical guidance based on previous experience. *Eur Heart J* 2020;41:4391–9.
 45. Mahler DA, Wells CK. Evaluation of clinical methods for rating dyspnea. *Chest* 1988;93:580–6.
 46. Finkelstein DM, Schoenfeld DA. Combining mortality and longitudinal measures in clinical trials. *Stat Med* 1999;18:1341–54.
 47. NYU Langone Health. Pulmonary embolism - thrombus removal with catheter-directed therapy (PE-TRACT). ClinicalTrials.gov identifier: NCT05591118. Updated October 24, 2022. [accessed 6 December 2022] <https://www.clinicaltrials.gov/ct2/show/NCT05591118>.
 48. Penumbra Inc. STORM-PE: a prospective, multicenter, randomized controlled trial evaluating anticoagulation alone vs anticoagulation plus mechanical aspiration with the Indigo® aspiration system for the treatment of intermediate high risk acute pulmonary embolism. ClinicalTrials.gov identifier: NCT05684796. Updated August 29, 2023. [accessed 31 August 2023] <https://classic.clinicaltrials.gov/ct2/show/NCT05684796>.
 49. Klok FA, Piazza G, Sharp ASP, et al. Ultrasound-facilitated, catheter-directed thrombolysis vs anticoagulation alone for acute intermediate-high-risk pulmonary embolism: Rationale and design of the HI-PEITHO study. *Am Heart J* 2022;251:43–53.