

University of Texas Rio Grande Valley

ScholarWorks @ UTRGV

Chemistry Faculty Publications and
Presentations

College of Sciences

6-2015

Internal dynamics in the molecular complex of CF₃CN and H₂O

Wei Lin

Anan Wu


Xin Lu

Xiao Tang

Daniel A. Obenchain

See next page for additional authors

Follow this and additional works at: https://scholarworks.utrgv.edu/chem_fac

 Part of the [Chemistry Commons](#)

Authors

Wei Lin, Anan Wu, Xin Lu, Xiao Tang, Daniel A. Obenchain, and Stewart E. Novick

PCCP

Accepted Manuscript



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

ARTICLE

Internal Dynamics in the Molecular Complex of CF_3CN and H_2O

Cite this: DOI: 10.1039/x0xx00000x

Wei Lin^{*a}, Anan Wu^{*b}, Xin Lu^b, Xiao Tang^b, Daniel A. Obenchain^c, and Stewart E. Novick^cReceived 00th March 2015,
Accepted 00th January 2015

DOI: 10.1039/x0xx00000x

www.rsc.org/

The rotational spectrum of trifluoroacetonitrile-water complex has been studied by pulsed-nozzle, Fourier transform microwave spectroscopy. Both *a*-type and *b*-type transitions have been observed. The rotational constants, centrifugal distortion constants, and the ^{14}N nuclear quadrupole coupling constants have been determined. The complex is T-shaped, with the oxygen atom from the water located 3.135 Å from the carbon atom of CF_3 of the trifluoroacetonitrile molecule.

Introduction

Trifluoroacetonitrile, CF_3CN , is a symmetric top molecule. The microwave spectrum, as well as millimeter-wave spectrum of the monomeric form CF_3CN have been extensively studied due to its large dipole moment and large thermal population in the excited vibrational states^{1,2,3,4,5,6,7,8,9,10,11,12,13}. We have reported the rotational analysis of the CF_3CN -Argon complex.¹⁴

The hydrogen bonding in small water containing clusters has received considerable attention. Water can serve as a proton donor, a proton acceptor, or in some cases as both. The types of hydrogen bonds involving water and its internal dynamics were summarized by Caminati *et al.*¹⁵ Microwave spectroscopy is a particularly suitable technique to study the structure and internal dynamics of water containing clusters. Several clusters formed by water and $\text{CF}_3\text{-X}$ ($\text{X} = \text{F}, \text{Cl}, \text{COOH}$) type molecules have been characterized. In the case of $\text{CF}_4\text{-H}_2\text{O}$,¹⁶ an anti-hydrogen bond which was first proposed by Novick *et al.*¹⁷ is formed due to the strong dipole-induced dipole interaction. Whereas Evangelisti *et al.*¹⁸ observed an O-Cl halogen bond in $\text{CF}_3\text{Cl-H}_2\text{O}$ that binds the water to CF_3Cl , as opposed to the expected O-H...Cl or O-H...F hydrogen bond. The symmetric top spectrum for $\text{CF}_3\text{Cl-H}_2\text{O}$ is rather surprising, indicating that the internal motion of water must convert the complex from an asymmetric to a symmetric top, similar to the case of $\text{CF}_4\text{-H}_2\text{O}$. This has also been recently found by Lovas *et al.*¹⁹ for $\text{CH}_3\text{CN-H}_2\text{O}$. One hydrogen of water bonds to the nitrogen along the C_{3v} axis of the acetonitrile; internal rotation of the other O-H of water about this C_{3v} axis turns this asymmetric top into an effective symmetric top. This does not happen in our analogous molecule $\text{CF}_3\text{CN-H}_2\text{O}$, whose spectrum is that of a true asymmetric top. Ouyang *et al.* studied $\text{CF}_3\text{COOH-H}_2\text{O}$,²⁰ in which water molecule forms a ring with the COOH by forming two hydrogen bonds. It will be interesting to see the effect of a cyanide group attaching to the CF_3 group. The simplest cyanide containing molecule, HCN, acts as a H atom donor when forming a complex with H_2O .^{21,22,23} Omron *et al.*²⁴ observed similar hydrogen bonded structure in $\text{HCCCN-H}_2\text{O}$. Lee *et al.*²⁵ studied the rotational spectra of $\text{NCCN-H}_2\text{O}$ and $\text{NCCN-D}_2\text{O}$ in which they derived a planar T-shaped asymmetric top structure

arising from dipole-dipole interaction. An early report of the Lovas result²⁶ on $\text{CH}_3\text{CN-H}_2\text{O}$ inspired this study. We started our project expecting that $\text{CF}_3\text{CN-H}_2\text{O}$, too, would behave as a symmetric top. The dominating force will still be hydrogen bonding after we replace CH_3 group with CF_3 group.


In this paper we report the rotational analysis of the $\text{CF}_3\text{CN-H}_2\text{O}$ complex with Fourier transform microwave spectroscopy as well as a detailed theoretical analysis of the orientation of the water molecule and the potential energy surface of the complex.

Experimental Section

The spectrum of trifluoroacetonitrile-water was measured using the Wesleyan pulsed-jet Fourier transform microwave spectrometer which has been described elsewhere^{27,28}. Trifluoroacetonitrile was prepared as a gas mixture of 1% in argon carrier gas. The sample gas was then bubbled through a reservoir of deionized water then expanded through a general valve nozzle into the vacuum chamber. The backing pressure is approximately 0.5 atm. The $\text{CF}_3\text{CN-H}_2\text{O}$ complexes were generated in the supersonic expansion. The spectrum was measured in the frequency range 16-19 GHz. Each line appears as a doublet due to the Doppler effect. The line position is calculated as the average frequencies of the two Doppler components. The full-width-half-maximum (fwhm) of the observed lines was ~5-7 kHz. With this fwhm, we can estimate peak centers to within 1 kHz.

Theoretical Calculations

Several factors play roles in the $\text{CF}_3\text{CN-H}_2\text{O}$ complex. It is hard to state either the hydrogen bonding or the dipole-dipole interaction is the predominant interaction. And, if the hydrogen bonding plays the leading role, both nitrogen and fluorine may be involved in the H bond. There is also the possibility that the halogen bond is preferred as in the cases of the Freons. To answer these questions, we performed *ab initio* calculations at MP2/6-311++G(3df,3pd) level of theory using the Gaussian09 program package.²⁹ Out of the three plausible conformers located from the calculation, the one dominated by dipole-dipole interactions, namely conf-1, as presented in Figure 1, is predicted to be the most stable conformation.

Figure 1. Three stable conformers of the CF₃CN-H₂O complex

Results and discussion

We scanned the region between 16 and 19 GHz following the prediction from the theoretical calculations. The ¹⁴N atom has a nuclear spin of 1, and the nuclear electric quadrupole coupling splits each rotational transition into a characteristic hyperfine pattern. The scan yielded almost two hundred transitions as expected from the hyperfine splitting. We first identified the transitions of the CF₃CN monomer, then identified the transitions of the CF₃CN-Argon complex based on the prediction from previous work. These CF₃CN-Argon transitions were outside of the previously scanned region. We combined these transitions with those from the previous work; the analysis is included in the following section. After removing these transitions, the remaining transitions were analysed using the predicted rotational constants and hyperfine coupling constants.

Rotational spectrum of the CF₃CN-Argon complex


A total of 18 transitions with 68 resolved hyperfine components have been identified as the transitions from the CF₃CN-Argon complex using the spectroscopic constants previously determined. Table S1 lists the newly measured frequencies of CF₃CN-Ar with their assignments.

These transitions were then combined with the previously measured transitions for the CF₃CN-Ar complex to obtain improved spectroscopic constants. The resulting spectroscopic constants are listed in Table S2 with the comparison to the literature. The rms error of the new fit is 2 kHz, which is consistent with the estimated experimental uncertainty.

Rotational spectrum of the CF₃CN-H₂O complex

The candidate transitions for the CF₃CN-H₂O complex were analysed with Pickett's least-squares fitting program, SPFIT,³⁰ using Watson's standard asymmetric rotor Hamiltonian³¹ (*A*-reduction, *J*^k representation) with an exact treatment of the nuclear quadrupole interaction. Both *a*-type and *b*-type transitions have been observed, allowing the determination of a complete set of rotational constants and centrifugal distortion constants. The nuclear quadrupole coupling constants of ¹⁴N are determined together with the rotational and centrifugal distortion constants. The coupling scheme used is **F** = **J** + **I**, where **J** is the molecular end-over-end angular momentum, **I** is ¹⁴N nuclear spin angular momentum where *I* = 1 is the nuclear

spin of ¹⁴N, and **F** is the total angular momentum. Each transition appeared as a doublet due to the tunnelling of the two equivalent forms of the water subunit within the dimer. Each of the hydrogen atom has a nuclear spin number of *I* = 1/2. The coupling of the two hydrogen nuclear spins results in two possible symmetry states. We will label the two symmetry states as A state (*para*-) for *I* = 0 and B state (*ortho*-) for *I* = 1 following reference 19. The intensity ratio of the two observed split transitions is approximately 1:3, which is in agreement with the (2*I* + 1) spin statistical weight. We assigned the weaker line of the two components to the A state. The splitting varies from transition to transition but is normally within a dozen MHz. Table S3 lists the measured frequencies and their assignment. Two hyperfine components of the *S*_{05-4₀₄ transition of the B state are shown in Figure 2. This spectrum was obtained with 64 averaging cycles. The doublets shown in Figure 2 for both hyperfine components are due to Doppler splitting which is because of the parallel setting of the molecular beam and the radiation direction.}

Figure 2. The *S*_{05-4₀₄ transition of the B state of the CF₃CN-H₂O complex. Lines are Doppler doubled, and the *F'*-*F''* quantum numbers are labelled for each transitions.}

The resulting spectroscopic constants are listed in Table 1. The rms error of the fit is 2 kHz, which is consistent with the estimated experimental uncertainty.

Table 1. Spectroscopic constants of CF₃CN-H₂O.

| | A state | B state |
|--|----------------|----------------|
| <i>A</i> /MHz | 3187.81379(48) | 3190.33856(43) |
| <i>B</i> /MHz | 1981.2828(15) | 1981.7161(14) |
| <i>C</i> /MHz | 1554.8307(18) | 1554.8797(17) |
| <i>Δ</i> _{<i>J</i>} /kHz | 2.743(32) | 2.709(29) |
| <i>Δ</i> _{<i>JK</i>} /kHz | -3.004(21) | -2.831(19) |
| <i>Δ</i> _{<i>K</i>} /kHz | 4.979(61) | 4.680(55) |
| <i>δ</i> _{<i>J</i>} /kHz | 0.7668(34) | 0.7732(32) |
| <i>δ</i> _{<i>K</i>} /kHz | 0.316(47) | 0.407(44) |
| <i>χ</i> _{<i>aa</i>} /MHz | 0.263(3) | 0.2737(25) |
| <i>χ</i> _{<i>bb</i>} - <i>χ</i> _{<i>cc</i>} /MHz | -5.0392(44) | -5.0528(40) |
| <i>N</i> | 49 | 45 |
| <i>σ</i> /kHz | 2.3 | 1.9 |

Structure Confirmation of the CF₃CN-H₂O complex

From the *ab initio* calculations at MP2/6-311++G(3df,3pd) level of theory three plausible conformers were located. Their energy, rotational constants and structures are given in Table 2 and Figure 1. We found that the conformer dominated by dipole-dipole interactions, namely conf-1, is predicted to be the most stable conformation. The calculated rotational constants from the equilibrium structure also agree well with experiment. The conformer with O-H...N H bond (conf-2) is about 0.32 kcal/mol less stable than conf-1. The conformer with halogen bond, conf-3, is significantly less stable by 2.13 kcal/mol. The dipole moments of CF₃CN and CH₃CN were determined to be 1.262 ± 0.010 D³² and 3.913 ± 0.002 D³³, respectively. The perfluorination of the acetonitrile significantly reduces the overall dipole moment of that moiety. Dipole-dipole interaction is expected to play an important role in their complexes with water. In the case of water-acetonitrile, the spectrum of a symmetric top molecule was observed¹⁹. This suggests that the oxygen atom of the water molecule is effectively on the axis of C≡N. The fitted structure of CH₃CN-H₂O complex suggests a hydrogen bonding distance of 2.075 Å between the nitrogen and hydrogen atoms. This conformation also favours the local dipole-dipole interaction of the C≡N group and O-H group. The nitrile group has much stronger polar effect than the carbon-fluorine bond, as is evidenced in the comparison of the dipole moments of acetonitrile (3.913 ± 0.002 D) and monofluoromethane (1.8572 ± 0.0010 D)³³. However, in the case of CF₃CN-H₂O complex, the local dipole moments of C-F bonds do play a more significant role than the C-H bonds in CH₃CN-H₂O complex. We speculate that molecular alignment in conf-1 may maximize the overall dipole-dipole interactions between local bonds, and by doing so stabilize conf-1 with respect to conf-2. This calls for further systematic investigation on the partially fluorinated acetonitriles. We also note that the *ab initio* rotational constants of conf-1 fit very well with the experimental values, with a maximum difference of about 1%, while those of conf. 2 and conf. 3 exhibit significant deviations. This indicates that the actual geometry of CF₃CN-H₂O is very close to that of the calculated conf. 1.

Table 2. *ab initio* (MP2/6-311++G(3df,3pd)) calculated results of energy, spectroscopic constants of the three stable conformers of the CF₃CN-H₂O complex.

| Type | Conf-1 | Conf-2 | Conf-3 | Exp. |
|---|---------|---------|---------|---------|
| <i>E</i> (kcal/mol) | 0.00 | 0.32 | 2.13 | |
| <i>E</i> ₀ (kcal/mol) ^a | 0.00 | 0.61 | 1.68 | |
| <i>A</i> /MHz | 3257.68 | 5628.49 | 5614.75 | 3228.85 |
| <i>B</i> /MHz | 2007.23 | 826.68 | 1365.78 | 2034.66 |
| <i>C</i> /MHz | 1587.25 | 824.83 | 1360.71 | 1598.49 |
| μ_a (Debye) ^b | 1.39 | 2.93 | -3.66 | |
| μ_b (Debye) ^b | 1.27 | -1.66 | 0.47 | |
| μ_c (Debye) ^b | 0.00 | 0.00 | 0.00 | |


[a] *E*₀ is the ZPE corrected relative energy of each conformer.

[b] μ_x are the dipole moment components.

Internal dynamics of the CF₃CN-H₂O complex


To rationalize the tunnelling motion of the CF₃CN-H₂O complex, we performed potential-energy surface scan using the coordinates defined in Figure 3. The angle α measures the bending amplitudes of the two constituent molecules. The angle θ reflects the rotation of water molecule around CF₃CN, and ϕ is defined as the angle between the norm of H₂O plane and the axis passing through two C atoms (see Fig. 3), which describe the internal rotation of H₂O with respect to CF₃CN.

Figure 3. Definition of geometrical parameters used to discuss the internal dynamics



To evaluate the potential curve for each geometrical parameter, an *ab initio* grid was calculated in steps of 2°. While the angle/dihedral angles were kept fixed at each step, the rest of geometrical parameters were fully optimized for each point. The results are plotted in Figure 4. As shown in Figure 4, the potential energy as a function of the angles α , θ and ϕ , which describe the bending motions of CF₃CN and water molecules, exhibits a single minimum, at $\alpha = 64^\circ$, $\theta = 0$, and $\phi = 90^\circ$, i. e., conf-1.

Figure 4. *ab initio* potential surface profile along α , θ , ϕ which are defined in Figure 3.




For the 3D potential surface scan, steps of 5° were employed. The results are plotted in Figure 5.

Conclusions

High resolution rotational spectrum of the CF₃CN-H₂O complex has been measured for the first time to produce rotational and centrifugal distortion constants, along with nuclear quadrupole coupling constants. We studied the internal dynamics of CF₃CN-H₂O complex using the rotational spectroscopic technique and the *ab initio* calculations. The observed conformation was characterized by a dipole-dipole interaction. The structure of CF₃CN-H₂O is completely different from that of CH₃CN-H₂O.

Figure 5. *ab initio* 3D potential surface with respect to θ , ϕ , which are defined in Figure 3.



Acknowledgements

This work was partially sponsored by National Science Foundation (NSF) under grant CHE-1011214 for S. E. N.. This work was partially sponsored by National Natural Science Foundation of China (NSFC) under grant Nos. 20525103, 20673088, 20973137, 20721001, 20423002, 21031004, 20703033, 21133004, 21273187 and 973 projects (Nos. 2007CB815301 and 2007CB815307) for X. L.. W. L. thanks the financial support from the Welch Foundation.

Notes and references

^a Department of Chemistry, University of Texas Rio Grande Valley, One West University Blvd., Brownsville, TX 78520, United States. E-mail: wei.lin@utrgv.edu.

^b State Key Laboratory of Physical Chemistry of Solid Surface, and Department of Chemistry, College of Chemistry and Chemical Engineering and School of Life Science, Xiamen University, Xiamen, 361005, China. E-mail: ananwu@xmu.edu.cn.

^c Department of Chemistry, Wesleyan University, 52 Lawn Ave., Middletown, CT 06459, United States

†Electronic Supplementary Information (ESI) available. See DOI: 10.1039/b000000x/

Citations here

¹ W. Kasten, H. Dreizler, B. E. Job, and J. Sheriden, *Z. Naturforsch A*, 1983, **38**, 1015.

² J. Sheridan and W. Gordy, *J. Chem. Phys.*, 1952, **20**, 591.

³ J. Sheridan and W. Gordy, *Phys. Rev.*, 1950, **77**, 292.

⁴ M. J. Whittle, J. G. Baker and G. Corbelli, *J. Mol. Spectrosc.*, 1971, **40**, 388.

⁵ C. A. Burrus and W. Gordy, *J. Chem. Phys.*, 1957, **26**, 391.

⁶ Pil Ja Seo, J. H. Carpenter and J. G. Smith, *J. Mol. Spectrosc.*, 1997, **184**, 362.

⁷ A. P. Cox, M. C. Ellis, A. C. Legon, A. Wallwork, *J. Chem. Soc., Faraday Trans.*, 1993, **89**, 2937.

⁸ J. H. Carpenter, M. Motamedi and J. G. Smith, *J. Mol. Spectrosc.*, 1995, **174**, 106.

⁹ M. Motamedi, J. H. Carpenter and J. G. Smith, *J. Mol. Spectrosc.*, 2003, **221**, 23.

¹⁰ J. H. Carpenter, M. Motamedi and J. G. Smith, *J. Mol. Spectrosc.*, 1995, **171**, 468.

¹¹ J. H. Carpenter, M. Motamedi, and J. G. Smith, *J. Mol. Spectrosc.*, 1996, **175**, 133.

¹² A. Friedrich, C. Gerke, H. Harder, H. Mader, J. Cosleou, G. Wlodarczak, and J. Demaison, *Mol. Phys.*, 1997, **91**, 697.

¹³ M. Motamedi and A. Haseli, *J. Mol. Spectrosc.*, 2006, **236**, 91.

¹⁴ W. Lin and S. E. Novick, *J. Mol. Spectrosc.*, 2007, **243**, 32.

¹⁵ L. Evangelisti and W. Caminati, *Phys. Chem. Chem. Phys.*, 2010, **12**, 14433.

¹⁶ W. Caminati, A. Maris, A. Dell'Erba, P. G. Favero, *Angew. Chem.*, 2006, **118**, 6863; *Angew. Chem. Int. Ed.*, 2006, **45**, 6711.

¹⁷ S. E. Novick, K. C. Janda, W. Klemperer, *J. Chem. Phys.*, 1976, **65**, 5115.

¹⁸ L. Evangelisti, G. Feng, P. Eciija, E. J. Cocinero, F. Castano, W. Caminati, *Angew. Chem. Int. Ed.*, 2011, **50**, 7807.

¹⁹ F. J. Lovas and J. Sobhanadri, *J. Mol. Spectrosc.*, 2015, **307**, 59

²⁰ B. Ouyang, T. G. Starkey, B. J. Howard, *J. Phys. Chem. A*, 2007, **111**, 6165

²¹ A. J. Fillery-Travis, A. C. Lego, L. C. Willoughby, *Chem. Phys. Lett.*, 1983, **98**, 369.

²² A. J. Fillery-Travis, A. C. Lego, L. C. Willoughby, *Proc. R. Soc. Lond. A*, 1984, **396**, 405

²³ H. S. Gutowsky, T. C. Germann, J. D. Augspurger, C. E. Dykstra, *J. Chem. Phys.*, 1992, **96**, 5808

²⁴ R. M. Omron, A. R. Hight Walker, G. Hilpert, G. T. Fraser, R. D. Suenram, *J. Mol. Spectrosc.*, 1996, **179**, 85

²⁵ S. Lee, I. I. Suni, W. Klemperer, *J. Chem. Phys.*, 1992, **96**, 5577

²⁶ F. J. Lovas and J. Sobhanadri, International Symposium on Molecular Spectroscopy, TC12

²⁷ A. R. Hight Walker, W. Chen, S. E. Novick, B. D. Bean, and M. D. Marshall, *J. Chem. Phys.*, 1995, **102**, 7298.

²⁸ W. Lin, L. Kang, and S. E. Novick, *J. Mol. Spectrosc.* 2005, **230**, 93.

²⁹ Gaussian 09, Revision **D.01**, M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, Ö. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski, and D. J. Fox, Gaussian, Inc., Wallingford CT, 2009.

³⁰ H. M. Pickett, *J. Mol. Spectrosc.*, 1991, **148**, 371.

³¹ J. K. G. Watson, *J. Chem. Phys.*, 1968, **45**, 4517.

³² P. B. Foreman, K. R. Chien, J. R. Williams, and S. G. Kukulich, *J. Mol. Spectrosc.*, 1974, **52**, 251.

³³ P. Alston Steiner, W. Gordy, *J. Mol. Spectrosc.*, 1966, **21**, 291.