



## Phosphoinositide-specific phospholipase C2, AtPLC2, in phosphoinositide metabolism and the endoplasmic reticulum stress response

著者	KANEHARA Kazue, YU Chao-Yuan , CHO Yueh , CHEONG Wei-Fun, TORTA Federico , SHUI Guanghou , WENK Markus R., NAKAMURA Yuki
journal or publication title	Proceedings of the annual meeting and Symposium, Japanese Society of Plant Physiologists
volume	57
page range	320-320
year	2016-03-11
URL	<a href="http://hdl.handle.net/10258/00009000">http://hdl.handle.net/10258/00009000</a>

## **Phosphoinositide-specific phospholipase C2, AtPLC2, in phosphoinositide metabolism and the endoplasmic reticulum stress response**

Kazue Kanehara, Chao-Yuan Yu, Yueh Cho, Wei-Fun Cheong, Federico Torta, Guanghou Shui, Markus R Wenk, Yuki Nakamura

Phosphoinositides are important lipid signals in plant developments and stress responses. However, multiple isoforms of the phosphoinositide biosynthetic genes hamper our understanding of the pivotal enzymes in each step of the pathway. In this study, we report that Arabidopsis phosphoinositide-specific phospholipase C2 (AtPLC2) is the primary phospholipase in phosphoinositide metabolism and is involved in the endoplasmic reticulum (ER) stress responses. Lipidomic profiling of multiple *plc* mutants showed that the *plc2-1* increased levels of its substrates phosphatidylinositol 4-phosphate and phosphatidylinositol 4,5-bisphosphate, suggesting that the major phosphoinositide metabolic pathway is impaired. AtPLC2 displayed a distinct tissue expression pattern and localized at the plasma membrane in different cell types. Moreover, the *plc2-1* showed hypersensitive response to ER stress as evidenced by changes in relevant phenotypes and gene expression profiles. Our results revealed the primary enzyme in phosphoinositide metabolism and an emerging link between phosphoinositides and the ER stress response in plants.