

Publication status: Preprint has been submitted for publication in journal

## BRAZILIAN HERNIA AND ABDOMINAL WALL SOCIETY STATEMENT ON LARGE HIATAL HERNIAS MANAGEMENT

André Brandalise, Fernando Augusto Mardiros Herbella, Renato Abrantes Luna, Sergio Szachnowicz, Rubens Antonio Aissar Sallum, Carlos Eduardo Domene, Paula Volpe, Leandro Totti Cavazzolla, Marcelo Lopes Furtado, Christiano Marlo Paggi Claus, José Francisco De Mattos Farah, Eduardo Crema in memorium

<https://doi.org/10.1590/0102-672020230069e17>

Submitted on: 2023-11-06

Posted on: 2023-11-06 (version 1)

(YYYY-MM-DD)

Position Paper, Arq. Bras. Cir. Dig. 36 • 2023

<https://doi.org/10.1590/0102-672020230069e1787>

André **BRANDALISE** <https://orcid.org/0000-0002-0285-2122>.

Fernando Augusto Mardiros **HERBELLA** <https://orcid.org/0000-0003-3594-5744>.

Renato Abrantes **LUNA** <https://orcid.org/0000-0003-2707-511X>

Sergio **SZACHNOWICZ** <https://orcid.org/0000-0002-6238-3532>,

Rubens Antonio Aissar **SALLUM** <https://orcid.org/0000-0003-1823-0042>,

Carlos Eduardo **DOMENE** <https://orcid.org/0000-0002-5115-569X>.

**PAULA VOLPE** <https://orcid.org/0000-0002-1586-5037>

Leandro Totti **CAVAZZOLLA** <https://orcid.org/https://orcid.org/0000-0003-2356-2789>

Marcelo Lopes Furtado <https://orcid.org/0000-0002-5103-5807>

Christiano Marlo Paggi **CLAUS** <https://orcid.org/0000-0003-1465-5430>

José Francisco de Mattos **FARAH** <https://orcid.org/0000-0002-3673-7341>

Eduardo **CREMA** (*in memoriam*) <https://orcid.org/0000-0002-6688-0819>

## **BRAZILIAN HERNIA AND ABDOMINAL WALL SOCIETY STATEMENT ON LARGE HIATAL HERNIAS MANAGEMENT**

Posição da Sociedade Brasileira de Hérnia e Parede Abdominal sobre o tratamento de grandes hérnias de hiato.

André **Brandalise**<sup>1</sup>, Fernando Augusto Mardiros **Herbella**<sup>2</sup>, Renato Abrantes **Luna**<sup>3</sup>, Sergio **Szachnowicz**<sup>4</sup>, Rubens Antonio Aissar **Sallum**<sup>5</sup>, Carlos Eduardo **Domene**<sup>6</sup>, Paula **Volpe**<sup>7</sup>, Leandro Totti **Cavazzolla**<sup>8</sup>, Marcelo Lopes **Furtado**<sup>9</sup>, Christiano Marlo Paggi **Claus**<sup>10</sup>, José Francisco De Mattos **Farah**<sup>11</sup>, Eduardo **Crema**<sup>12</sup>.

From <sup>1</sup> Robotic Surgery Program Coordinator, Centro Médico de Campinas Foundation, Campinas (SP), Brazil; <sup>2</sup> Department of Surgery, Federal University of Sao Paulo, São Paulo (SP), Brazil ; <sup>3</sup> Faculty of Medicine and Surgery, Servidores do Estado Federal Hospital, Universidade Federal do Estado do Rio de Janeiro, Rio de Janeiro (RJ), Brazil ; <sup>4</sup> Department of Gastroenterology, Universidade de Sao Paulo, São Paulo (SP), Brazil ; <sup>5</sup> Department of Gastroenterology, Universidade de Sao

Paulo, São Paulo (SP), Brazil; <sup>6</sup> Integrated Center for Advanced Medicine, Surgery – CIMA, President of Brazilian Society of Laparoscopic and Robotic Surgery, Sao Paulo (SP), Brazil ; <sup>7</sup> Integrated Center for Advanced Medicine, Surgery – CIMA, Sao Paulo (SP), Brazil ; <sup>8</sup> General Surgery Unit, University Hospital, Universidade Federal do Rio Grande do Sul, Porto Alegre (RS), Brazil; <sup>9</sup> Videolaparoscopic Surgery Service, Pintagueras Hospital, President of the Brazilian Hernia and Abdominal Wall Society, Jundiaí, Sao Paulo (SP), Brazil. ; <sup>10</sup> Department of Surgical Clinic, Postgraduate Program in Minimally Invasive Surgery, Universidade Positivo, Curitiba (PR), Brazil ; <sup>11</sup> Department of General and Oncological Surgery, Servidor Público Estadual de São Paulo Hospital, Sao Paulo (SP), Brazil; <sup>12</sup> Digestive Surgical Unit, Universidade Federal do Triangulo Mineiro, Uberaba (MG), Brazil (*in memorium*)

**How to cite this article:** Brandalise A, Herbella FAM, Luna RA, Szachnowicz S, Sallum RAA, Domene CE, Volpe P, Cavazzolla LT, Furtado ML, Claus CMP, Farah JFM, Crema E. ABCD Arq Bras Cir Dig. 2023;36e1787. <https://doi.org/10.1590/0102-672020230069e1787>

---

Author's contributions: All co-authors contributed to the bibliographic review, drafting the conclusions of the article, as well as the final draft of the article.

---

**Correspondence:** André Brandalise. Email: [andre@gastrosite.com.br](mailto:andre@gastrosite.com.br).

## **ABSTRACT**

Large hiatal hernias (LHH) besides being more prevalent in the elderly, have different clinical presentation: fewer reflux, more mechanical symptoms and a greater possibility of acute, life-threatening complications such as gastric volvulus, ischemia and visceral mediastinal perforation. Thus, surgical indications are distinct from gastroesophageal reflux disease (GERD-related), sliding hiatal hernias. Heartburn

tends to be less intense, while symptoms of chest pain, cough, discomfort, and tiredness are reported more frequently. Complaints of vomiting and dysphagia may suggest the presence of associated gastric volvulus. Signs of iron deficiency and anemia are found. Surgical indication is still controversial and was previously based on high mortality reported in emergency surgeries for gastric volvulus. Postoperative mortality is especially related to three factors: body mass index (BMI above 35), age over 70 years and presence of comorbidity.

Minimally invasive elective surgery should be offered to symptomatic individuals with good or reasonable performance status, regardless of age group. In asymptomatic and oligosymptomatic patients, besides obviously identifying the patient's desire, case-by-case analysis of surgical risk factors such as age, obesity and comorbidities, should be taken under consideration. One should also pay attention to situations with greater technical difficulty and risks of acute migration due to increased abdominal pressure (abdominoplasty, manual workers, spastic diseases). Technical alternatives such as partial fundoplication and anterior gastropexy can be considered.

We emphasize the importance of performing surgical procedures in cases of LHH in high-volume centers, with experienced surgeons.

**KEYWORDS:** Hernia, Hiatal. General Surgery. Fundoplication. Laparoscopy. Robotic Surgery. Surgical Mesh.

## **RESUMO**

As grandes hérnias de hiato (HHG), além de serem mais prevalentes em idosos, têm apresentação clínica diferente: menos refluxo, mais sintomas mecânicos e maior possibilidade de complicações agudas e potencialmente fatais, como vólvulo gástrico, isquemia e perfuração mediastinal visceral. Assim, as indicações cirúrgicas são distintas das hérnias de hiato por deslizamento, relacionadas à doença do refluxo gastroesofágico (DRGE). A azia tende a ser menos intensa, enquanto os sintomas de dor no peito, tosse, desconforto e cansaço são relatados com maior frequência. Queixas de vômitos e disfagia podem sugerir a presença de volvo gástrico associado.

São encontrados sinais de deficiência de ferro e anemia. A indicação cirúrgica ainda é controversa e foi anteriormente baseada na alta mortalidade relatada em cirurgias de emergência para volvo gástrico. A mortalidade pós-operatória está especialmente relacionada a três fatores: índice de massa corporal (IMC acima de 35), idade superior a 70 anos e presença de comorbidades.

A cirurgia eletiva minimamente invasiva deve ser oferecida a indivíduos sintomáticos, com desempenho bom ou razoável, independentemente da faixa etária. Em pacientes assintomáticos e oligossintomáticos, além de obviamente identificar o desejo do paciente, deve-se levar em consideração a análise caso a caso dos fatores de risco cirúrgico, como idade, obesidade e comorbidades. Deve-se atentar também para situações de maior dificuldade técnica e riscos de migração aguda por aumento da pressão abdominal (abdominoplastia, trabalhos manuais, doenças espásticas). Alternativas técnicas como funduplicatura parcial e gastropexia anterior podem ser consideradas.

Ressaltamos a importância da realização de procedimentos cirúrgicos nos casos de GHH em centros de grande volume, com cirurgiões experientes.

**DESCRITORES:** Hérnia Hiatal. Cirurgia Geral. Funduplicatura. Laparoscopia. Cirurgia Robótica. Telas Cirúrgicas.

### **Central message**

Large hiatal hernias (LHH) management, also known as intrathoracic stomach or giant hiatal hernias, is a controversial topic. The Brazilian Hernia and Abdominal Wall Society did a consensus meeting on the area based on the experience of national expert surgeons and current literature. The main topics was arranged regarding the management of LHH in questions, which were presented by two experts and discussed among the others until a consensus was reached.

## **Perspectives**

Minimally invasive elective surgery should be offered to symptomatic individuals with good or reasonable performance status, regardless of age group. In asymptomatic and oligosymptomatic patients, besides obviously identifying the patient's desire, case-by-case analysis of surgical risk factors such as age, obesity and comorbidities, should be taken under consideration. One should also pay attention to situations with greater technical difficulty and risks of acute migration due to increased abdominal pressure (abdominoplasty, manual workers, spastic diseases). Technical alternatives such as partial fundoplication and anterior gastropexy can be considered. We emphasize the importance of performing surgical procedures in cases of LHH in high-volume centers, with experienced surgeons.

Financial Source: None

Conflicts of Interest: None

Received: 10/13/2013

Accepted: 10/21/2023

## **1. INTRODUCTION**

Large hiatal hernias (LHH) management, also known as intrathoracic stomach or giant hiatal hernias, is a controversial topic. Hiatal hernias may be classified as: type I (sliding); type II (paraesophageal hernia); type III (mixed sliding and paraesophageal); and type IV (associated to the herniation of other organs) <sup>29</sup>.

The Brazilian Hernia and Abdominal Wall Society did a consensus meeting on the area based on the experience of national expert surgeons and current literature.

We have arranged the main topics regarding the management of LHH in questions, which were presented by two experts and discussed among the others until a consensus was reached.

## **2. DEFINITION**

The group considered for this statement only LHH type II, III and IV, defined as those with more than 5cm or one third of herniated stomach into the mediastinum.

### **3. ROUTINE SURGICAL THERAPY FOR ASYMPTOMATIC OR OLIGOSYMPTOMATIC PATIENTS**

- **Large hiatal hernias (LHH) are most commonly found after the seventh decade of life.**
- **The vast majority is asymptomatic, and the chances of acute complications are low.**
- **When complications occur, especially gastric volvulus, mortality rates are high.**

Large hiatal hernias (LHH) besides being more prevalent in the elderly, have different clinical presentation - less reflux, more mechanical symptoms and a greater possibility of acute, life-threatening complications such as gastric volvulus, ischemia and visceral mediastinal perforation. Thus, surgical indications are distinct from GERD-related, sliding hiatal hernias <sup>18,20,27</sup>.

Heartburn tends to be less intense, while symptoms of chest pain, cough, discomfort, and tiredness are reported more frequently. Complaints of vomiting and dysphagia may suggest the presence of associated gastric volvulus. Signs of iron deficiency and anemia are found in almost 20% of cases <sup>9</sup>.

Surgical indication in patients with large LHH still controversial and was previously based on high mortality reported in emergency surgeries for gastric volvulus <sup>7,18,28</sup>.

In the 1980s and 1990s, publications with analysis of large populations, demonstrated that acute complications, although severe and of high mortality, are less frequent than was once imagined and asymptomatic or oligosymptomatic patients may be followed up without surgical intervention. Watch and wait is an acceptable approach, especially based on the findings that the overall mortality was 5.4% and the

frequency of 1.1% per year of asymptomatic cases required surgery <sup>45</sup>. More recent studies highlight the possibility of observation in asymptomatic patients, although surgery can be safely offered and good results in patients with controlled comorbidities regardless of age group <sup>32</sup>.

On the other hand, acute complications, such as gastric volvulus with incarceration or vascular suffering remains one of the most dramatic urgent situations, especially in the elderly <sup>10</sup>. A recent publication compared a series of urgent surgeries in LHH and showed a 1.7-fold increase in morbidity and a 2.7-fold increase in mortality when compared to elective surgery, even after age adjustment and comorbidities <sup>14</sup>. These authors recommend elective, minimally invasive surgery for symptomatic and oligosymptomatic patients considered fit for the procedure. This onset of paradigm shift (contrary to observation in asymptomatic patients) can be considered an effect of the results of minimally invasive surgery, which showed complication and mortality rates lower than 1% in large series <sup>10,18</sup>.

Postoperative mortality is especially related to three factors: body mass index (BMI above 35), age over 70 years and presence of comorbidity (Charlson index greater than 3) <sup>2</sup>. Age-related mortality in patients under 60 years of age or 60-69 is 0%, between 70-79 is 0.9% and in patients over 80 years the mortality of 7.8% <sup>28,29</sup>. In general, mortality between 0.9 and 4.6% is described after elective correction of LHH <sup>13</sup>.

## **Recommendations**

Minimally invasive elective surgery should be offered to symptomatic individuals with good or reasonable performance status, regardless of age group.

In asymptomatic and oligosymptomatic patients, besides obviously identifying the patient's desire, case-by-case analysis of surgical risk factors such as age, obesity and comorbidities, should be taken under consideration. One should also pay attention to situations with greater technical difficulty and risks of acute migration due to increased abdominal pressure (abdominoplasty, manual workers, spastic diseases).

We emphasize the importance of performing surgical procedures in cases of LHH in high-volume centers, with experienced surgeons.



#### **4. PRE-OPERATIVE WORK-UP**

- **The commonly used work-up in LHHs includes: upper digestive endoscopy, barium upper GI series, prolonged esophageal pH monitoring (pH-metry), esophageal manometry, computed tomography.**

##### **Upper GI endoscopy**

Most commonly performed when the patient's complaint is reflux, regurgitation or dysphagia. The examination is challenging in very large hernias, with alteration in gastric anatomy and esophageal tortuosity. Sometimes endoscopy fails to estimate the size of the hernia or the length of the esophagus, and should always be associated with an imaging test <sup>27</sup>.

In emergency cases endoscopy can decompress an obstructed stomach and evaluate the mucosal integrity.

##### **Upper GI Barium Series**

It's a non-invasive and low-cost test, and gives us an overview of the hernia, which sometimes becomes difficult by endoscopy.

It is possible to identify the size of the hernia, location of the esophagogastric junction in relation to the diaphragm and the shape of the herniated stomach, whether anatomical or with some type of volvulus <sup>47</sup>.

Esophageal shortening or stenosis are also very well evaluated. In cases when a motility test is not feasible, it shows, especially when a video barium swallow test is performed, esophageal emptying.

##### **CT scan**

Nowadays we have seen many patients referred to a Computed tomography (CT) scan for investigation of thoracic symptoms that discovers a LHH. CT scan allows a good anatomical vision of the hernia and identify other herniated organs.

CT scan is especially useful in acute cases, where a complication of hiatal hernia, such as volvulus or perforation, is suspected. The use of venous contrast may evaluate perfusion changes in the stomach or other herniated organs.

Due to the ability to make sagittal, coronal and axial cuts and three-dimensional reconstructions, it is also possible to evaluate the diameter of the hiatal defect at its rest position and determine the herniated volume to try predicting cases of greater complexity<sup>24,26</sup>.

### **Esophageal Manometry**

Many patients with LHH have dysphagia as part of their symptoms and to evaluate esophageal motility is important, not for the diagnosis of the disease, but to plan the surgical procedure<sup>15</sup>.

The test itself bares greater technical difficulty since frequently it isn't possible to transpose the GEJ with the probe. If this is the case, it can be guided by endoscopy before the examination or it can be studied only the esophageal body to assess whether there is adequate motility for a fundoplication.

Although up to 53% of motor alterations are found in patients with LHH, those may be secondary to the hernia and would not have a negative influence at the results of surgeries in relation to patients without motor alterations detected<sup>52</sup>.

Manometry can be conventional or high resolution, and there is no clear advantage of one over the other.

### **Prolonged pH or pH-impedance monitoring**

Tests known as "gold standard" for the diagnosis of GERD, they are not important in patients with LHH. Should be requested if there is doubt whether the

patient's symptoms or main complaint to decide about having surgery is related to gastroesophageal reflux<sup>15</sup>.

## **Recommendations**

Both upper digestive endoscopy and upper GI series are important for the diagnosis and classification of LHH and should be requested to all patients with this type of hernia, being complementary to each other.

CT scan helps in the identification of multivisceral herniations (type IV) and are recommended in more complex hernias.

Esophageal manometry is recommended in patients complaining of dysphagia or when a fundoplication is planned.

Prolonged pH monitoring study is unnecessary in most of the cases.

## **5. CONSIDERATIONS ON SURGICAL TECHNIQUE**

### **5.1 Hernia sac management**

- **Finding the appropriate dissection plan and removing the hernia sac from the mediastinum is an important surgical step**
- **Partial resection of the hernia sac with complete disconnection from the crura may be an alternative, especially in redos.**

The hernia sac is of main importance in the surgical treatment of LHH. Often the viscera, in particular the stomach and esophagogastric junction can't be reduced by traction back to the abdominal cavity because the entire posterior part of the sac covers the esophagus and is adhered to it.

The recommended surgical technique for LHH should be finding a dissection plan between the hernial sac and the mediastinal structures and pleura, which is practically avascular, completely reducing the hernia sac to the abdominal cavity and thereby the EGJ and stomach return to their anatomical position <sup>16,51</sup>.

In recurrent LHH, this correct plane is more fibrotic and adherent to the pleura, leading to higher risks of pleural injury. In the event of a capnotorax, within a few moments, chest and abdominal pressures are equalized and the surgery can be continued with minor adjustments to ventilatory parameters by the anesthesiologist, associated with a reduction of the pneumoperitoneum pressure if needed.

It is strongly recommended that both vagal trunks are identified and preserved. In LHH, the posterior vagus nerve may be further away from the esophagus, increasing the risk of its injury.

After complete reduction of the hernia sac it can be removed or transected to adequately expose the Angle of His, thus establishing an anatomical landmark for the EGJ to make sure an adequate length of abdominal esophagus was achieved.

## **Recommendations**

The hernia sac must always be completely disconnected from the hiatus. Complete resection is optional.

It is strongly recommended to actively search and identify the anterior and posterior vagal trunks, due to a higher risk of iatrogenic injury.

### 5.2. How cranial should the thoracic esophagus be dissected?

- **Achieving an adequate abdominal length is of main importance on the surgical treatment of LHH**
- **Mediastinal dissection must be as cranial as possible to reduce axial tension (cranial traction) so the esophagus maintains its abdominal position at the end of surgery, without traction**

The existence of an intra-abdominal segment of the esophagus is a natural antireflux mechanism since abdominal positive pressure collapses the esophageal walls by external pressure. Also, for decades it has been demonstrated that reflux control is directly related to the length of the abdominal segment of the esophagus <sup>15</sup>.

Thus, one of the objectives of antireflux surgery is to restore of an adequate abdominal esophageal length, especially in patients with hiatal hernia. In addition, inadequate length of intra-abdominal esophagus is associated with recurrence of hernias.

Restoration of abdominal esophagus length is achieved through circular dissection of the esophagus in the cranial direction. Therefore, the thoracic esophagus should be dissected to sufficient extent to obtain satisfactory abdominal esophageal length <sup>17,21</sup>.

There are no studies comparing the benefits of different final lengths of the abdominal esophagus, and the *recommendation of experts* is to obtain 3 cm of extension, without caudal traction <sup>17,41</sup>.

In some cases, even after extensive dissection, adequate abdominal esophageal length may not be obtained, defining the so-called "short esophagus". In such situations, other measures should be taken and will be discussed later on.

## **Recommendations:**

The thoracic esophagus should be dissected in an extension sufficient to obtain 3cm of abdominal esophagus, without traction.

### **5.3 – Can adjuvant techniques improve the results?**

Axial tension from the upward traction forces of the esophagus may be involved in recurrence. Some techniques are described to try to reduce it.

#### **5.3.1 – Collis gastroplasty:**

Collis gastroplasty is a technical surgical step used to reduce axial tension when at least 3 cm of abdominal esophagus isn't obtained after extensive mediastinal dissection.

Although efficient and potentially reducing hernia recurrences, this technique has some drawbacks.

Firstly, the morbidity of the operation increases, and fistula is a rare but possible occurrence. Secondly, the longer the stomach segment required in gastroplasty, the greater the number of parietal cells above the fundoplication, leading to pathological reflux in up to 50% of cases. Another concern is the presence of a gastric segment without peristalsis above the fundoplication, which may lead to dysphagia <sup>23,48</sup>.

Despite these potential problems, the quality of life reported in patients undergoing this technique is similar to those undergoing traditional fundoplication <sup>33</sup>.

### **5.3.2 Vagotomy;**

Truncal vagotomy was described by Oeschlager et al. <sup>37</sup> as a simple maneuver to obtain EGJ reduction and thus avoid Collis gastroplasty. This is a solitary study, showing that when EGJ was not reduced to more than 2 cm in the abdomen without tension, performing truncal vagotomy led to achieve adequate abdominal length without the association of potential known adverse effects related to this maneuver (delayed gastric emptying or dumping).

With scarcity of data in the literature, this should be a maneuver used with extreme caution since it is irreversible and with potential adverse effects.

### **5.3.2 Gastropexy or gastrostomy:**

Gastropexy is an attempt to fix the stomach in the abdominal cavity, and some forms have been proposed: fixation to the crura, Hill's gastropexy (where the small curvature is fixed to the arcuate ligament), anterior gastropexy with nonabsorbable sutures and finally fixation by gastrostomy.

All these techniques of anterior gastropexies are described in the literature and presented as case series. There is insufficient evidence for a definitive conclusion. However, the frequency of recurrences described in those series is relatively low - reaching 15% <sup>1,3,39</sup>.

## **Recommendations:**

Collis gastroplasty should be part of the tactical armamentarium of the surgeon who deals with complex cases of hiatal hernia, but its use should be restricted to exceptional situations where it is not possible to reduce the GEJ to the abdominal cavity.

Different forms of gastropexy can be used, given the low rates of complications reported, but they do not serve as substitutes for an adequate mobilization and reduction of the JEG to the abdominal cavity.

### **5.4 – How should the hiatus be closed?**

- **Adequate hiatal closure around the esophagus is of main importance in the proper treatment of LHH.**
- **There is consensus in the literature that nonabsorbable sutures must be used, but suture technique (simple stitches, figure-of-eight, mattress or continuous suture) and types of suture (multifilament, monofilament, barbed) are still the surgeon's preference**

Axial and radial tensions to which EGJ and diaphragmatic pillars are submitted, contribute to the higher rate of recurrences seen in LHH. In this topic we are interested in the radial tension on the crura.

Recent literature on the subject demonstrates that radial tension is proportionally related to the width and the area of the hiatus, and that the surgeon's clinical impression does not always correspond to the tension objectively measured in the pillars. In addition, the shape of the hiatus (slit, drop, D or oval) is also related to the closing tension of the pillars, which are larger in the D and oval configurations<sup>4,34</sup>.

In large, D-shaped or oval hiatus, when the tension is high, lateral sutures approaching the left crus to itself (at 2 o'clock position, in analogy) may be necessary to adequate closure around the esophagus.

Regarding the closure itself, nonabsorbable sutures should be used, with no difference between continuous suture, separate, simple stitches, in X or U. The use of Teflon pledgets is popular among many services, but to this day has not proven to be beneficial in this scenario, prevailing here the preference of the surgeon.

Usually, the hiatal closure begins posteriorly, where the radial tension is lower, and continued anteriorly. Excessive anterior angulation of the esophagus may cause dysphagia and should be avoided. Placing sutures anterior to the esophagus may be performed, when necessary, without compromising the efficiency of the repair.

In very large defects, some experts are performing diaphragmatic relaxing incisions. By making an opening, on the right or left diaphragm, reduction of the tension on the crural sutures is achieved. It is reported a reduction of 46% at the tension with relaxing incision alone, and 56% reduction when associated with pleurotomy <sup>4</sup>. Such an incision produces a weak area at the incision site and must be fixed with an inlay, non-absorbable mesh, placed away from the esophagus or fundoplication.

One large series describes the use of the relaxing incision in 11% of paraesophageal hernia cases (in the so-called more severe cases of suture tension). With this tactic authors report recurrence similar to that observed in cases of lower severity. They draw attention to the need to associate a prosthesis to cover the defect created by the incision. Although they did not report complications related to the prosthesis, there is a risk of iatrogenic injury due its fixation next to the vena cava or thoracic duct, conditioning the use of this tactic to surgeons with a high affinity with the technique <sup>12</sup>.

Another tactic maneuver described to help complete hiatal closure around the esophagus is pleurotomy. The intentional creation of a capnothorax (usually at the left side), results in a “flattening” of the diaphragm, which reduces the tension at the hiatal sutures <sup>4</sup>.

### **Recommendations:**



Complete closure of the hiatus around the esophagus with non-absorbable sutures must be obtained. An intraesophageal bougie should be used to avoid tight closure or anterior angulation of the esophagus.

Regarding adjuvant maneuvers to help hiatal closure, such as pleurotomy and relaxing incisions, the lack of consistent studies suggests that their use should be based on a case-by-case evaluation, highly depending on the surgeon's expertise.

### **5.5 – Can the use of mesh can improve results?**

- **The use of mesh as a method to reduce recurrences of LHH is based on the success they had in other abdominal wall hernias, but this positive effect is still matter of great debate among specialists.**
- **The use of nonabsorbable prostheses has a good effect in preventing anatomical recurrence in short-term follow-up, but without long-term evidence. Its use can potentially cause severe complications, requiring complex surgeries to treat, associated with high morbidity and mortality rates.**
- **Absorbable meshes can reduce short- and medium-term recurrences, but the long-term follow-up is still under investigation**

Prospective and randomized studies in the literature have conflicting results on the reduction of recurrences using prostheses to reinforce hiatoplasty. A major limitation is that the few randomized studies have analyzed different types of meshes, varying in relation to the material (polypropylene, PTFE, biological, absorbable synthetic), shape (U-shaped, C-, rectangular, circumferential), size (1 x 3 cm, 3 x 4 cm), fixation methods (points, staples, glues) and application position (anterior or posterior to the esophagus). In addition to having different definitions as to what would large hernias be <sup>11,12,30,40</sup>.

The use of permanent prosthesis is described to be associated with severe complications, during the surgery due to accidents with tacks fixation and later on the follow-up, such as intra luminal erosion (esophagus or stomach) and stenosis at the

EGJ due to retraction and dense fibrosis, which usually require complex visceral resections to treat <sup>44</sup>.

These risks, despite being isolated case reports, led to reluctance in applying permanent prosthesis on the hiatus and the search for biological prostheses, which would be absorbed after a certain time. However, a randomized controlled trial, showed long term recurrence rates similar on mesh versus non-mesh groups <sup>35,36</sup>.

Recently, absorbable synthetic meshes were introduced, aiming to induce local proliferation of fibroblasts, favoring the formation of a more resistant scar and causing a local inflammatory reaction that would cause greater fixation of the esophagus/fundoplication complex in the hiatus region <sup>40</sup>.

A recent meta-analysis comparing synthetic absorbable meshes to no mesh, reported a significantly lower recurrence rate compared to the non-mesh group (8% vs. 18%; 95%CI 0.08–0.17). Long term results of these types of mesh are still unknown <sup>11</sup>.

### **Recommendations:**

Based on recent literature, there is no data that can contraindicate or strongly recommend the use of prosthesis as an absolute way to reduce recurrences in LHH.

The use prosthesis to reinforce hiatoplasty remains a surgeon's personal choice, based on experience, always considering the potential risks of complications in the short and long term.

We recommend that, if a mesh is to be applied, it should be resorbable and fixed with suture or glue.

Mesh must not be placed without complete approximation of the hiatus around the esophagus. There is no indication of using meshes as a bridge or inlay repair.

### **5.6 – Is the fundoplication necessary?**

- **Fundoplication is recommended by the majority of experts during surgical management of LHH**
- **Fundoplication lowers the occurrence of GERD , which is very common after correction of the hernia.**
- **Despite lacking strong evidence in the literature, it's suggested that fundoplication should also lower the risk of LHH recurrence.**

The addition of a fundoplication to LHH repair may be controversial. It prevents gastroesophageal reflux disease that may occur even if it was not present previously due to extensive dissection of the esophagogastric junction and disruption of natural antireflux mechanisms. Moreover, the wrap may help keep the stomach in abdominal position.

On the other hand, some authors recommend only selective performance of fundoplication, that is, only in those patients with evidence of reflux, due to the fact that fundoplication may be associated with post operative complications such as dysphagia and gas bloat,

In a recent study, 60 patients who underwent surgery for LHH were divided into 35 patients with fundoplication versus 25 without. In patients where fundoplication was not performed, despite not having reflux previously, there was a 28% incidence of esophagitis and 39% of abnormal acid exposure <sup>19</sup>.

Upon decided that a fundoplication will be part of the procedure, whether it should be a total or partial fundoplication is still a matter of debate. Each one has its own pros and cons, but the general consensus is that if the aim of the fundoplication is preventing long term reflux, it favors a total fundoplication. If the main goal is preventing gas bloat and dysphagia, a partial fundoplication should be the choice.

## **Recommendations**

A fundoplication should be added routinely to the LHH repair. Besides restoring the anatomy after extended EGJ dissection, helps keeping the stomach in the abdominal situation, preventing hiatal hernia and GERD recurrence.

We recommend a total fundoplication but a partial may also be used specially in the setting of old age, dysphagia or dysmotility.

### **5.7. Robotic platform**

Robotic is a stable platform with many ergonomic advantages to the surgeon. For LHH repair, the stability of optics and the ability of the robotic arms to work parallel to the esophagus within the restricted space of the mediastinum, may aid achieving an easier and higher dissection of the proximal esophagus.

In longer surgeries, specially LHH in a redo situation, the comfort delivered for the surgeon in the console, favors better surgical technique.

There are not yet high quality prospective studies, clinical trials or metanalysis of results to compare robotic LHH repair to laparoscopy. A low quality systematic review disclosed similarity of results <sup>49</sup>. A populational study with almost 170,000 individuals; however, showed a higher index of complication for robotic surgery, especially pulmonary and esophageal perforations, even when matched for surgeon's expertise <sup>50</sup>.

Some studies evaluated robotic surgery in special situations. The repair of recurrent HH showed similar results to the index operation in 51 patients <sup>42</sup>. Robotic surgery was superior in emergency cases according to a multicenter study; however, the procedure was performed only by the most experienced teams in the emergency <sup>22</sup>.

It is undeniable that robotic platforms are a huge advance in technology applied to minimally invasive surgery. Adds ergonomics, stability of the camera and precision of dissection, but it is still waiting consistent data to conclude whether it contributes to better results than laparoscopic surgery, both in teams with experience in complex LHH surgeries.

## **Recommendations**

Robotic platforms may add stability of dissection and ergonomic advantages in LHH surgeries. There is no clear advantages in outcomes or complications when compared to laparoscopic approach, when both are performed by experienced surgeons.

## **6. SPECIAL SITUATIONS**

Specific clinical scenarios may demand a different approach when managing LHH.

### **6.1 – Acute complication (emergency scenario)**

Emergent situations may occur in the setting of LHH. Upper digestive hemorrhage is a rare but possible situation. Fortunately, it is generally controlled by clinical and endoscopic methods <sup>25</sup>. The most common and morbid condition is obstruction usually followed by ischemia of the stomach that comprises 98% of the cases of emergent operations. Almost 30% of the patients with acute volvulus may present with ischemia leading to 50% mortality rate <sup>6</sup>. Obstruction is caused by gastric volvulus. Classically, it is clinically represented by the Borchardt triad: thoracic (or upper abdominal) pain, impossibility to vomit and inability to pass a nasogastric tube<sup>6</sup>.

An upper digestive endoscopy must be the initial approach in the emergency to try emptying the stomach and locate a nasogastric tube for decompression. These acts may prevent ischemia and eventually devolvulate the stomach or postpone surgical therapy to allow improvement of the clinical condition.

Emergency surgery becomes the only option in cases of failure of endoscopic decompression or signs of ischemia and/or sepsis. Complex operations may be anticipated and gastric resection may be necessary. HH should be treated and an alternative feeding route provided. The access may through laparotomy or minimally invasive depending on the expertise of the team. A recent study showed that matched

groups had double risk for sever complications and three times more mortality for emergency operation even if age and comorbidities are considered <sup>46</sup>.

## **6.2 – Obesity**

Obesity is a risk factor for HH. One third of the obese population may have HH and 5% will present a LHH <sup>8</sup>. Moreover, obesity is linked to higher rates of recurrence <sup>29,31</sup>.

Obese patients, as well as patients with increased intra-abdominal pressure due to other causes such as abdominoplasty, have worse long-term results following hiatal hernia repairs and should be also managed with caution in cases of LHH.

In cases of obesity grades I and II without comorbidities, weight loss guidance before the surgical procedure is common in clinical practice, despite presenting irregular results. In morbidly obese patients, there is already a consensus that a bariatric procedure should be added to the hernia correction. Roux-en-Y gastric by-pass is the recommended technique because it presents better results in treating reflux, with good weight loss, control of comorbidities, and reduction of HHG recurrences <sup>43</sup>.

Sleeve gastrectomy is associated with the development of postoperative reflux, presents a greater chance of herniation and should be avoided.

## **6.3 – Old age**

LHH are frequently found in old age. The average age is between 65-75 years. Safety, quality of life and recurrence do not seem to be affected by age, even considering a group of octogenarians. Time of hospitalization, need for intensive care, and mechanical ventilation were; however, longer in the oldest <sup>38</sup>.

## **Recommendations**

Endoscopic therapy should always be tried as a first tentative to decompress the stomach and locate a nasogastric tube at least to prepare the patient to a definite surgical treatment in better clinical conditions in the emergency setting. Cases of ischemia and sepsis should be treated by surgical therapy. Minimally invasive procedures are the preferential route if possible and sepsis control and a feeding route are mandatory.

A bariatric procedure should be added in the morbidly obese. Roux-en-Y gastric bypass is the procedure of choice.

Treatment of LHH in elderly patients should be performed in symptomatic patients electively with results and similar safety to non-elderly patients. Technical alternatives such as partial fundoplication and anterior gastropexy can be considered.

## REFERENCES

1. Aye RW, Swanstrom LL, Kapur S, Buduhan G, Dunst CM, Knight A, et al. A randomized multiinstitution comparison of the laparoscopic Nissen and Hill repairs. *Ann Thorac Surg.* 2012;94(3):951-7; discussion 957-8. doi: 10.1016/j.athoracsur.2012.04.083.
2. Ballian N, Luketich JD, Levy RM, Awais O, Winger D, Weksler B, et al. A clinical prediction rule for perioperative mortality and major morbidity after laparoscopic giant paraesophageal hernia repair. *J Thorac Cardiovasc Surg.* 2013;145(3):721-9. doi: 10.1016/j.jtcvs.2012.12.026.
3. Bellevue OC, Louie BE, Jutric Z, Farivar AS, Aye RW. A Hill Gastropexy Combined with Nissen Fundoplication Appears Equivalent to a Collis-Nissen in the Management of Short Esophagus. *J Gastrointest Surg.* 2018;22(3):389-395. doi: 10.1007/s11605-017-3598-4.
4. Bradley DD, Louie BE, Farivar AS, Wilshire CL, Baik PU, Aye RW. Assessment and reduction of diaphragmatic tension during hiatal hernia repair. *Surg Endosc.* 2015;29(4):796-804. doi: 10.1007/s00464-014-3744-y.

5. Brandalise A, Aranha NC, Brandalise NA. The polypropylene mesh in the laparoscopic repair of large hiatal hernias: technical aspects. *Arq Bras Cir Dig.* 2012; 25(4):224–228. <https://doi.org/10.1590/S0102-67202012000400003>
6. Carter R, Brewer LA 3rd, Hinshaw DB. Acute gastric volvulus. A study of 25 cases. *Am J Surg.* 1980;140(1):99-106. doi: 10.1016/0002-9610(80)90424-9.
7. Chan EG, Sarkaria IS, Luketich JD, Levy R. Laparoscopic Approach to Paraesophageal Hernia Repair. *Thorac Surg Clin.* 2019;29(4):395-403. doi: 10.1016/j.thorsurg.2019.07.002.
8. Che F, Nguyen B, Cohen A, Nguyen NT. Prevalence of hiatal hernia in the morbidly obese. *Surg Obes Relat Dis.* 2013;9(6):920-4. doi: 10.1016/j.soard.2013.03.013.
9. Cheverie JN, Lam J, Neki K, Broderick RC, Lee AM, Matsuzaki T, et al. Paraesophageal hernia repair: a curative consideration for chronic anemia? *Surg Endosc.* 2020;34(5):2243-2247. doi: 10.1007/s00464-019-07014-3.
10. Choi S, Tang A, Murthy S, Raja S. Preoperative Evaluation and Clinical Decision Making for Giant Paraesophageal Hernias: Who Gets an Operation? *Thorac Surg Clin.* 2019;29(4):415-419. doi: 10.1016/j.thorsurg.2019.07.006.
11. Clapp B, Kara AM, Nguyen-Lee PJ, Annabi HM, Alvarado L, Marr JD, et al. Does bioabsorbable mesh reduce hiatal hernia recurrence rates? A meta-analysis. *Surg Endosc.* 2023;37(3):2295-2303. doi: 10.1007/s00464-022-09514-1.
12. Crespín OM, Yates RB, Martin AV, Pellegrini CA, Oelschlager BK. The use of crural relaxing incisions with biologic mesh reinforcement during laparoscopic repair of complex hiatal hernias. *Surg Endosc.* 2016;30(6):2179-85. doi: 10.1007/s00464-015-4522-1.
13. Damani T, Ray JJ, Farag M, Shah PC. Elective paraesophageal hernia repair in elderly patients: an analysis of ACS-NSQIP database for contemporary morbidity and mortality. *Surg Endosc.* 2022;36(2):1407-1413. doi: 10.1007/s00464-021-08425-x.
14. Dellaportas D, Papaconstantinou I, Nastos C, Karamanolis G, Theodosopoulos T. Large Paraesophageal Hiatus Hernia: Is Surgery Mandatory? *Chirurgia (Bucur).* 2018;113(6):765-771. doi: 10.21614/chirurgia.113.6.765.
15. DeMeester TR, Wernly JA, Bryant GH, Little AG, Skinner DB. Clinical and in vitro analysis of determinants of gastroesophageal competence. A study of the



- principles of antireflux surgery. *Am J Surg.* 1979;137(1):39-46. doi: 10.1016/0002-9610(79)90008-4.
16. Edey M, Salky B, Posner A, Fierer A. Sac excision is essential to adequate laparoscopic repair of paraesophageal hernia. *Surg Endosc.* 1998;12(10):1259-63. doi: 10.1007/s004649900832.
  17. Flores LE, Armijo PR, Xu T, Otten M, Hennings D, Krause C, et al. How high is too high? Extensive mediastinal dissection in patients with hiatal hernia repair. *Surg Endosc.* 2021;35(5):2332-2338. doi: 10.1007/s00464-020-07647-9.
  18. Fullum TM, Oyetunji TA, Ortega G, Tran DD, Woods IM, Obayomi-Davies O, et al. Open versus laparoscopic hiatal hernia repair. *JLS.* 2013;17(1):23-9. doi: 10.4293/108680812X13517013316951.
  19. Furnée EJ, Draaisma WA, Gooszen HG, Hazebroek EJ, Smout AJ, Broeders IA. Tailored or routine addition of an antireflux fundoplication in laparoscopic large hiatal hernia repair: a comparative cohort study. *World J Surg.* 2011;35(1):78-84. doi: 10.1007/s00268-010-0814-8.
  20. Grubnik VV, Malynovskyy AV. Laparoscopic repair of hiatal hernias: new classification supported by long-term results. *Surg Endosc.* 2013;27(11):4337-46. doi: 10.1007/s00464-013-3069-2.
  21. Herbella FA, Del Grande JC, Colleoni R. Short esophagus or bad dissected esophagus? An experimental cadaveric study. *J Gastrointest Surg.* 2003;7(6):721-5. doi: 10.1016/s1091-255x(03)00077-5.
  22. Hosein S, Carlson T, Flores L, Armijo PR, Oleynikov D. Minimally invasive approach to hiatal hernia repair is superior to open, even in the emergent setting: a large national database analysis. *Surg Endosc.* 2021;35(1):423-428. doi: 10.1007/s00464-020-07404-y.
  23. Jobe BA, Horvath KD, Swanstrom LL. Postoperative function following laparoscopic collis gastroplasty for shortened esophagus. *Arch Surg.* 1998;133(8):867-74. doi: 10.1001/archsurg.133.8.867.
  24. Kao AM, Ross SW, Otero J, Maloney SR, Prasad T, Augenstein VA, et al. Use of computed tomography volumetric measurements to predict operative techniques in paraesophageal hernia repair. *Surg Endosc.* 2020;34(4):1785-1794. doi: 10.1007/s00464-019-06930-8.

25. Kapadia S, Jagroop S, Kumar A. Cameron ulcers: an atypical source for a massive upper gastrointestinal bleed. *World J Gastroenterol.* 2012;18(35):4959-61. doi: 10.3748/wjg.v18.i35.4959.
26. Kavic SM, Segan RD, George IM, Turner PL, Roth JS, Park A. Classification of hiatal hernias using dynamic three-dimensional reconstruction. *Surg Innov.* 2006;13(1):49-52. doi: 10.1177/155335060601300108.
27. Lopes LR, Chaim FHM, Santos IGG, Coelho Neto JS, Terciotti V Jr, Ferrer JAP, Andreollo NA. Video-Laparoscopic Treatment of Intrathoracic Gastric Volvulus. *JLS.* 2020;24(4):e2020.00061. doi: 10.4293/JLS.2020.0006
28. Luketich JD, Raja S, Fernando HC, Campbell W, Christie NA, Buenaventura PO, et al. Laparoscopic repair of giant paraesophageal hernia: 100 consecutive cases. *Ann Surg.* 2000;232(4):608-18. doi: 10.1097/00000658-200010000-00016.
29. Luna RA, Peixoto EM, Carvalho CFA, Velasque LS. Impact of body mass index on perioperative outcomes for complex hiatus hernia by videolaparoscopy. *Arq Bras Cir Dig.* 2022;35:e1672. doi: 10.1590/0102-672020220002e1672.
30. Memon MA, Siddaiah-Subramanya M, Yunus RM, Memon B, Khan S. Suture Cruroplasty Versus Mesh Hiatal Herniorrhaphy for Large Hiatal Hernias (HHs): An Updated Meta-Analysis and Systematic Review of Randomized Controlled Trials. *Surg Laparosc Endosc Percutan Tech.* 2019;29(4):221-232. doi: 10.1097/SLE.0000000000000655.
31. Morgenthal CB, Lin E, Shane MD, Hunter JG, Smith CD. Who will fail laparoscopic Nissen fundoplication? Preoperative prediction of long-term outcomes. *Surg Endosc.* 2007;21(11):1978-84. doi: 10.1007/s00464-007-9490-7.
32. Morrow EH, Chen J, Patel R, Bellows B, Nirula R, Glasgow R, et al. Watchful waiting versus elective repair for asymptomatic and minimally symptomatic paraesophageal hernias: A cost-effectiveness analysis. *Am J Surg.* 2018;216(4):760-763. doi: 10.1016/j.amjsurg.2018.07.037.
33. Nason KS, Luketich JD, Awais O, Abbas G, Pennathur A, Landreneau RJ, et al. Quality of life after collis gastroplasty for short esophagus in patients with paraesophageal hernia. *Ann Thorac Surg.* 2011;92(5):1854-60; discussion 1860-1. doi: 10.1016/j.athoracsur.2011.06.030.

34. Navaratne L, Ashrafian H, Martínez-Isla A. Quantifying tension in tension-free hiatal hernia repair: a new intra-operative technique. *Surg Endosc.* 2019;33(9):3040-3049. doi: 10.1007/s00464-019-06843-6.
35. Oelschlager BK, Pellegrini CA, Hunter J, Soper N, Brunt M, Sheppard B, et al. Biologic prosthesis reduces recurrence after laparoscopic paraesophageal hernia repair: a multicenter, prospective, randomized trial. *Ann Surg.* 2006;244(4):481-90. doi: 10.1097/01.sla.0000237759.42831.03.
36. Oelschlager BK, Pellegrini CA, Hunter JG, Brunt ML, Soper NJ, Sheppard BC, et al. Biologic prosthesis to prevent recurrence after laparoscopic paraesophageal hernia repair: long-term follow-up from a multicenter, prospective, randomized trial. *J Am Coll Surg.* 2011;213(4):461-8. doi: 10.1016/j.jamcollsurg.2011.05.017.
37. Oelschlager BK, Yamamoto K, Woltman T, Pellegrini C. Vagotomy during hiatal hernia repair: a benign esophageal lengthening procedure. *J Gastrointest Surg.* 2008;12(7):1155-62. doi: 10.1007/s11605-008-0520-0.
38. Parker DM, Rambhajan AA, Horsley RD, Johanson K, Gabrielsen JD, Petrick AT. Laparoscopic paraesophageal hernia repair is safe in elderly patients. *Surg Endosc.* 2017;31(3):1186-1191. doi: 10.1007/s00464-016-5089-1.
39. Poncet G, Robert M, Roman S, Boulez JC. Laparoscopic repair of large hiatal hernia without prosthetic reinforcement: late results and relevance of anterior gastropexy. *J Gastrointest Surg.* 2010;14(12):1910-6. doi: 10.1007/s11605-010-1308-6.
40. Saad AR, Velanovich V. Laparoscopic antireflux surgery: are old questions answered? Mesh hernioplasty. *Arq Bras Cir Dig.* 2023;35:e1710. doi: 10.1590/0102-672020220002e1710.
41. Soper NJ, Teitelbaum EN. Laparoscopic paraesophageal hernia repair: current controversies. *Surg Laparosc Endosc Percutan Tech.* 2013;23(5):442-5. doi: 10.1097/SLE.0b013e3182a12680.
42. Sowards KJ, Holton NF, Elliott EG, Hall J, Bajwa KS, Snyder BE, et al. Safety of robotic assisted laparoscopic recurrent paraesophageal hernia repair: insights from a large single institution experience. *Surg Endosc.* 2020;34(6):2560-2566. doi: 10.1007/s00464-019-07291-y.

43. Spann MD, Harrison NJ, English WJ, Bolduc AR, Aher CV, Williams DB, et al. Efficacy and Safety of Recurrent Paraesophageal Hernia Repair with Roux-en-Y Gastric Bypass. *Am Surg*. 2020;86(3):250-255. PMID: 32223806.
44. Stadlhuber RJ, Sherif AE, Mittal SK, Fitzgibbons RJ Jr, Michael Brunt L, Hunter JG, et al. Mesh complications after prosthetic reinforcement of hiatal closure: a 28-case series. *Surg Endosc*. 2009;23(6):1219-26. doi: 10.1007/s00464-008-0205-5.
45. Stylopoulos N, Gazelle GS, Rattner DW. Paraesophageal hernias: operation or observation? *Ann Surg*. 2002;236(4):492-500; discussion 500-1. doi: 10.1097/00000658-200210000-00012.
46. Tam V, Luketich JD, Winger DG, Sarkaria IS, Levy RM, Christie NA, et al. Non- Elective Paraesophageal Hernia Repair Portends Worse Outcomes in Comparable Patients: a Propensity-Adjusted Analysis. *J Gastrointest Surg*. 2017;21(1):137-145. doi: 10.1007/s11605-016-3231-y.
47. Tatum JM, Samakar K, Bowdish ME, Mack WJ, Bildzukewicz N, Lipham JC. Videoesophagography *versus* Endoscopy for Prediction of Intraoperative Hiatal Hernia Size. *Am Surg*. 2018;84(3):387-391. PMID: 29559053.
48. Terry ML, Vernon A, Hunter JG. Stapled-wedge Collis gastroplasty for the shortened esophagus. *Am J Surg*. 2004;188(2):195-9. doi: 10.1016/j.amjsurg.2003.12.069.
49. Vasudevan V, Reusche R, Nelson E, Kaza S. Robotic paraesophageal hernia repair: a single-center experience and systematic review. *J Robot Surg*. 2018;12(1):81-86. doi: 10.1007/s11701-017-0697-x.
50. Ward MA, Hasan SS, Sanchez CE, Whitfield EP, Ogola GO, Leeds SG. Complications Following Robotic Hiatal Hernia Repair Are Higher Compared to Laparoscopy. *J Gastrointest Surg*. 2021;25(12):3049-3055. doi: 10.1007/s11605-021-05005-1.
51. Watson DI, Davies N, Devitt PG, Jamieson GG. Importance of dissection of the hernial sac in laparoscopic surgery for large hiatal hernias. *Arch Surg*. 1999;134(10):1069-73. doi: 10.1001/archsurg.134.10.1069.
52. Wirsching A, Zhang Q, McCormick SE, Hubka M, Low DE. Abnormal High-Resolution Manometry Findings and Outcomes after Paraesophageal Hernia

Repair. J Am Coll Surg. 2018;227(2):181-188.e2. doi:  
10.1016/j.jamcollsurg.2018.03.033.

This preprint was submitted under the following conditions:

- The authors declare that they are aware that they are solely responsible for the content of the preprint and that the deposit in SciELO Preprints does not mean any commitment on the part of SciELO, except its preservation and dissemination.
- The authors declare that the necessary Terms of Free and Informed Consent of participants or patients in the research were obtained and are described in the manuscript, when applicable.
- The authors declare that the preparation of the manuscript followed the ethical norms of scientific communication.
- The authors declare that the data, applications, and other content underlying the manuscript are referenced.
- The deposited manuscript is in PDF format.
- The authors declare that the research that originated the manuscript followed good ethical practices and that the necessary approvals from research ethics committees, when applicable, are described in the manuscript.
- The authors declare that once a manuscript is posted on the SciELO Preprints server, it can only be taken down on request to the SciELO Preprints server Editorial Secretariat, who will post a retraction notice in its place.
- The authors agree that the approved manuscript will be made available under a [Creative Commons CC-BY](#) license.
- The submitting author declares that the contributions of all authors and conflict of interest statement are included explicitly and in specific sections of the manuscript.
- The authors declare that the manuscript was not deposited and/or previously made available on another preprint server or published by a journal.
- If the manuscript is being reviewed or being prepared for publishing but not yet published by a journal, the authors declare that they have received authorization from the journal to make this deposit.
- The submitting author declares that all authors of the manuscript agree with the submission to SciELO Preprints.