



Effect of Gamma Irradiation on the Pasting, Functional and Microbial Properties of Flours from white Yam (*Dioscorea rotundata*) and Water Yam (*Diocorea alata*)

Ugbedejo, A. E., Momoh, C., Dickson, G., & Idoko, F. A. (2020). Effect of Gamma Irradiation on the Pasting, Functional and Microbial Properties of Flours from white Yam (*Dioscorea rotundata*) and Water Yam (*Diocorea alata*). *Global Journal of Medical Research*, 20(L2), 1-6. Advance online publication. https://medicalresearchjournal.org/index.php/GJMR/article/view/2330/1-Effect-of-Gamma-Irradiation_JATS_NLM_xml

[Link to publication record in Ulster University Research Portal](#)

Publication Status:

Published online: 15/01/2020

Document Version

Publisher's PDF, also known as Version of record

General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.



GLOBAL JOURNAL OF MEDICAL RESEARCH: L
NUTRITION & FOOD SCIENCE
Volume 20 Issue 2 Version 1.0 Year 2020
Type: Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals
Online ISSN: 2249-4618 & Print ISSN: 0975-5888

Effect of Gamma Irradiation on the Pasting, Functional and Microbial Properties of Flours from white Yam (*Dioscorea rotundata*) and Water Yam (*Diocorea alata*)

By Ameh Elijah Ugbedeajo, Momoh Clement Owoicho, Gerna Dickson & Felicia Adizaldoko
University of Agriculture

Abstract- In this study, flours prepared from white yam and water yam tubers were treated using gamma-irradiation at 0, 0.15 kGy and 2 kGy dosages, giving rise to four samples. The samples were analysed for functional, pasting and microbiological properties using standard analytical methods. The functional properties of the flour samples showed significant differences ($p < 0.05$) for irradiated samples. The pasting properties of the irradiated water yam and white yam flour sample ranged from 206.11 ± 1.99 RVU to 162.44 ± 2.12 RVU for doses of 0.15 kGy and 2 kGy (peak viscosity), trough ranged from 108.54 ± 1.98 to 90.68 ± 1.56 RVU, breakdown ranged 102.18 ± 1.00 to 62.67 ± 1.87 RVU, while the pasting temperature ranged from 83.10 ± 1.10 to 77.29 RVU.

Keywords: gamma, irradiation and yam.

GJMR-L Classification: NLMC Code: WB 400



Strictly as per the compliance and regulations of:



© 2020. Ameh Elijah Ugbedeajo, Momoh Clement Owoicho, Gerna Dickson & Felicia Adizaldoko. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Effect of Gamma Irradiation on the Pasting, Functional and Microbial Properties of Flours from white Yam (*Dioscorea rotundata*) and Water Yam (*Diocoreaalata*)

Ameh Elijah Ugbedeajo ^α, Momoh Clement Owoicho ^σ, Gerna Dickson ^ρ & Felicia Adizaldoko ^ω

Abstract- In this study, flours prepared from white yam and water yam tubers were treated using gamma-irradiation at 0, 0.15 kGy and 2 kGy dosages, giving rise to four samples. The samples were analysed for functional, pasting and microbiological properties using standard analytical methods. The functional properties of the flour samples showed significant differences ($p < 0.05$) for irradiated samples. The pasting properties of the irradiated water yam and white yam flour sample ranged from 206.11 ± 1.99 RVU to 162.44 ± 2.12 RVU for doses of 0.15 kGy and 2 kGy (peak viscosity), trough ranged from 108.54 ± 1.98 to 90.68 ± 1.56 RVU, breakdown ranged 102.18 ± 1.00 to 62.67 ± 1.87 RVU, while the pasting temperature ranged from 83.10 ± 1.10 to 77.29 RVU. The microbial content of the white yam and water yam flours decreased with increase in irradiation does ranging from 1.46×10^2 cfu/g to 1.00×10^2 cfu/g for the white yam flour and 1.39×10^2 cfu/g to 1.10×10^2 cfu/g for the water yam flour, for the aerobic plate count with the un-irradiated sample value of 240×10^3 cfu/g and 2.10×10^3 cfu/g for white yam and water yam flours were obtained. The fungal counts for un-irradiated samples were 1.32×10^2 cfu/g and 1.20×10^2 cfu/g for the white yam and water yam flour respectively. Microbial content of the un-irradiated flours differed significantly ($p < 0.05$) from irradiated flours.

Keywords: gamma, irradiation and yam.

I. INTRODUCTION

Yam is the second most important tuber crop in Africa next to cassava. Nigeria is the main producer of yam globally with 71% of world production (FAO, 2009). Yams are annual or perennial tuber-bearing and climbing plants with over 600 species in which only few are cultivated for food and medicine (IITA, 2006). The most cultivated species in Nigeria are the white yam (*Dioscorea. rotundata*), yellow yam (*Dioscoreacayenensis*), water yam (*Dioscorea. alata*) and trifoliate yam (*Dioscorea. dumetorum*) (Amusa *et al.*, 2003). The crop is of major importance in the diet and economic life of people in West Africa (Giranidin, *et al.*, 1998). Yam is an elite crop, preferred over other root and tuber crops of West Africa and a choice during

ceremonies and festivities. *Dioscorea. alata* is also referred to as greater yam. It is more important as food in West Africa and the Caribbean than in Asia and the Americas where it originated, and has been competing with the most important native species, *Dioscorea rotundata*. *Dioscoreaalata* is also known for its high nutritional content (Osagie, 1992). *Dioscoreaalata* tubers have variable shapes, the majority being cylindrical. The flesh of the tuber ranges in colour from white to purplish (FAO, 2009). The texture of its flesh is usually not as firm as that of white yam and less suitable compared to other species for the preparation of the most popular food products from yam (fufu and pounded yam especially) in the West Africa region.

The production of the yam is seasonal, so storage is necessary before subsequent planting or for use as food. Fresh yam tubers are often difficult to store and are subject to deterioration by sprouting and microbial rot during storage. Post-harvest losses usually range from 25 to 60% (Afoakwa and Sefa-dede, 2001).

Loss of yam in storage is very high. Several inhibitory chemical growth regulators such as maleic hydrazide, Tetrachloronitro benzene, acetic acid, naphthalene have been used to retard sprouting in stored yam tubers. Apart from unavailability of the right type of chemicals and their toxicity nature, widespread adulteration of the available ones especially in Nigeria is a serious problem (David, 2009). Processing of yams into various products including flour is a viable alternative for checking postharvest losses and promotes year round availability. Radiation technology can complement existing technologies to ensure food security and safety (ICGFI, 1999).

Radiation processing could be used for anti-infestation and spoilage of food grains, flours and pulses; inhibition of sprouting in onions, potatoes, garlic, yam and ginger, preventing microbial contamination of spices; extending shelf-life under recommended conditions of storage of flours; and overcoming quarantine barriers in international trade. Ionizing radiations have the potentials of reducing considerable storage losses through inhibition of sprouting, inactivation of food spoilage micro-organism, control of

Author α σ ρ : Department of Food Science & Technology, University of Agriculture, Makurdi, Benue State, Nigeria.

Author ω : Department of Food Science & Technology Kaduna Polytechnic, Kaduna. e-mails: momohclement86@gmail.com, amehelijah@gmail.com

insects and sterilization of food crops. Irradiation technology is easy to apply, clean, and environment friendly. It is a direct, simple and efficient on-line process (Bansa and Appiah, 1999).

Unlike most previous studies which have focused on other species or cultivars of root and tuber crops, this study investigated the potential of applying gamma irradiation for improving the quality of two yam tuber varieties in Makurdi, Benue State. It was carried out to determine the effect of gamma irradiation on the physico-chemical and microbial properties of flours from *Dioscorea rotundata* and *Dioscorea alata*

II. MATERIALS AND METHODS


a) Procurement of Materials

White yam (*Dioscorea rotundata*), and Water yam (*Dioscorea alata*), were purchased from Modern market in Makurdi, Benue State, Nigeria.

b) Sample Preparation

Fresh White yam (*Dioscorea rotundata*), and water yam (*Dioscorea alata*) were properly cleaned and sorted to remove dirt, and other extraneous matters, before they were processed into flour. The method used was a modification of the method described by Enwere, (1998). The yam tuber was washed, peeled, sliced, heated and dried, which was then milled into flour. The flow diagram for the production of yam flours is shown in Figure 1.

The yam tubers were processed into flour and then subjected to gamma irradiation doses, with one un-irradiated batch to serve as control. Irradiation of samples were done using ⁶⁰Co source. Target doses were 150 kGy and 2 kGy



Source: Modified Enwere (1998)

Figure 1: Yam flour processing flow diagram

III. ANALYTICAL METHODS

Pasting properties, functional properties and microbiological properties were done using standard methods as described by AOAC, 2012.

a) Statistical Analysis

Statistical Package for Social Sciences (SPSS) V23 computer software was used to analyze the data. Means and Standard deviation were calculated where appropriate. One way Analysis of variance (ANOVA) was used to determine the treatment that was different from others in the various parameters tested; differences were considered significant at 95% ($p < 0.05$) significant level and 99% ($p < 0.01$) significant level where mentioned.

IV. RESULT AND DISCUSSION

a) Effect of Gamma Irradiation on the Pasting Properties of White Yam Flour and Water Yam Flour

Peak viscosity and trough viscosity of the two varieties of yam flours were relatively low. The values of the peak viscosity were 162.44 RVU for the white yam (*D. rotundata*) flour at irradiation dose of 2 KGy and 198.88 RVU for the control sample. Peak viscosity is the maximum viscosity attained by the paste during the heating cycle (i.e from 50 to 95°C) due to starch granules swelling and leaching out of the soluble components into the solution. It reflects the ability of starch granules to swell freely before their physical breakdown (Sing *et al.*, 2003) and often correlates with product quality. Peak viscosity had values of 198.33 and 225.56 RVU for white yam flour and 162.44 and 198.88 RVU for water yam flour. The relatively longer time taken for most (*D. alata*) flour to paste further indicates stronger bonding forces in their starch granules. Yam flours generally have some level of resistance to swelling which in this case is more pronounced in (*D. alata*). Richard *et al.* (1991) reported that cassava starch has a high peak viscosity because it exhibits a high degree of swelling. The low value of the peak viscosity in water yam (*D. alata*) and white yam (*D. rotundata*) flour indicates strong internal forces between their starch granules which resulted in lower swelling power pasting viscosity. High amylose content has been linked to low swelling power due to greater reinforcement of the internal network by amylose molecules (Hoover, 2001). Highly associated starch granules with an extensive and strongly bonded structure also exhibit resistance towards swelling. According to Richard *et al.*, (1991) pasting viscosity and swelling are positively related. The higher the swelling power of a flour sample, the higher the pasting viscosity.

Kulkanir (1996), observed that (*D. alata*) flour has relatively low viscosity but high gel strength when compared with *D. rotundata* which was in agreement with the present study under irradiation. Therefore, *D.*

alata flour may be more suitable for products which require high gel strength and a low viscosity or they could be parboiled and processed to products such as amala. Flour with a higher peak viscosity is also required for making food products such as jelly or binder while those with lower viscosity are desirable for preparing weaning foods and lighter gruels (Kulkani *et al.*, 1996).

Trough is considered as a measure of the breakdown of hot paste. The ability of a paste to withstand the heating and shear stress is an important factor for most food processing operations and is also a factor in describing the quality of starch gels (Madens and Christensen, 1996). High paste stability is a requirement for industrial users of starch (Bainbridge *et al.*, 1996). This is because drastic changes in paste during and after processing could lead to textural changes that may be undesirable. Trough values obtained were 96.38 RVU and 140.40 RVU for white yam (*D. rotundata*) flour also 90.68RVU and 146.21RVU for water yam (*D. alata*) flour. The *D. alata* flour had the highest trough and lowest breakdown values, which indicates greater ability to withstand shear at high temperatures and higher cooked paste stability (Farhat *et al.*, 1999). Flour with low trough values would have greater need for cross-linking than one with high value, *D. alata* flour could therefore be targeted for industrial uses because of its hot paste stability.

The breakdown viscosity increased with the increasing radiation dose in the white yam flour (*D. rotundata*) and water yam (*D. alata*) flour. The values ranged from 95.82 ± 0.70 to 103.68 ± 1.32 RVU for the white yam flour, while the water yam flour ranged from 89.92 ± 0.50 to 109.89 ± 0.02 RVU. Mohd *et al.*, (2009) observed significant changes in breakdown viscosity by the radiation dose indicates considerable disruption or weakening of the bonding forces (hydrogen bonds) in the starch granules by the radiation. The lower breakdown of viscosity indicates good stability of the starch granules, Celik *et al.* (2004) observed a reduction in the cooking period of irradiated legumes.

The viscosity after cooling cooked paste to 50 °C is the final viscosity. Anonymous (1990) reported that starch paste increases in viscosity when cooled. The increase in viscosity is not only caused by simple kinetic effect of cooling but also by re-association of molecules (particularly amylose). Final viscosity is the most commonly used parameter to determine the quality of starch-based samples because it indicates starch/flour ability to form a gel after cooking.

Yam flour is thixotropic and, as observed in this study, has a higher cooled paste viscosity than hot paste viscosity. The observation supports the general fact that yam pastes form firm gels rather than viscous gels after cooking and cooling. This has implications for the kind of products yam flour could be used for, such as weaning diets or crackers. Yam flour is noted for high retro gradation during cooling and this might have

accounted for the increase in final viscosities during cooling. This is brought about by the high degree of association between the starch-water systems and their high ability to re-crystallize, resulting in progressively higher viscosities during cooling of yam starches (Anonymous, 1990). Ayernor (2012) reported that the rate at which the development of rigidity occurs in yam flour is dependent on the degree of starch-water binding which can be affected through processes that influence the interaction between the flour particles and water.

Setback shows the cold paste viscosity's tendency to retro- grade. Therefore, the decreased viscosity of the irradiated samples reveals the lack of ability of amylopectin to hold the granule during the imbibition of water (Mohd *et al.*, 2009). Therefore, from the values obtained for the white and water yam flour decreasing as the irradiation dose increase show lack of ability of amylopectin to hold granules during imbibition of water.

Setback viscosity is an important factor for starch used as a food ingredient in processing and preservation, because the quality of the food's texture and physical properties deteriorate due to retro gradation as time passes (Nunoo, 2009). A decrease in setback viscosity was observed with increase in the irradiation dose. The lower setback viscosity of the paste indicated that the reassociation of the starch polymers is not as great as in the control. Pomeranz (1991) established that high setback (retrogradation) will cause undesirable gel texture or high freeze-thaw stability. This means the protein and starch in the flour obtained from the two irradiated yam flour would be unable to form strong gel matrices or networks, perhaps, due to poor ionic interactions between the individual molecules. Gamma irradiation is capable of hydrolyzing chemical bonds, there by cleaving large molecules of starch into smaller fragments of dextrin that maybe either electrically charged or uncharged as free radicals (Wu *et al.*, 2002). These changes may affect the physical and rheological properties of irradiated starches, resulting in changes in properties of starch paste (Mohd *et al.*, 2009). Decreases in pasting properties, such as breakdown and setback values, may result primarily from irradiation-induced starch degradation and may present opportunities such as ease of cooking and reduced starch retrogradation, respectively (Sabularse *et al.*, 1991).

There was a reduction in the peak temperature among the two varieties of yam flour, although the reduction was not dose dependent as there were little variation in the values.

The peak temperature for the white yam (*D. rotundata*) flour with irradiation doses of 0.15 kGy and 2 kGy had a reduced value whereas in the water yam (*D. alata*) flour, the minimum peak temperature was observed in the 2 KGy irradiated samples. It can be deduced that irradiation caused the reduction. The

reduction will result in low temperature and time of cooking of the flours from irradiated yam flour and temperature may be prepared when all other properties are equal.

Decreases in pasting properties, such as breakdown and setback values, may result primarily from irradiation-induced starch degradation and may present opportunities such as ease of cooking and reduced starch retro-gradation, respectively (Sabularse *et al.*, 1991).

b) Effect of Gamma Irradiation on the Functional Properties of White Yam flour and Water Yam Flour

In this study, it was observed that water absorption capacity (WAC) and oil absorption capacity (OAC) of the two varieties of yam flours were significantly ($p < 0.05$) influenced by the irradiation.

Significant increase in water absorption capacity in the irradiated white yam (*D. alata*) and water yam (*D. rotundata*) was observed, the control of white yam flour had the lowest value for the oil absorption capacity and the irradiated water yam flour at 2 kGy had the highest value of water absorption capacity 1.93ml/g. Abu *et al.*, (2006a,b) observed similar trends. This suggests that the irradiation increases water affinity of resulting flours. The increase in the WAC may in part be due to irradiation-induced damage or degradation of cowpea starch to simpler molecules such as dextrans and sugar that have higher affinity for water than starch (Abu *et al.*, 2006a,b). High WAC of flour is advantageous in the preparation of food items like bread and sausages to maintain freshness and for easy handling. Oshodi and Adeladun (1993) reported that high WAC of seed flour is useful in the preparation of soups and gravies. The phenomenon could be attributed to a degree of cross-linking, which may have occurred simultaneously with chain scission at the doses (1.0 and 1.5 KGy) hence the observed results. Similar results were reported by Abu *et al.*, (2006b).

OAC is an important functional trait in food industries for retention of flavor and improvement of shelf-life and palatability (Bhat *et al.*, 2008). The OAC of the flour obtained from the irradiated samples of yam flours of the white yam (*D. rotundata*) and water yam (*D. alata*) were higher than the control samples. This was in agreement as reported by Abu *et al.* (2006a,b) that OAC of cowpea irradiated increased as doses increased. The increase in the OAC may be related to the radiation. The mechanism of the increase in the OAC upon irradiation involves the increased ability of degraded and/or cross-linked starch to physically entrap more oil and the high affinity of non-polar protein side chains for lipids (Salbe *et al.*, 1982). As reported by Urbein, (1986) increase in OAC of seed flour has been attributed to the unmasking of non-polar protein residues as a result of irradiation-induced denaturation. White yam (*D. rotundata*) and

water yam (*D. alata*) flour had relatively significant lowest OAC at 0.15 kGy with 1.11 ± 0.01 ml/g.

Water solubility index of the irradiated samples was significantly lower than the control across the two varieties of yam flour. This is an indication that the flours obtained from irradiated yam flour samples were less soluble in water to an extent. The irradiation may have caused the soluble substances in the flour to idolize making their binding sites for water inaccessible. Medoua *et al.*, (2005) reported that the decrease in water solubility could be due to the mobilization of soluble substances. The water solubility of the yam flours for both white yam (*D. rotundata*) and water yam (*D. alata*) decreased significantly with the radiation doses but was not dose dependent. The values for control for both yam flours ranged from 8.58 ± 0.02 ml/g and 6.15 ± 0.03 ml/g as compared to the irradiated samples with values of 7.5604 ± 1.04 ml/g for white yam flour and 5.13 ± 1.31 ml/g for water yam flour. This result was similar to those observed by Azimet *et al.* (2009). Abu *et al.* (2006a,b), however, observed reduction in WSI with increasing radiation dose.

Bulk density of the white yam (*D. rotundata*) and water yam (*D. alata*) flour of irradiated or non-irradiated were not significantly different ($p < 0.05$) from each other, even at 2 kGy the bulk density was not significantly different, which may not be attributed to the irradiation. This means that the bulk density of the yam flour was not significantly ($p < 0.05$) affected by the irradiation. Abu *et al.*, 2006 a, b. Azim *et al.* (2009) reported that the bulk density of groundnut flour was not affected significantly by gamma irradiation.

Reconstitution index of the yam flour samples were not significantly ($p < 0.05$) different from the irradiated and non-irradiated from each other. The control values for the white yam (*D. alata*) and water yam (*D. rotundata*) flour ranged between 6.76 ± 0.02 ml/g and 6.09 ± 0.02 ml/g as compared with the irradiated samples with values of 6.40 ± 0.01 ml/g and 5.58 ± 0.01 ml/g. though statistical differences were observed in the values obtained for reconstitution index in the white yam flour, their functionality in sample may not differ.

c) Effect of Gamma Irradiation on the Microbial Content of White Yam Flour and Water Yam Flour

The microbial analysis for the white yam sample for aerobic plate count for the control was 1.93×10^3 cfu/ml and that for water yam sample was 2.4×10^3 . The bacteria present in the samples before irradiation would have been picked up during processing (Jay *et al.*, 2005), but the low moisture content (due to the flour nature) might have prevented the growth of some bacteria even before irradiation and when there is low moisture in stored products, *Bacillus* species are usually the one ones that grow (Jay *et al.*

2005); this might have accounted for the low presence of *Bacillus* species in the flour samples after irradiation. Most of the bacteria isolated from the flour samples (white yam and water yam) before irradiation have been obtained by previous workers (Ojokoh and Gabriel, 2010). Report of occurrence of many *Lactobacillus* and *Bacillus* species were found to be dominant during the production of a fermented cassava product (Achi and Akomas, 2006). The same situation was obtained for the fungi whereby the samples contained less fungi qualitative after irradiation than before irradiation. *Aspergillus niger* which was the most predominant fungus in the samples before irradiation. This should be desirable because *A. niger* is one of the aflatoxin-producing moulds (Frazier and Westhoff, 2004) whose presence in foods will constitute health hazard to consumers. *A. flavus* and *A. niger* which are sporadic (Adegoke, 2004). Toxicogenic strains of *A. flavus* have been known to produce aflatoxin, a potent hepatotoxic and carcinogenic agent (Uraih and Ogbadu, 2019).

V. CONCLUSION

The pasting temperature was not significantly affected by gamma irradiation which indicates the ability to withstand shear stress and high temperature. Gamma irradiation reduced significantly bacteria and fungi count in the two varieties of yam flour, thereby lowering the activity of micro-organism. The use of gamma irradiation in the treatment of water yam flour and white yam flour showed that irradiation can preserve and maintain the quality of yam flour

REFERENCES RÉFÉRENCES REFERENCIAS

1. Abu, J. O., Duodu, K. G., Minnaar, A. (2006a). Effects of gamma irradiation on some physico-chemical and thermal properties of cowpea (*Vigna unguiculata* L. Walp) starch. *Food Chem.* 95, 386–393.
2. Abu, J. O., Klaus.M., Kwaku, G.D, Duodu, G. and Amanda, M. (2005) Functional properties of cowpea (*Vigna unguiculata* L. Walp) flours and pastes as affected by γ -irradiation. *Food chemistry*, 93: 103-111.
3. Abu, J. O., Muller, K., Duodu, K. G., Minnaar, A. (2006b). Gamma irradiation of cowpea (*Vigna unguiculata* L. Walp) flours and pastes: effects on functional, thermal and molecular properties of isolated proteins. *Food Chem.* 95, 138–147.
4. Adegoke, G.O., 2004. Understanding Food Microbiology. 2nd Edn., Alleluia Ventures Ltd., Ibadan, Nigeria
5. Afoakwa E. O. and Sefa-Dedeh, S. (2001). Chemical composition and quality changes occurring in *Dioscorea rotundata* tubers after harvest. *Food Chem.* 75: 85-91.
6. Amusa, N. A., Adegite, A. A., Mohammed, S and Baiyewu, R. A. (2003). Yam disease and its management in Nigeria. *African Journal of Biotechnology* 2(12):497 – 502.
7. Ayernor, G.S. (2012). The Yam (*Dioscorea*) Starches. In: *Advances in Yam research*. 1st ed. Godson Osuji. Anambra State University of Technology Publishers. Nigeria., pp.79-88.
8. Bainbridge, Z., K. Tomlins, K. Wellings, and A. Westby. 1996. Methods for assessing quality characteristics of non-grains starch (Part 3. Laboratory methods). *Natural Resources Institute*, Chalthom, U.K.
9. Bansa, D and V. Appiah (1999). The effect of irradiation on the storage yam in Ghana. *Journal the Ghana Science Association*. Vol. 1 (3) pp. 100-103.
10. Bhat, R., Sridhar, R. K., Young, C. C., Bhagwath, A. A. and Ganesh, S. (2008). Composition and functional properties of raw and electron beam-irradiated *Mucuna pruriens* seeds. *Int. J. Food Sci. Technol.* 43, 1338–1351.
11. Celik, S., Yalcin, E., Basman, A., and Koksel, H. (2004). Effects of irradiation on protein electrophoretic properties, water absorption and cooking quality of lentils. *Int. J. Food Sci. Nutr.* 55, 641–648
12. David, M (2009). Nigeria: Food and Agricultural Import and Standards. USAID Foreign Agricultural Services Gain Report. Global Agricultural Information Network. Pp 1-9.
13. Enwere, N. J. (1998). *Food of Plant Origin*. Afro-Orbis Publication Limited, Nsukka.
14. F. A. O. (2009), "Food Loss Prevention in Perishable Crops", *Corporate Document Repository*, pp. 220-231.
15. Farhat I. A., Oguntona T., and Neale R. J. (1999). Characterization of starches from West African yams. *J. Sci. Food Agric.* 79:2105–2112.
16. Frazier, W. C. and Westhoff, D. C. (2004). *Food Microbiology* 4th edition. Tata Mc Graw-Hill Publishing Company Ltd., New Delhi.
17. Girardin, O., Nindjin, C., Farah, Z., Escher, F., Stamp, P., Otokore, D. (1998). Use of gibberellic acid to prolong dormancy and reduce losses during traditional storage of yams. *Journal of the Science of Food and Agriculture*. Vol. 77(2).
18. Hoover, R. (2001) composition, molecular structure and physiochemical properties of 165 tuber and root starches. *Carbohydrate Polymers* 45:253-847.
19. International Consultative Group on Food Irradiation (ICGFI). Facts about food irradiation. 1999. [Cited 10 March 2008] Available from: <http://www.cdc.gov/ncidod/dbmd/diseaseinfo/foodirradiation.ht>
20. International Institute of Tropical Agriculture (IITA) (1993). Crop Improvement Division/Tuber root improvement program archival reports (1989–

- 1993). Part III yam (*Dioscorea* spp.) Ibadan, Nigeria pp. 20 – 85.
21. Jay, J. M., Leossner, M. J. and Golden, D. A. (2005). *Modern Food Microbiology*. 7th edition. Springer Science and Business Inc. London. pp. 45-54.
22. Kulkarni J, de Castella A, Smith D, Taffe J, Keks N, Copolov D (1996) A clinical trial of the effects of estrogen in acutely psychotic women. *Schizophr Res* 20:247–252.
23. Madsen MH, Christensen DH (1996) Changes in viscosity properties of potato starch during growth. *Starch/Starke* 48:245–249.
24. Medoua, G. N., Mbome, I. L., Agbor-Egbe, T. and Mbofung, C. M. F. (2005). Physicochemical changes occurring during post-harvest hardening of trifoliate yam (*Dioscorea dumetorum*) tubers. *Food Chem.* 90, 597–601.
25. Medoua, G. N., Mbome, I. L., Agbor-Egbe, T. and Mbofung, C. M. F. (2005). Physicochemical changes occurring during post-harvest hardening of trifoliate yam (*Dioscorea dumetorum*) tubers. *Food Chem.* 90, 597–601.
26. Mohd, A. N., Mat, H. D., Muhammad, K., Rahman, A. R., Ghazali, Z. and Hashim, K., (2009). Pasting and leaching properties of irradiated starches from various botanical sources. *Int. Food Res. J.* 16, 415–429.
27. Nunoo, A. P. (2009). Pasting of G-irradiated proteins from *Vignasubterranea* in native starch models and the surface functional properties of the proteins. M. Sc. Thesis. Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.
28. Osagie, A. U. (1992). The yam tuber in storage. Postharvest research Unit, Department of Biochemistry, University of Benin, Benin-City, Nigeria. Pp 146 –149.
29. Oshodi, A. A. and Adeladun, A. O. (1993). Proximate composition, some valuable minerals Functional properties of three varieties of Lima bean flour. *Inter Jour. Food Sci and Nutri.* 43: 175-181
30. Pomeranz Y. *Functional properties of food components*. 2nd edition, Academic Press. Inc. New York. 1991; pp. 24-78.
31. Richard J. R, Asauka M, A, Blanshard J. N. V (1991). The Physiochemical properties of cassava starch. *Trop. Sci.* 31:189-207
32. Sabularse VC, Liuzzo JA, Rao RM, Grodner RM (1991) Cooking quality of brown rice as influenced by gamma irradiation, variety and storage. *J Food Sci* 56: 96–98, 108.
33. Salbe, G. O., Taylor, J. R. N., and Minnaar, A. (2001) Effect of irradiation, with and without cooking of maize and kidney bean flours, on porridge viscosity and in vitro starch digestibility. *Journal of the Science of Food and Agriculture*, 81: 497–502.
34. Uraih, W and Ogbadu, G (1980): Incidence of Aflatoxin in Nigerian Sorghum. *Microbios*.10:
35. Urbain, W. M. (1986) *Food irradiation*. (pp. 37–81) New York: Academic Press.
36. Wu, D., Shu, Q., Wang, Z. and Xia, Y. 2002. Effect of gamma irradiation on starch viscosity and physicochemical properties of different rice. *Radiation Physics and Chemistry* 65(1): 79-86.