



Examining the inter-relationships between social isolation and loneliness and their correlates among older British adults before and during the COVID-19 lockdown: Evidence from four British longitudinal studies

Mansfield, R., Di Gessa, G., Patel, K., McElroy, E., Wels, J., Henderson, M., Maddock, J., Stafford, J., Steptoe, A., Richards, M., & Patalay, P. (2023). Examining the inter-relationships between social isolation and loneliness and their correlates among older British adults before and during the COVID-19 lockdown: Evidence from four British longitudinal studies. *Innovation in Aging*. Advance online publication. <https://doi.org/10.1093/geroni/igad126>

[Link to publication record in Ulster University Research Portal](#)

Published in:
Innovation in Aging

Publication Status:
Published online: 07/11/2023

DOI:
[10.1093/geroni/igad126](https://doi.org/10.1093/geroni/igad126)

Document Version
Author Accepted version

General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

Examining the inter-relationships between social isolation and loneliness and their correlates among older British adults before and during the COVID-19 lockdown: Evidence from four British longitudinal studies

Rosie Mansfield PhD^{1*+}, Giorgio Di Gessa PhD²⁺, Kishan Patel PhD³⁺, Eoin McElroy PhD⁴⁺, Jaques Wels PhD^{3,5+}, Morag Henderson PhD¹, Jane Maddock PhD³, Jean Stafford PhD³, Andrew Steptoe PhD⁶, Marcus Richards PhD³, and Praveetha Patalay PhD^{1,3}

¹ Centre for Longitudinal Studies, University College London, London, UK

² Department of Epidemiology and Public Health, University College London, London, UK

³ MRC Unit for Lifelong Health and Ageing, University College London, London, UK

⁴ School of Psychology, Ulster University, Coleraine, Northern Ireland, UK

⁵ Centre Metices, Université libre de Bruxelles, Brussels, Belgium

⁶ Department of Behavioural Science and Health, University College London, London, UK

+ R. Mansfield, G. Di Gessa, K. Patel, E. McElroy, and J. Wels served as joint first authors for this study.

*Address correspondence to: Rosie Mansfield. Email: r.mansfield@ucl.ac.uk

© The Author(s) 2023. Published by Oxford University Press on behalf of The Gerontological Society of America.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background and Objectives: Unprecedented social restrictions during the COVID-19 pandemic have provided a new lens for considering the inter-relationship between social isolation and loneliness in later life. We present these inter-relationships before and during the COVID-19 restrictions and investigate to what extent demographic, socio-economic, and health factors associated with such experiences differed during the pandemic.

Research Design and Method: We used data from four British longitudinal population-based studies (1946 NSHD, 1958 NCDS, 1970 BCS, and ELSA, N=12,129). Rates, co-occurrences, and correlates of social isolation and loneliness are presented prior to and during the early stage of the COVID-19 pandemic and the inter-relationships between these experiences are elucidated in both periods.

Results: Across the four studies, pre-pandemic proportions reporting social isolation ranged from 15 to 54%, with higher rates in older ages (e.g., 32% of 70-79 and 54% of those over 80). During the pandemic, the percentage of older people reporting both social isolation and loneliness and isolation only slightly increased. The inter-relationship between social isolation and loneliness did not change. Associations between socio-demographic and health characteristics and social isolation and loneliness also remained consistent, with greater burden among those with higher economic precarity (females, non-homeowners, unemployed, illness and greater financial stress).

Discussion and Implications: There were already large inequalities in experiences of social isolation and loneliness and the pandemic had a small impact on worsening extent and inequalities in these. The concepts of loneliness and social isolation are not interchangeable, and clarity is needed in how they are conceptualised, operationalised, and interpreted. Given many older adults experience high levels of social isolation, there should be greater emphasis on reducing social isolation and the inequalities observed in who experiences greater isolation and loneliness.

Keywords: Conceptualisation, Overlap, Networks, Inequalities

Translational Significance

There is a conflation of “loneliness” and “social isolation” in research and policy, and over-emphasis on loneliness in UK policy (e.g., loneliness ministers). Being female, economic precarity, and long-standing illness consistently correlated with isolation and loneliness; structural changes and policies designed to reduce these inequalities are needed. Given many older adults experience high levels of social isolation, there should be greater emphasis on reducing social isolation. The increases in social isolation observed in the pandemic highlight the need for efforts to support older people to (re)start hobbies, volunteer, and socialise to reduce isolation and its consequences for health and wellbeing.

Accepted Manuscript

Background and Objectives

Prior to the COVID-19 pandemic, loneliness was identified as a significant public health concern (Jeste et al., 2019): for example, in the UK, a Minister for Loneliness was appointed in 2018 and the 'Campaign to End Loneliness' was launched. Despite increased policy interest, there remains a need to better understand the conceptualisation and measurement of social isolation and loneliness as they are often inconsistently applied and interchangeably referred to across research, policy, and practice (Wigfield et al., 2022). With the implementation of social distancing and quarantine measures due to the COVID-19 pandemic and several national lockdowns across Britain, social isolation – and its relationship with loneliness – were brought into even sharper focus. By carefully considering the operationalisation of these complex concepts, this study has the dual aim to 1) explore the conceptual and empirical inter-relationships between social isolation and loneliness, and 2) identify overlapping and independent correlates of social isolation and loneliness prior to and during the COVID-19 pandemic. Using data from four British longitudinal studies, we investigate associations between social isolation and loneliness under normal circumstances and during COVID-19 restrictions and examine whether the factors associated with such experiences differ due to the pandemic.

As an objective condition, social isolation can be quantified by a person's network size, composition, and frequency of contact (Holt-Lunstad et al., 2015; Huisman & van Tilburg, 2021). On the other hand, the way in which an individual perceives and experiences their social circumstances includes qualitative assessments of the value, function, and meaning ascribed to relationships. Loneliness arises as a negative feeling associated with the perception of an inadequate quantity and or quality of social relationships (Zavaleta et al., 2017). It can therefore be experienced in the absence of isolation and *vice versa*, i.e., those who are socially isolated may

not experience loneliness (Dykstra, 2006; Perlman & Letitia Anne Peplau, 1984; J. E. M. Power et al., 2019)

Data from the European Social Survey collected before the pandemic indicated that 8.6 percent of the adult population had frequent feelings of loneliness whereas 20.8 percent were socially isolated (d'Hombres et al., 2021). It is clear that one experience can exist without the other, with only a moderate association observed between social isolation and loneliness (Hughes et al., 2004). Both constructs have also been found to independently predict poorer health, wellbeing, cognitive capability, and mortality in older age through different mechanisms (Coyle & Dugan, 2012; Golden et al., 2009; Steptoe, Shankar, et al., 2013), providing empirical evidence for a conceptual distinction between these two constructs. However, the investigation of objective indicators of social isolation is often neglected, with few studies examining the interaction between isolation and loneliness (Holt-Lunstad & Steptoe, 2022).

The burden of these experiences is also not equally shared and, although overlapping, the socio-demographic factors associated with social isolation and loneliness are varied. Common risk factors include low economic position and poor health; however, older age is associated with increased social isolation but not loneliness (d'Hombres et al., 2021). Despite being more likely to live alone in later life, recent findings indicate that women are less lonely than men (Barreto et al., 2021; Esteve et al., 2020). Investigations of the interaction between age and other socio-demographic factors provide a more nuanced picture. For example, low education level, deprivation, and female gender were only associated with loneliness in adolescence and early adulthood in a large Danish population-based study (Lasgaard et al., 2016). In the UK, lower levels of loneliness were associated with the number of social interactions in early adult life and relationship quality in midlife (Victor & Yang, 2012).

Although several cross-sectional studies have indicated high levels of social isolation and loneliness during the COVID-19 restrictions, it is difficult to infer causality in the absence of pre-pandemic scores (Killgore et al., 2020). The first large-scale, population-based study investigating the psychological impact of the pandemic was based on data collected in the first Understanding Society COVID-19 survey (Li & Wang, 2020). Over a third of the sample reported feeling lonely sometimes or often during the pandemic. Young people, women, and those with COVID-19 symptoms were more likely to report loneliness and mental health difficulties, while those in employment and with a cohabiting partner were found to report less loneliness. Low income, not being married or cohabiting, smaller household size (adults only), higher depressive symptoms, living in an urban area, and lower number of close friends and social support were also associated with loneliness (Bu et al., 2020; Groarke et al., 2020).

Limited to the unique experience of lockdown, few of these studies tell us much about the stability of demographic, socio-economic, and health characteristics associated with social isolation and loneliness before and during the pandemic and the strength of these associations. Were the people at risk of social isolation and loneliness prior to COVID-19 more likely to have these experiences during lockdown? Or were new groups disproportionately affected by the drastic changes to their lifestyle? To answer these questions in relation to loneliness, Bu et al., (2020) conducted a cross-cohort study using data from the UK Household Longitudinal Study (Understanding Society) (2017-2019) and the COVID-19 Social Study. Different groups of individuals were identified, including those whose risk of loneliness remained the same during the pandemic (e.g., women, urban residents, and those living alone), those who experienced heightened risk (e.g., those with low income and young people), and those who emerged as high-risk groups during the pandemic (e.g., students). Due to the use of different cohorts, this study is unable to directly compare the

experiences of individuals over time and therefore cannot make conclusions about changes due to the pandemic. The use of multiple successive birth cohorts in the current study, alongside a multi-generational ageing cohort, provides a further opportunity to examine cross-generational differences in experiences during the pandemic and tease apart age or cohort effects from period effects related to the pandemic.

This article aims to overcome some of the existing methodological limitations in the field and has two main aims. First, we describe levels of social isolation and loneliness prior to and during the COVID-19 restrictions, testing associations between social isolation and loneliness under normal circumstances and during the early stage of COVID-19 restrictions. By following the same individuals before and during the pandemic, we also provide better evidence for the differential impacts of restrictions on social isolation, loneliness, and their intersection. Second, we compare the correlates of social isolation and loneliness, and the relationship between them, pre- and during the COVID-19 restrictions to answer the following research questions:

Research Questions

1. What were the levels of social isolation and loneliness, and what proportion of the sample was classified into different groups e.g., isolated, and/or lonely prior to and during the COVID-19 restrictions?
2. What were the inter-relationships between social isolation and loneliness indicators prior to and during the COVID-19 restrictions?
3. To what extent were demographic, socio-economic factors, and physical and mental health associated with social isolation and loneliness prior to and during the COVID-19 restrictions?

Research Design and Methods

Data sources

Data collected at two time points prior to and during the COVID-19 pandemic were utilised from four UK population-based studies. Three are longitudinal birth cohort studies with samples born within a single week across England, Scotland, and Wales: the 1946 MRC National Survey of Health and Development (NSHD) (Kuh et al., 2016; Wadsworth et al., 2006), the 1958 National Child Development Study (NCDS) (Power & Elliott, 2006), and the 1970 British Cohort Study (BCS) (Elliott & Shepherd, 2006; Sullivan et al., 2022)). In contrast, the English Longitudinal Study of Ageing (ELSA) is a panel study following individuals aged ≥ 50 years biennially since 2002 (Stephens, Breeze, et al., 2013).

Table 1 provides details of the studies including participants' age and timing of data collection across surveys. It also provides details of the survey designs, sampling frames, response rates, and analytic sample sizes.

Participants

The analytic sample for each cohort included those who were alive, living in Great Britain, who took part in the study at two time points prior to and one wave during the early stages of the pandemic, and who completed the pre-pandemic and wave 1 COVID-19 surveys including outcomes of interest (social isolation and loneliness). Across all cohorts, we further restricted the sample to those who directly participated in the surveys (i.e., we excluded proxy respondents). Age bands that mapped onto the age homogenous cohorts during the COVID-19 pandemic (e.g., 50-59, 60-69, 70-79, and 80+) were generated for ELSA participants to enable analysis to differentiate between age or cohort and period effects related to the pandemic. Participants' demographic, socio-economic, and health characteristics are summarised in [supplementary file S1](#).

Sampling Design Weights and Accounting for Non-Response Bias

To account for sampling design and non-response biases, weights were applied to studies to improve representativeness of their target populations i.e., the general population of mid to older age adults in Great Britain/England. For NSHD, design weights were included in the generation of non-response weights for the wave 1 COVID-19 survey. Only survey non-response weights for the wave 1 COVID-19 timepoint were included for NCDS, BCS, and ELSA. Given that certain groups of individuals are more likely to discontinue participation in longitudinal surveys (e.g., males and those disadvantaged and less healthy), accounting for non-response in analyses ensures that data from these participants are given more weight, resulting in a more representative sample.

Measures

This section provides details of the measurement of social isolation and loneliness prior to and during the COVID-19 pandemic, and an overview of demographic, socio-economic, and health characteristics being investigated in the current study. Full details of the original items, harmonisation, and re-coding are included in [supplementary file S2](#).

Social isolation

Self-reported indicators of social isolation were identified in the four studies prior to and during the COVID-19 restrictions. Relevant items were organised by their relational context (e.g., household, community) and by the domain of social isolation assessed (e.g., network size, frequency of contact). For example, items were generated for isolation within the household (i.e., living alone), family network (i.e., partnership and children), frequency of contact with friends and relatives, education and employment status, frequency of contact with people in the community (e.g., frequency of attending community groups/organisations), and volunteering. To compare study members who were isolated across contexts prior to and during the COVID-19 restrictions,

items were reduced to those which could be harmonised across time points. An overall social isolation score was generated with a total index score (maximum value of six) indicating the number of contexts an individual was isolated across. This variable was also recoded as a binary variable, for estimating proportions and visualising overlaps with loneliness, where a score greater than three (indicating isolation in at least three contexts) was used as a binary indicator of social isolation.

Loneliness

Prior to the COVID-19 pandemic, not all cohorts asked participants a full measure of loneliness such as the UCLA Loneliness Scale. However, items included ‘I feel left out of things’ and related to the extent to which cohort members had been feeling close to others e.g., ‘I’ve been feeling close to other people’. To make variables comparable across cohorts, items were recoded as binary, indicating those that were lonely and not lonely. Across all four cohorts, the UCLA Loneliness Scale (3-item) (Hughes et al., 2004) was included in the COVID-19 survey along with an overall item ‘How often do you feel lonely?’. The short version of the Revised UCLA Loneliness Scale (R-UCLA; (Russell et al., 1980) consists of 3 items relating to lacking companionship, feeling left out, and feeling isolated from others, with simplified response options (‘hardly ever’ = 1, ‘some of the time’ = 2, or ‘often’ = 3). For cohorts that did not include the UCLA Loneliness Scale prior to the COVID-19 pandemic, the best-matched item was selected during COVID-19 to generate the loneliness indicator. For example, as can be seen in [supplementary file S2](#), in NCDS there was only 1 item relating to feeling left out collected prior to the COVID-19 pandemic; this was therefore matched with the UCLA item during COVID-19 relating to feeling left out to generate the most comparable loneliness indicators. Designed for large-scale social surveys, the three-item UCLA Loneliness Scale provides a reliable assessment of loneliness. Only a modest relationship

was found between this measure of loneliness and objective social isolation (Hughes et al., 2004) supporting the conceptual distinction in the current study.

Demographic, socio-economic and health characteristics

Demographic variables included sex as well as age and ethnicity (in ELSA only). Cohort members' highest level of educational achievement (degree vs. no degree) was also included. Socio-economic indicators were self-reported financial difficulties, home ownership, and occupational social class. A binary variable was generated to indicate those with 'poor/fair' health (vs good, very good, or excellent health). In addition, whether cohort members report a limiting longstanding illness or health problem was included as a binary variable. Continuous measures of psychological distress and life satisfaction were also included as indicators of mental health and wellbeing (see [Supplementary File S2](#) for details).

Analysis strategy

To answer the first research question and investigate the percentage of participants experiencing social isolation and loneliness prior to and during the first COVID-19 lockdown, binary variables were generated to identify those in each group. Using these indicator variables, the percentage of the sample reporting both, one, or neither are summarised prior to and during the COVID-19 restrictions. Analyses were stratified using age bands that mapped onto the other cohorts during the wave 1 COVID-19 timepoint for ELSA (50-59, 60-69, 70-79, 80+). We also present Venn diagrams depicting the proportion of the cohort experiencing social isolation and loneliness prior to and during the pandemic and the extent of overlap between these experiences.

Our second research question was addressed by examining the associations between individual indicators of social isolation and loneliness prior to and during the first COVID-19 lockdown using tetrachoric correlations. Within each cohort, we estimated tetrachoric correlation matrices using

data collected prior to the pandemic, and again during the COVID-19 restrictions. We then visualised these matrices as networks, using the using ‘qgraph’ package (Epskamp et al., 2018) in R. Our nodes were all the binary indicators of social isolation and loneliness that were available in each cohort, and the edges in the networks represented the strength of the tetrachoric correlations. We chose to present bivariate relationships in the networks rather than partial correlation coefficients to avoid introducing biased or spurious connections due to inappropriate statistical control (Wysocki et al., 2022).

The extent to which demographic, socio-economic and health characteristics were associated with social isolation and loneliness prior to and during the COVID-19 pandemic was examined using two modified Poisson regression models (Zou, 2004; Zou & Donner, 2013) for each outcome of interest and within each cohort study i.e., with social isolation and loneliness as dependent variables. For ELSA, models were stratified by age bands to ensure any age effects identified in the other cohort studies were not masked by ELSA’s age heterogenous sample.

Results

RQ1: Descriptives and overlap of social isolation and loneliness prior to and during the COVID-19 restrictions.

Table 2 provides the count and weighted percentages of the sample reporting different indicators of social isolation and loneliness prior to and during the COVID-19 restrictions. Overall rates of social isolation and loneliness are also presented for each cohort and, for ELSA, presented by age band. In addition, Table 3 summarises the count and weighted percentages of the sample reporting both social isolation and loneliness, social isolation only, loneliness only, or neither for the periods prior to and during the COVID-19 restrictions. By matching the appropriate ELSA age band with

each of the other birth cohorts, Figure 1 also offers the opportunity to differentiate between age or cohort and period effects related to the pandemic.

Table 3 shows that the percentages of respondents reporting social isolation and loneliness is generally higher in older ages in ELSA. We observe that 8.6, 10.7, 8.9 and 15.7 percent of those aged respectively 50-59, 60-69, 70-79 and 80+ reported being both lonely and isolated in the pre-Covid-19 ELSA sample. Figures slightly increased in the Covid-19 wave with, respectively, 11.3, 12.5, 11.5 and 16.4 percent of the sample. Looking at BCS, NCDS and NSHD, percentages were 4.8, 1.2 and 3.3 in the pre-Covid-19 sweeps against 5.1, 2.9 and 6.2 during the pandemic.

The same trend is observed when looking at being socially isolated in ELSA: 14.2, 18.1, 22.,7 and 38.4 percent of those aged 50-59, 60-69, 70-7,9 and 80+ respectively reported being socially isolated in the pre-pandemic wave against 14.9, 20, 25.1 and 40.8 during the pandemic. Differences between pre-pandemic and pandemic times are of the same nature in BCS (17.8 and 17.5 percent) and NSHD (14.2 and 16 percent) but are slightly higher in NCDS (21.6 and 32 percent)

The percentages of those reporting being lonely but not socially isolated and neither socially isolated nor lonely decreased by age and did not change noticeably during the pandemic.

Finally, the percentages of those reporting being neither lonely nor socially isolated tend to be lower among older age groups. In ELSA, we can observe that 63 percent of those aged 50-59 reported neither social isolation nor loneliness against 59.5 percent during the pandemic. By contrast, figures for those aged 80+ were 39.7 percent before the pandemic and 38.2 percent during the pandemic. Percentages from BCS, NCDS and NSHD are higher but show a similar pattern with respectively 70.3, 76.3 and 73.5 percent in pre-pandemic times and 73.8, 63 and 69.7 percent during the pandemic restrictions.

The Venn diagrams in [Figure 1](#) illustrate the overlap between isolation and loneliness for each age-group and study as well as before and during pandemic restrictions. The left side of the figures shows the proportions for each age-band within ELSA while their corresponding age-bands in BCS, NCDS and NSHD are shown on the right side. Three main observations flow from these figures. First, the size of the circles representing social isolation and loneliness combined as well as their intersections tend to be bigger within ELSA than within the birth cohorts due to higher percentage of respondents reporting neither loneliness nor social isolation within the birth cohorts. Second, the size of the intersections (i.e., those reporting being lonely and isolated) has not noticeably changed during the pandemic restrictions. Finally, it can be observed that the share of the population aged 80+ reporting being only lonely (and not isolated) is small compared with other age-bands. For those aged 80 and over, social isolation seems to be associated with loneliness more than in the other age-groups.

RQ2: inter-relationships between social isolation and loneliness indicators prior to and during the COVID-19 restrictions

Tetrachoric correlations between all indicators of social isolation and loneliness in the NCDS and BCS are presented as networks in [Figure 2](#), and networks by age-bands in ELSA are presented in the [supplementary file S3](#). Networks could not be estimated in NSHD due to a non-positive definite correlation matrix, likely due to a tetrachoric correlation of 1 between living and partner status (i.e., all cohort members who lived alone also had no partner – likely a result of the advanced age of the cohort). As such, NSHD was excluded from this portion of the analyses. Within the NCDS and BCS cohorts, and also within ELSA age-bands, the networks were broadly similar pre- and during COVID, particularly for the strongest edges. In NCDS and BCS, loneliness was directly correlated with all measures of social isolation prior-to and during the pandemic. Prior to the

pandemic, loneliness was most strongly associated with being out of work/education (NEET), living alone and having less than weekly contact with friends in the NCDS. During the pandemic, having no partner was most strongly associated with loneliness in this cohort, followed by living alone and NEET. For the BCS, the strongest pre-pandemic correlates of loneliness were living alone, NEET, and no community engagement. Living alone, having no partner, and NEET were the strongest correlates of loneliness in BCS during the pandemic.

In ELSA, the four indicators of self-reported loneliness/social isolation formed a strongly connected cluster of nodes. However, these clusters had many connections with the objective indicators of social isolation. Both prior to and during the pandemic, the strongest bridges between objective and subjective indicators of social isolation were between the ‘lives alone’, ‘has no partner’, ‘lacks companionship’, and ‘feels lonely’ nodes. The tetrachoric correlation matrices used to create these networks are available in the [supplementary file S4](#).

RQ3: Predictors of social isolation and loneliness prior to and during the COVID-19 restrictions

[Tables 4 and 5](#) present the results from the modified Poisson regression models for social isolation and loneliness prior to and during the COVID-19 pandemic. Demographic, socio-economic and health variables were added into the model in blocks.

Correlates of greater social isolation included female gender, manual occupational social class, self-reported financial difficulties, not being a homeowner, longstanding illness, and lower life satisfaction. There were no notable differences in the effect sizes of these associations in the pre-pandemic and lockdown periods.

Correlates of greater loneliness included female gender, not having degree level education, manual occupational social class, self-reported financial difficulties, not being a homeowner, longstanding

illness and greater psychological distress and lower life satisfaction. There were no notable differences in the effect sizes of these associations in the pre-pandemic and lockdown periods.

Discussion and Implications

This study provides a conceptual and empirical contribution, presenting the inter-relationship between social isolation and loneliness before and during the pandemic, with its unprecedented social restrictions. By using data from multiple successive birth cohorts, alongside several age-bands derived from a multi-generational ageing cohort, we were also able to examine cross-cohort differences in experiences during the early stages of COVID-19 and distinguish age or cohort effects from period effects related to the pandemic. Furthermore, we present the demographic, socio-economic, and health factors associated with experiences of social isolation and loneliness prior to and during the lockdown restrictions, adding to the literature that highlights the unequal burden of these experiences for females, those with greater economic precarity, including not owning a home, manual occupational social class, and greater financial stress.

We show support for these experiences as independent but related constructs (Hughes et al., 2004); however, our examination of demographic, socio-economic and health characteristics suggests mostly common correlates for these outcomes. Across all the datasets examined, and during both pre-pandemic and restricted periods of restrictions, more people reported being socially isolated than lonely based on our cut-off points. This maps onto findings from the European Social Survey collected before the pandemic, which indicated that 8.6 percent of the adult population had frequent feelings of loneliness whereas 20.8 percent were socially isolated (d'Hombres et al., 2021). When examining age-based differences, we observe higher levels of isolation at older ages. However, levels of loneliness were more stable across later life. When comparing different age

groups, we see that the overlap between social isolation and loneliness is fairly consistent pre- and during the COVID-19 restrictions.

Examining the associations at the indicator-level, we find that the inter-relationships between indicators during the lockdown were less strongly connected compared to before the pandemic. This analysis suggests that there might have been an impact on how these concepts relate during restrictions, but these differences were not marked and are unlikely to indicate any fundamental differences in the conceptual links between loneliness and social isolation indicators during lockdown. The COVID-19 pandemic restrictions had specific effects on the prevalence of some of these indicators. For instance, remote contact increased and likely partially compensated for reductions in in-person contacts (Wels et al., 2023). There were also substantially fewer individuals in education and employment (Wels et al., 2022).

The study has several strengths, including the use of multiple data sources with slightly different designs and measures that permit the examination of consistency and replication of findings. However, there are limitations to note including the lack of availability of detailed measures of loneliness in the three birth cohorts before the pandemic, and data availability for relevant measures at certain timepoints (e.g., ELSA did not collect measures of face-to-face contact during the pandemic). We used data from participants who had responded before and during the pandemic surveys; this maintains a comparable sample, although those who dropped out might have been more isolated and in poorer health, potentially leading to some underestimation in the observed associations despite the use of sample and non-response weights in analyses.

By comparing data prior to and during enforced pandemic restrictions, we were able to provide insight into how these associations might vary in different contexts. In both these periods, we find similar associations between indicators of social isolation and loneliness, and with regard to other

demographic, socioeconomic, and health correlates in both these periods. The number of study members reporting only isolation or loneliness, and the moderate overlap between older adults reporting both experiences indicate that these concepts are not interchangeable, and clarity in how they are conceptualised, operationalised and interpreted in quantitative research is necessary. This, in turn, will contribute to a better understanding of the role and consequences of social isolation and loneliness in older age, and inform how interventions might support different aspects of these outcomes in older adults.

The increases in social isolation observed in the pandemic highlight the need for efforts to support older people to (re)start hobbies, volunteer, and schedule time to meet up with friends and neighbours, as these activities can also lead to health and other psychological benefits (Fancourt et al., 2022). The findings that being female, economic precarity, and long-standing illness are stable correlates of isolation and loneliness indicate a need for structural changes and policies designed to reduce these inequalities in experiences of isolation and loneliness.

We would also like to highlight the implications of our findings for the currently widespread conflation of the two terms “loneliness” and “social isolation” in policy, and the over emphasis on loneliness within the UK context (e.g., campaign to end loneliness, loneliness ministers). Given many older adults experience high levels of social isolation, there should be greater emphasis on reducing social isolation through policy intervention rather than focusing on reducing individuals’ feelings of loneliness. Investigation of objective social isolation shifts the focus away from individuals and towards structural factors contributing greater isolation and the inequities in these experiences and contributing factors (Umberson et al., 2022). This can help to identify areas that are modifiable through targeted policy and intervention.

Funding

This research was primarily supported by grants from the UK Economic and Social Council (ES/V012789/1 and ES/T007575/1). We also acknowledge the support of the MRC for the National Core Study in Longitudinal Health and Wellbeing (MC_PC_20059) and MRC Unit for Lifelong Health and Ageing (MC_UU_00019/1 and MC_UU_00019/3). In addition, JS, JM and PP are supported by Alzheimer's Society (Ref:469). JW is funded by the Belgian FNRS CQ research grant (40010931).

Conflict of Interest

None.

Data Availability

The data used for this project is available to all researchers from the UK Data Service: <https://ukdataservice.ac.uk/> and NSHD: <https://nshd.mrc.ac.uk/>

Acknowledgements

We are grateful to all the cohort members who took part in the NSHD, NCDS, BCS and ELSA cohorts analysed in this paper. We are also grateful to the survey and data management teams of the included longitudinal cohorts. We also acknowledge the input of the What Works Centre for Wellbeing, Campaign to End Loneliness and our Stakeholder Advisory Group.

References

- Barreto, M., Victor, C., Hammond, C., Eccles, A., Richins, M. T., & Qualter, P. (2021). Loneliness around the world: Age, gender, and cultural differences in loneliness. *Personality and Individual Differences, 169*, 110066. <https://doi.org/10.1016/j.paid.2020.110066>
- Bu, F., Steptoe, A., & Fancourt, D. (2020). Loneliness during a strict lockdown: Trajectories and predictors during the COVID-19 pandemic in 38,217 United Kingdom adults. *Social Science and Medicine, 265*. <https://doi.org/10.1016/j.socscimed.2020.113521>
- Coyle, C. E., & Dugan, E. (2012). Social isolation, loneliness and health among older adults. *Journal of Aging and Health, 24*(8), 1346–1363. <https://doi.org/10.1177/0898264312460275>
- d’Hombres, B., Barjaková, M., & Schnepf, S. V. (2021). *Loneliness and Social Isolation: An unequally shared burden in Europe. 14245, 23.*
- Dykstra, J. de J. G. T. van T. P. A. (2006). *Loneliness and Social Isolation.*
- Elliott, J., & Shepherd, P. (2006). Cohort profile: 1970 British Birth Cohort (BCS70). *International Journal of Epidemiology, 35*(4), 836–843. <https://doi.org/10.1093/ije/dy1174>
- Epskamp, S., Borsboom, D., & Fried, E. I. (2018). Estimating psychological networks and their accuracy: A tutorial paper. *Behavior Research Methods, 50*(1), 195–212. <https://doi.org/10.3758/s13428-017-0862-1>
- Esteve, A., Reher, D. S., Treviño, R., Zueras, P., & Turu, A. (2020). Living Alone over the Life Course: Cross-National Variations on an Emerging Issue. *Population and Development Review, 46*(1), 169–189. <https://doi.org/10.1111/padr.12311>

- Fancourt, D., Mak, H. W., Noguchi, T., Bone, J., Wels, J., Gao, Q., Kondo, K., & Saito, T. (2022). *Psychological benefits of hobby engagement in older age: a longitudinal cross-country analysis of 93,263 older adults in 16 countries*. <https://doi.org/10.21203/rs.3.rs-2349781/v1>
- Golden, J., Conroy, R. M., Bruce, I., Denihan, A., Greene, E., Kirby, M., & Lawlor, B. A. (2009). Loneliness, social support networks, mood and wellbeing in community-dwelling elderly. *International Journal of Geriatric Psychiatry*, *24*(7), 694–700. <https://doi.org/10.1002/gps.2181>
- Groarke, J. M., Berry, E., Graham-Wisener, L., McKenna-Plumley, P. E., McGlinchey, E., & Armour, C. (2020). Loneliness in the UK during the COVID-19 pandemic: Cross-sectional results from the COVID-19 Psychological Wellbeing Study. *PLoS ONE*, *15*(9 September), 1–18. <https://doi.org/10.1371/journal.pone.0239698>
- Holt-Lunstad, J., Smith, T. B., Baker, M., Harris, T., & Stephenson, D. (2015). Loneliness and Social Isolation as Risk Factors for Mortality: A Meta-Analytic Review. *Perspectives on Psychological Science*, *10*(2), 227–237. <https://doi.org/10.1177/1745691614568352>
- Holt-Lunstad, J., & Steptoe, A. (2022). Social isolation: An underappreciated determinant of physical health. *Current Opinion in Psychology*, *43*, 232–237. <https://doi.org/10.1016/j.copsyc.2021.07.012>
- Hughes, M. E., Waite, L. J., Hawkey, L. C., & Cacioppo, J. T. (2004). A short scale for measuring loneliness in large surveys: Results from two population-based studies. *Research on Aging*, *26*(6), 655–672. <https://doi.org/10.1177/0164027504268574>

- Huisman, M., & van Tilburg, T. G. (2021). Social exclusion and social isolation in later life. In *Handbook of Aging and the Social Sciences*. Academic Press. <https://doi.org/10.1016/b978-0-12-815970-5.00007-3>
- Jeste, D. V., Lee, E. E., & Cacioppo, S. (2019). Battling the Modern Behavioral Epidemic of Loneliness: Suggestions for Research and Interventions. *Physiology & Behavior*, *176*(3), 139–148. <https://doi.org/10.1001/jamapsychiatry.2020.0027>.
- Killgore, W. D. S., Cloonan, S. A., Taylor, E. C., & Dailey, N. S. (2020). Loneliness: A signature mental health concern in the era of COVID-19. *Psychiatry Research Journal*, *290*. <https://doi.org/10.1016/j.psychres.2020.113117>.
- Kuh, D., Wong, A., Shah, I., Moore, A., Popham, M., Curran, P., Davis, D., Sharma, N., Richards, M., Stafford, M., Hardy, R., & Cooper, R. (2016). The MRC National Survey of Health and Development reaches age 70: maintaining participation at older ages in a birth cohort study. *European Journal of Epidemiology*, *31*(11), 1135–1147. <https://doi.org/10.1007/s10654-016-0217-8>
- Lasgaard, M., Friis, K., & Shevlin, M. (2016). “Where are all the lonely people?” *Social Psychiatry and Psychiatric Epidemiology*, *51*(10), 1373–1384. <https://doi.org/10.1007/s00127-016-1279-3>
- Li, L. Z., & Wang, S. (2020). Prevalence and predictors of general psychiatric disorders and loneliness during COVID-19 in the United Kingdom. *Psychiatry Research*, *291*(June), 113267. <https://doi.org/10.1016/j.psychres.2020.113267>
- Perlman, D., & Letitia Anne Peplau. (1984). Loneliness research: A survey of empirical findings. In L. A. Peplau & S. E. Goldston (Eds.), *Preventing the harmful consequences of severe and persistent loneliness* (pp. 13–46). National Institute of Mental Health.

- Power, C., & Elliott, J. (2006). Cohort profile: 1958 British birth cohort (National Child Development Study). *International Journal of Epidemiology*, 35(1), 34–41.
<https://doi.org/10.1093/ije/dyi183>
- Power, J. E. M., Sjöberg, L., Kee, F., Kenny, R. A., & Lawlor, B. (2019). Comparisons of the discrepancy between loneliness and social isolation across Ireland and Sweden: findings from TILDA and SNAC-K. *Social Psychiatry and Psychiatric Epidemiology*, 54(9), 1079–1088. <https://doi.org/10.1007/s00127-019-01679-w>
- Russell, D., Peplau, L. A., & Cutrona, C. E. (1980). The revised UCLA Loneliness Scale: Concurrent and discriminant validity evidence. *Journal of Personality and Social Psychology*, 39(3), 472–480. <https://doi.org/10.1037//0022-3514.39.3.472>
- Step toe, A., Breeze, E., Banks, J., & Nazroo, J. (2013). Cohort profile: The English Longitudinal Study of Ageing. *International Journal of Epidemiology*, 42(6), 1640–1648.
<https://doi.org/10.1093/ije/dys168>
- Step toe, A., Shankar, A., Demakakos, P., & Wardle, J. (2013). Social isolation, loneliness, and all-cause mortality in older men and women. *Proceedings of the National Academy of Sciences of the United States of America*, 110(15), 5797–5801.
<https://doi.org/10.1073/pnas.1219686110>
- Sullivan, A., Brown, M., Hamer, M., & Ploubidis, G. B. (2022). Cohort Profile Update: The 1970 British Cohort Study (BCS70). *International Journal of Epidemiology*, 1–8.
<https://doi.org/10.1093/ije/dyac148>
- Umberson, D., Lin, Z., & Cha, H. (2022). Gender and Social Isolation across the Life Course. *Journal of Health and Social Behavior*, 63(3), 319–335.
<https://doi.org/10.1177/00221465221109634>

- Victor, C. R., & Yang, K. (2012). The prevalence of loneliness among adults: A case study of the United Kingdom. *Journal of Psychology, 146*(1–2), 85–104.
<https://doi.org/10.1080/00223980.2011.613875>
- Wadsworth, M., Kuh, D., Richards, M., & Hardy, R. (2006). Cohort profile: The 1946 National Birth Cohort (MRC National Survey of Health and Development). *International Journal of Epidemiology, 35*(1), 49–54. <https://doi.org/10.1093/ije/dyi201>
- Wels, J., Booth, C., Wielgoszewska, B., Green, M. J., Di Gessa, G., Huggins, C. F., Griffith, G. J., Kwong, A. S. F., Bowyer, R. C. E., Maddock, J., Patalay, P., Silverwood, R. J., Fitzsimons, E., Shaw, R., Thompson, E. J., Steptoe, A., Hughes, A., Chaturvedi, N., Steves, C. J., ... Ploubidis, G. B. (2022). Mental and social wellbeing and the UK coronavirus job retention scheme: Evidence from nine longitudinal studies. *Social Science & Medicine, 308*, 115226. <https://doi.org/10.1016/j.socscimed.2022.115226>
- Wels, J., Wielgoszewska, B., Moltrecht, B., Booth, C., Green, M. J., Hamilton, O. K., Demou, E., Di Gessa, G., Huggins, C., Zhu, J., Santorelli, G., Silverwood, R. J., Kopasker, D., Shaw, R. J., Hughes, A., Patalay, P., Steves, C., Chaturvedi, N., Porteous, D. J., ... Ploubidis, G. B. (2023). Home working and social and mental wellbeing at different stages of the COVID-19 pandemic in the UK: Evidence from 7 longitudinal population surveys. *PLOS Medicine, 20*(4), e1004214. <https://doi.org/10.1371/journal.pmed.1004214>
- Wigfield, A., Turner, R., Alden, S., Green, M., & Karania, V. (2022). Developing a New Conceptual Framework of Meaningful Interaction for Understanding Social Isolation and Loneliness. *Social Policy and Society, 21*(2), 172-193.
<https://doi.org/10.1017/S147474642000055X>

Wysocki, A. C., Lawson, K. M., & Rhemtulla, M. (2022). Statistical Control Requires Causal Justification. *Advances in Methods and Practices in Psychological Science*, 5(2).

<https://doi.org/10.1177/25152459221095823>

Zavaleta, D., Samuel, K., & Mills, C. T. (2017). Measures of Social Isolation. *Social Indicators Research*, 131(1), 367–391. <https://doi.org/10.1007/s11205-016-1252-2>

Zou, G. (2004). A Modified Poisson Regression Approach to Prospective Studies with Binary Data. *American Journal of Epidemiology*, 159(7), 702–706.

<https://doi.org/10.1093/aje/kwh090>

Zou, G., & Donner, A. (2013). Extension of the modified Poisson regression model to prospective studies with correlated binary data. *Statistical Methods in Medical Research*, 22(6), 661–670. <https://doi.org/10.1177/0962280211427759>

Accepted Manuscript

Table 1. Details of the cohort studies including participants' age and timing of data collection, survey designs, sampling frames, response rates, and analytic sample sizes

Study Population	Design and Sample Frame	2020 Age Range in years	Most Recent Pre-Pandemic Survey	Details of wave 1 COVID-19 Survey (Response Rate)	Analytic N
<i>Age Homogenous Cohorts</i>					
BCS70: British Cohort Study 1970	Cohort of all children born in Great Britain (i.e., England, Wales & Scotland) in one week in 1970, with regular follow-up surveys from birth.	50	2016	Feb-Mar (40%)	3,075
NCDS: National Child Development Study	Cohort of all children born in Great Britain (i.e., England, Wales & Scotland) in one week in 1958, with regular follow-up surveys from birth.	62	2008	Feb-Mar (52%)	3,851
NSHD: National Survey of Health and Development	Cohort of all children born in Great Britain (i.e., England, Wales & Scotland) in one week in 1946, with regular follow-up surveys from birth.	74	2015	Feb-Mar (90%)	763
<i>Age Heterogeneous Studies</i>					
ELSA: English Longitudinal Study of Aging	A nationally representative population study of individuals aged 50+ living in England, with biennial surveys and periodic refreshing of the sample to maintain representativeness.	52-90+	2018-2019	Jun-Jul (75%)	4,440

Table 2. Count and percentage of the sample reporting social isolation and loneliness prior to and during the COVID-19 restrictions for each cohort

Variable	NSHD	NCDS	BCS	ELSA			
	70	50	46	50-59 (n=236)	60-69 (n=1,501)	70-79 (n=1,887)	80+ (n=816)
Pre-COVID-19							
Social Isolation % (N)							
Living alone	14.3 (109)	9.8 (292)	8.3 (160)	16.6 (42)	19.8 (282)	22.8 (456)	42.6 (340)
No children	9.6 (73)	16.2 (587)	16.6 (489)	18.9 (50)	16.7 (259)	10.7 (216)	8.3 (74)
No partner	16.0 (122)	18.0 (581)	15.9 (359)	25.8 (59)	27.9 (369)	26.4 (510)	50.2 (383)
No weekly in person contact with friends/relatives	18.6 (142)	57.7 (2,288)	23.2 (766)	-	-	-	-
No weekly remote contact with friends/relatives		-	-	5.1 (8)	3.8 (56)	4.3 (79)	4.4 (35)
In neither education nor employment	78.8 (601)	13.0 (372)	13.2 (269)	23.4 (53)	53.6 (930)	92.0 (18740)	99.2 (806)
No community engagement i.e., not a member of a club/organisation or volunteer	72.7 (555)	77.2 (2,877)	76.1 (2,174)	90.0 (209)	82.8 (1196)	77.6 (1414)	85.8 (672)
Social Isolation Indicator – High Score (≥ 3)	17.4 (133)	22.8 (734)	15.1 (346)	22.8 (54)	28.7 (415)	31.6 (607)	54.1 (418)
Loneliness % (N)							
Feeling lack of companionship – often	4.6 (35)	-	-	8.4 (17)	6.7 (93)	5.1 (92)	5.7 (41)
Feeling left out – often	2.5 (19)	2.2 (58)	-	7.1 (12)	3.5 (52)	3.7 (62)	4.3 (25)
Feeling isolated from others - often	2.8 (21)	-	11.9 (269)	7.5 (16)	4.7 (67)	3.1 (53)	3.9 (27)
Feeling lonely - often	-	-	-	9.7 (21)	7.2 (98)	4.9 (86)	5.0 (35)
UCLA total mean score (items 1-3) (3-9 range)	3.74	-	-	4.28	4.13	3.98	4.20
Loneliness Indicator - High score	12.3 (94)	2.2 (58)	11.9 (269)	22.8 (50)	21.1 (281)	17.3 (313)	22.0 (156)
COVID-19							
Social Isolation % (N)							
Living alone	24.4 (186)	23.4 (834)	13.8 (305)	18.1 (43)	22.2 (325)	25.2 (495)	45.7 (366)
No children	5.5 (42)	21.4 (765)	18.9 (534)	18.9 (50)	16.7 (259)	10.7 (216)	8.3 (74)
No partner	24.4 (186)	16.7 (557)	15.2 (345)	28.0 (65)	30.0 (414)	30.6 (583)	54.1 (422)
No weekly in person contact with friends/relatives	0.5 (4)	57.8 (2,217)	59.5 (1,848)	-	-	-	-
No weekly remote contact with friends/relatives	0.3 (2)	-	-	3.0 (6)	4.0 (54)	4.2 (70)	5.4 (38)
In neither education nor employment	94.8 (723)	66.9 (2,523)	39.1 (1,034)	43.5 (101)	71.4 (1171)	96.7 (1826)	99.1 (806)
No community engagement i.e., not a member of a club/organisation or volunteer	36.4 (278)	32.2 (1,139)	26.3 (702)	83.3 (193)	79.6 (1191)	85.0 (1586)	88.1 (708)
Social Isolation Indicator – High Score (≥ 3)	22.2 (169)	34.9 (1,226)	22.6 (531)	26.2 (63)	32.5 (477)	36.5 (693)	57.2 (448)
Loneliness % (N)							

Feeling lack of companionship – often	4.5 (34)	-	-	8.1 (20)	7.1 (96)	5.2 (100)	8.5 (64)
Feeling left out – often	2.2 (17)	5.0 (156)	-	6.2 (13)	3.9 (57)	3.4 (61)	4.7 (32)
Feeling isolated from others - often	9.0 (69)	-	8.7 (205)	8.3 (22)	7.5 (102)	6.2 (111)	9.1 (64)
Feeling lonely - often	3.7 (28)	-	-	9.5 (20)	5.8 (78)	5.2 (90)	4.9 (36)
UCLA total mean score (items 1-3) (3-9 range)	3.98	-	-	4.42	4.20	4.14	4.25
Loneliness Indicator - High score	14.3 (109)	5.0 (156)	8.7 (205)	25.6 (60)	23.0 (320)	20.4 (385)	21.0 (165)

Note. NSHD = 1946 MRC National Survey of Health and Development; NCDS = 1958 National Child Development Study; BCS = 1970 British Cohort Study; ELSA = English Longitudinal Study of Ageing; UCLA = UCLA Loneliness Scale. Weighted %. Social isolation indicator is calculated by giving a score of 1 for every type of social isolation and generating a score out of 6. Those with scores ≥ 3 indicate social isolation. Loneliness indicator is generated differently for NCDS and BCS where the full UCLA Loneliness Scale is not available. For these cohorts, loneliness = a response of 'often' to the loneliness item available.

Table 3. Count and % of the sample reporting possible combinations of social isolation and loneliness prior to and during the COVID-19 restrictions for each cohort and for the samples combined

Outcome Variable	Pre-COVID-19							COVID-19						
	NSHD 70-79	NCDS 60-69	BCS 50-59	ELSA				NSHD 70-79	NCDS 60-69	BCS 50-59	ELSA			
				50-59	60-69	70-79	80+				50-59	60-69	70-79	80+
Both (social isolated and lonely)	3.3 (25)	1.2 (26)	4.8 (76)	8.6 (20)	10.7 (144)	8.9 (165)	15.7 (110)	6.2 (47)	2.9 (80)	5.1 (81)	11.3 (28)	12.5 (170)	11.4 (217)	16.4 (126)
Only socially isolated	14.2 (108)	21.6 (708)	17.8 (455)	14.2 (34)	18.1 (271)	22.7 (442)	38.4 (308)	16.0 (122)	32.0 (1,146)	17.5 (450)	14.9 (35)	20.0 (307)	25.1 (476)	40.8 (322)
Only lonely	9.0 (69)	1.0 (32)	7.2 (193)	14.2 (30)	10.4 (137)	8.4 (148)	6.2 (46)	8.1 (62)	2.1 (76)	3.7 (124)	14.3 (32)	10.5 (150)	9.0 (169)	4.6 (39)
Neither	73.5 (561)	76.2 (3,085)	70.3 (2,351)	63.0 (152)	60.9 (949)	60.0 (1,132)	39.7 (352)	69.7 (532)	63.0 (2,549)	73.8 (2,420)	59.5 (141)	57.0 (874)	54.5 (1,026)	38.2 (329)

Note. NSHD = 1946 MRC National Survey of Health and Development; NCDS = 1958 National Child Development Study; BCS = 1970 British Cohort Study; ELSA = English Longitudinal Study of Ageing; UCLA = UCLA Loneliness Scale. Weighted percentages.

Table 4. Results from modified Poisson regression models for social isolation prior to and during the COVID-19 pandemic including relative risk (RR) and 95% Confidence Intervals (95% CI)

Cohort	NSHD		NCDS		BCS		ELSA							
	Pre-COVID-19	COVID-19	Pre-COVID-19	COVID-19	Pre-COVID-19	COVID-19	50-59		60-69		70-79		80+	
							Pre-COVID-19	COVID-19	Pre-COVID-19	COVID-19	Pre-COVID-19	COVID-19	Pre-COVID-19	COVID-19
<i>Demographic</i>														
Gender (woman)	1.64 0.92-2.91	1.25 0.61-2.56	1.02 .86-1.20	1.05 .94-1.18	.65*** .48-.88	.83 .67-1.04	1.21 0.70-2.06	1.22 0.75-1.98	1.11 0.89-1.36	1.18 0.97-1.43	1.30 1.12-1.51	1.29 1.13-1.48	1.76 1.48-2.09	1.63 0.38-1.90
Black, Asian, and Minority Ethnic (if ethnic minority)	-	-	-	-	-	-	0.44 0.13-1.50	0.75 0.26-2.19	1.02 0.54-1.87	1.03 0.59-1.80	0.98 0.55-1.72	1.20 0.77-1.87	1.09 0.79-1.50	1.03 0.74-1.44
<i>Education/Socio-economic</i>														
Degree (yes)	1.40 0.81-2.41	0.96 0.38-2.44	.94 .71-1.25	.91 .74-1.12	1.15 .80-1.67	1.01 .77-1.33	0.69 0.35-1.35	0.98 0.56-1.71	1.09 0.83-1.42	0.92 0.73-1.15	0.94 0.77-1.15	0.82 0.69-0.98	0.86 0.67-1.11	0.91 0.72-1.15
Self-reported financial difficulties (yes)	1.28 0.54-3.04	1.25 0.48-3.25	1.41** 1.06-1.88	1.10 .86-1.40	1.03 .69-1.53	1.21 .91-1.61	1.05 0.60-1.81	1.37 0.80-2.37	1.25 1.00-1.57	1.50 1.21-1.86	1.16 0.96-1.40	1.17 0.99-1.38	0.88 0.73-1.06	1.00 0.83-1.21
Homeowner (yes)	1.91 0.40-9.06	0.81 0.25-2.61	.67*** .54-.83	.81** .68-.97	.41*** .31-.54	.57*** .45-.73	0.47 0.27-0.83	0.60 0.33-1.07	0.46 0.37-0.57	0.52 0.42-0.63	0.52 0.44-0.63	0.51 0.44-0.59	0.67 0.57-0.79	0.72 0.61-0.83
<i>Occupational Social Class^a</i>														
Intermediate (if yes)	1.02 0.54-1.92	2.41 0.95-6.11	1.05 .88-1.25	1.03 .91-1.17	.95 .67-1.34	1.32 .98-1.78	3.72 1.07-13.0	1.97 0.74-5.23	1.13 0.86-1.49	1.01 0.79-1.28	1.03 0.86-1.24	0.90 0.77-1.06	0.94 0.79-1.13	0.93 0.77-1.10
Manual (if yes)	0.65 0.22-1.91	2.72 0.67-10.93	.85 .63-1.15	.91 .09-.76	.78 .54-1.14	1.47** 1.09-1.97	2.22 0.66-7.56	1.86 0.79-3.43	0.99 0.76-1.29	0.91 0.72-1.14	0.96 0.80-1.15	0.85 0.72-0.99	0.88 0.74-1.04	0.92 0.78-1.08
Other (if yes)	0.96 0.31-2.97	2.60 0.72-9.43	1.54*** 1.19-2.00	1.28** 1.02-1.62	2.56*** 1.79-3.65	2.81*** 2.08-3.79	2.81 0.99-7.98	1.79 0.85-3.79	1.08 0.76-1.53	1.06 0.77-1.44	0.49 0.11-2.14	0.37 0.08-1.73	1.00 0.71-1.42	1.07 0.77-1.48
Managerial (ref.)														
<i>Physical and Mental Health</i>														
Self-reported health	0.90 0.44-1.85	0.29 0.06-1.27	1.01 .80-1.28	1.14 .95-1.38	1.27 .93-1.74	1.01 .74-1.39	0.95 0.42-2.17	1.12 0.64-1.92	0.90 0.67-1.21	0.85 0.67-1.09	1.03 0.84-1.26	1.08 0.90-1.28	0.97 0.81-1.15	1.03 0.88-1.21
Limiting long-standing illness or health problem (if yes)	1.12 0.65-1.92	1.83 0.98-3.41	1.10 .86-1.42	1.11 .91-1.35	1.17 .85-1.60	.94 .67-1.32	2.09 0.98-4.43	2.17 1.21-3.87	1.43 1.11-1.83	1.14 0.91-1.42	0.98 0.82-1.17	1.01 0.87-1.17	0.89 0.75-1.04	0.97 0.84-1.11

Psychological distress	1.29 0.81-2.04	0.96 0.55-1.66	1.01 .94-1.09	.96 .89-1.02	1.02 .91-1.15	1.05 .95-1.17	0.98 0.77-1.24	1.13 0.90-1.41	1.06 0.96-1.17	1.00 0.92-1.08	1.05 0.97-1.14	0.99 0.89-1.06	1.08 1.01-1.15	1.06 1.00-1.13
Life satisfaction	0.74 0.66-0.84	0.70 0.60-0.81	.91*** .87-.95	.93*** .89-.96	.92** .86-.98	.92*** .86-.98	0.93 0.79-1.09	0.94 0.84-1.03	0.92 0.88-0.97	0.93 0.90-0.97	0.91 0.87-0.94	0.91 0.89-0.94	0.98 0.94-1.01	0.96 0.93-0.99

Notes. NSHD = 1946 MRC National Survey of Health and Development; NCDS = 1958 National Child Development Study; BCS = 1970 British Cohort Study; ELSA = English Longitudinal Study of Ageing; UCLA = UCLA Loneliness Scale.

^a. Occupational Social Class refers to NS-SEC, the National Statistics Socio-Economic Classification, and is derived from the current or most recent (when the person is not working) job. Respondents who have never worked or for whom no NS-SEC information was collected are coded as 'Other'.

** $p < .05$; *** $p < .01$.

Table 5. Results from modified Poisson regression models for loneliness prior to and during the COVID-19 pandemic including relative risk (RR) and 95% Confidence Intervals (95% CI)

Cohort	NSHD		NCDS		BCS		ELSA							
	Pre-COVID-19	COVID-19	Pre-COVID-19	COVID-19	Pre-COVID-19	COVID-19	50-59		60-69		70-79		80+	
							Pre-COVID-19	COVID-19	Pre-COVID-19	COVID-19	Pre-COVID-19	COVID-19	Pre-COVID-19	COVID-19
Gender (if woman)	2.36 1.37-4.07	1.94 0.99-3.82	.60 .33-1.12	1.47** .96-2.25	.54*** .39-.75	1.04 .65-1.67	1.08 0.62-1.86	1.79 1.05-3.06	1.39 1.07-1.81	1.75 1.35-2.27	1.32 1.05-1.65	1.69 1.37-2.10	1.39 0.99-1.95	1.62 1.14-2.30
Black, Asian, and Minority Ethnic (if ethnic minority)	-	-	-	-	-	-	2.47 1.22-5.00	2.28 1.12-4.64	2.21 1.46-3.35	1.52 0.94-2.46	0.96 0.39-2.13	0.54 0.17-2.10	0.92 0.28-3.02	1.32 0.54-3.26
Education/Socio-economic														
Degree (if yes)	1.38 0.84-2.27	1.01 0.51-1.99	.46 .11-1.91	.70 .32-1.51	.98 .64-1.49	.86 .52-1.41	0.73 0.37-1.40	0.61 0.32-1.14	0.47 0.32-0.70	1.04 0.79-1.37	0.63 0.45-0.89	0.99 0.78-1.27	0.79 0.47-1.32	0.91 0.58-1.43
Self-reported financial difficulties (if yes)	0.63 0.16-2.51	1.92 0.69-5.39	4.04*** 1.82-8.96	2.15** 1.14-4.05	2.19*** 1.40-3.43	2.05** 1.19-3.54	1.74 0.98-3.07	1.92 1.09-3.37	1.53 1.17-2.02	2.05 1.58-2.67	1.66 1.28-2.15	1.91 1.52-2.38	1.49 1.05-2.10	1.65 1.13-2.41
Homeowner (if yes)	0.90 0.30-2.71	0.74 0.28-1.94	1.58 .62-4.02	.57 .31-1.05	.77 .55-1.09	.42*** .29-.62	0.70 0.40-1.21	0.95 0.51-1.76	0.80 0.58-1.09	0.70 0.52-0.93	0.72 0.52-1.00	0.74 0.55-1.00	0.70 0.47-1.04	1.00 0.62-1.63
Occupational Social Class ^a														
Intermediate (if yes)	0.71 0.39-1.27	0.86 0.39-1.88	1.46 .72-2.95	.77 .44-1.35	1.34 .90-2.00	.77 .46-1.28	2.20 0.66-7.31	1.48 0.59-3.69	0.84 0.57-1.24	0.82 0.59-1.44	1.01 0.76-1.35	1.04 0.80-1.34	0.95 0.61-1.48	0.55 0.36-0.83
Manual (if yes)	0.57 0.22-1.53	1.28 0.42-3.96	2.26 .69-7.39	.87 .49-1.54	1.93*** 1.33-2.80	1.29 .81-2.04	1.64 0.60-4.47	1.08 0.51-2.27	1.32 0.95-1.83	0.99 0.73-1.34	0.97 0.74-1.28	1.11 0.87-1.41	0.91 0.63-1.32	0.78 0.55-1.11
Other (if yes)	0.45 0.15-1.34	0.49 0.15-1.61	5.76*** 2.61-12.70	1.98** 1.04-3.77	2.37*** 1.47-3.83	3.13*** 1.93-5.07	2.03 0.83-4.96	1.01 0.53-1.91	1.19 0.76-1.85	1.18 0.79-1.75	0.87 0.22-3.43	1.77 0.57-5.47	0.19 0.02-1.69	-- ²
Managerial (ref.)														
Physical and Mental Health														
Self-reported health	0.97 0.55-1.69	1.64 0.93-2.90	1.04 .58-1.85	.77 .41-1.45	1.52 .99-2.34	.91 .58-1.41	1.00 0.46-2.17	1.27 0.74-2.16	0.76 0.54-1.06	0.76 0.54-1.04	0.62 0.46-0.82	0.89 0.69-1.13	1.28 0.88-1.86	0.75 0.51-1.11
Limiting Long-standing illness or health problem (if yes)	1.10 0.71-1.71	1.16 0.71-1.90	1.02 .51-2.06	1.89** 1.10-3.25	.90 .59-1.38	1.07 .71-1.61	1.57 0.76-3.26	1.59 0.86-2.97	1.22 0.90-1.65	0.87 0.66-1.17	0.76 0.58-0.97	1.09 0.87-1.36	1.36 0.95-1.96	0.88 0.61-1.26
Psychological distress	0.89 0.72-1.11	0.90 0.67-1.22	1.56*** 1.30-1.88	1.60*** 1.35-1.91	1.20*** 1.06-1.36	1.33*** 1.15-1.52	1.06 0.83-1.33	1.36 1.04-1.78	1.24 1.09-1.41	1.15 1.03-1.28	1.34 1.23-1.46	1.17 1.08-1.27	1.40 1.23-1.58	1.30 1.14-1.48

Life satisfaction	0.65 0.55-0.76	0.65 0.53-0.79	.77*** .68-.86	.86*** .77-.95	.81*** .75-.86	.80*** .74-.87	0.76 0.66-0.86	0.90 0.79-1.03	0.84 0.79-0.90	0.85 0.80-0.90	0.81 0.77-0.86	0.83 0.80-0.88	0.90 0.85-0.96	0.87 0.81-0.92
-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------	-------------------

Notes. NSHD = 1946 MRC National Survey of Health and Development; NCDS = 1958 National Child Development Study; BCS = 1970 British Cohort Study; ELSA = English Longitudinal Study of Ageing; UCLA = UCLA Loneliness Scale.

^a. Occupational Social Class refers to NS-SEC, the National Statistics Socio-Economic Classification, and is derived from the current or most recent (when the person is not working) job. Respondents who have never worked or for whom no NS-SEC information was collected are coded as 'Other'.

^b. Too few cases.

** $p < .05$; *** $p < .01$.

Figure 1. Figure 1. Venn diagrams showing overlap in social isolation and loneliness across cohorts.

Figure 2. Tetrachoric correlation networks between indicators of social isolation and loneliness in NCDS and BCS.

Accepted Manuscript

Figure 1





Figure 2



Accepted Manuscript