

Welcome to **FENS** Forum 2022



E-Book of Abstracts

fens.org/2022

To search in the book please download it and use control + F

BOARD NUMBER: S05-250

AGE-DEPENDENT ROLE OF NMDA RECEPTORS IN EXPERIMENTAL AUTOIMMUNE ENCEPHALOMYELITIS

POSTER SESSION 05 - SECTION: NEUROINFLAMMATION AND AUTOIMMUNITY

Ivana Ćuruvija¹, Biljana Bufan², Emilija Đorović³, Veljko Blagojević¹, Jelica Grujić-Milanović⁴, Milica Marković³, Jasmina Djuretić³

¹Institute of Virology, Vaccines and Sera "Torlak", Immunology Research Centre "branišlav Janković", Belgrade, Serbia, ²University of Belgrade - Faculty of Pharmacy, Department Of Microbiology And Immunology, Belgrade, Serbia, ³University of Belgrade - Faculty of Pharmacy, Department Of Pathobiology, Belgrade, Serbia, ⁴Institute for Medical Research, National Institute of Republic of Serbia, Department For Cardiovascular Research, Laboratory For Experimental Hypertension, Belgrade, Serbia

Aims Ageing affects *N*-methyl-D-aspartate receptors (NMDARs), their expression and function in neuronal and non-neuronal cells. Contribution of NMDARs to pathogenesis of experimental autoimmune encephalomyelitis (EAE) has been investigated but further study is still needed. The aim of this study was to determine whether ageing affects the role of NMDARs in EAE. **Methods** Memantine, a non-competitive NMDAR antagonist which limits pathological activity of NMDARs while sparing normal synaptic activity, was administered orally from day 7 after immunization to 3- and 24-month-old female Dark Agouti rats. The animals were sacrificed at the peak of the disease. Spinal cord mononuclear cells were analyzed by flow cytometry. Brain tissue was collected for biochemical analysis of redox status and RT-qPCR. **Results** Semiprophylactic administration of memantine ameliorated clinical disease course, with greater effect in aged rats. Memantine reduced the number, frequency, and reactivation of CD4⁺ T lymphocytes and increased the relative percentage of CX3CR1-expressing microglia in spinal cord, but to a greater extent in aged rats. Additionally, analysis of brain redox status parameters showed that memantine was more effective in reducing superoxide anion radical, malondialdehyde and advanced oxidation protein products in aged rats than in young ones. In accordance with previous findings, NMDAR inhibition by memantine decreased NADPH oxidase and IL-1 β expression and increased the nuclear factor erythroid 2-related factor 2 and heme oxygenase-1 expression, to a greater extent in aged rats. **Conclusions** The involvement of NMDARs in the pathogenesis of EAE was age-dependent, being more pronounced in aged than in young rats.