

Combined dandelion extract and all-trans retinoic acid induces cytotoxicity in human breast cancer cells

Rezaie, H., Alipanah-Moghadam, R., Jeddi, F., Clark, C. C. T., Aghamohammadi, V. & Nemati, A

Published PDF deposited in Coventry University's Repository

Original citation:

Rezaie, H, Alipanah-Moghadam, R, Jeddi, F, Clark, CCT, Aghamohammadi, V & Nemati, A 2023, 'Combined dandelion extract and all-trans retinoic acid induces cytotoxicity in human breast cancer cells', Scientific Reports, vol. 13, no. 1, 15074. <https://doi.org/10.1038/s41598-023-42177-z>

DOI 10.1038/s41598-023-42177-z

ESSN 2045-2322

Publisher: Nature Research

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.



OPEN

Combined dandelion extract and all-trans retinoic acid induces cytotoxicity in human breast cancer cells

Hamed Rezaie^{1,5}, Reza Alipanah-Moghadam^{1,5}, Farhad Jeddi², Cain C. T. Clark³, Vahideh Aghamohammadi⁴ & Ali Nemat¹

Breast cancer is one of the most prevalent and deadly cancers among women worldwide. Recently, natural compounds have been widely used for the treatment of breast cancer. Present study evaluated antiproliferative and anti-metastasis activities of two natural compounds of dandelion and all-trans-retinoic acid (ATRA) in human MCF-7 and MDA-MB231 breast cancer cells. We also evaluated the expression of MMP-2, MMP-9, IL-1 β , p53, NM23 and KAI1 genes. Data showed a clear additive cytotoxic effect in concentrations of 40 μ M ATRA with 1.5 and 4 mg/ml of dandelion extract in MCF-7 and MDA-MB231 cells, respectively. In both cell lines, compared with the untreated cells, the expression levels of MMP-9 and IL-1 β were significantly decreased while p53 and KAI1 expression levels were increased. Besides, MMP-2 and NM23 had different expressions in the two studied cell lines. In conclusion, dandelion/ATRA co-treatment, in addition to having strong cytotoxic effects, has putative effects on the expression of anti-metastatic genes in both breast cancer cells.

Abbreviations

ATRA	All-trans retinoic acid
DEPC	Diethyl pyro carbonate
DMSO	Dimethylsulfoxide
FBS	Fetal Bovine Serum
IC50	Fifty percent inhibitory concentration
KAI1 (CD82)	Kangai-1 Protein
IL-1 β	Interleukin-1beta
MDM2	Murine double minute 2
MMP-2	Matrix Metalloproteinase -2
MMP-9	Matrix Metalloproteinase- 9
NF- κ B	Nuclear factor kappa B
NHPs	Natural health products
PBS	Phosphate buffered saline

Breast cancer is one of the most prevalent and deadly cancers among women worldwide, in both developed and developing countries¹. The incidence of breast cancer patients is predicted to reach 3,059,829 people by 2040², and every year, 2,300,000 new cases are diagnosed, with approximately 700,000 deaths yearly, accounting for nearly 6.9% of all cancer deaths³. The prevalence of breast cancer varies in different regions of the world, based on race and nationality. Various factors, such as genetics, age, diet, obesity, physical activity, environmental carcinogens, and sex hormones are involved in the etiology of this cancer^{4,5}. Breast cancer is characterized by the rapid proliferation of cancer cells and its invasion into healthy tissues, which can reduce patients' survivability⁴. Metastasis is the main leading cause of mortality in patients suffering from this cancer; thus, one of the main

¹Department of Clinical Biochemistry, School of Medicine, Ardabil University of Medical Sciences, Ardabil, Iran. ²Department of Genetics and Pathology, School of Medicine, Ardabil University of Medical Sciences, Ardabil, Iran. ³Centre for Intelligent Healthcare, Coventry University, Coventry CV1 5FB, UK. ⁴Department of Nutrition, Khalkhal University of Medical Sciences, Khalkhal, Iran. ⁵These authors contributed equally: Hamed Rezaie and Reza Alipanah-Moghadam. ✉email: alipanahreza9@gmail.com; r.alipanah@arums.ac.ir; farhadjeddi@gmail.com; f.jeddi@arums.ac.ir; v_agamohammadi@yahoo.com

defences against breast cancer is the controlling of cancer cells proliferation and prevention of its tissue invasion to identify novel targets for treatment^{6,7}. Various chemical drugs have been developed for breast cancer treatment, which have severe side effects, such as cardiac toxicity, nephrotoxicity, and neurotoxicity as well as drug resistance^{8–11}. Recently, the use of natural health products (NHPs), as well as plant extracts, as therapeutic agents to ameliorate a wide range of diseases, including cancer, has been considered. Moreover, 60% of the global population is habituated to the consumption of vegetables, fruits, herbs, and vitamins to prevent and/or remedy non-communicable diseases^{12,13}. Indeed, natural compounds, in addition to having very strong therapeutic effects, have far fewer side effects compared to chemical anticancer drugs^{14,15}. All-trans retinoic acid (ATRA), a derivative of vitamin A, is known as a strong anticancer agent causing growth inhibition, differentiation and apoptosis induction in several human cancers¹⁶. Moreover, several studies have identified an additive cytotoxic effect of ATRA combined with other natural compounds, such as caffeic acid, genistein, and curcumin, against various cancer cells^{17–19}.

Taraxacum officinale (Dandelion), belonging to Asteraceae family, has received a lot of attention in recent years. All dandelion organs, i.e. the roots, leaves, and flowers contain significant amounts of flavonoid compounds and L-chicoric acid, which have potent antioxidant, anti-inflammatory, and anti-cancer activities^{20,21}. Dandelion is used as traditional folk medicine throughout Asia, Europe and North America to improve outcomes in several diseases²². Empirical studies have shown that dandelion extract can inhibit the proliferation of cancer cells and induce death in them^{13,23,24}, and has strong anti-metastasis effects on cancer cells such as glioblastoma cells²⁵. In addition to breast cancer cells, it has been shown that dandelion potentially inhibit proliferation and migration of other cancer cells such as gastric cancer cells²⁶. Several studies have also shown that polyphenolic compounds in plant extracts can apply potential anticancer effects in combination with retinoic acid^{27–29}. Therefore, this study was designed to evaluate the cytotoxic effect of dandelion/ATRA combination. To achieve this goal, we used MCF-7 and MDA-MB231 well-established breast cancer cell lines which have different phenotypic/genotypic features such as progesterone and estrogen receptor status, different aspects of epithelial or mesenchymal type, multidrug resistance and invasiveness^{30,31}. Additionally, we aimed to evaluate the expression level of IL-1 β , as a gene that plays a role in cell proliferation, in human breast cancer cells. Thus, along with IL-1 β , its downstream genes, MMP-2 and MMP-9, expressions were assessed. To evaluate the anti-metastasis effects of dandelion/ATRA co-treatment, the expression of p53, NM23, and KAI1 genes, which are known as tumor metastasis suppressors, were measured.

Results

Effect of dandelion extract, ATRA, or dandelion/ATRA co-treatment on cell viability. MCF-7 cells were cultured in DMEM medium with 10% FBS and seeded in 96-well plates, after 24 h and when cell density reached 80–90%, the cells were treated as triplicates with previously mentioned concentrations of dandelion extract, ATRA, or the combination of dandelion extract and ATRA. After 48 h of treatment, cytoviability was assessed by the MTT test. Exposure of MCF-7 cells to all three compounds inhibited cell survival in a dose-dependent manner. The lowest cytotoxic concentrations of dandelion extract and ATRA on cell viability occurred in concentrations of 1 mg/ml and 1 μ M, respectively, as well as a combination of these concentrations (Fig. 1). Also, the highest cytotoxic concentrations of dandelion extract and ATRA on cell viability occurred in concentrations of 10 mg/ml and 100 μ M, respectively, as well as a combination of these concentrations. According to the dose–response curve, the IC₅₀ values for dandelion extract and ATRA were 1.69 mg/ml and 48 μ M, respectively. The concentrations of dandelion extract and ATRA used for the final treatment were considered at 1.5 mg/ml and 40 μ M, respectively (Fig. 1).

Similarly, the effect of dandelion extract, ATRA, or the combination of dandelion extract and ATRA, on cell viability of MDA-MB231 cells was studied. Based on the results of the MTT test, the lowest cytotoxic concentrations of dandelion extract and ATRA on cell viability occurred in concentrations of 1 mg/ml and 1 μ M, respectively, as well as a combination of these concentrations (Fig. 2). Furthermore, the highest cytotoxic concentrations of dandelion extract and ATRA on cell viability occurred in concentrations of 10 mg/ml and 200 μ M, respectively, as well as a combination of these concentrations. Using Sigma Plot software, the IC₅₀ values for dandelion extract and ATRA were calculated at 5.9 mg/ml and 149 μ M, respectively. The concentrations of 4 mg/ml dandelion extract and 40 μ M ATRA were used to evaluate their individual or combined effects on MDA-MB231 cells (Fig. 2).

Treatment concentrations for gene expression. In MCF-7 and MDA-MB231 cells, gene expression were assessed in cells treated with the concentrations of 1.5 and 4 mg/ml of dandelion extract, respectively, and constant concentration 40 μ M of ATRA. To determine gene expression in dandelion/ATRA co-treated cells, the concentrations of 1.5 mg/ml dandelion with 40 μ M ATRA and 4 mg/ml dandelion with 40 μ M ATRA were used in MCF-7 and MDA-MB231 cells, respectively.

Effect of dandelion extract or ATRA and dandelion/ATRA co-treatment on MMP-2 expression. In the MCF-7 cells, in all treated groups, MMP-2 expression level was significantly higher compared to the untreated control group ($p < 0.05$) (Fig. 3A). The highest MMP-2 mRNA expression was found in the dandelion extract-treated group, compared to the other groups ($p < 0.05$). Also, the MMP-2 expression level in the combined group was higher than the ATRA-treated group ($p < 0.05$). However, in the MDA-MB231 cells, the MMP-2 expression was shown to decrease significantly in all the treated groups, unlike in the untreated control group ($p < 0.05$). There is no significant difference in the MMP-2 mRNA expression among the treatment groups ($p > 0.05$) (Fig. 4A).






Figure 1. The effect of dandelion or ATRA and dandelion/ATRA co-treatment on MCF-7 cell viability. Viability was assessed with the MTT method after the exposure to the increasing concentrations of components for 48 h. The data are the mean \pm SD values of triplicate individual determinations. The lowest cytotoxic concentrations of dandelion extract and ATRA were observed in concentrations of 1 mg/ml and 1 μ M, respectively. Also, the highest cytotoxic concentrations of dandelion extract and ATRA on cell viability occurred in concentrations of 10 mg/ml and 100 μ M, respectively. To evaluate the effect of dandelion/ATRA co-treatment on cell viability, the concentration of ATRA was fixed at 40 μ M and the concentration of dandelion was considered variable between 0 and 1.5 mg/ml (* $P < 0.05$).




Figure 2. The effect of dandelion or ATRA and dandelion/ATRA co-treatment on MDA-MB231 cell viability. Viability was assessed with the MTT method after the exposure to the increasing concentrations of components for 48 h. The data are the mean \pm SD values of triplicate individual determinations. The minimum cytotoxic effects of dandelion and ATRA were seen in 1 mg/ml and 1 μ M concentrations, respectively. Moreover, the maximum cytotoxic effects of dandelion and ATRA were observed in 10 mg/ml and 200 μ M concentrations, respectively. To estimate the effect of dandelion/ATRA co-treatment on cell viability, the concentration of dandelion was considered variable between 0 to 4 mg/ml and the ATRA concentration was fixed at 40 μ M. (*P < 0.05).




Figure 3. The Effect of dandelion or ATRA and dandelion/ATRA co-treatment on gene expression in MCF-7 breast cancer cells. (A–F) Real time PCR method was done to calculate the expressions of MMP-2, MMP-9, IL-1 β , p53, NM23, and KAI1. The data is represented as the mean \pm SD of 3 independent determinations. (* $P < 0.05$).

Effect of dandelion extract, ATRA, or a combination of dandelion extract and ATRA, on MMP-9 expression. Our results indicated that MMP-9 expression in both MCF-7 and MDA-MB231 cells




Figure 4. The Effect of dandelion or ATRA and dandelion/ATRA co-treatment on gene expression in MDA-MB231 breast cancer cells. (A–F) The mRNA expressions of MMP-2, MMP-9, IL-1 β , p53, NM23, and KAI1 were carried out by quantitative real time PCR. The data are the mean \pm SD values of triplicate individual determinations. (* $P < 0.05$).

had a significant decrease in the treated groups compared to the untreated control group ($p < 0.05$). However, the expression of MMP-9 declined in the combination group in MCF-7 cells compared to the dandelion group

($p > 0.05$), but its association with the ATRA group was not significant ($p > 0.05$) (Fig. 3B). Likewise, in the MDA-MB231 cell line, the MMP-9 mRNA expression level was not significantly different between the treatment groups ($p > 0.05$) (Fig. 4B).

Effect of dandelion extract, ATRA, or a combination of dandelion extract and ATRA on IL-1 β expression. Based on our results, IL-1 β expression in both cell lines were decreased in the treatment groups with dandelion extract, ATRA, or their combination, compared to the untreated control group ($p < 0.05$). Interestingly, the reduction in IL-1 β expression level in MCF-7 cells in dandelion and combined groups was much greater than in the ATRA group ($p < 0.05$). However, dandelion elicited a further reduction in expression of IL-1 β compared to the combined group, although this was not significant ($p > 0.05$) (Fig. 3C). IL-1 β gene expression in the MDA-MB231 cell line was significantly reduced in the combined group compared to the ATRA and dandelion groups, moreover, ATRA elicited a further reduction in expression of IL-1 β compared to the dandelion group ($p < 0.05$) (Fig. 4C).

Effect of dandelion extract, ATRA, or a combination of dandelion extract and ATRA on p53 expression. Based on analysis of variance, p53 mRNA expression significantly increased in MCF-7 cells after incubation with dandelion extract, ATRA, or a combination of both, compared to the untreated control group ($p < 0.05$). Surprisingly, a significant increase in the p53 gene expression was detected in the combined group compared to both the dandelion and ATRA groups ($p < 0.05$). There were significant changes in the p53 mRNA levels between dandelion and ATRA groups ($p < 0.05$) (Fig. 3D). In MDA-MB231 cells, the rate of p53 mRNA expression levels was significantly increased in all treatment groups compared to the untreated control group ($p < 0.05$), however, the p53 expression level in dandelion and combined groups was similarly higher (Fig. 4D).

Effect of dandelion extract, ATRA, or a combination of dandelion extract and ATRA on NM23 expression. Our results showed that NM23 expression in MCF-7 cells, after 48 h of exposure to dandelion extract, ATRA, or a combination of both, decreased significantly compared to the untreated control group control group ($p < 0.05$). In addition, the dandelion and dandelion/ATRA co-treatment significantly reduced NM23 expression in MCF-7 cells compared to the ATRA group ($p < 0.05$). However, there was no significant difference in the NM23 mRNA levels between dandelion and combined groups ($p > 0.05$) (Fig. 3E). NM23 expression in MDA-MB231 cells decreased significantly in ATRA and dandelion/ATRA co-treatment, as compared to the untreated control group ($p < 0.05$). Additionally, dandelion significantly increased NM23 expression in MDA-MB231 cells compared to other groups ($p < 0.05$) (Fig. 4E).

Effect of dandelion extract, ATRA, or a combination of dandelion extract and ATRA on KAI1 expression. The results of the present study showed that the expression of KAI1 was increased in MDA-MB231 and MCF-7 cells after 48 h in treated groups compared to the untreated control group ($p < 0.05$). Furthermore, a significant increase in KAI1 expression was detected in MCF-7 cells after treatment by dandelion, compared to the ATRA and combined groups ($p < 0.05$). In MCF-7 cells no significant correlation was observed between ATRA and combined groups ($p > 0.05$) (Fig. 3F). In MDA-MB231 cells, the ATRA group significantly increased KAI1 expression, as compared to the dandelion and dandelion/ATRA co-treatment ($p < 0.05$). In addition, the combination of ATRA with dandelion significantly increased KAI1 expression in MDA-MB231 cells compared to the dandelion group ($p < 0.05$) (Fig. 4F).

As confirmed in our study, the combination of dandelion and ATRA elicited cell cytotoxicity following overexpression of p53 in both MCF-7 and MDA-MB231 breast cancer cells. The p53 gene, as a cell guard, is critically involved in programmed cell death, cell cycle regulation, and DNA repair, and, in addition to stimulating the apoptotic pathway, the p53 protein suppresses the expression of the anti-apoptotic Bcl-2 and Bcl-XL genes in its normal function³². Choi et al. studied the proapoptotic effect of ethanol extracts acquired from the dandelion flower on human ovarian cancer SK-OV-3 cells and stated that apoptosis induction by dandelion involved p53 activation, bax upregulation, and downregulation of Bcl-2³³. In the study by Chen et al. administration of dandelion leaf polysaccharide in H22 tumor-bearing mice directly induced tumor cell apoptosis via upregulation of p53 and pro-apoptotic protein Bax expression³⁴. George and Abrahamse used *Rubus fairholmanianus* derivatives against MCF-7 breast cancer cells and reported that overexpression of p53 in MCF-7 cells treated with these derivatives induced apoptosis in MCF-7 cells³⁵. Furthermore, Motadi et al. indicated the anticancer effects of *Tulbaghia violacea* extract in several cancer cell lines, such as MCF-7 and MDA-MB231 following p53 overexpression. The authors concluded that the p53-dependent pathway may be responsible for cell death³⁶. In addition, Won et al., using Sanggenol L, as a natural flavonoid, showed apoptosis in human prostate cancer cells can be directed via suppression of PI3K/Akt/mTOR signaling and cell cycle arrest via activation of p53³⁷. Previous studies have also evaluated the combination of ATRA with other anti-cancer agents. Wang et al. indicated that ATRA and decitabine in combination synergistically induce apoptosis in higher-risk myelodysplastic syndromes and elderly acute myeloid leukaemia cells³⁸. Another study also showed that there was an additive effect in leukemic cells when treated with a combination of ATRA and epigallocatechin-3-gallate (EGCG), as active polyphenolic compounds of green tea. The authors concluded that the cytotoxic effects of EGCG were enhanced by ATRA³⁹. In this study, we present a novel finding that shows an additive cytotoxic effect of the combination of dandelion with ATRA on breast cancer cells followed by overexpression of p53, which indicates the possible role of p53-related pathways in this process.

Along with the increase in the expression of p53, we observed a marked decrease in the expression of IL-1 β in MCF-7 and MDA-MB231 cells treated with dandelion extract, ATRA, or combination of both. It has been

shown that IL-1 β level is associated with the progression and stage of breast cancer^{40,41}. In this regard, it has been reported that IL-1 β progresses tumorigenesis through attenuation of p53 protein expression and promotion of the growth of different types of cancer cells by nuclear factor kappa B (NF- κ B) activation^{42–44}. Suppression of IL-1 β expression decrease tumor growth and inhibit perfusion of tumor cells from the primary location into circulation^{45–48}. Therefore, regarding central role of IL-1 β in growth and proliferation of breast cancer cells, our findings may be defined as a part of the mechanism of cytotoxicity by dandelion extract, ATRA, or the combination of both, on breast cancer cells, MCF-7, and MDA-MB231. Several studies indicated the anti-inflammatory effects of dandelion extract^{49,50}. In the study by Liu et al. dandelion extract significantly decreased the expression of IL-1 β , IL-6, and TNF- α , and reduced the LPS-induced inflammatory response⁵¹. Studies have indicated that some of compounds in dandelion extract such as phenolic acids, flavonoids, triterpenes and sterols have potential therapeutic effects in inflammation^{51–55}. Moreover, anti-inflammatory effect of ATRA has been demonstrated in previous studies^{56,57}. However, Guo et al. reported that ATRA may stimulate the expression of IL-1 β by activating NF- κ B signalling and caspase-1 in macrophages⁵⁸. Furthermore, a study indicated that the anti-proliferative effect of ATRA is associated with selective stimulation of IL-1 β , a cytokine that directly prevents growth of lung cancer cells⁵⁹.

Based on our results, dandelion, alone or combined with ATRA, reduced MMP-2 and MMP-9 expressions in MDA-MB231 cells, while also reducing MMP-9 expression in MCF-7 cells, and increasing MMP-2 expression in these cells. Despite that different genes contribute to the control of metastasis, the role of matrix metalloproteinases (MMPs) are more pronounced. MMPs are a group of zinc-containing enzymes that play a key role in various processes, such as inflammation, angiogenesis, tissue regeneration, and invasion of cancer cells into other tissues. Among the MMPs, MMP-2 and 9 play the most prominent role in metastasis. Several studies have also shown that MMP-2 and MMP-9 play a key role in the invasion of cancer cells, and their increased expression is directly related to the malignancy and aggressiveness of cancer cells^{60,61}. Furthermore, it has been shown that the attenuation of MMP-2 and 9 expressions can greatly reduce the invasiveness of cancer cells. Moreover, it has been found that MMP-9 is expressed in small amounts by healthy breast tissue, but it notably increases in malignant breast cancer cells as well as in the metastatic stage of breast cancer^{62,63}. On the other hand, MMP-9 expression reduction can inhibit the aggressive malignancy of cancer cells⁶⁴. Moreover, multiple studies emphasized the role of MMP-2 in metastasis in various cancers, including breast cancer^{65,66}. Furthermore, other natural products, such as curcumin, quercetin, and polyphenol, inhibit MMP-2 and MMP-9 expressions and tumor metastasis⁶⁷. There are several studies confirming our results regarding MMP-9 expression, such as Liang et al. and Liu et al., who showed that ATRA decreased the expression of MMP-9 in glioma and MDA-MB-231 cells^{68,69}. However, consistent with the results obtained in our study in MCF-7 cells, which showed that the expression of MMP-2 was increased in the treatment groups, Malmira et al. showed that colorectal cancer HCT-116 cells treated using ginger extract, significantly increased MMP-2 expression level in a dose-dependent manner⁷⁰. Furthermore, it has been shown that ATRA has an effective role in the regulation of MMP-2 or MMP-9 expression in different cells⁷¹. Although, there are conflicting reports in the literature regarding the effect of ATRA on the expression of MMP-2 and 9, such that, in some studies, the expression of MMP-2 and MMP-9 was inhibited by ATRA. However, in other studies, the expression of MMP-9 was reduced and the expression of MMP-2 remained unchanged, whilst another study reported that both MMP-2 and MMP-9 expression were elevated^{68,72,73}. Consistent with the results obtained in our study in MCF-7 cells, Vu et al. showed that ATRA increases MMP-2 expression and secretion in human myeloid leukemia THP-1 cells⁷⁴. Also, Kim et al., by investigating the effect of ATRA on the expression of MMP-2 and 9 in human dental pulp cells (HDPCs), showed that ATRA decreases the expression of MMP-2 in these cells, but has no effect on the expression of MMP-9⁷⁵. The amplification effect of ATRA, in combination with other natural products including curcumin and vitamin D, on cell death and the invasion of cancer cells has also been investigated in various studies^{19,76}. Overall, the effect of dandelion, or its combination with ATRA, on the expressions of MMP-2 and MMP-9 in MDA-MB231 and MCF-7 cells should be considered a novel finding that can be used in cancer treatment strategies by conducting further research. Another finding in our study is the significant increase in KAI1 expression in both MCF-7 and MDA-MB231 cells in all treatments consisting of dandelion, ATRA, or the combination of ATRA with dandelion. The KAI1 or CD82 gene is a major tumor metastasis suppressor that primarily inhibits cancer cell motility and invasiveness in a wide variety of solid malignant tumors, such as prostate, stomach, colon, hepatocarcinoma, thyroid, pancreas, and breast cancers. KAI1/CD82 may be regarded as a main prognostic biomarker in anticipating metastatic development and maybe a hopeful candidate for therapeutic purposes in cancers^{77,78}. The expression of KAI1 is abundant in normal breast tissue, but its expression is reduced in primary tumors, as well as in the metastatic stage of breast cancer⁷⁹. KAI1 inhibits cancer cell migration and adhesion, in addition to apoptosis, by preventing integrin-mediated signalling⁸⁰. Moreover, KAI1 upregulates tissue inhibitor of metalloproteinase 1 (TIMP1), which causes the inactivation of MMP-9. Restored KAI1 expression is significantly associated with loss of different cancer cell proliferation, such as ovary, stomach, pancreas, and breast^{77,79}. Thus, the increase in the expression of KAI1 in our treatment groups can indicate the effect of dandelion and ATRA on reducing the aggressiveness of breast cancer cells. Also, interestingly, our results showed a reducing effect of ATRA, or its combination with dandelion, on NM23 expression in both MCF-7 and MDA-MB231 cells. NM23 is considered as a potential anti-invasive and tumor-metastasis suppressor, moreover, it has been shown that low NM23 expression may be associated with a poor prognosis in breast cancer⁸¹. However, in our study, elevated expression in NM23 was observed only in the dandelion-treated group in MDA-MB-231 cells, as compared to the control group, which should be studied so to elucidate possible mechanistic actions. Overall, it appears that dandelion, or its combination with ATRA, can be very effective in inducing cell death and controlling the main agents involved in the invasion of breast cancer cells.

The present study highlighted that dandelion extract, in combination with ATRA, in addition to having strong cytotoxic effects on breast cancer cells, also has potential effects on the expression of genes involved in the proliferation and invasion of MCF-7 and MDA-MB231 breast cancer cells.

Materials and methods

Materials. MCF-7 and MDA-MB231 human breast cancer cell lines were obtained from the Pasteur Institute of Iran, Tehran. DMEM (Dulbecco's Modified Eagle Medium) (Cat. No: 12800-116), FBS (Fetal Bovine Serum cell culture) (Cat. No: 10270), penicillin/streptomycin (Cat. No: 15140), Glutamax (Cat. No: 25030) and non-essential amino acids (Cat. No: 11140050) were purchased from Thermo Fisher Scientific Inc. (UK). Cell Proliferation Kit (MTT) (Cat. No: M2128), DMSO (Dimethylsulfoxide) (Cat. No: D4540), and ATRA (Cat. No: R2625-50 Mg) were purchased from Sigma-Aldrich, USA. Trypsin (X0930) was purchased from Biowest, France. Dandelion root was prepared from the Namin region of Ardabil province, Iran (Latitude of 38°25' 28.04" N, Longitude 48°28' 41.67" E). Plant material was collected in accordance with applicable national and international guidelines⁸². Plant specimens were identified by the Research Institute of Forests and Rangelands (RIFR), Tehran, Iran. Voucher specimens were deposited in the Research Institute of Forests and Rangelands (RIFR), Tehran, Iran with voucher Numbers: TARRI 107642.

Preparation of hydro alcoholic extract of dandelion root. Briefly, the root of the plant was washed with water and dried in air. After complete drying, it was gently powdered by an electric grinder. Then, obtained powder was mixed with 70% ethanol in a ratio of 1:10 w/v. The mixture was then shaken in room temperature for 72 h. The mixture was then filtered using Whatman No. 42 filter paper and dried at 50 °C. The extract obtained were then stored in the dark in a freezer at – 70 °C until use.

Cell culture. MCF-7 and MDA-MB231 cells were grown and maintained in DMEM supplemented with 10% FBS, which added penicillin/streptomycin (P/S), and kept in a humid atmosphere at 37 °C with 5% CO₂.

Tetrazolium (MTT) assay for measuring cellular viability and drug treatment. 3-(4,5)-Dimethylimidazole (-Zyl)-3, 5-diphenyltetrazolium bromide used for assessment of cell proliferation and cell viability. In this method, the metabolic activity of the cells is determined by reducing the yellow tetrazole to purple formazan in living cells by the action of the mitochondrial succinate dehydrogenase enzyme. MCF-7 and MDA-MB231 cells were cultured in a 96-well plate with a density of 5000 cells per well. After 24 h, MCF-7 cells were exposed to increasing concentrations of dandelion extract (0–10 mg/ml), ATRA (0–100 μM), or the combination of dandelion extract and ATRA (in the same concentration ranges) to determine cell viability. Similarly, MDA-MB231 cells were exposed to increasing concentrations of dandelion extract (0–10 mg/ml), ATRA (0–200 μM) or the combination of dandelion extract and ATRA (in the same concentration ranges) to determine cell viability. Next, plates containing cells were incubated for 48 h at 37 °C in a CO₂ atmosphere. After 48 h, the medium was removed from the wells and culture medium containing 5 mg/ml of MTT solution was added to each well and cells were incubated for 4 h at 37 °C with 5% CO₂. DMSO was then added to the wells to dissolve formazan crystals by shaking them for 15 min. The absorbance was measured at 570 nm on an ELISA reader (Synergy HT, USA Biotech, USA). Cell viability at all concentrations was calculated relative to untreated control and as a percentage. It was repeated three times for all groups. Fifty percent inhibitory concentration (IC₅₀) was calculated using Sigma Plot Software version 14. In this study, different concentrations of dandelion and ATRA were evaluated in combined therapy, and based on the Bliss independence equation the results showed that the most effective combined dose was the concentration lower than IC₅₀.⁸³

Real-time PCR. Briefly, RNA isolation and qPCR studies were conducted according to the MIQE guideline standards⁸⁴. Total RNA was extracted from cultured cells, after 48 h treatment, using Trizol solution as recommended by the manufacturer (Biobasic, Canada). RNA integrity was confirmed by gel electrophoresis on a denaturing 1.5% agarose gel (28S/18S rRNA ratio). Nucleic acid purity and quantity was evaluated on a

Gene	Gen ID	Primer sequences (5' to 3')	Product size (bp)
MMP2	4313	F: GGC GAT GGA TAC CCC TTT GAC	114
		R: TCC CAA GGT CCA TAG CTC	
MMP9	4318	F: CTG GGC AGA TTC CAA ACC TTT	91
		R: GCG GCA AGT CTT CCG AGT AG	
IL-1β	3553	F: CTG GTA CAT CAG CAC CTC TCA	134
		R: GCA CAG GAC TCT CTG GGT ACA	
p53	7157	F: CACAGCACATGACGGAGGTT	70
		R: GCCAGACCATCGCTATCTGA	
NM23	4830	F: TCATGCTCGGGGAGACCAA	126
		R: CGATCTCCTTCTCTGCACTCT	
KAI1	3732	F: GGGAAGAGGACAACAGCCTTT	112
		R: ATGCAGCCCTCCTGGTACAC	
β-actin	60	F: ACAGAGCCTCGCCTTTGC	76
		R: CGCGGCGATATCATCATCCA	

Table 1. Primers and sequencing information.

NanoDrop (Thermo Scientific Nanodrop 2000c, USA) where the A260/A280 & A260/A230 ratios were acquired. RNAs with a ratio of 1.7–2.1 were used for gene expression analysis. Three microgram total RNA was reverse transcribed using SMOBIO Kit (RP1300, Taiwan) in 20 µl volume according to the manufacturer's procedure. The cDNA was diluted 1:5 in nuclease free water and then real-time PCR was performed by using SYBR Green PCR master mix (Takara TB Green Premix Ex Taq II, RR820Q, Japan). All experiments were conducted on a Real-time PCR Detection System (Ruche light cyclor 96, Germany). The PCR reaction was performed in a total volume of 20 µl (10 µl master mix, 1 µl cDNA, 1 µl primer Forward, 1 µl primer Reverse, and 8 µl nuclease free water). The real-time PCR protocol for evaluating each gene expression is provided in the Supplementary Table S1. Oligo Primer Analysis Software (version 7.60) was used for primer designing. Primers and sequencing information are shown in Table 1. For all target and housekeeping genes considered, standard curves were calculated with serial tenfold dilutions of control cDNA to determine amplification *efficiency*. Product specificity was assessed by melting curve analyses and the non-template control (NTC) and genomic DNA (gDNA) samples. Only primer pairs which did not result in more than one melting peak or any sigmoid curve in the NTC and gDNA samples were used for final analyses. Beta-actin (ACTB), as a housekeeping gene, was used as an internal control for normalization. The relative mRNA expression levels were obtained according to the $2^{-\Delta\Delta Ct}$ method⁸⁵. Each measurement was performed in triplicate.

Data availability

All data and materials are fully presented in the manuscript.

Received: 3 April 2023; Accepted: 6 September 2023

Published online: 12 September 2023

References

- Wang, Z. *et al.* DNER promotes epithelial–mesenchymal transition and prevents chemosensitivity through the Wnt/ β -catenin pathway in breast cancer. *Cell Death Dis.* **11**, 1–16 (2020).
- Hashemi, S.-M. *et al.* Health-related quality of life of breast cancer patients in the Eastern Mediterranean region: A systematic review and meta-analysis. *Breast Cancer Res. Treat.* **174**, 585–596 (2019).
- Sung, H. *et al.* Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA* **71**, 209–249 (2021).
- Ullah, M. F. Breast cancer: Current perspectives on the disease status. In *Breast Cancer Metastasis and Drug Resistance* 51–64 (Springer, 2019).
- Momenimovahed, Z. & Salehiniya, H. Epidemiological characteristics of and risk factors for breast cancer in the world. *Breast Cancer* **11**, 151 (2019).
- Wen, X., Chen, Y. & Fang, X. Overexpression of HOXD8 inhibits the proliferation, migration and invasion of breast cancer cells by downregulating ILP2 expression. *Exp. Ther. Med.* **22**, 1–9 (2021).
- Qi, L., Sun, B., Yang, B. & Lu, S. circHIPK3 (hsa_circ_0000284) promotes proliferation, migration and invasion of breast cancer cells via miR-326. *Oncol. Targets. Ther.* **14**, 3671 (2021).
- Alkahtani, S. *et al.* Poly lactic-co-glycolic acid-(PLGA-) loaded nanoformulation of cisplatin as a therapeutic approach for breast cancers. *Oxid. Med. Cell. Longev.* **2021**, 1–8 (2021).
- Najjary, S. *et al.* Role of miR-21 as an authentic oncogene in mediating drug resistance in breast cancer. *Gene* **738**, 144453 (2020).
- Hasna, J. *et al.* Orai3 calcium channel and resistance to chemotherapy in breast cancer cells: The p53 connection. *Cell Death Differ.* **25**, 693–707 (2018).
- Muley, H., Fadó, R., Rodríguez-Rodríguez, R. & Casals, N. Drug uptake-based chemoresistance in breast cancer treatment. *Biochem. Pharmacol.* **177**, 113959 (2020).
- Mishra, A. & Meherotra, R. Head and neck cancer: Global burden and regional trends in India. *Asian Pac. J. Cancer Prev.* **15**, 537–550 (2014).
- Rehman, G. *et al.* Effect of methanolic extract of dandelion roots on cancer cell lines and AMP-activated protein kinase pathway. *Front. Pharmacol.* **8**, 875 (2017).
- Akram, R. *et al.* In *Medicinal Plants for Lung Diseases* 229–256 (Springer, 2021).
- Jiang, W. *et al.* Remodeling the epigenetic landscape of cancer—application potential of flavonoids in the prevention and treatment of cancer. *Front. Oncol.* **11**, 2331 (2021).
- Patrad, E., Niapour, A., Farassati, F. & Amani, M. Combination treatment of all-trans retinoic acid (ATRA) and γ -secretase inhibitor (DAPT) cause growth inhibition and apoptosis induction in the human gastric cancer cell line. *Cytotechnology* **70**, 865–877 (2018).
- Maity, S. *et al.* Caffeic acid, a dietary polyphenol, as a promising candidate for combination therapy. *Chem. Pap.* **76**, 1271–1283 (2022).
- Dastjerdi, M. N., Zamani, S., Mardani, M. & Beni, B. H. All-trans retinoic acid and genistein induce cell apoptosis in OVCAR-3 cells by increasing the P14 tumor suppressor gene. *Res. Pharm. Sci.* **11**, 505 (2016).
- Thulasiraman, P., McAndrews, D. J. & Mohiudddin, I. Q. Curcumin restores sensitivity to retinoic acid in triple negative breast cancer cells. *BMC Cancer* **14**, 1–14 (2014).
- Lis, B., Jedrejek, D., Moldoch, J., Stochmal, A. & Olas, B. The anti-oxidative and hemostasis-related multifunctionality of L-chicoric acid, the main component of dandelion: An in vitro study of its cellular safety, antioxidant and anti-platelet properties, and effect on coagulation. *J. Funct. Foods* **62**, 103524 (2019).
- Liu, N. *et al.* The effects of solid-state fermentation on the content, composition and in vitro antioxidant activity of flavonoids from dandelion. *PLoS ONE* **15**, e0239076 (2020).
- Kania-Dobrowolska, M. & Baraniak, J. Dandelion (*Taraxacum officinale* L.) as a source of biologically active compounds supporting the therapy of co-existing diseases in metabolic syndrome. *Foods* **11**, 2858 (2022).
- Nguyen, C. *et al.* Dandelion root and lemongrass extracts induce apoptosis, enhance chemotherapeutic efficacy, and reduce tumour xenograft growth in vivo in prostate cancer. *Evid.-Based Complement. Altern. Med.* **2019**, 1–12 (2019).
- Husieva, O. & Baranovsky, M. The impact of *Taraxacum officinale* aqueous extracts on WISH cancer cell line. *Problems of Environmental Biotechnology* (2019).
- Rashidi, I. *et al.* Anti-invasion and anti-metastasis effects of dandelion (*Taraxacum officinale*) hydroalcoholic extract on glioblastoma multiforme cell line model. *WCRJ* **9**, e2361 (2022).
- Zhu, H. *et al.* Dandelion root extract suppressed gastric cancer cells proliferation and migration through targeting lncRNA-CCAT1. *Biomed. Pharmacother.* **93**, 1010–1017 (2017).

27. Lee, J. H., Kishikawa, M., Kumazoe, M., Yamada, K. & Tachibana, H. Vitamin A enhances antitumor effect of a green tea polyphenol on melanoma by upregulating the polyphenol sensing molecule 67-kDa laminin receptor. *PLoS ONE* **5**, e11051 (2010).
28. Chen, M.-C., Hsu, S.-L., Lin, H. & Yang, T.-Y. Retinoic acid and cancer treatment. *BioMedicine* **4**, 22. <https://doi.org/10.7603/s40681-014-0022-1> (2014).
29. Panda, A. K., Chakraborty, D., Sarkar, I., Khan, T. & Sa, G. New insights into therapeutic activity and anticancer properties of curcumin. *J. Exp. Pharmacol.* **9**, 31–45 (2017).
30. Theodossiou, T. A. *et al.* Simultaneous defeat of MCF7 and MDA-MB-231 resistances by a hypericin PDT–tamoxifen hybrid therapy. *npj Breast Cancer* **5**, 13. <https://doi.org/10.1038/s41523-019-0108-8> (2019).
31. Chen, J., Lu, L., Wang, H., Dai, L. & Zhang, P. PKD2 mediates multi-drug resistance in breast cancer cells through modulation of P-glycoprotein expression. *Cancer Lett.* **300**, 48–56 (2011).
32. Wu, G. S. & Ding, Z. Caspase 9 is required for p53-dependent apoptosis and chemosensitivity in a human ovarian cancer cell line. *Oncogene* **21**, 1–8 (2002).
33. Choi, E. J. & Kim, G.-H. Dandelion (*Taraxacum officinale*) flower ethanol extract inhibits cell proliferation and induces apoptosis in human ovarian cancer SK-OV-3 cells. *Food Sci. Biotechnol.* **18**, 552–555 (2009).
34. Chen, P. *et al.* Protective effect of the polysaccharides from taraxacum mongolicum leaf by modulating the p53 signaling pathway in H22 tumor-bearing mice. *Foods* **11**, 3340 (2022).
35. George, B. P. & Abrahamse, H. Increased oxidative stress induced by rubus bioactive compounds induce apoptotic cell death in human breast cancer cells. *Oxid. Med. Cell. Longev.* **2019**, 1–18 (2019).
36. Motadi, L. R., Choene, M. S. & Mthembu, N. N. Anticancer properties of Tulbaghia violacea regulate the expression of p53-dependent mechanisms in cancer cell lines. *Sci. Rep.* **10**, 1–11 (2020).
37. Won, Y.-S. & Seo, K.-I. Sanggenol L induces apoptosis and cell cycle arrest via activation of p53 and suppression of PI3K/Akt/mTOR signaling in human prostate cancer cells. *Nutrients* **12**, 488 (2020).
38. Wang, L. *et al.* All-trans retinoic acid enhances the cytotoxic effect of decitabine on myelodysplastic syndromes and acute myeloid leukaemia by activating the RAR α -Nrf2 complex. *Br. J. Cancer* **128**, 691–701 (2022).
39. Ly, B. T. K. & Chi, H. T. Combined effect of (-)-epigallocatechin-3-gallate and all-trans retinoic acid in FLT3-mutated cell lines. *Biomed. Rep.* **13**, 1–1 (2020).
40. Diep, S., Maddukuri, M., Yamauchi, S., Geshow, G. & Delk, N. A. Interleukin-1 and nuclear factor kappa B signaling promote breast cancer progression and treatment resistance. *Cells* **11**, 1673 (2022).
41. Rébé, C. & Ghiringhelli, F. Interleukin-1 β and cancer. *Cancers* **12**, 1791 (2020).
42. Li, L. & Hong, Z. IL-1 β /NF- κ B signaling promotes colorectal cancer cell growth through miR-181a/PTEN axis. *Arch. Biochem. Biophys.* **604**, 20–26 (2016).
43. Schauer, I. G. *et al.* Interleukin-1 β promotes ovarian tumorigenesis through a p53/NF- κ B-mediated inflammatory response in stromal fibroblasts. *Neoplasia* **15**, 409–418 (2013).
44. Mürköster, S. S. *et al.* Acquired chemoresistance in pancreatic carcinoma cells: Induced secretion of IL-1 β and NO lead to inactivation of caspases. *Oncogene* **25**, 3973–3981 (2006).
45. Xia, Y. *et al.* Piperine inhibits IL-1 β -induced IL-6 expression by suppressing p38 MAPK and STAT3 activation in gastric cancer cells. *Mol. Cell. Biochem.* **398**, 147–156 (2015).
46. Jin, H., Jin, X., Cao, B. & Wang, W. Berberine affects osteosarcoma via downregulating the caspase-1/IL-1 β signaling axis. *Oncol. Rep.* **37**, 729–736 (2017).
47. Zhang, J. & Veeramachaneni, N. Targeting interleukin-1 β and inflammation in lung cancer. *Biomark. Res.* **10**, 1–9 (2022).
48. Safari, H. *et al.* Decrease of intracellular ROS by arbutin is associated with apoptosis induction and downregulation of IL-1 β and TNF- α in LNCaP; prostate cancer. *J. Food Biochem.* **44**, e13360 (2020).
49. Seo, S.-W. *et al.* Taraxacum officinale protects against cholecystokinin-induced acute pancreatitis in rats. *World J. Gastroenterol.* **WJG** **11**, 597 (2005).
50. Badr, A., Fouad, D. & Attia, H. Insights into protective mechanisms of dandelion leaf extract against cisplatin-induced nephrotoxicity in rats: Role of inhibitory effect on inflammatory and apoptotic pathways. *Dose-Response* **17**, 1559325819874897 (2019).
51. Liu, Q. *et al.* Effects of dandelion extract on the proliferation of rat skeletal muscle cells and the inhibition of a lipopolysaccharide-induced inflammatory reaction. *Chin. Med. J.* **131**, 1724–1731 (2018).
52. Park, C. M., Park, J. Y. & Song, Y. S. Luteolin and chicoric acid, two major constituents of dandelion leaf, inhibit nitric oxide and lipid peroxide formation in lipopolysaccharide-stimulated RAW 264.7 cells. *J. Food Sci. Nutr.* **15**, 92–97 (2010).
53. Budzianowski, J. Coumarins, caffeoyltartaric acids and their artifactual methyl esters from Taraxacum officinale leaves. *Planta Med.* **63**, 288–288 (1997).
54. Shan, J. *et al.* Chlorogenic acid inhibits lipopolysaccharide-induced cyclooxygenase-2 expression in RAW264.7 cells through suppressing NF- κ B and JNK/AP-1 activation. *Int. Immunopharmacol.* **9**, 1042–1048 (2009).
55. Banno, N. *et al.* Anti-inflammatory activities of the triterpene acids from the resin of Boswellia carteri. *J. Ethnopharmacol.* **107**, 249–253 (2006).
56. Li, S., Lei, Y., Lei, J. & Li, H. All-trans retinoic acid promotes macrophage phagocytosis and decreases inflammation via inhibiting CD14/TLR4 in acute lung injury. *Mol. Med. Rep.* **24**, 1–11 (2021).
57. He, T.-R. *et al.* All-trans retinoic acid-incorporated glycol chitosan nanoparticles regulate macrophage polarization in Pg-LPS-induced inflammation. *Curr. Med. Sci.* **42**, 974–980 (2022).
58. Li, G., Yan, Z., Wen-ping, L., Tian-yu, Z. & De-qin, Y. Regulatory effect of all-trans retinoic acid on the expression of IL-1 β in macrophages and the mechanisms involved. *J. Sichuan Univ.* **53**, 444–451 (2022).
59. Ross, H. J. The antiproliferative effect of trans-retinoic acid is associated with selective induction of interleukin-1 beta, a cytokine that directly inhibits growth of lung cancer cells. *Oncol. Res.* **8**, 171–178 (1996).
60. Das, S., Amin, S. A. & Jha, T. Inhibitors of gelatinases (MMP-2 and MMP-9) for the management of hematological malignancies. *Eur. J. Med. Chem.* **223**, 113623 (2021).
61. Wang, R. *et al.* GOLPH3 overexpression is closely correlated with poor prognosis in human non-small cell lung cancer and mediates its metastasis through upregulating MMP-2 and MMP-9. *Cell. Physiol. Biochem.* **35**, 969–982 (2015).
62. Winkler, J., Abisoye-Ogunniyan, A., Metcalf, K. J. & Werb, Z. Concepts of extracellular matrix remodelling in tumour progression and metastasis. *Nat. Commun.* **11**, 1–19 (2020).
63. Huang, H. Matrix metalloproteinase-9 (MMP-9) as a cancer biomarker and MMP-9 biosensors: Recent advances. *Sensors* **18**, 3249 (2018).
64. Yin, P. *et al.* MMP-9 knockdown inhibits oral squamous cell carcinoma lymph node metastasis in the nude mouse tongue-xenografted model through the RhoC/Src pathway. *Anal. Cell. Pathol.* **2021**, 1–16 (2021).
65. Maybee, D. V., Ink, N. L. & Ali, M. A. Novel roles of MT1-MMP and MMP-2: Beyond the extracellular milieu. *Int. J. Mol. Sci.* **23**, 9513 (2022).
66. Nakopoulou, L. *et al.* MMP-2 protein in invasive breast cancer and the impact of MMP-2/TIMP-2 phenotype on overall survival. *Breast Cancer Res. Treat.* **77**, 145–155 (2003).
67. Li, L.-H. *et al.* Methanol extract of Elaeagnus glabra, a Korean medicinal plant, inhibits HT1080 tumor cell invasion. *Oncol. Rep.* **21**, 559–563 (2009).

68. Liu, H., Zang, C., Fenner, M., Possinger, K. & Elstner, E. PPAR γ ligands and ATRA inhibit the invasion of human breast cancer cells in vitro. *Breast Cancer Res. Treat.* **79**, 63–74 (2003).
69. Liang, C., Yang, L. & Guo, S. All-trans retinoic acid inhibits migration, invasion and proliferation, and promotes apoptosis in glioma cells in vitro. *Oncol. Lett.* **9**, 2833–2838 (2015).
70. Malmir, S., Ebrahimi, A. & Mahjoubi, F. Effect of ginger extracts on colorectal cancer HCT-116 cell line in the expression of MMP-2 and KRAS. *Gene Rep.* **21**, 100824 (2020).
71. Zhou, T.-B. & Qin, Y.-H. The potential mechanism for the different expressions of gelatinases induced by all-trans retinoic acid in different cells. *J. Recept. Signal Transduct.* **32**, 129–133 (2012).
72. Lei, F.-Y., Zhou, T.-B., Qin, Y.-H., Chen, X.-P. & Li, Z.-Y. Potential signal pathway of all-trans retinoic acid for MMP-2 and MMP-9 expression in injury podocyte induced by adriamycin. *J. Recept. Signal Transduct.* **34**, 378–385 (2014).
73. Wu, G.-D. *et al.* The role of ZFP580 in the regulation of rat VSMCs migration by ATRA and its mechanism. *Zhongguo Ying Yong Sheng li xue za zhi Zhongguo Yingyong Shenglixue Zazhi Chin. J. Appl. Physiol.* **36**, 45–50 (2020).
74. Vu, H. T., Hoang, T. X. & Kim, J. Y. All-trans retinoic acid enhances matrix metalloproteinase 2 expression and secretion in human myeloid leukemia THP-1 cells. *BioMed Res. Int.* **2018**, 1–10 (2018).
75. Kim, J. M. *et al.* Inhibition of matrix metalloproteinases expression in human dental pulp cells by all-trans retinoic acid. *Int. J. Oral Sci.* **6**, 150–153 (2014).
76. Elmaci, I., Ozpinar, A., Ozpinar, A., Perez, J. L. & Altinoz, M. A. From epidemiology and neurometabolism to treatment: Vitamin D in pathogenesis of glioblastoma Multiforme (GBM) and a proposal for Vitamin D+ all-trans retinoic acid+ Temozolomide combination in treatment of GBM. *Metab. Brain Dis.* **34**, 687–704 (2019).
77. Liu, W. M. & Zhang, X. A. KAI1/CD82, a tumor metastasis suppressor. *Cancer Lett.* **240**, 183–194 (2006).
78. Yan, W., Huang, J., Zhang, Q. & Zhang, J. Role of metastasis suppressor KAI1/CD82 in different cancers. *J. Oncol.* **2021**, 1–8 (2021).
79. Feng, J. *et al.* Tetraspanin CD82: A suppressor of solid tumors and a modulator of membrane heterogeneity. *Cancer Metastasis Rev.* **34**, 619–633 (2015).
80. He, X. *et al.* The peptide mimicking small extracellular ring domain of CD82 inhibits tumor cell migration in vitro and metastasis in vivo. *J. Cancer Res. Clin. Oncol.* **147**, 1927–1934 (2021).
81. Mao, H. *et al.* Loss of nm23 expression predicts distal metastases and poorer survival for breast cancer. *Int. J. Oncol.* **18**, 587–591 (2001).
82. IUCN. (IUCN, Gland and Cambridge, 2001).
83. Fouquier, J. & Guedj, M. Analysis of drug combinations: Current methodological landscape. *Pharmacol. Res. Perspect.* **3**, e00149 (2015).
84. Bustin, S. A. *et al.* (Oxford University Press, 2009).
85. Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time quantitative PCR and the 2 $^{-\Delta\Delta CT}$ method. *Methods* **25**, 402–408 (2001).

Acknowledgements

We hereby express our gratitude to the Ardabil University of Medical Sciences for funding this research.

Author contributions

R.A.M., F.J. and H.R. contributed to the conception and design of the study and the study protocol. R.A.M., F.J. and H.R. managed the running of the study. R.A.M., F.J. and A.N. conducted data analysis and all authors helped with data interpretation. R.A.M., C.C., and V.A. wrote this manuscript with input from all co-authors. All authors read and approved the final version of the manuscript.

Funding

This study is financially supported by the Ardabil university of medical sciences.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-023-42177-z>.

Correspondence and requests for materials should be addressed to R.A.-M., F.J. or V.A.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023