



岐阜大学機関リポジトリ

Gifu University Institutional Repository

Title	4-Hydroxy-4-methyl-2,6-diphenyl-5,6-dihydro-4H-1,3-thiazine(本文(Fulltext))
Author(s)	KOKETSU, Mamoru; EBIHARA, Masahiro; ISHIHARA, Hideharu
Citation	[Acta Crystallographica Section E-Structure Reports Online] vol.[62] no.[Part4] p.[O1218]-[O1220]
Issue Date	2006-04
Rights	International Union of Crystallography
Version	出版社版 (publisher version) postprint
URL	http://hdl.handle.net/20.500.12099/28358

この資料の著作権は、各資料の著者・学協会・出版社等に帰属します。

Mamoru Koketsu,^{a*} Masahiro
Ebihara^b and Hideharu Ishihara^b^aDivision of Instrumental Analysis, Life Science
Research Centre, Gifu University, Yanagido,
Gifu 501-1193, Japan, and ^bDepartment of
Chemistry, Faculty of Engineering, Gifu
University, Yanagido, Gifu 501-1193, JapanCorrespondence e-mail:
koketsu@cc.gifu-u.ac.jp

Key indicators


Single-crystal X-ray study
 $T = 190$ K
Mean $\sigma(C-C) = 0.003$ Å
 R factor = 0.046
 wR factor = 0.102
Data-to-parameter ratio = 14.5For details of how these key indicators were
automatically derived from the article, see
<http://journals.iucr.org/e>.4-Hydroxy-4-methyl-2,6-diphenyl-5,6-
dihydro-4*H*-1,3-thiazineIn the title compound, $C_{17}H_{17}NOS$, the phenyl ring at position
6 of the thiazine ring is *trans* to the hydroxy group. The
thiazine ring is in a sofa conformation.

Received 24 February 2006

Accepted 27 February 2006

Comment

Mycobacterium tuberculosis is the main causative agent of a chronic and often fatal condition in humans known as tuberculosis (TB). It is estimated that one third of the world's population is TB-infected, with about eight million new cases annually, and of these, 3.1 million die annually. TB is currently the leading killer of youths, women, and AIDS patients in the world (Snider *et al.*, 1994). The antimycobacterial activities of 5,6-dihydro-4*H*-1,3-thiazine derivatives have been investigated. They show activity against *Mycobacterium tuberculosis* H37Rv (ATCC 27294) (Koketsu *et al.*, 2002). Confirmation of the conformation of the thiazine ring is essential for the study of structure–biological activity relationships of thiazine derivatives. 4,6-Disubstituted-4-hydroxy-5,6-dihydro-4*H*-1,3-thiazines are synthesized by the $BF_3 \cdot Et_2O$ -catalysed reaction of a primary thioamide with an α,β -unsaturated ketone (Koketsu *et al.*, 1999, 2002). They are obtained as diastereomers resulting from the asymmetric centres at positions 4 and 6 of the thiazine ring. In the present diastereomer, (I), the relationship between the OH group at position 4 and the phenyl group at position 6 is *trans*.



The molecular structure of (I) is shown in Fig. 1. The thiazine ring of (I) adopts a sofa conformation, with atom C3 deviating by 0.686 (3) Å from the plane of the remaining five atoms which lie in a common plane (r.m.s. deviation 0.007 Å). There are intermolecular O—H...N hydrogen bonds between neighbouring molecules (Table 2 and Fig. 2).

Experimental

4-Phenyl-3-buten-2-one (0.58 g, 4.0 mmol) was added to a solution of thiobenzamide (0.55 g, 4.0 mmol) in dry dichloromethane (40 ml) at room temperature under an argon atmosphere. To this solution was added $BF_3 \cdot Et_2O$ (1.2 mmol). The reaction mixture was stirred for 2 h, quenched with saturated sodium carbonate solution, and extracted with dichloromethane. The extracts were dried (Na_2SO_4) and

evaporated to dryness. Recrystallizations from ether/hexane (4:6) gave (I) as crystals (yield 0.65 g, 69%; m.p. 399.6–400.0 K). IR (KBr) 3188, 1590 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3 , δ , p.p.m.): 1.49 (3H, s, CH_3), 1.89 (1H, t, $J = 13.2$ Hz), 2.30 (1H, d, $J = 17.2$ Hz), 3.63 (1H, br, OH), 4.52 (1H, d, $J = 17.2$ Hz, C6), 7.29–7.47 (8H, m, Ar), 7.81–7.84 (2H, m, Ar); ^{13}C NMR (100 MHz, CDCl_3 , δ , p.p.m.): = 26.2, 38.6, 44.2, 85.8, 126.7, 127.7, 128.2, 128.8, 130.8, 138.2, 139.8, 158.1; MS (CI): $m/z = 284$ [$M^+ + 1$].

Crystal data

$\text{C}_{17}\text{H}_{17}\text{NOS}$	Mo $K\alpha$ radiation
$M_r = 283.39$	Cell parameters from 10470 reflections
Orthorhombic, $Pbca$	$\theta = 3.0\text{--}25.0^\circ$
$a = 12.1452$ (2) \AA	$\mu = 0.21$ mm^{-1}
$b = 10.6688$ (2) \AA	$T = 190$ (2) K
$c = 23.1573$ (6) \AA	Prism, colourless
$V = 3000.60$ (11) \AA^3	$0.13 \times 0.13 \times 0.02$ mm
$Z = 8$	
$D_x = 1.255$ Mg m^{-3}	

Data collection

Nonius KappaCCD diffractometer	$R_{\text{int}} = 0.120$
φ or ω scans	$\theta_{\text{max}} = 25.1^\circ$
Absorption correction: none	$h = -14 \rightarrow 14$
45044 measured reflections	$k = -12 \rightarrow 12$
2663 independent reflections	$l = -27 \rightarrow 27$
1718 reflections with $I > 2\sigma(I)$	

Refinement

Refinement on F^2	$w = 1/[\sigma^2(F_o^2) + (0.045P)^2 + 0.1974P]$
$R[F^2 > 2\sigma(F^2)] = 0.046$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.102$	$(\Delta/\sigma)_{\text{max}} < 0.001$
$S = 1.02$	$\Delta\rho_{\text{max}} = 0.18$ e \AA^{-3}
2663 reflections	$\Delta\rho_{\text{min}} = -0.28$ e \AA^{-3}
184 parameters	
H-atom parameters constrained	

Table 1

Selected geometric parameters (\AA , $^\circ$).

S1–C1	1.762 (2)	C2–O1	1.425 (3)
S1–C4	1.826 (2)	C2–C11	1.516 (3)
C1–N1	1.283 (3)	C2–C3	1.522 (3)
C1–C5	1.483 (3)	C3–C4	1.522 (3)
N1–C2	1.479 (3)	C4–C12	1.509 (3)
C1–S1–C4	102.84 (10)	O1–C2–C3	106.48 (18)
N1–C1–C5	118.51 (19)	N1–C2–C3	113.38 (18)
N1–C1–S1	128.66 (17)	C11–C2–C3	110.97 (18)
C5–C1–S1	112.80 (16)	C2–C3–C4	113.10 (18)
C1–N1–C2	121.39 (18)	C12–C4–C3	113.88 (17)
O1–C2–N1	108.00 (17)	C12–C4–S1	107.72 (15)
O1–C2–C11	111.09 (18)	C3–C4–S1	109.60 (15)
N1–C2–C11	106.93 (18)		

Table 2

Hydrogen-bond geometry (\AA , $^\circ$).

$D\text{--}H\cdots A$	$D\text{--}H$	$H\cdots A$	$D\cdots A$	$D\text{--}H\cdots A$
$\text{O1--H9}\cdots\text{N1}^i$	0.84	2.05	2.888 (2)	173

Symmetry code: (i) $-x + 1, -y + 1, -z$.

All H atoms were placed in idealized positions and refined with fixed individual displacement parameters [$U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl C})$], with $\text{C--H} = 0.95\text{--}0.99$ \AA and $\text{O--H} = 0.84$ \AA . The




Figure 1

The molecular structure of (I), showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as small spheres of arbitrary radii.




Figure 2

Hydrogen-bonded (dashed lines) dimeric structure of (I). Displacement ellipsoids are drawn at the 30% probability level and H atoms are shown as small spheres of arbitrary radii.

displacement parameter of the hydroxy H atom was refined. In addition, the torsion angles involving the methyl and hydroxyl group were refined.

Data collection: *COLLECT* (Nonius, 2000); cell refinement: *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *SCALEPACK* and *DENZO* (Otwinowski & Minor, 1997); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEPII* (Johnson, 1976); software used to prepare material for publication: *SHELXL97*.

This work was supported by a Grant-in-Aid for Science Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan (Nos. 15550030 and 17550099), for which we are grateful.

References

- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
Koketsu, M., Senda, T., Yoshimura, K. & Ishihara H. (1999). *J. Chem. Soc. Perkin Trans. 1*, pp. 453–456.

- Koketsu, M., Tanaka, K., Takenaka, Y., Kwong, C. D. & Ishihara, H. (2002). *Eur. J. Pharm. Sci.* **15**, 307–310.
- Nonius (2000). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Snider, D. E. Jr, Raviglione, M. & Kochi, A. (1994). *Global Burden of Tuberculosis*, in *Tuberculosis: Pathogenesis, Protection, and Control*, edited by B. Bloom, pp. 3–11. Washington, DC: ASM Press.