



岐阜大学機関リポジトリ

Gifu University Institutional Repository

Title	Relationship Between Functional Exercise Capacity and Functional Stenosis in Patients With Stable Angina and Intermediate Coronary Stenosis(本文(Fulltext))
Author(s)	TANAKA, Shinichiro; NODA, Toshiyuki; SEGAWA, Tomonori; MINAGAWA, Taro; WATANABE, Sachiro; MINATOBUCHI, Shinya
Citation	[Circulation journal : official journal of the Japanese Circulation Society] vol.[73] no.[12] p.[2308]-[2314]
Issue Date	2009-11-20
Rights	The Japanese Circulation Society (社団法人日本循環器学会)
Version	出版社版 (publisher version) postprint
URL	http://hdl.handle.net/20.500.12099/38580

この資料の著作権は、各資料の著者・学協会・出版社等に帰属します。

Relationship Between Functional Exercise Capacity and Functional Stenosis in Patients With Stable Angina and Intermediate Coronary Stenosis

Shinichiro Tanaka, MD; Toshiyuki Noda, MD; Tomonori Segawa, MD;
Taro Minagawa, MD*; Sachiro Watanabe, MD; Shinya Minatoguchi, MD**

Background: Some stable angina patients with significant coronary function have low exercise capacity, whereas some have high exercise capacity. The aim of the present study was to determine whether coronary pressure-derived fractional flow reserve (FFR_{myo}), a functional index of coronary stenosis, is a better indicator of exercise capacity than angiographic stenosis.

Methods and Results: The 15 male (65.8±8.9 years old) subjects with stable angina and 75% angiographic stenosis underwent a cardiopulmonary exercise test (CPX), and peak oxygen uptake (Peak $\dot{V}O_2$) and oxygen uptake at anaerobic threshold (AT) were measured. The relationship between FFR_{myo} and CPX values was assessed. The left anterior descending artery was affected in 8 patients, the left circumflex artery in 5, and the right coronary artery in 2. Percent diameter stenosis (%DS) was 61.7±9.1% by quantitative coronary angiography. Mean FFR_{myo}, Peak $\dot{V}O_2$, and AT was 0.84±0.66, 17.1±3.2 ml·kg⁻¹·min⁻¹, and 11.1±2.0 ml·kg⁻¹·min⁻¹, respectively. There was no significant correlation between %DS and FFR_{myo}, Peak $\dot{V}O_2$, or AT (r=0.12, -0.051, and -0.013, respectively; P=NS), but FFR_{myo} had a significant positive correlation with Peak $\dot{V}O_2$ and AT (r=0.534 and 0.542, respectively; P<0.05).

Conclusions: Exercise capacity reflects functional stenosis in stable angina patients. (Circ J 2009; 73: 2308–2314)

Key Words: Fractional flow reserve; Functional exercise capacity; Ischemic heart disease

Low peak oxygen uptake (Peak $\dot{V}O_2$) and poor exercise capacity are generally considered to be predictors of mortality in patients with ischemic heart disease.^{1–4} Patients presenting with these symptoms benefit from revascularization via percutaneous coronary intervention (PCI). However, there were some patients with higher functional capacity who already have oxygen consumption at high levels⁵ and often get no benefit from PCI.^{3,4} For example, it has been reported that patients with low functional capacity, low peak $\dot{V}O_2$, and poor oxygen uptake at the anaerobic threshold (AT) during a cardiopulmonary exercise test (CPX) show significant improvement after PCI, suggesting that myocardial impairment induced by intermittent ischemia is reversible, whereas the functional capacity of those with higher peak $\dot{V}O_2$ is not significantly improved after PCI.³ Thus, it is suggested that the efficacy of PCI is influenced by the baseline condition of the patient.

Although angiographic stenosis is often associated with reduced exercise capacity and functional stenosis, the relationship between angiographic stenosis and functionality varies widely.^{6,7} Coronary pressure-derived fractional flow reserve (FFR_{myo}), which is calculated from coronary pressure measurements, is a reliable index of the functional severity of coronary stenosis, and an FFR_{myo} value of 0.75

distinguishes stenoses that are associated with inducible ischemia from those that are not.^{8,9} Indeed, in patients with stable chest pain, the most important prognostic factor is the occurrence of myocardial ischemia reflected by an FFR_{myo} value <0.75. Even after treatment with PCI, the clinical outcomes of such patients are found to be significantly worse than those of patients with functionally insignificant stenoses (FFR_{myo} ≥0.75).^{6,7} We hypothesized that there is an association between exercise capacity and functional stenosis, as reflected by FFR_{myo}, in patients with stable coronary heart disease and we tested this hypothesis using pressure wires and CPX tests.

Methods

Study Design and Participants

The present study was conducted according to the principles of the Declaration of Helsinki and was approved by the Ethics Committee of the Gifu Prefectural General Medical Center. All study participants gave written informed consent prior to enrollment in the study.

Between November 2006 and September 2008, 15 patients were enrolled. All of them showed 75% angiographic stenosis (classified according to the American Heart

Received April 14, 2009; accepted August 20, 2009; released online October 17, 2009

Department of Cardiology, Gifu Prefectural General Medical Center, *Gifu Cardiovascular Institution, **Second Department of Medicine, Gifu University Graduate School of Medicine, Gifu, Japan

Mailing address: Shinichiro Tanaka, MD, Department of Cardiology, Gifu Prefectural General Medical Center, 4-6-1 Noisshiki, Gifu 500-8717, Japan. E-mail: shin19711015@yahoo.co.jp

All rights are reserved to the Japanese Circulation Society. For permissions, please e-mail: cj@j-circ.or.jp

Table 2. Angiographic Data and FFRmyo

Patient no.	Affected artery	QCA				FFRmyo
		RD (mm)	MLD (mm)	Stenosis (%)	Lesion length (mm)	
1	LCX	4.29	1.66	61.4	16.2	0.88
2	LCX	2.91	1.27	56.3	10.8	0.83
3	LAD	2.06	0.73	64.4	17.2	0.78
4	LCX	3.59	2.06	42.7	6.7	0.85
5	LAD	2.39	0.76	68.4	7.9	0.86
6	LAD	2.42	1.08	55.3	6	0.83
7	LAD	2.89	0.91	68.4	7.8	0.86
8	LCX	2.56	0.61	76.4	22.9	0.88
9	LAD	2.46	0.74	70.0	22	0.83
10	LAD	3.58	1.43	60.18	8.76	0.86
11	LAD	2.46	0.74	70.0	21.9	0.83
12	LCX	2.86	1.50	47.5	10.7	0.9
13	LAD	2.52	1.01	60.1	22.5	0.69
14	RCA	3.22	1.39	56.9	13.1	0.77
15	RCA	5.45	1.77	67.5	9.2	0.99
Mean		3.04	1.12	61.7	13.6	0.84
SD		0.89	0.45	9.1	6.3	0.66

FFRmyo, myocardial fractional flow reserve; QCA, quantitative coronary angiography; RD, reference diameter; MLD, minimum luminal diameter; LCX, left circumflex coronary artery; LAD, left anterior descending coronary artery; RCA, right coronary artery.

Antianginal Thrombotic Treatment

All the patients received oral therapy comprising 75 mg of clopidogrel or 200 mg of ticlopidine plus 81 mg of aspirin daily. Anti-ischemic therapy included a long-acting β -blocker, calcium-channel blocker, nitrates and nicorandil, alone or in combination with statins, and either an angiotensin II receptor blocker or an angiotensin-converting enzyme inhibitor for standard secondary prevention.¹³

Statistical Analysis

All analyses were performed using the intention-to-treat principle. Continuous data are expressed as means \pm SD and proportions, and categorical data are expressed as frequencies and percentages. Pearson's correlation coefficient analysis and simple regression analysis were used to assess the relationships between subjects. Values of $P < 0.05$ were considered significant. All analyses were performed using SAS software version 5.0 (SAS Institute, Inc, Cary, NC, USA).

Results

Characteristics of the Patients and their Lesions

The clinical characteristics of the 15 study participants are shown in **Table 1**. Their mean age was 66 years (range, 54–76); 53% of the patients had hypertension, 47% had hyperlipidemia, and 67% had diabetes. Although 10 patients had angina, all recorded normal ECGs while resting. Ergometry revealed myocardial ischemia (eg, ST-segment depression) in 5 patients, and 6 patients were positive on stress thallium scintigraphy.

Angiographic characteristics are summarized in **Table 2**. Stenotic lesions were localized in the left anterior descending artery in 8 patients, in the left circumflex artery in 5 patients, and in the right coronary artery in 2 patients. The average reference vessel diameter was 3.04 mm, and the mean lesion length was 13.6 mm. Quantitative coronary angiography revealed that the mean minimal lumen diameter was 1.12 ± 0.45 mm and the %DS was $61.8 \pm 9.4\%$. Mean FFRmyo was 0.84 (range, 0.69–0.99).

Functional Capacity Assessed by CPX

Table 3 shows the results of the CPX tests. The average

maximal exercise tolerance was 96.3 W, and the Peak $\dot{V}O_2$ was 16.7 ± 2.9 ml \cdot kg⁻¹ \cdot min⁻¹ at 96.3 W. All the $\dot{V}O_2$ curves were observed to plateau during CPX. Peak $\dot{V}O_2$ reflected the maximum $\dot{V}O_2$ value. Mean AT value was 11.1 ± 2.0 ml \cdot kg⁻¹ \cdot min⁻¹, and the mean load at that AT level was 53.1 W. The endpoint of the CPX test was shortness of breath in 4 patients and leg fatigue in 6 patients. Target heart rate was nearly achieved in 4 patients: 90% of maximum heart rate. CPX test was terminated in 1 patient because of elevated diastolic blood pressure (>120 mmHg).

Correlations Among Functional Capacity, Angiographic Stenosis, and Functional Stenosis

There was no correlation between FFRmyo and %DS (**Figure 1**). Likewise, neither Peak $\dot{V}O_2$ nor AT correlated with %DS (**Figure 2**). There was no correlation between the double products and FFRmyo or %DS (**Figure 3**). On the other hand, there was a positive correlation between oxygen uptake and the double products: $r = 0.761$ ($P < 0.01$) at peak and $r = 0.775$ ($P < 0.01$) at the AT. In addition, a significant positive correlation was observed between FFRmyo and Peak $\dot{V}O_2$ ($r = 0.534$, $P < 0.05$) and between FFRmyo and the AT ($r = 0.542$, $P < 0.05$) (**Figure 4**).

Discussion

The results of this study showed (1) a positive correlation between FFRmyo and Peak $\dot{V}O_2$ [$r = 0.534$ ($P < 0.05$)] and between FFRmyo and the AT [$r = 0.542$ ($P < 0.05$)], (2) no significant correlation between %DS and FFRmyo, and (3) no correlation at all between %DS and Peak $\dot{V}O_2$ or between %DS and the AT.

The diagnostic accuracy of FFRmyo for assessing functional stenosis is more than 90%, which is higher than for any other invasive or noninvasive test.^{8,9} When FFRmyo was used to divide patients into a group in which stenosis was most likely to be physiologically significant (FFRmyo < 0.75) and a group in which it was not, the overlap between the 2 groups for angiographic severity was so large that it suggested angiography could not be used to predict the absence or presence of inducible ischemia in individual patients.^{6,7} Our finding that there was no correlation between

Table 3. Anaerobic Threshold, O₂ Uptake, and Hemodynamic Data at Rest and During Exercise

Patient no.	Peak $\dot{V}O_2$ (ml·kg ⁻¹ ·min ⁻¹)	Rest					AT					Maximum					Endpoint of CPX test
		AT		Rest			AT		Maximum			AT		Maximum			
		(ml·kg ⁻¹ ·min ⁻¹)	(ml·kg ⁻¹ ·min ⁻¹)	SBP (mmHg)	DBP (mmHg)	HR (beats/min)	Double product (mmHg·beats/min)	SBP (mmHg)	DBP (mmHg)	HR (beats/min)	Double product (mmHg·beats/min)	SBP (mmHg)	DBP (mmHg)	HR (beats/min)	Double product (mmHg·beats/min)	W	
1	16	8.4	94	47	85	7,990	107	66	101	10,807	135	69	131	17,685	65	Target HR	
2	13.2	9.0	112	74	70	7,840	120	85	92	11,040	152	78	109	16,568	75	Shortness of breath	
3	12.8	8.7	118	73	71	8,378	126	82	91	11,466	154	80	107	16,478	75	Shortness of breath	
4	14.2	10.3	136	69	70	9,520	134	67	81	10,854	116	53	94	10,904	65	Shortness of breath	
5	19.9	13.6	161	110	80	12,880	193	103	120	23,160	206	110	140	28,840	125	Target HR	
6	19.4	13.3	165	106	76	12,540	198	116	122	24,156	202	121	131	26,462	100	High pressure of DBP	
7	17.5	12.0	134	63	78	10,452	116	53	112	12,992	142	60	131	18,602	100	Leg fatigue	
8	15	10.2	152	78	62	9,424	174	73	92	16,008	181	82	114	20,634	90	Leg fatigue	
9	16.5	11.9	134	63	68	9,112	130	68	96	12,480	142	60	102	14,484	90	Leg fatigue	
10	22.2	11.7	126	76	52	6,552	144	86	81	11,664	181	94	116	20,996	130	Leg fatigue	
11	14.2	10.3	142	66	69	9,798	126	50	82	10,332	134	57	93	12,462	65	Shortness of breath	
12	20.8	14.2	163	102	79	12,877	184	109	121	22,264	210	106	144	30,240	125	Target HR	
13	14.7	9.7	148	70	64	9,472	168	66	90	15,120	181	79	106	19,186	85	Leg fatigue	
14	18	9.8	126	78	78	9,828	146	78	94	13,724	220	97	128	28,160	125	Leg fatigue	
15	22	14.0	134	89	63	8,442	198	104	92	18,216	215	112	130	27,950	130	Target HR	
Mean	17.1	11.1	136.3	77.6	71	9,674	151	80.4	97.8	14,952	171.4	83.9	118	20,643	96.3		
SD	3.2	2	19.8	17.4	8.6	1,807	31.9	20.2	14.3	4,636	34.2	21.9	16.3	6,066	25.1		

peak $\dot{V}O_2$, oxygen uptake at peak exercise; AT, anaerobic threshold at the corresponding oxygen uptake; HR, heart rate; SBP, systolic blood pressure; DBP, diastolic blood pressure; Target HR, 90% of maximum HR.




Figure 1. Relationship between fractional flow reserve and percent stenosis.

%DS calculated from quantitative coronary angiography and the 3 functional parameters, namely, FFR_{myo}, peak $\dot{V}O_2$, and AT, is consistent with earlier studies.

FFR_{myo} reflects the maximum achievable blood flow to the myocardium supplied by a stenotic artery as a fraction of normal maximum flow, and is an accurate and specific index of the severity of epicardial stenosis.^{8,9} Generally, one of the most important determinants of myocardial ischemia is endothelial function.^{1,2,14} In patients with a higher maximal exercise tolerance, sufficient myocardial perfusion is achieved because of improved endothelium-dependent vasodilatation and collateralization.¹⁵ In addition, it has been reported that an increase in functional capacity is associated with an increase in coronary flow reserve (coronary flow velocity response to adenosine) and that impaired overall functional capacity is independently associated with coronary microvascular dysfunction.¹ Thus, a reduction in coronary flow reserve apparently reflects both epicardial and microvascular disease.^{16,17} Other studies have shown that exercise training improves endothelium-dependent vasodilatation in both epicardial coronary vessels and resistance vessels in patients with coronary artery disease.² Exercise brings about a change in the relationship between myocardial blood flow and coronary pressure in patients with moderate stenosis; the linear relationship between perfusion pressure and blood flow at exercise in cases of a moderate stenosis of the epicardial artery shifted upward and were closer to those found with hyperemia than those for the sub-endocardial artery.¹⁸ Such results are consistent with those of our study, indicating that there is a positive correlation between FFR_{myo} and functional capacity and little, if any, relationship between %DS and functional capacity. The reason why FFR_{myo} does not correlate with the maximum double product appears to be associated with the endpoint of the CPX test. In the present study, none of the patients experienced anginal pain or significant ST depression during the CPX test. Because the maximal double product is important as an angina threshold, it should correlate with FFR_{myo} when exercise is terminated because of myocardial ischemia. In the absence of myocardial ischemia, which was the case in the present study, peak $\dot{V}O_2$ may be a more appropriate index for measuring exercise capacity. In addition, exercise increases the levels of catecholamines, which may be responsible for the improvement in blood flow and




Figure 2. Relationship between percent stenosis and parameters reflecting exercise capacity. (A) Peak $\dot{V}O_2$ vs percent stenosis; (B) AT vs percent stenosis. Peak $\dot{V}O_2$, oxygen uptake at peak exercise; AT, anaerobic threshold at the corresponding oxygen uptake.




Figure 3. Relationship between double product and percent stenosis or functional stenosis. (A) Double product at peak exercise vs fractional flow reserve; (B) double product at peak exercise vs percent stenosis; (C) double product at AT vs fractional flow reserve; (D) double product at AT vs percent stenosis. AT, anaerobic threshold; bpm, beats/min.

oxygen delivery to the skeletal muscles.¹⁹ This may be one of reasons why FFR_{myo}, which was measured during hyperemia induced by adenosine, showed a significant correlation with oxygen uptake but not with the double product.

When treating patients with coronary artery disease and

stable angina, the goal of treatment is complete, or nearly complete, elimination of anginal chest pain and a return to normal activity with good functional capacity.^{20,21} Although PCI has become a common initial management strategy for patients with stable coronary artery disease, studies have




Figure 4. Relationship between fractional flow reserve and parameters reflecting exercise capacity. (A) Peak $\dot{V}O_2$ vs fractional flow reserve; (B) AT vs fractional flow reserve. Fractional flow reserve showed a positive correlation with exercise capacity. Peak $\dot{V}O_2$, oxygen uptake at peak exercise; AT, anaerobic threshold at the corresponding oxygen uptake.

shown that it does not reduce the long-term rates of death risk, myocardial infarction or other major cardiovascular events when combined with optimal medical therapy.¹³ Nonetheless, for approximately 24 months, PCI accompanied by optimal medical therapy does relieve angina and improve the self-assessed health status to a greater degree than that achieved by optimal medical therapy alone. In addition, a greater benefit of PCI is observed in patients with more severe and more frequent angina. Therefore, the importance of PCI in terms of rapid relief of symptoms must be critically weighed against the inherent risks and the higher cost of the procedure.⁴ From that viewpoint, FFR_{myo} can be used for immediate decision-making in the catheterization laboratory. However, under other circumstances, a noninvasive procedure would be preferable. Assessment of exercise capacity using a CPX test is not invasive and can be used together with metabolic equivalent values to serve as an index of daily activity. CPX testing is useful over self-assessed health status, which is subjective and can vary among patients depending upon their symptom complexity and their unique perceptions, expectations, and preferences. For instance, patients with more severe angina and several coexisting medical problems may be satisfied with some reduction in the symptoms that enables them to perform only limited daily life activities.^{20,21} Another investigation has indicated that 6,500–8,500 steps/day should be considered as the minimal goal and optimal level of physical activity for secondary prevention of cardiovascular disease.²² For assessment of daily activity or for secondary prevention, CPX testing is a noninvasive guide to functional capacity and treatment efficacy.

PCI significantly increased AT and peak $\dot{V}O_2$ in patients with peak $\dot{V}O_2 < 15$ ml·kg⁻¹·min⁻¹, whereas it yielded no significant improvement in those with peak $\dot{V}O_2 > 15$ ml·kg⁻¹·min⁻¹.³ It is noteworthy that an FFR_{myo} value of 0.75 reflects stenoses associated with inducible ischemia and hence appears to be similar to a peak $\dot{V}O_2$ value of 15.0.

A previous study showed that routine FFR_{myo} measurement with angiographic guidance resulted in a significant reduction in major adverse cardiac effects, even when patients underwent PCI with drug-eluting stents.²³ Cardiac rehabilitation is an important strategy for secondary prevention, provided there is no risk of inducible ischemia.^{5,15,22}

Exercise training can improve myocardial perfusion, as revealed by myocardial scintigraphy, as well as maximal exercise capacity.¹⁶ Therefore, it may be possible to improve FFR_{myo} by increasing exercise capacity through training.

Study Limitations

This study has several important limitations. Our study evaluated a small group of Japanese men with stable angina warranting referral for coronary angiography, which limits generalization of the results. We measured FFR_{myo} at only one point in time, so unknown or unmeasured factors that could potentially alter the responses at other time points were not studied. In addition, it was a nonrandomized study in which the study population was small, and the results reflected the experience of a single center only. Therefore, the study lacks the obvious advantages of a larger, multi-center, multinational randomized study. Inclusion criteria for this study were restrictive and hence limited the number of study patients, even though intermediate coronary heart disease and the resultant myocardial ischemia are known to be important factors that affect the daily activity of numerous individuals. This study included a very high proportion of diabetic patients (67%). Although glucose levels were efficiently controlled by administration of only α -glucosidase inhibitors or diet therapy alone, our study did not reflect the vascular functional changes that are induced by hyperglycemia. A larger, multicenter, multinational randomized study is warranted to clarify these issues.

Conclusions

Japanese men with reduced FFR_{myo} are likely to have reduced functional capacity as assessed by a CPX test. CPX testing may offer a noninvasive measure of myocardial ischemia that is associated with the fractional flow reserve.

Acknowledgments

Reiko Matsuoka, MD; Shintaro Abe, MD; Yoshiaki Goto, MD; Takashi Kato, MD; Tomoko Hirose, MD; Shunnichiro Warita, MD; Tai Kojima, MD; Takeshi Hirose, MD; Makoto Iwama, MD; Koji Ono, MD and Haruki Takahashi, MD helped in collecting the data for this study.

We thanked Toshihiko Hirose, Yuko Ishida, and Masashi Nishiwaki for assisting with the CPX tests and with the preparation of the manuscript.

Disclosure

This study was not financially supported by any organization. And all authors have no conflict of interest with this study.

References

- Handberg E, Johnson BD, Arant CB, Wessel TR, Kerensky RA, von Mering G, et al. Impaired coronary vascular reactivity and functional capacity in women: Results from the NHLBI Women's Ischemia Syndrome Evaluation (WISE) Study. *J Am Coll Cardiol* 2006; **47**: S44–S49.
- Hambrecht R, Wolf A, Gielen S, Linke A, Hofer J, Erbs S, et al. Effect of exercise on coronary endothelial function in patients with coronary artery disease. *N Engl J Med* 2000; **342**: 454–460.
- Barmeyer A, Meinertz T. Anaerobic threshold and maximal oxygen uptake in patients with coronary artery disease and stable angina before and after percutaneous transluminal coronary angioplasty. *Cardiology* 2002; **98**: 127–131.
- Weintraub WS, Spertus JA, Kolm P, Maron DJ, Zhang Z, Jurkovicz C, et al. Effect of PCI on quality of life in patients with stable coronary disease. *N Engl J Med* 2008; **359**: 677–687.
- Seki E, Watanabe Y, Shimada K, Sunayama S, Onishi T, Kawakami K, et al. Effects of a phase III cardiac rehabilitation program on physical status and lipid profiles in elderly patients with coronary artery disease: Juntendo Cardiac Rehabilitation Program (J-CARP). *Circ J* 2008; **72**: 1230–1234.
- Bech GJ, De Bruyne B, Pijls NH, de Muinck ED, Hoorntje JC, Escaned J, et al. Fractional flow reserve to determine the appropriateness of angioplasty in moderate coronary stenosis: A randomized trial. *Circulation* 2001; **103**: 2928–2934.
- Pijls NH, van Schaardenburgh P, Manoharan G, Boersma E, Bech JW, van't Veer M, et al. Percutaneous coronary intervention of functionally nonsignificant stenosis: 5-year follow-up of the DEFER Study. *J Am Coll Cardiol* 2007; **49**: 2105–2111.
- Pijls NH, De Bruyne B, Peels K, Van Der Voort PH, Bonnier HJ, Bartunek J, et al. Measurement of fractional flow reserve to assess the functional severity of coronary-artery stenoses. *N Engl J Med* 1996; **334**: 1703–1708.
- Pijls NH, Van Gelder B, Van der Voort P, Peels K, Bracke FA, Bonnier HJ, et al. Fractional flow reserve: A useful index to evaluate the influence of an epicardial coronary stenosis on myocardial blood flow. *Circulation* 1995; **92**: 3183–3193.
- Yokoyama M. Guidelines for diagnostic evaluation of patients with chronic ischemic heart disease (JCS 2005). Available at http://www.j-circ.or.jp/guideline/pdf/JCS2005_yokoyama_h.pdf
- Lopez-Palop R, Saura D, Pinar E, Lozano I, Pérez-Lorente F, Picó F, et al. Adequate intracoronary adenosine doses to achieve maximum hyperaemia in coronary functional studies by pressure derived fractional flow reserve: A dose response study. *Heart* 2004; **90**: 95–96.
- Wilson RF, Wyche K, Christensen BV, Zimmer S, Laxson DD. Effects of adenosine on human coronary arterial circulation. *Circulation* 1990; **82**: 1595–1606.
- Boden WE, O'Rourke RA, Teo KK, Hartigan PM, Maron DJ, Kostuk WJ, et al. Optimal medical therapy with or without PCI for stable coronary disease. *N Engl J Med* 2007; **356**: 1503–1516.
- Hambrecht R, Adams V, Erbs S, Linke A, Kränkel N, Shu Y, et al. Regular physical activity improves endothelial function in patients with coronary artery disease by increasing phosphorylation of endothelial nitric oxide synthase. *Circulation* 2003; **107**: 3152–3158.
- Hambrecht R, Walther C, Möbius-Winkler S, Gielen S, Linke A, Conradi K, et al. Percutaneous coronary angioplasty compared with exercise training in patients with stable coronary artery disease: A randomized trial. *Circulation* 2004; **109**: 1371–1378.
- Albertal M, Regar E, Van Langenhove G, Carlier SG, Serrano P, Boersma E, et al. Flow velocity and predictors of a suboptimal coronary flow velocity reserve after coronary balloon angioplasty. *Eur Heart J* 2002; **23**: 133–138.
- Albertal M, Voskuil M, Piek JJ, de Bruyne B, Van Langenhove G, Kay PI, et al. Doppler Endpoints Balloon Angioplasty Trial Europe (DEBATE) II Study Group: Coronary flow velocity reserve after percutaneous interventions is predictive of periprocedural outcome. *Circulation* 2002; **105**: 1573–1578.
- Duncker DJ, Bache RJ. Regulation of coronary blood flow during exercise. *Physiol Rev* 2008; **88**: 1009–1086.
- Riley M, Machara K, Pórszász J, Engelen MP, Bartstow TJ, Tanaka H, et al. Association between the anaerobic threshold and the breakpoint in the double product/work rate relationship. *Eur J Appl Physiol Occup Physiol* 1997; **75**: 14–21.
- Fraker TD Jr, Fihn SD, Gibbons RJ, Abrams J, Chatterjee K, Daley J, et al. 2007 chronic angina focused update of the ACC/AHA 2002 guidelines for the management of patients with chronic stable angina: A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines Writing Group to develop the focused update of the 2002 guidelines for the management of patients with chronic stable angina. *J Am Coll Cardiol* 2007; **50**: 2264–2274.
- Tanaka S, Yokoyama C, Kawamura I, Takasugi N, Kubota T, Ushikoshi H, et al. Conservative medication follow-up for over 20 years of a patient with ischemic heart disease after diagnosis of chronic total occlusion of the 3 main coronary arteries. *Circ J* 2008; **72**: 1205–1209.
- Ayabe M, Brubaker PH, Dobrosielski D, Miller HS, Kiyonaga A, Shindo M, et al. Target step count for the secondary prevention of cardiovascular disease. *Circ J* 2008; **72**: 299–303.
- Tonino PA, De Bruyne B, Pijls NH, Siebert U, Ikeno F, van't Veer M, et al. Fractional flow reserve versus angiography for guiding percutaneous coronary intervention. *N Engl J Med* 2009; **360**: 213–224.