

University of Kentucky

UKnowledge

Sanders-Brown Center on Aging Faculty
Publications

Aging

1-5-2022

Sensory Processing Abnormalities in Community-Dwelling Older Adults with Cognitive Impairment: A Mixed Methods Study

Elizabeth K. Rhodus
University of Kentucky

Elizabeth G. Hunter
University of Kentucky

Graham D. Rowles
University of Kentucky

Shoshana H. Bardach
University of Kentucky

Kelly Parsons
University of Kentucky

See next page for additional authors

Follow this and additional works at: https://uknowledge.uky.edu/sbcoa_facpub



Part of the [Geriatrics Commons](#), and the [Neurosciences Commons](#)

[Right click to open a feedback form in a new tab to let us know how this document benefits you.](#)

Repository Citation

Rhodus, Elizabeth K.; Hunter, Elizabeth G.; Rowles, Graham D.; Bardach, Shoshana H.; Parsons, Kelly; Barber, Justin M.; Thompson, MaryEllen; and Jicha, Gregory A., "Sensory Processing Abnormalities in Community-Dwelling Older Adults with Cognitive Impairment: A Mixed Methods Study" (2022). *Sanders-Brown Center on Aging Faculty Publications*. 205.
https://uknowledge.uky.edu/sbcoa_facpub/205

This Article is brought to you for free and open access by the Aging at UKnowledge. It has been accepted for inclusion in Sanders-Brown Center on Aging Faculty Publications by an authorized administrator of UKnowledge. For more information, please contact UKnowledge@lsv.uky.edu.

Sensory Processing Abnormalities in Community-Dwelling Older Adults with Cognitive Impairment: A Mixed Methods Study


Digital Object Identifier (DOI)


<https://doi.org/10.1177/23337214211068290>

Authors

Elizabeth K. Rhodus, Elizabeth G. Hunter, Graham D. Rowles, Shoshana H. Bardach, Kelly Parsons, Justin M. Barber, MaryEllen Thompson, and Gregory A. Jicha

Sensory Processing Abnormalities in Community-Dwelling Older Adults with Cognitive Impairment: A Mixed Methods Study

Gerontology & Geriatric Medicine
Volume 8: 1–11
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/23337214211068290
journals.sagepub.com/home/ggm


Elizabeth K. Rhodus, PhD, OTR/L^{1,2,3} , Elizabeth G. Hunter, PhD, OTR/L⁴,
Graham D. Rowles, PhD⁴, Shoshana H. Bardach, PhD⁵, Kelly Parsons, MSW, CSW¹,
Justin Barber, MS¹, MaryEllen Thompson, PhD, OTR/L⁶ and Gregory A. Jicha, MD, PhD^{1,2,7}

Abstract

Mild cognitive impairment (MCI) or dementia often leads to behavioral and psychiatric symptoms of dementia (BPSD). Sensory processing abnormalities may be associated with BPSD. The purpose of this study was to explore relationships among sensory processing, behavior, and environmental features within the homes of people with MCI or dementia. This project used mixed methods to assess participants' sensory processing, care partner perspectives on behaviors, and *in situ* observations of the home environment. Nine participants with cognitive impairment (MCI $n = 8$, early dementia = 1) and their care partners were included. Seven participants with cognitive impairment were reported to have abnormal sensory processing. Findings suggest that unique environmental adaptations, tailored to personal and sensory preferences for each participant, were associated with a decreased level of behavioral disruption during the observation periods. Implementing sensory-based approaches to maximize environment adaptation may be beneficial in reducing disruptive behaviors for adults with cognitive impairment.

Manuscript received: July 30, 2021; **final revision received:** November 30, 2021; **accepted:** December 2, 2021.

Introduction

One in nine individuals over the age of 65 have mild cognitive impairment (MCI) or Alzheimer's disease or related dementias (ADRD), and as many as one in three over the age of 85 have these conditions (Alzheimer's Association, 2020). Including persons with MCI, nearly 90% of individuals with these diagnoses demonstrate behavioral and psychiatric symptoms of dementia (BPSD) (Gerlach & Kales, 2020; Zhao et al., 2016). Often BPSD, such as agitation, anxiety, apathy, and depression, present in early stages of cognitive impairment including MCI and can be a precursor to severity of ADRD (Sun et al., 2021; Wise et al., 2019; Zhao et al., 2016). These symptoms interfere with optimal care provision, decrease functional independence, result in inappropriate use of antipsychotic medications, and reduce quality of life for both persons with cognitive impairment and their care partners (Kales et al., 2014).

¹Sanders-Brown Center on Aging, University of Kentucky, Lexington, KY, USA

²Department of Behavioral Science, University of Kentucky, Lexington, KY, USA

³Center for Health Equity Transformation, University of Kentucky, Lexington, KY, USA

⁴Graduate Center for Gerontology, University of Kentucky, Lexington, KY, USA

⁵The Dartmouth Institute for Health Policy & Clinical Practice, Dartmouth University, Lebanon, NH, USA

⁶Department of Occupational Science and Occupational Therapy, Eastern Kentucky University, Richmond, KY, USA

⁷Department of Neurology, University of Kentucky, Lexington, KY, USA

Corresponding Author:

Elizabeth K. Rhodus, PhD, OTR/L, Sanders-Brown Center on Aging, University of Kentucky, 463 Healthy Kentucky Research Building, 760 Press Avenue, Lexington, KY 40508, USA.
Email: Elizabeth.rhodus@uky.edu



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE

and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).


Table 1. Adult Sensory Profile Questionnaire Domains, Behaviors, and Examples.

Sensory Domain	Associated Behavior	Sample Statements from ASP
Low registration	Decreased awareness to sensory information and need for high intensity stimulation	I trip or bump into things. I do not seem to notice when my face or hands are dirty.
Sensory seeking	Behaviors that seek out sensory stimuli	I add spice/herbs to my food. I wear colorful clothing.
Sensory sensitivity	Likely to respond and be sensitive to low levels of sensory stimuli	I feel dizzy easily. I am distracted when there is a lot of noise around.
Sensation avoiding	Tendency to control or avoid overwhelming sensory stimuli	I only eat familiar foods. I stay away from crowds.

The Science of Behavior Change program (Nielsen et al., 2018), endorsed by the National Institutes of Health, suggests that behavioral scientists need to identify mechanisms of action which influence behavior and to determine appropriate assessment of such mechanisms. Identification of underlying mechanisms contributing to BPSD is needed to improve care and behavior science (Topaz et al., 2020; Nielsen et al., 2018). National and international agencies recognize a strong need for nonpharmacological interventions, including environmental assessment and modification, as first-line treatment for BPSD in ADRD (Gitlin et al., 2012; Zeisel et al., 2003). However, lack of assessments and interventions to guide and apply such approaches severely limit real-world implementation (Padilla, 2011). To fill this gap, innovative methodologies are needed to define mechanisms of BPSD.

The current study was influenced by mechanisms established by environmental gerontologists who have long recognized behavior as a product of environmental interaction. In 1936, Lewin defined (B)ehavior as a (f)unction of the (P)erson and (E)nvironment: $B = f(P, E)$ (Lewin, 1936). Building on this, Lawton and Nahemow (1973) developed the Ecological Model of Aging which depicts zones of adaptive behavior when environmental factors/demands are aligned with a person's capability. This model identifies zones of maximum comfort and performance when environmental factors are consonant with individual ability. Person-environment fit, as identified in this model, informs pragmatic development of assessment and intervention for community-residing adults with ADRD. However, there is sparse evidence for the combined effects of environmental factors on behavioral regulation and performance for people with ADRD (Jensen & Padilla, 2017; Magasi et al., 2015).

Neurobehavioral disruptions, including BPSD, may be a manifestation of sensory impairment and/or sensory processing abnormalities (Gerlach & Kales, 2020; Luo et al., 2018; Pinto et al., 2017; Sánchez et al., 2016). For example, agitation may occur in response to overstimulation, and wandering may be a self-initiated attempt to increase sensory stimulation (Ravn et al., 2018). Behaviors related to sensory processing are unique to the individual and reflect personal preferences, prior experiences, and neurological needs (Dunn, 1997; Kovach

**Figure 1.** Recruitment and data collection.

et al., 2004). Such behaviors have been described as patterns which reflect sensory processing abilities and have been classified within four categories: low registration; sensation seeking; sensory sensitivity; and sensation avoiding (see Table 1) (Brown & Dunn, 2002; Dunn, 1997).

Utilization of sensory processing considered in environmental context as an innovative approach for understanding BPSD may be appropriate and effective for persons with ADRD and their caregivers (Champagne, 2018; Ravn et al., 2018).

The aim of this study was to assess the presence of sensory processing abnormalities in older adults with cognitive impairment and evaluate the relationship among these abnormalities, environmental features, and behavior in their home environment.

Methods

Design

We used a mixed method approach involving descriptive quantitative evaluation of cognition and sensory processing

and qualitative exploration of community-dwelling older adults with cognitive impairment through interviews with caregivers and *in situ* observation of behavior and environmental features in the home setting (see Figure 1). This approach was selected due to the complexity of the situational context. Using these methods increases rigor through triangulation of data sources and provides a pragmatic approach to investigating and interpretation of behavior in the context of which it occurs (Creswell, 2013; Morgan, 2007). Written consent for participation was obtained from care partners and the legally authorized representative of the person with cognitive impairment. Verbal assent for participation was obtained from those with cognitive impairment. All procedures were approved by the University Institutional Review Board.

Participants

All participants were enrolled in an Alzheimer's Disease Research Center (ADRC) cohort (Schmitt et al., 2012). Purposeful sampling was used to recruit participants from the ADRC if they had a clinical diagnosis of MCI or dementia, were aged 65 years or older, were deemed appropriate for participation by a medical clinician or social worker from the ADRC, lived at home with a care partner who was willing to participate in this study, were able to speak and understand English, and resided in the same county as the research institution. Residence in the same county of the research institution was necessary to maintain feasible travel for in-home observations.

Diagnosis

The diagnosis of MCI was determined according to the consensus guidelines developed by the Second International Working Group on MCI (Winblad et al., 2004) and adopted by the National Institute on Aging-Alzheimer's Association Workgroup on Diagnostic Guidelines for Alzheimer's Disease as follows: (1) a cognitive complaint by the subject or informant, or evidence for longitudinal decline on cognitive test performance (at least 1.5 standard deviation decline); (2) generally intact global cognition; (3) no or minimal functional impairment; and (4) not demented according to DSM-IV criteria. Diagnosis of dementia was based on criteria set forth by the Diagnostic and Statistical Manual of Mental Disorders Fourth Edition (DSM-IV) (American Psychiatric Association, 2013). Additional guidance by the National Institute for Aging and the Alzheimer's Association (Albert et al., 2011) was used for MCI and ADRC diagnoses. All participants' diagnoses were confirmed with a neurologist.

Quantitative Data Collection

Quantitative data was assembled from cross-sectional cognitive test scores and measurement of sensory processing.

Cognitive test scores were collected from the ADRC database (test date within 24 months of observation) to supplement clinical diagnoses with methods described elsewhere (Schmitt et al., 2012). Measurement of sensory processing was collected at the time of participant observation. These descriptive quantitative measures are presented to provide a diagnostic description of the participants; statistical analyses, including power analysis and sample size calculation, were not conducted.

Cognitive assessment. The Clinical Dementia Rating (CDR) (Morris, 1993) scores from the most recent ADRC annual visit of each participant were extracted from the ADRC database and used to identify cognitive impairment severity to supplement the clinical diagnosis. CDR scores reflect participants' status at the most recent ADRC visit (within 24 months) from the date of study enrollment. The CDR has high reliability and validity for assessment of cognitive function and serves as a primary tool in psychometric analysis of functional cognition of ADRC (Hughes et al., 1982; Morris, 1993). Six cognitive domains were assessed via caregiver interview during the CDR (memory, orientation, judgment and problem solving, community affairs, home and hobbies, and personal care). Following the interview, each domain is rated on a 5-point scale indicating the level of impairment (0 = none, 0.5 = questionable, 1 = mild, 2 = moderate, and 3 = severe) based on standard rules.

Sensory processing. The Adult Sensory Profile (ASP) questionnaire was used to measure sensory processing (Brown et al., 2001). Participants or their primary care partner completed the 60-item ASP questionnaire, which is validated and reliable for use in older adults, including those with MCI and dementia (Chung, 2006; Engel-Yeger & Rosenblum, 2021). Each item describes a behavior attributed to sensory processing and results present sensory processing abilities among four categories (see Table 1). For the respondent, the ASP items are presented in groups, by sensory system (number of items): gustatory/olfactory (8), proprioceptive (8), visual (10), haptic (13), and auditory (11). The 10 remaining items are classified as activity level. Participants rate the frequency with which they engage in each behavior or they observe it in the participant, in the case of caregivers responding. Ratings for all items are summed within each sensory processing pattern to obtain summary scores (Brown et al., 2001). The summary score of each category is compared to normative data established via a national sample of same-aged adults (Dunn & Westman, 1997). Identification of more than one category outside of normative results was indicative of sensory processing abnormalities.

Qualitative Data Collection

A qualitative descriptive approach was used to describe the situation of each study participant. Prior to data collection, the

Table 2. Observation Chart Prompts Used for Data Collection.

Assessment	Behavioral Signs
Neuropsychiatric Inventory (NPI)	Delusions/hallucinations, irritability, anxiety/agitation, depression/apathy, disinhibition, aberrant motor behaviors, appetite/eating changes, nighttime behaviors
In-Home Occupational Performance Evaluation (I-HOPE)	Lighting/sounds, smells/temperature, visual aids/distractions, position (seated/standing/mobile), inside/outside, time of day, access within and surrounding home, performance within environment (including decision making)
Social context	Caregiver present, interactions (empathy, theory of mind, facial recognition), lives with, alertness, communication (figurative language, nonverbal), response to social contact, participant's mood, routine interactions

first author used principles of bracketing and reflexivity to acknowledge their background as an occupational therapist which may introduce biases, such as emphasis on occupations and functional behaviors (Creswell, 2013). Avenues to ensure trustworthiness and consistency of findings included peer debriefing with AD researchers, constant comparison of emergent findings, and triangulation (Creswell, 2013).

Care partner interviews. Semi-structured interviews took place with the primary care partner (interview prompts included in [Supplementary file](#)). These interviews were designed to gain information regarding care partner observations of behaviors and environmental interaction. All interviews were audio recorded and transcribed verbatim by the first author.

Behavioral observation. In-home observations were undertaken after semi-structured interviews with care partners to discuss their perceptions of behaviors and interactions within the home environment by the person with cognitive impairment. The first author acted as a participant-observer, meaning that verbal interaction between the participant and observer occurred, if appropriate, but was not solicited. Second home visits were observation-only and were conducted at a different time of day than the first visit (morning vs. afternoon) to promote representation of behaviors throughout daylight hours.

In-home observations were documented via field notes and observation charts (see [Table 2](#)). These charts were developed from select items of the Neuropsychiatric Inventory (NPI) (Cummings, 1997) and In-Home Occupational Performance Evaluation (I-HOPE) (Stark et al., 2010). Items from the NPI were used to identify behaviors specific to cognitive impairment, including observation of the sleeping environment which might have contributed to nighttime behaviors. The NPI is a commonly used behavior assessment tool for persons with cognitive impairment (Cummings, 1997). Items from the I-HOPE were selected for observation data acquisition of environmental features that might influence behaviors.

Analysis

Mixed methods were used for data analysis. Quantitative measures were scored and used for descriptive purposes in

combination with qualitative findings. Qualitative analyses used descriptive, case study analysis allowing for illustration of the situational context of the participants' living environment in relation to their sensory processing and behavior. Data analysis began following the first interview and used a constant comparison approach with appropriate modifications following each field visit (Boeije, 2002). Peer debriefing with ADRC researchers and expert consulting with the ADRC clinical director (senior author) was used during data collection and analysis to ensure validity and rigor of data. Transcriptions were coded using HyperResearch 4.0.2 (Researchware, 2018).

Findings

Twenty-four individuals with cognitive impairment who were enrolled in the ADRC met inclusion criteria and were selected for recruitment. Of these, 11 were unable to be contacted and four declined participation. Common reasons for exclusion of potential participants included: recent move into facility-based care, death, and care partners' refusal to participate.

Nine participants with cognitive impairment and their care partners agreed to study enrollment ([Table 3](#); pseudonyms are used). All participants were in stable condition at the time of study participation without recent (4 weeks) exacerbations or onset of cognitive, behavioral, or medical conditions. Two participants (Charlotte and Ronald) were married, both with diagnosis of MCI, and both participated as cognitively impaired participants and simultaneously responded as one another's care partner. Both were legally and cognitively able to make consent decisions for this study, and their family was aware of their participation. For purposes of data collection and presentation of findings, their personal experiences as a participant with cognitive impairment as well as insights from a care partner perspective were collected, and observations occurred simultaneously.

Sixteen observations took place within participants' homes. ASP summary scores indicated at least one sensory processing category deviated from the normative data of older adults for all nine participants ([Table 4](#)). Seven of the nine participants had two categories outside of normative data. Abnormalities with low registration were reported most

Table 3. Participant Characteristics.

Name	Age	Sex	Education (Years)	Race	Diagnosis	Global CDR	Care Partner	Length of Relationship (Years)
Betty	73	F	18	White	MCI	0	Samuel	50
Chris	93	M	21	White	MCI	0	Karen	50
Walter	82	M	21	Black	MCI	0.5	Laura	48
Charlotte	88	F	13	White	MCI	0.5	Ronald	58
Ronald	83	M	13	White	MCI	0.5	Charlotte	58
Adam	85	M	21	White	MCI	0.5	Ellen	20
Pamela	67	F	12	Black	MCI	0.5	Alex	50
Ricky	81	M	16	White	MCI	0.5	Janelle	10
Vicky	81	F	12	White	Dementia	1	John	61

Notes. M = male; F = female; MCI = mild cognitive impairment; CDR = cognitive dementia rating.

Table 4. Adult Sensory Profile results for participants with cognitive impairment compared to normative data for persons aged 65 years and older.

Participant	Low Registration	Sensation Seeking	Sensory Sensitivity	Sensation Avoiding
Ronald	+	=	=	+
Pamela	+	=	=	=
Walter	+	=	=	=
Charlotte	+	=	+	+
Vicky	=	+	-	=
Adam	+	=	=	++
Betty	+	=	=	+
Ricky	+	+	=	=
Chris	=	-	=	-

Notes. Symbols representative of normative data for specific age group as reported on the Adult Sensory Profile:¹⁹ (-) Less than most people; (=) similar to most people; (+) more than most people; (++) much more than most people.

frequently in 78% of participants ($n = 7$). Scores outside of normal range were also common in sensory avoiding, noted in 56% of participants ($n = 5$).

During interviews, all participants and their care partners described elements of their environment associated with their sensory processing patterns and behavior. These responses were unique for each participant. Some described self-initiated modifications of the environment to adjust sensory stimuli in the home. Vignettes below describe participants' sensory profiles, care partner discussion of behavioral symptoms, and environmental features to illustrate interrelationships among sensory processing, home environment elements, and behavior, whereby illustrating the potential mechanism of action related to behavioral expression for this population.

Chris

Chris, age 93 had a diagnosis of MCI, and lived with his wife, Karen. She described his temperament as very laid-back. He went with the flow of the day and said he was not bothered by anything within his environment. He enjoyed weekly outings to have coffee with the "guys" and to exercise at a local boxing club.

Sensory profile. Chris was "less than most" in sensation seeking and sensation avoiding, meaning his sensory processing system sought out sensory stimulation less than others of his age, and he also actively avoided stimulation less than others. That behaviors were passive in relation to environmental stimuli was evident by minimal influence from the environment impacting his behavioral state. A sensory profile indicating passivity in behavioral reactions to stimulation correlates with his self-described laid-back personality. For example, he was indifferent regarding the noise level in their home when friends or family visited, and he did not actively attempt to clean or reorganize their belongings as Karen packed things to donate.

Environment. Chris and Karen's home was quiet, well-lit, and without clutter. Chris stayed on the primary floor, rather than descending stairs to the basement. For this reason, Karen completed the laundry. Chris slept in the living room in an electric, reclining chair. Karen helped him with bathing and dressing, but she said the bathrooms were well-equipped with grab bars and bathroom stools to support his decreased balance. In addition to Karen, Chris had regular help with transportation from friends and neighbors. Their children visited monthly.

Behavior. Chris' sensory profile and home environment characteristics appeared to be a good match as Chris did not present with negative behaviors or limited accessibility within his home. While he did describe some frustration with any clutter of mail and/or and medicine bottles on the dining room table which caused him to have difficulty differentiating or remembering which medicine to take, he had no other comments or complaints of discomfort within the home.

Ricky

Ricky, age 81 with a diagnosis of MCI, lived with his wife of 10 years, Janelle. He described himself as the shopper for the family because he enjoyed going to the grocery store and driving his car.

Sensory profile. Ricky's sensory profile revealed he was "*more than most*" in areas of low registration and sensation seeking. More than most in low registration indicates a difficulty with sensory discrimination and a need for increased intensity of sensory stimulation for perception. More than most in sensation seeking indicates a drive to seek out sensory stimulation, such as driving or seeking a cluttered space. His sensory profile scores correspond with his interest in shopping, driving, and need for high levels of visual stimulation in his office.

Environment. Ricky's home was well-kept with one room dedicated as his office. In his office, the walls were covered with pictures and shelves were filled with books. He mentioned he felt comfortable in this space with *his* belongings. They had moved into their patio home in a small retirement community a few years prior to the interview. Ricky and his wife describe a rich social life with their neighbors, including frequent visits from friends, neighborhood socials, and game night with friends.

Behavior. Ricky was pleasant and friendly throughout observations. He appeared to have good environmental fit within his home, as it was clean with limited distractions to the completion of tasks. He and his wife described times of agitation and frustration when he was cognitively challenged. For example, Ricky asked Janelle to make decisions regularly. When in the community, she navigated as he drove; she made the grocery list and weekly meals, and he shopped. When he did not have her guidance in making decisions, he became overwhelmed and agitated. They were both close with their church leadership and described church support groups which aided Ricky's ability to manage his agitation.

Betty

Betty, age 73 with a diagnosis of MCI, was an active member of her community and church as a volunteer. She lived with

her husband, Samuel, in a home where they had lived for the majority of their adulthood.

Sensory profile. Betty's ASP results identified her as "*more than most*" in areas of low registration and sensation avoiding. More than most in low registration indicates a need for increased intensity in stimulation for perception. Also, more than most in sensation avoiding indicates a tendency to avoid unpredictable sensory environments. These are evident by her need for increased attention to detail and self-report of need to control aspects of her environment. Those who are sensory avoiding are often more accepting of environmental stimuli they can control. For example, Betty said she loved to sing but often avoided loud sounds. She slept in a separate room from her husband due to his loud snoring.

Environment. Betty and Samuel lived in a two-story home which had undergone major renovation and expansion to accommodate their needs as they aged. They were committed to living in their home for as long as possible. She and her husband laughed about her desire to organize everything in their home and keen attention to details in all her activities.

Behavior. Betty was friendly and inviting during observations. She and Samuel had no concerns regarding her behaviors and moods. She did discuss the need to decrease her involvement in community organizations in preparation for worsening cognition in the coming years, but she was accepting of this transition and had adapted her home and daily routines to support her needs.

Adam

Adam, aged 85 with a diagnosis of MCI, had lived in his home for decades, and he and his wife planned renovations to their home to enhance access and safety.

Sensory profile. Adam's profile indicated that he was considered "*more than most*" others of his age in the area of low registration. This indicated that he needed a higher level of sensory input to perceive stimuli. His wife accommodated this by adding large signs in their home to remind him to drink water and find his keys.

Environment. Adam described times when he felt overwhelmed by sensory stimuli from his environment which caused some avoidant and agitated behaviors. He preferred a quiet environment and actively sought out decreasing the volume when it was too high. He also required control over stimulation. For example, he enjoyed bright lights as long as he was in control of when they were on and where. He gave examples of becoming agitated when there was too much stimulation in his environment. His wife told a story of a time when he was asked to leave a grocery store because he felt out

of control of the noise level in the store and had lashed out at store staff.

Behavior. Adam demonstrated persistent behaviors to align his environment with his sensory preferences. He and his wife shared examples of increased stress and agitation when he was not able to accommodate his environment to meet his sensory preferences and processing abilities, for example, he became agitated when the television was on while he talked on the phone, or he was restless when the lighting was not as he preferred when falling asleep. However, he was active in adjusting elements of the environment as needed (i.e., turning the television off).

Vicky

Vicky, aged 81 with a diagnosis of dementia, lived with her husband of 61 years. She enjoyed looking at family photos and visiting with her children and grandchildren.

Sensory profile. Vicky's ASP indicated that she sought sensation "*more than most*" and had sensory sensitivity "*less than most*" others within the normative data. These indicate that she preferred sensory stimuli and tolerated higher levels of sensory stimuli. These were evident in her desire for regular car rides and pictures covering the majority of the wall space within her home.

Environment. Vicky and her husband described a daily routine with prayer in the morning, television shows mid-morning, a drive and eating lunch out of the home, resting in the afternoon and a light meal for an early dinner each day. Vicky's husband described her desire to look through photo albums daily and routine requests to go out for a drive with him. He easily accommodated these requests and did not report any concerns regarding her behavior. She was pleasant and quiet throughout the interview and observations. Vicky said she liked to sweep and help in the kitchen, but her husband stated he preferred she did not do these tasks due to previous lower extremity joint replacements and back pain.

Behavior. Vicky was soft-spoken and kind throughout observations. Her husband described her as a wonderful wife and stated that she never acted out or had difficulty with behaviors. She navigated her home environment with ease and appeared to have good environmental fit within her home.

Charlotte

Charlotte, aged 88 with a diagnosis of MCI, was a care provider for her husband, *Ronald*, described below.

Sensory profile. Her ASP results indicated that she had sensory sensitivity, low registration, and sensation avoidance "*more than most*" for her age. Similar to Betty, Charlotte

needed to be in control of her environment to help accommodate a need to avoid sensation. She said she often spent an entire day in bed after a busy day prior, which may be related to sensory sensitivity and sensation avoiding tendencies. She also described situations, such as trouble with cooking and bumping into things, which are evident in those with low registration.

Environment. Charlotte said she enjoyed rearranging her home but felt that care for the home was beginning to be burdensome. She discussed increasing reliance on supports within her environment such as needing assistance for shopping trips and new troubles related to cooking (i.e., burning food and improper seasoning). She mentioned that she frequently bumped into items in her home and needed extra equipment (shower chair and grab bars) in the bathroom for stability. Charlotte required continual oxygen through a nasal canula and had long electrical cords throughout her living space.

Behavior. She described bouts of depression and explained that she would spend days in bed. She said she felt depressed and lonely and had little enthusiasm to do things outside of her home. Her husband described her as a social person, but she no longer had regular contact with her friends.

Ronald

Ronald, aged 83 with a diagnosis of MCI, the husband of Charlotte, was a retired salesman who enjoyed driving.

Sensory profile. Ronald's sensory profile indicated three domains that were different than most others his age. He had low registration "*less than most*," sensory sensitivity "*less than most*," and sensation avoiding "*much more than most*." Low registration indicates a need for increased sensory stimuli for perception. Lowered sensory sensitivity also indicates that he is not sensitive to increased sensory input. These are consistent with a tendency to interrupt in conversation and speak loudly despite normal hearing. Despite these areas being lowered, he has increased tendencies to be sensory avoiding, which indicated he did not prefer high amounts of stimuli. This was evident in his reluctance to grocery shop, resistance to moving or trying new situations, and a work history of traveling alone.

Environment. Ronald is happy in his home and hopes to remain living there for as long as possible. He said he was satisfied with the home and did not want to make changes to this living environment. Socially, he participated in several community activities, including a weekly breakfast with the guys at McDonalds, YMCA exercise classes, church, and offering transportation to disabled friends. Ronald continues to enjoy driving, an activity he did throughout his career as a regional sales representative which he reports would cause

him to be away from home at least three nights per week. Because of his love for driving, he would often help his friends if they needed transportation. For tasks in their home, Charlotte managed all finances, grocery shopping, and cooking. Ronald drove for the couple, did the laundry, and had a decreasing role in outside yard work, as they hired additional help.

Behavior. Ronald was welcoming and open during observations. He did tend to dominate conversations, and was, at times insistent that he was correct. Charlotte said they often bickered back and forth but got along, overall. He did describe times of becoming agitated when in highly stimulating environments, such as the grocery store, and so he avoided such places. Ronald was able to maintain appropriate environmental contexts to meet his needs and sensory preferences.

Pamela

Pamela was an active 67-year-old living with MCI. She lived with Alex, her husband of 50 years. She enjoyed community service and housework.

Sensory profile. Pamela's sensory profile results indicated that she has low registration "more than most" others her age, and the other three domains were "similar to most" others of her age group. During observation, she talked about the need for strong sensory input, including her desire to have candles in her home and preference to have the television on for sound. Her craft room was also full of supplies but organized. She discussed avoidance of large box stores, and she frequently stays home while her husband runs errands.

Environment. Pamela and Alex had lived in their current home for most of their marriage. Pamela said she felt safe in her home, and that she enjoyed working on projects in her space. They participated in regular exercise together three to 4 days per week. Pamela and Alex engaged in weekly volunteering with meal delivery for their church. Pamela said she liked to help navigate, and her husband drives. They both attend the local senior citizen's center for social engagement. Alex manages all finances and grocery shopping, and Pamela helps with cooking, cleaning, and other housework.

Behavior. Pamela was friendly and inviting during observations. She presented with some perseveration during conversation and daily tasks, but this did not seem to interfere with her daily function. Pamela described a lifetime of challenges with psychiatric conditions that often caused her "fly off the handle." She said she often became agitated and anxious over minute details. Her husband stated that he remained calm and "let her blow off steam." These abrupt changes may have been related to lack of

environmental fit or her inability to maintain a good fit as elements of her environment changed, such as noise level or disorganization.

Walter

Walter, age 82 with a diagnosis of MCI, lives with Laura, his wife of 48 years.

Sensory profile. Low registration was "more than most" others his age. All other areas of sensory processing and patterns were reported as "similar to most" others. Low registration is evident by his tolerance of clutter throughout his home and yard, along with frequent small car accidents, as he may not register all needed environmental precautions while driving.

Environment. Walter and Laura lived in a large three-story home. They had limited outside support or help from their family or social groups. The home was cluttered and needed home improvements, such as painting and small repairs. They both agreed that the house had become difficult to care for and maintain. There was clutter throughout the house that presented numerous tripping hazards. Laura said Walter liked to get started on house projects but was not likely to finish or often made bigger problems in the home. She gave the example of when he attempted to re-tile the bathroom wall, and she had to call in a contractor to repair damage he inadvertently caused.

Behavior. Walter had a passion for cars and driving, despite recent accidents and growing concerns from Laura for his safety. She talked about struggles in their partnership in terms of communication. During observations, he talked loudly, often over Laura, was agitated, and had difficulty with transition between topics. Laura said he became aggressive at times. Walter's situation is an example of an environment not supporting his needs and preferences, whereby impacting his behavior.

Discussion

This study explored sensory processing of community-dwelling adults with cognitive impairment as a potential mechanism to better understand behavioral presentation in the home environment. Seven of the nine cognitively impaired participants presented with sensory processing abnormalities. Several individuals employed self-identified environmental adaptations that were in line with their sensory processing preferences. These environmental adaptations led to greater behavioral regulation in the home context; a novel perspective for assessment and intervention for BPSD.

Findings of this study highlight the unique contributions of sensory processing as a potential mechanism of behavior for older adults with cognitive impairment (Kovach et al., 2004).

Prevalence and implications of sensory processing impairment in older adults, especially those with cognitive impairment, has not been fully explored in existing literature (Champagne, 2018). However, evidence of sensory processing abnormalities creates a realm of assessment and interventions that may prove beneficial for BPSD in the earliest stages of cognitive impairment. For example, Adam often became agitated when overstimulated by sound. Without identification of this sensory preference, his behavioral response could manifest as a “problem behavior,” causing additional distress for him and his spouse and potentially leading to unnecessary or inappropriate intervention. Awareness of sensory processing abilities and preferences has the potential to inform small but significant modifications and adjustments to the environment to decrease behavioral symptoms for older adults with cognitive impairment (Crowe, 2014).

Several participants engaged in self-initiated and/or care partner-initiated environmental adaptations to accommodate for sensory processing ability and preference. Those who implemented environmental adaptations demonstrated and reported less behavioral disturbance in the home when compared to participants who did not engage in such adaptations (i.e., clutter throughout home, lack of control over noise volume). A person-centered approach toward tailoring environmental adaptation to support behavior may be most advantageous due to the unique complexities involved in the situational contexts of the home environment (Hung et al., 2016; Wang et al., 2019). Further, communication of such preference, if relocation to institutional care is warranted, may improve transition and integration into the new living environment (Ashbourne et al., 2021).

Use of sensory systems to regulate behavior involves identifying environment and sensory-based techniques to target behavior for persons with cognitive impairment (American Occupational Therapy Association, 2017). Expertise, such as that offered by occupational therapy, could provide support beyond what individuals can do on their own to adapt and facilitate a supportive home environment for management of BPSD over time (Dunn & Bennett, 2002). Despite practice guideline recommendations, there are limited clinical tools for rigorous assessment and intervention related to sensory processing and sensory modulation for persons with ADRD (Champagne, 2018).

Our findings add to growing evidence of the need for supporting sensory-based assessment and intervention in late-life neurodegenerative disorders as a potential mechanism of action in behavioral expression for persons with MCI, as well as need for increased methods for evaluation of the situational context related to BPSD (Butler et al., 2020; Gitlin et al., 2012). Although replication with larger, more diverse samples is necessary, these findings offer preliminary description of sensory processing as it relates to behavior in the home environment. Future studies are needed to understand the nature of sensory processing and subtleties of environmental

adaptation in relation to BPSD, as well as potential avenues for treatment, such as nonpharmacological, occupational therapy interventions, to improve quality of life and enable prolonged residence in familiar settings for older adults with cognitive impairment.

Limitations

This study has limitations. The authors attempted to recruit participants with a dementia diagnosis; however, care partners of those with advanced dementia declined to participate in the current study. This limited observation of the breadth of cognitive impairment provides continued opportunity for investigation. Future research is needed to confirm behavioral phenotypes and the relationship to sensory processing in later stages of cognitive impairment.

Conclusion

Onset and progression of altered sensory processing as a symptom of neurodegenerative cognitive impairment may have significant implications for behavior and environmental interaction. Findings of this study offer evidence of reduced sensory processing abilities as a potential mechanism of behavior in the home environment of adults with MCI. A rigorous, nonpharmacological, therapeutic approach to assessment and intervention of sensory processing as it relates to behaviors of individuals living with cognitive impairment could provide opportunity for improved quality of life for all individuals involved. Additional research is needed to define subtleties of environmental adaptation to support older adults with MCI as they age in their home environment.

Acknowledgments

This study utilized University of Kentucky Alzheimer’s Disease Research Center. The authors acknowledge and thank all participants.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Funding for the longitudinal cohort is provided by NIH/NIA under Grant NIH/NIA P30 AG072946; and the first author is funded by the NIH under Grant NIH T32 AG057461: “Training in Translational Research in Alzheimer’s and Related Dementias (TRIAD).”

Ethical Approval

University of Kentucky IRB Protocol: 46351.

ORCID iD

Elizabeth K. Rhodus  <https://orcid.org/0000-0001-5108-0766>

Supplemental Material

Supplemental material for this article is available online.

References

- Albert, M. S., DeKosky, S. T., Dickson, D., Dubois, B., Feldman, H. H., Fox, N. C., Gamst, A., Holtzman, D. M., Jagust, W. J., Petersen, R. C., Snyder, P. J., Carrillo, M. C., Thies, B., & Phelps, C. H. (2011). The diagnosis of mild cognitive impairment due to Alzheimer's disease: Recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimer's & Dementia: The Journal of the Alzheimer's Association*, 7(3), 270–279. <https://doi.org/10.1016/j.jalz.2011.03.008>
- Alzheimer's Association. (2020). 2020 Alzheimer's disease facts and figures. *Alzheimer's & Dementia*, 16(3), 391–460. <https://doi.org/https://doi.org/10.1002/alz.12068>
- American Occupational Therapy Association. (2017). *Occupational therapy practice guidelines for adults with Alzheimer's Disease and related major neurocognitive disorders*. AOTA Press.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). American Psychiatric Association.
- Ashbourne, J., Boscart, V., Meyer, S., Tong, C. E., & Stolee, P. (2021). Health care transitions for persons living with dementia and their caregivers. *BMC Geriatrics*, 21(1), 285. <https://doi.org/10.1186/s12877-021-02235-5>
- Boeije, H. (2002). A purposeful approach to the constant comparative Method in the analysis of qualitative interviews. *Quality and Quantity*, 36(4), 391–409. <https://doi.org/10.1023/A:1020909529486>
- Brown, C., & Dunn, W. (2002). *Adolescent/Adult Sensory Profile user's manual*. Pearson.
- Brown, C., Tollefson, N., Dunn, W., Cromwell, R., & Filion, D. (2001). The adult sensory profile: Measuring patterns of sensory processing. *American Journal of Occupational Therapy*, 55(1), 75–82. <https://doi.org/10.5014/ajot.55.1.75>
- Butler, M., Gaugler, J., Talley, K., Abdi, H., Desai, P., Duval, S., Forte, M., Nelson, V., Ng, W., Ouellette, J., Ratner, E., Saha, J., Shippee, T., Wagner, B., Wilt, T., & Yesli, L. (2020). *Care interventions for people living with dementia and their caregivers. Comparative Effectiveness Review No. 231*. Agency for Healthcare Research and Quality. AHRQ Publication No. 20-EHC023 <https://doi.org/10.23970/AHRQEPCCER231>
- Champagne, T. (2018). *Sensory modulation in dementia care: Assessment and activities for sensory enriched care*. Jessica Kingsley Publishers.
- Chung, J. C. (2006). Measuring sensory processing patterns of older Chinese people: Psychometric validation of the adult sensory profile. *Aging & Mental Health*, 10(6), 648–655. <https://doi.org/10.1080/13607860600648080>
- Creswell, J. W. (2013). *Qualitative inquiry & research design: Choosing among five approaches* (3rd ed.). SAGE.
- Crowe, J. (2014). Professional reasoning and occupational therapist's use of a multi-sensory environment for clients with dementia. *Physical & Occupational Therapy In Geriatrics*, 32(3), 198–209. <https://doi.org/10.3109/02703181.2014.927947>
- Cummings, J. L. (1997). The Neuropsychiatric Inventory: Assessing psychopathology in dementia patients. *Neurology*, 48(5 Suppl 6), S10–S16. https://doi.org/10.1212/wnl.48.5_suppl_6.10s
- Dunn, W. (1997). The impact of sensory processing abilities on the daily lives of young children and their families: A Conceptual Model. *Infants & Young Children*, 9(4), 23–35. <https://doi.org/10.1097/00001163-199704000-00005>
- Dunn, W., & Bennett, D. (2002). Patterns of sensory processing in children with attention deficit hyperactivity disorder. *OTJR: Occupation, Participation and Health*, 22(1), 4–15. <https://doi.org/10.1177/153944920202200102>
- Dunn, W., & Westman, K. (1997). The sensory profile: The performance of a national sample of children without disabilities. *American Journal of Occupational Therapy*, 51(1), 25–34. <https://doi.org/10.5014/ajot.51.1.25>
- Engel-Yeger, B., & Rosenblum, S. (2021). Executive dysfunctions mediate between altered sensory processing and daily activity performance in older adults. *BMC Geriatrics*, 21(1), 132. <https://doi.org/10.1186/s12877-021-02032-0>
- Gerlach, L. B., & Kales, H. C. (2020). Managing behavioral and psychological symptoms of dementia. *Clinics in Geriatric Medicine*, 36(2), 315–327. <https://doi.org/10.1016/j.cger.2019.11.010>
- Gitlin, L. N., Kales, H. C., & Lyketsos, C. G. (2012). Non-pharmacologic management of behavioral symptoms in dementia. *JAMA*, 308(19), 2020–2029. <https://doi.org/10.1001/jama.2012.36918>
- Hughes, C. P., Berg, L., Danziger, W. L., Coben, L. A., & Martin, R. L. (1982). A new clinical scale for the staging of dementia. *British Journal of Psychiatry*, 140, 566–572. <https://doi.org/10.1192/bjp.140.6.566>
- Hung, L., Chaudhury, H., & Rust, T. (2016). The effect of dining room physical environmental renovations on person-centered care practice and residents' dining experiences in long-term care facilities. *Journal of Applied Gerontology*, 35(12), 1279–1301. <https://doi.org/10.1177/0733464815574094>
- Jensen, L., & Padilla, R. (2017). Effectiveness of environment-based interventions that address behavior, perception, and falls in people with Alzheimer's disease and related major neurocognitive disorders: A systematic review. *American Journal of Occupational Therapy*, 71(5), 7105180030p7105180031–7105180030p7105180010. <https://doi.org/10.5014/ajot.2017.027409>
- Kales, H. C., Gitlin, L. N., & Lyketsos, C. G. (2014). Management of neuropsychiatric symptoms of dementia in clinical settings: Recommendations from a multidisciplinary expert panel. *Journal of American Geriatric Society*, 62(4), 762–769. <https://doi.org/10.1111/jgs.12730>

- Kovach, C. R., Taneli, Y., Dohearty, P., Schlidt, A. M., Cashin, S., & Silva-Smith, A. L. (2004). Effect of the BACE intervention on agitation of people with dementia. *The Gerontologist*, 44(6), 797–806. <https://doi.org/10.1093/geront/44.6.797>
- Lawton, M. P., & Nahemow, L. (1973). Ecology and the aging process. In *The psychology of adult development and aging*. American Psychological Association. <https://doi.org/10.1037/10044-020>
- Lewin, K. (1936). *Principles of topological psychology*. McGraw-Hill.
- Luo, Y., He, P., Guo, C., Chen, G., Li, N., & Zheng, X. (2018). Association between sensory impairment and dementia in older adults: Evidence from China. *Journal of American Geriatric Society*, 66(3), 480–486. <https://doi.org/10.1111/jgs.15202>
- Magasi, S., Wong, A., Gray, D. B., Hammel, J., Baum, C., Wang, C. C., & Heinemann, A. W. (2015). Theoretical foundations for the measurement of environmental factors and their impact on participation among people with disabilities. *Archives of Physical Medicine and Rehabilitation*, 96(4), 569–577. <https://doi.org/10.1016/j.apmr.2014.12.002>
- Morgan, D. L. (2007). Paradigms lost and pragmatism regained: Methodological implications of combining qualitative and quantitative methods. *Journal of Mixed Methods Research*, 1(1), 48–76. <https://doi.org/10.1177/2345678906292462>
- Morris, J. C. (1993, November). The clinical dementia rating (CDR): Current version and scoring rules. *Neurology*, 43(11), 2412–2414. <https://doi.org/10.1212/wnl.43.11.2412-a>
- Nielsen, L., Riddle, M., King, J. W., , NIH Science of Behavior Change Implementation Team Aklin, W. M., Chen, W., Clark, D., Collier, E., Czajkowski, S., Esposito, L., Ferrer, R., Green, P., Hunter, C., Kehl, K., King, R., Onken, L., Simmons, J. M., Stoeckel, L., Stoney, C., & Weber, W. (2018). The NIH science of behavior change program: Transforming the science through a focus on mechanisms of change. *Behaviour Research and Therapy*, 101, 3–11. <https://doi.org/10.1016/j.brat.2017.07.002>
- Padilla, R. (2011). Effectiveness of environment-based interventions for people with Alzheimer's disease and related dementias. *American Journal of Occupational Therapy*, 65(5), 514–522. <https://doi.org/10.5014/ajot.2011.002600>
- Pinto, J. M., Wroblewski, K. E., Huisingh-Scheetz, M., Correia, C., Lopez, K. J., Chen, R. C., Kern, D. W., Schumm, P. L., Dale, W., & McClintock, M. K. (2017). Global sensory impairment predicts morbidity and mortality in older U.S. adults. *Journal of American Geriatric Society*, 65(12), 2587–2595. <https://doi.org/10.1111/jgs.15031>
- Ravn, M. B., Klingberg, T., & Petersen, K. S. (2018). The Adult Sensory Profile™ in care homes targeting people diagnosed with dementia: A qualitative study from the care provider perspective. *Rehabilitation research and practice*, 2018, 5091643. <https://doi.org/10.1155/2018/5091643>
- Researchware, I. (2018). *HyperRESEARCH 4.0.2*. [Computer Software] <http://www.researchware.com/>
- Sánchez, A., Marante-Moar, M. P., Sarabia, C., de Labra, C., Lorenzo, T., Maseda, A., & Millán-Calenti, J. C. (2016). Multisensory stimulation as an intervention strategy for elderly patients with severe dementia: A Pilot Randomized Controlled Trial. *Am J Alzheimers Disease Other Dementias*, 31(4), 341–350. <https://doi.org/10.1177/1533317515618801>
- Schmitt, F. A., Nelson, P. T., Abner, E., Scheff, S., Jicha, G. A., Smith, C., Cooper, G., Mendiondo, M., Danner, D. D., Van Eldik, L. J., Caban-Holt, A., Lovell, M. A., & Kryscio, R. J. (2012). University of Kentucky Sanders-Brown healthy brain aging volunteers: donor characteristics, procedures and neuropathology. *Current Alzheimer's Research*, 9(6), 724–733. <https://doi.org/10.2174/156720512801322591>
- Stark, S. L., Somerville, E. K., & Morris, J. C. (2010). In-home occupational performance evaluation (I-HOPE). *American Occupational Therapy Association*, 64(4), 580–589. <https://doi.org/10.5014/ajot.2010.08065>
- Sun, Y., Xu, W., Chen, K.-L., Shen, X.-N., Tan, L., & Yu, J.-T., the Alzheimer's Disease Neuroimaging, I. (2021). Mild behavioral impairment correlates of cognitive impairments in older adults without dementia: Mediation by amyloid pathology. *Translational Psychiatry*, 11(1), 577. <https://doi.org/10.1038/s41398-021-01675-2>
- Topaz, M., Adams, V., Wilson, P., Woo, K., & Ryvicker, M. (2020). Free-text documentation of dementia symptoms in home healthcare: A natural language processing study. *Gerontology and Geriatric Medicine*. <https://doi.org/10.1177/2333721420959861>
- Wang, J., Wu, B., Bowers, B. J., Lepore, M. J., Ding, D., McConnell, E. S., & Corazzini, K. N. (2019). Person-centered dementia care in China: A Bilingual literature review. *Gerontology and Geriatric Medicine*. <https://doi.org/10.1177/2333721419844349>
- Winblad, B., Palmer, K., Kivipelto, M., Jelic, V., Fratiglioni, L., Wahlund, L.-O., Nordberg, A., Bäckman, L., Albert, M., Almkvist, O., Arai, H., Basun, H., Blennow, K., de Leon, M., DeCarli, C., Erkinjuntti, T., Giacobini, E., Graff, C., Hardy, J., & Petersen, R. (2004). Mild cognitive impairment - Beyond controversies, towards a consensus: Report of the International working group on mild cognitive impairment. *Journal of Internal Medicine*, 256(3), 240–246. <https://doi.org/10.1111/j.1365-2796.2004.01380.x>
- Wise, E. A., Rosenberg, P. B., Lyketsos, C. G., & Leoutsakos, J. M. (2019). Time course of neuropsychiatric symptoms and cognitive diagnosis in National Alzheimer's Coordinating Centers volunteers. *Alzheimer's & Dementia: Diagnosis, Assessment & Disease Monitoring*, 11, 333–339. <https://doi.org/https://doi.org/10.1016/j.dadm.2019.02.006>
- Zeisel, J., Silverstein, N. M., Hyde, J., Levkoff, S., Lawton, M. P., & Holmes, W. (2003). Environmental correlates to behavioral health outcomes in Alzheimer's special care units. *The Gerontologist*, 43(5), 697–711. <https://doi.org/10.1093/geront/43.5.697>
- Zhao, Q. F., Tan, L., Wang, H. F., Jiang, T., Tan, M. S., Tan, L., Xu, W., Li, J. Q., Wang, J., Lai, T. J., & Yu, J. T. (2016). The prevalence of neuropsychiatric symptoms in Alzheimer's disease: Systematic review and meta-analysis. *Journal of Affective Disorders*, 190, 264–271. <https://doi.org/10.1016/j.jad.2015.09.069>