



Please cite the Published Version

El Mohtadi, Mohamed, Vagg-Whitehead, Kathryn , Belboul, Amina, Fadel, Abdulmannan and Ashworth, Jason  (2023) Estrogen stimulates the clearance of wound bacteria by monocyte-derived macrophages via activation of estrogen-receptor alpha and actin cytoskeleton reorganization. In: Cold Spring Harbor Laboratory (CSHL): Microbial Pathogenesis and Host Response, 11 September 2023 - 15 September 2023, Cold Spring Harbor Laboratory, New York, USA. (Unpublished)

Version: Accepted Version

Downloaded from: <https://e-space.mmu.ac.uk/633340/>

Usage rights:  In Copyright

Additional Information: This is an abstract of a poster presentation given at Cold Spring Harbor Laboratory (CSHL): Microbial Pathogenesis and Host Response

Enquiries:

If you have questions about this document, contact rsl@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Estrogen stimulates the clearance of wound bacteria by monocyte-derived macrophages via activation of estrogen-receptor alpha and actin cytoskeleton reorganisation

Mohamed El Mohtadi¹, Kathryn Whitehead², Amina Belboul², Abdulmannan Fadel³ and Jason Ashworth².

¹ Department of Biology, Edge Hill University, Ormskirk, United Kingdom.

² Centre for Bioscience, Manchester Metropolitan University, Manchester, United Kingdom.

³ School of Sport and Exercise Sciences, John Moores University, Liverpool, United Kingdom.

Mohamed El Mohtadi (elmohtam@edgehill.ac.uk)

Kathryn Whitehead (k.a.Whitehead@mmu.ac.uk)

Amina Belboul (amina.belboul@stu.mmu.ac.uk)

Abdulmannan Fadel (A.Fadel@ljmu.ac.uk)

Jason Ashworth (J.Ashworth@mmu.ac.uk)

Abstract

Background:

Chronic wounds in the elderly often become infected, leading to substantial morbidity and mortality. Age-related impaired healing is mediated by age-related changes in steroid hormones, particularly declining levels of estrogen with increasing age. Although the anti-inflammatory activity of estrogen has been defined, little is known about the effects of estrogen deprivation on bacterial clearance. The aim of this study was to determine the effect of ageing (estrogen deprivation) on the ability of human monocyte-derived macrophages to eliminate bacteria via phagocytosis.

Materials/methods:

Host-pathogen assays were used to measure macrophage-mediated phagocytosis of two major wound pathogens, methicillin-resistant *Staphylococcus aureus* (MRSA) and *Pseudomonas aeruginosa*, under *in vitro* and *ex vivo* conditions that model estrogen levels in the elderly, young adults and following estrogen supplementation. Fluorescence and scanning electron microscopy (SEM) were used to visualise host-pathogen interactions and protein mediators of phagocytosis were measured by immunoblotting.

Results:

Estrogen at concentrations typical of youth or supraphysiological levels significantly ($P < 0.05$) increased the phagocytosis of MRSA and *P. aeruginosa* in a dose-dependent manner compared to estrogen deprivation with significantly enhanced clearance of bacteria by M1 macrophages compared to M2 macrophages. Epifluorescence, confocal and SEM confirmed estrogen increases co-localisation of fluorescent GFP-*S. aureus* or mCherry-*P. aeruginosa* within macrophages and promotes bacterial internalisation. Activation of estrogen receptor-alpha (ER- α) mirrored the stimulatory effect of estrogen on phagocytosis whilst ER- α antagonism significantly ($P < 0.01$) blocked the phagocytic effect of estrogen. In contrast, activation of ER-beta (ER- β) had no significant effect on phagocytosis, confirming estrogen mediates bacterial clearance via ER- α . Immunoblotting analysis demonstrated that estrogen-enhanced phagocytosis is associated with altered levels of mediators involved in the actin cytoskeleton of phagocytes including increased levels of FAK, Rac1, Cdc42 and RhoG, but reduced levels of RhoA.

Conclusion:

Findings suggest estrogen may promote the resolution of wound infections during youth but this protection is lost as estrogen levels decline with increasing age, resulting in increased propensity and progression of age-related wound infections. Thus, novel wound dressings providing estrogen supplementation or selective activation of ER- α and/or specific targeting of downstream mediators of the actin cytoskeleton may provide effective treatment options for infected wounds in the elderly.