

Citation for published version:

Mccluskey, A, Coles, S & Morgan, B 2023, 'Accurate Estimation of Diffusion Coefficients and their Uncertainties

from Computer Simulation', Simulation Meets Molecular Spectroscopy (SiMol) Meeting 2023, Abingdon, UK United Kingdom, 30/10/23 - 1/11/23.
Publication date: 2023
Link to publication
Publisher Rights CC BY

University of Bath

Alternative formats

If you require this document in an alternative format, please contact: openaccess@bath.ac.uk

General rightsCopyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy
If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


Download date: 04. Dec. 2023


ACCURATE ESTIMATION OF DIFFUSION COEFFICIENTS AND THEIR UNCERTAINTIES FROM COMPUTER SIMULATION


ANDREW R. MCCLUSKEY1,2, SAMUEL W. COLES3,4, BENJAMIN J. MORGAN3,4

1. SCHOOL OF CHEMISTRY, UNIVERSITY OF BRISTOL, CANTOCK'S CLOSE, BRISTOL BS8 1TS, UK. 2. EUROPEAN SPALLATION SOURCE ERIC, OLE MAALØES VEJ 3, 2200 KØBENHAVN N, DK. 3. DEPARTMENT OF CHEMISTRY, UNIVERSITY OF BATH, CLAVERTON DOWN, BATH, BA2 7AY, UK. 4. THE FARADAY INSTITUTION, QUAD ONE, HARWELL SCIENCE AND INNOVATION CAMPUS, DIDCOT, OX11 ORA, UK. EMAIL: ANDREW.MCCLUSKEY@BRISTÓL.AC.UK


THE OPTIMAL SCHEME TO ESTIMATE THE DIFFUSION COEFFICIENT REQUIRES **BOTH** VARIANCE AND COVARIANCE TO BE CONSIDERED.










A COVARIANT MULTI-DIMENSIONAL NORMAL DISTRIBUTION CAN BE USED AS A GENERATIVE MODEL FOR MEAN-SQUARED DISPLACEMENT...






DATA FROM MOLECULAR DYNAMICS SIMULATION CAN BE USED TO PARAMETERISE THE MODEL.







WE CAN SAMPLE THE PARAMETERISED DISTRIBUTION LIKELIHOOD WITH A HEAVISIDE PRIOR TO GET AN ESTIMATE OF THE DIFFUSION COEFFICIENT DISTRIBUTION, WHICH WOULD TYPICALLY REQUIRE **MANY 1000**≤ **OF SIMULATIONS**.

