

University of Groningen

Physical Demands of Tennis Across the Different Court Surfaces, Performance Levels and Sexes

Pluim, Babette M.; Jansen, Marleen G.T.; Williamson, Samuel; Berry, Cain; Camporesi, Silvia; Fagher, Kristina; Heron, Neil; van Rensburg, Dina C.Janse; Moreno-Pérez, Víctor; Murray, Andrew

Published in:
Sports Medicine

DOI:
[10.1007/s40279-022-01807-8](https://doi.org/10.1007/s40279-022-01807-8)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2023

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Pluim, B. M., Jansen, M. G. T., Williamson, S., Berry, C., Camporesi, S., Fagher, K., Heron, N., van Rensburg, D. C. J., Moreno-Pérez, V., Murray, A., O'Connor, S. R., de Oliveira, F. C. L., Reid, M., van Reijen, M., Saueressig, T., Schoonmade, L. J., Thornton, J. S., Webborn, N., & Arden, C. L. (2023). Physical Demands of Tennis Across the Different Court Surfaces, Performance Levels and Sexes: A Systematic Review with Meta-analysis. *Sports Medicine*, 53(4), 807-836. <https://doi.org/10.1007/s40279-022-01807-8>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Physical Demands of Tennis Across the Different Court Surfaces, Performance Levels and Sexes: A Systematic Review with Meta-analysis

Babette M. Pluim^{1,2,3,29,30} · Marleen G. T. Jansen^{4,5} · Samuel Williamson⁶ · Cain Berry⁷ · Silvia Camporesi⁸ · Kristina Fagher⁹ · Neil Heron^{10,11} · Dina C. Janse van Rensburg^{1,12} · Víctor Moreno-Pérez¹³ · Andrew Murray^{14,15} · Seán R. O'Connor¹⁶ · Fábio C. L. de Oliveira¹⁷ · Machar Reid¹⁸ · Miriam van Reijen^{4,19} · Tobias Saueressig²⁰ · Linda J. Schoonmade²¹ · Jane S. Thornton^{22,23} · Nick Webborn^{24,25,26} · Clare L. Ardern^{27,28}

Accepted: 21 December 2022 / Published online: 8 February 2023
© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2023

Abstract

Background Tennis is a multidirectional high-intensity intermittent sport for male and female individuals played across multiple surfaces. Although several studies have attempted to characterise the physical demands of tennis, a meta-analysis is still lacking.

Objective We aimed to describe and synthesise the physical demands of tennis across the different court surfaces, performance levels and sexes.

Methods PubMed, Embase, CINAHL and SPORTDiscus were searched from inception to 19 April, 2022. A backward citation search was conducted for included articles using Scopus. The PECOS framework was used to formulate eligibility criteria. Population: tennis players of regional, national or international playing levels (juniors and adults). Exposure: singles match play. Comparison: sex (male/female), court surface (hard, clay, grass). Outcome: duration of play, on-court movement and stroke performance. Study design: cross-sectional, longitudinal. Pooled means or mean differences with 95% confidence intervals were calculated. A random-effects meta-analysis with robust variance estimation was performed. The measures of heterogeneity were Cochrane Q and 95% prediction intervals. Subgroup analysis was used for different court surfaces.

Results The literature search generated 7736 references; 64 articles were included for qualitative and 42 for quantitative review. Mean [95% confidence interval] rally duration, strokes per rally and effective playing time on all surfaces were 5.5 s [4.9, 6.3], 4.1 [3.4, 5.0] and 18.6% [15.8, 21.7] for international male players and 6.4 s [5.4, 7.6], 3.9 [2.4, 6.2] and 20% [17.3, 23.3] for international female players. Mean running distances per point, set and match were 9.6 m [7.6, 12.2], 607 m [443, 832] and 2292 m [1767, 2973] (best-of-5) for international male players and 8.2 m [4.4, 15.2], 574 m [373, 883] and 1249 m [767, 2035] for international female players. Mean first- and second-serve speeds were 182 km·h⁻¹ [178, 187] and 149 km·h⁻¹ [135, 164] for international male players and 156 km·h⁻¹ 95% confidence interval [151, 161] and 134 km·h⁻¹ [107, 168] for international female players.

Conclusions The findings from this study provide a comprehensive summary of the physical demands of tennis. These results may guide tennis-specific training programmes. We recommend more consistent measuring and reporting of data to enable future meta-analysts to pool meaningful data.

Clinical Trial Registration The protocol for this systematic review was registered a priori at the Open Science Framework (Registration DOI <https://doi.org/10.17605/OSF.IO/MDWFY>).

Key Points

The mean match duration (best-of-3 sets) was around 1.5 h for international male and female players (all surfaces) and approximately 75 min for junior national male and female players (hard court).

The mean rally duration of international players was shortest on grass court with 4.3 s 95% confidence interval [3.1, 5.9] for male players and 5.7 s [4.8, 6.7] for female players, followed by hard court with 5.6 s [4.9, 6.5] and 6.4 s [5.0, 8.1 s], and clay court with 7.1 s confidence interval [6.2, 8.1] and 8.8 s [5.2, 15.0].

The mean effective playing time was around one fifth of the actual playing time in international male and female players and roughly one fourth in junior national players of both sexes.

The mean first- and second-serve speeds were higher in international male players (182 km·h⁻¹ [178, 187] and 149 km·h⁻¹ [135, 164]) than in female players (156 km·h⁻¹ 95% confidence interval [151, 161] and 134 km·h⁻¹ [107, 168]), and mean first-serve speeds were higher than second-serve speeds.

1 Introduction

Tennis is a multidirectional high-intensity intermittent sport for male and female individuals played across multiple indoor and outdoor surfaces. Globally, over 87 million people play tennis, of whom 59% are male and 41% are female [1]. The three main playing surfaces in tennis are hard court, clay court and grass court [2].

Tennis requires technical and tactical proficiency, mental skills and high physical performance levels for players to excel [3–5]. Elite tennis players need to perform both endurance and strength training to produce powerful strokes and handle the high intensities of play, including the many rapid accelerations, decelerations and changes of directions that occur over the length of a match (one or more hours) [3, 6].

Understanding the physical demands of tennis during match play is vital for developing young players and preparing elite players for competition [3]. However, the external loads are hard to generalise due to the unpredictable and varied nature of the sport and match play. For example, the duration of a tennis match is a crucial variable [7] but can vary greatly. Many factors influence match duration, including the scoring system (e.g. no-advantage scoring) [8, 9], match format (e.g. best-of-3 or best-of-5 sets), the quality of the opponent, and the playing surface and type of balls used [10]. These factors likely influence the physical demands of tennis.

Advances in technology, including visual tracking systems and wearable sensors, have made it easier to analyse the specific demands of tennis [11]. Match duration, the number of strokes and tennis-specific loads (e.g. acceleration, deceleration, change of direction, stroke speed) can now be quantified. These parameters allow a comprehensive overview of the external loads of tennis. Although there are substantial differences between matches, synthesising the data using a meta-analysis can provide valuable insights into the specific loads players encounter and manage.

Therefore, we aimed to describe and synthesise the physical demands of tennis. We reviewed the data across the different playing surfaces (clay, hard and grass court), performance levels (regional, national and international) and sex of tennis players (male and female).

2 Methods

The Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) statement [12] guided the reporting of this systematic review (Electronic Supplementary Material [ESM]). We registered the protocol for this systematic review a priori at the Open Science Framework (Registration DOI: <https://doi.org/10.17605/OSF.IO/MDWFY>).

2.1 Eligibility Criteria

The PECOS (Population, Exposure, Comparison, Outcome, and Study design) framework was used to define inclusion and exclusion criteria.

2.1.1 Inclusion Criteria

- (i) Population: male and female tennis players of regional, national or international playing level; juniors (aged ≤ 18 years) and adults.
- (ii) Intervention: singles tennis match play according to the International Tennis Federation (ITF) Association of Tennis Professionals (ATP), or Women's Tennis Association (WTA) rules.
- (iii) Comparison: sex (male/female), court surface (hard, clay, grass).
- (iv) Outcome: data collected included at least one parameter related to the duration of play (e.g. strokes, rallies, games, sets and matches), on-court movement (e.g. accelerations, decelerations, changes of direction, distance covered and running speed) or stroke performance (count and speed).
- (v) Study designs: cross-sectional, longitudinal.

Exclusion criteria:

- (i) Editorials, notes, letters, case reports and reviews.
- (ii) Articles on tennis during match play with modified rules (e.g. time-limited matches).
- (iii) Articles reporting on wheelchair tennis or doubles tennis.
- (iv) Articles providing physiological or biomechanical variables only.

2.2 Data Sources and Searches

A systematic literature search was performed in the bibliographic databases PubMed, Embase, CINAHL (via Ebsco) and SPORTDiscus (via Ebsco) from inception to 19 April, 2022. The search syntax was designed by a medical information specialist (LJS) with input from two authors (MGTJ and BMP). Search terms included controlled terms (MeSH in PubMed and Emtree in Embase, CINAHL Headings in CINAHL and Thesaurus terms in SPORTDiscus) and free-text terms. The following terms were used (including synonyms and closely related words) as index terms or free-text words: 'tennis' and search terms comprising 'athletic performance'. The search was performed without date or language restrictions. Duplicate articles were excluded by LS using Endnote X20.0.1 (Clarivate™), following the Amsterdam Efficient Deduplication method [13] and the Bramer method [14]. Additionally, a backward citation search was conducted for included articles using Scopus. The entire search strategy for all databases is detailed in the ESM.

2.3 Study Selection

Four pairs of reviewers (MGTJ and ML, MGTJ and BMP, BMP and SW, and SW and ML) independently screened all potentially relevant titles and abstracts for eligibility using Rayyan [15]. If deemed appropriate, the full-text article was checked for the eligibility criteria. Differences in judgement were resolved through a consensus procedure, and an independent reviewer (FCLO) was available to make a final decision if the reviewer pairs did not reach a consensus. The same four teams of independent reviewers conducted a full-text review of the screened articles to confirm eligibility. Disagreements regarding inclusion were resolved through discussion between reviewers, and once again, an independent reviewer (FCLO) was available to make final decisions if needed. The reviewers documented the reasons for exclusion at each stage.

2.4 Data Extraction

Two reviewers (BMP and MGTJ) independently extracted the following data related to the characteristics of the

included studies: name of the first author; the year of publication; country where the study was carried out; study design; study aim; study population; sample size (participants and matches); age; sex (% male); playing level; court surface; the assessment tool; comparison; and an overview of the outcome parameters of each study. Playing level was determined by the level of the tournaments the player participated in, i.e. regional, national or international, or by their ranking. Court surface was defined by the playing surface of the matches, i.e. hard, clay or grass.

Table 1 presents the data on the outcome parameters extracted from the articles. The relevant authors were contacted if data were missing or required clarification.

2.5 Methodological Quality Assessment

Two reviewers (BMP and SW) independently assessed the methodological quality of all included studies using the Joanna Briggs Institute checklist for analytical cross-sectional studies [17]. The non-analytical nature of the included studies required minor modifications to the checklist with the removal of item three ("Was the exposure measured in a valid and reliable way?") and the separation of item 2 ("Were the study subjects and the setting described in detail?") to assess descriptions of study participants and study settings separately. Therefore, the modified checklist comprised eight items, including questions on study inclusion criteria, participants, the setting, the condition, confounding factors, validity and reliability of the measurement technique, and appropriate statistical analysis. Each question was rated as 'yes', 'no' or 'unclear'. The reviewers discussed differences until they reached a consensus. The quality assessment outcome was not used to determine study inclusion or perform a sub-group analysis based on methodological quality or risk of bias.

2.6 Data Synthesis

The outcomes were analysed based on three overall categories: "male", "female" and "male vs female". We separated our analysis based on experience level, "national" or "international", to ensure reasonable (statistical) homogeneity of the studies.

For quantitative statistical analysis, pooled means or mean differences with 95% confidence intervals (CIs) were calculated for movement variables reported by three or more studies. A random-effects meta-analysis was performed with robust variance estimation to account for the dependence of the study means [18, 19]. We calculated a pooled mean or mean difference from the studies using the inverse variance method for pooling [20]. Measures of means were log-transformed for analysis and then back-transformed to ensure no implausible (i.e. negative) estimates were obtained [21]. The

Table 1 Outcome parameters for duration, on-court movement and stroke performance

Duration	Stroke performance (speed)
Match duration (min)	Peak serve speed (km·h ⁻¹)
Rally/stroke duration (s)	First-serve speed (km·h ⁻¹)
Effective playing time ^a (%)	Second-serve speed (km·h ⁻¹)
Work-to-rest ratio ^b	Groundstroke speed (km·h ⁻¹)
Points per game/set/match (#)	Forehand speed (km·h ⁻¹)
Games per set/match (#)	Backhand speed (km·h ⁻¹)
Sets per match (#)	
On-court movement	Stroke performance (count)
Accelerations per minute/speed zone/match (#)	Strokes per rally/game/set/match/second (#)
Decelerations per minute/speed zone/match (#)	Serves per game/set/match (#)
Direction changes per rally/match (#)	First serves per game/set/match (#)
Distance covered per point/game/set/match (m)	Second serves per game/set/match (#)
Distance covered per speed zone (m)	Forehands per game/set/match (#)
Distance covered per minute (m)	Backhands per game/set/match (#)
Distance covered per speed zone per minute (m)	Forehand volleys per game/set/match (#)
Average running speed (m·s ⁻¹)	Backhand volleys per game/set/match (#)
Peak running speed (m·s ⁻¹)	Overheads per game/set/match (#)

^aEffective playing time is the percentage of the total playing times against the real match time. It consists of the playing time, all the breaks between points, games and sets, and the rest periods during any other breaks (e.g. injury time out, heat rule, replay by Hawk-Eye [16] and discussion with chair umpire)

^bThe work-to-rest ratio is the ratio between rally duration and the rest periods between rallies and expressed as a quotient

standard error of the log-transformed mean was calculated with the formula $\sqrt{\frac{SD^2}{n \cdot \text{mean}^2}}$ [22].

The measures of heterogeneity used were Cochrane Q and the resulting chi-squared statistic, I^2 statistic and 95% prediction intervals (PIs). A 95% PI estimate is where the actual effects are expected for 95% of similar studies that may be conducted in the future. The estimate of the PI is imprecise if the number of studies is low [23].

We assumed a correlation value of $\rho=0.8$ for all analyses if correlation values were missing. We performed all calculations and graphics with the software R [24] and the extension packages ‘metafor’ [25] and ‘robumeta’ [19].

2.7 Qualitative Synthesis

We summarised the descriptive characteristics of each study and presented these in summary tables and text.

2.7.1 Subgroup Analysis

We considered the impact of the three main court surfaces used in tennis: hard court, grass court and clay court.

2.7.2 Sensitivity Analysis

As correlation values were unknown, a sensitivity analysis with a range of different correlation parameters was performed ($\rho=0, 0.2, 0.4, 0.6, 1.0$). Missing standard deviations were imputed as the median value of the included standard deviations in the corresponding analysis [26]. Sensitivity analyses were calculated without the studies with the imputed standard deviation.

3 Results

3.1 Search

The literature search generated 7736 references: 1925 in PubMed, 2056 in Embase, 1149 in CINAHL and 2606 in SPORTDiscus. After removing duplicates, 4162 papers remained. Based on reviewing the titles and abstracts, 155 articles that dealt with participants playing singles tennis matches and reported on the external loads of tennis (duration of play, on-court movement or stroke performance) were selected. After the full-text screening, a further 91 articles were excluded, leaving 64 articles for the review,




Fig. 1 Flow chart of the screening process. *ATP* Association of Tennis Professionals, *ITF* International Tennis Federation, *WTA* Women's Tennis Association

of which 42 articles were included in the quantitative synthesis (meta-analysis) (Fig. 1).

3.2 Characteristics of the Included Studies

The characteristics of the included articles are presented in the ESM. Most articles were either analytical ($n = 46$ studies) or descriptive ($n = 10$) cross-sectional studies. There were three pre-post studies, two cross-over trials, two repeated-measures experimental designs and one randomised controlled trial. Most research came from Spain ($n = 20$), followed by Australia ($n = 11$), the UK ($n = 8$) and the Czech Republic ($n = 5$). The sample size ranged from 8 to 1188 players and 7 to 6984 matches, with one

large study analysing 3715 best-of-5-set and 16,246 best-of-3-set matches. Most studies included international-level players ($n = 44$), followed by national-level ($n = 18$) and regional-level ($n = 2$) players. International-level players were primarily adults, whereas national-level players were mainly adolescents. More studies included male players ($n = 30$) than female players ($n = 7$), and 27 had both sexes.

Most studies assessed physical demands on hard court ($n = 48$), followed by clay court ($n = 26$), grass court ($n = 19$) and wood ($n = 1$). Regarding duration, rally and match duration were studied most often ($n = 25$), followed by the work-to-rest ratio ($n = 14$), effective playing time ($n = 13$) and rally pace ($n = 6$). From the on-court movements, distances covered (per match, set, game, point or minute) were studied most often ($n = 26$), followed by

Table 2 (Modified) JBI checklist score (%) of the studies included in this review ($n=64$)

Author(s) (year)	Item number and corresponding score								Yes	No	Unclear	JBI checklist score
	1	2a	2b	4	5	6	7	8				
Bergeron et al. (2007) [27]	N	Y	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Brown and O'Donoghue (2008) [28]	N	N	Y	Y	Y	Y	Y	N	5	3	0	62.5
Brown (2021) [29]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Carboch (2017) [30]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Carboch and Plachá (2018) [31]	N	Y	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Carboch et al. (2018) [32]	N	Y	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Carboch et al. (2019) [33]	N	Y	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Carboch et al. (2020) [34]	N	Y	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Cui et al. (2017) [35]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Cui et al. (2018) [36]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Cui et al. (2020a) [37]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Cui et al. (2020b) [38]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Fernández-Elias et al. (2020) [39]	N	Y	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Fernandez-Fernandez et al. (2007) [40]	N	Y	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Fernandez-Fernandez et al. (2008) [41]	N	Y	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Filipicic et al. (2015) [42]	Y	N	Y	Y	Y	N	Y	Y	6	2	0	75.0
Filipicic et al. (2021) [43]	N	Y	Y	Y	Y	N	Y	Y	6	2	0	7.0
Fitzpatrick et al. (2019) [44]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Galé-Ansodi et al. (2014) [45]	Y	U	Y	Y	N	Y	Y	N	5	2	1	6.5
Galé-Ansodi et al. (2016) [46]	N	N	Y	Y	N	N	U	Y	3	4	1	37.5
Galé-Ansodi et al. (2017a) [47]	N	Y	Y	Y	Y	Y	U	U	5	1	2	62.5
Galé-Ansodi et al. (2017b) [48]	N	N	Y	Y	Y	Y	Y	Y	6	2	0	75.0
Galé-Ansodi et al. (2018) [49]	N	Y	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Gallo-Salazar et al. (2015) [50]	Y	Y	Y	Y	Y	Y	Y	Y	8	0	0	100.0
Gallo-Salazar et al. (2019) [51]	Y	Y	Y	Y	Y	Y	Y	Y	8	0	0	100.0
Giles et al. (2021) [52]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Hoppe et al. (2014) [53]	Y	Y	Y	U	Y	Y	Y	Y	7	0	1	87.5
Hoppe et al. (2016) [54]	N	Y	Y	U	Y	Y	Y	Y	6	1	1	75.0
Hornery et al. (2007) [55]	N	Y	Y	Y	Y	Y	Y	U	6	1	1	75.0
Johnson and McHugh (2005) [56]	N	N	Y	Y	N	N	Y	N	3	5	0	37.5
Kilit and Arslan (2017) [57]	N	Y	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Kilit and Arslan (2018) [58]	N	Y	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Klaassen et al. (1998) [59]	N	N	Y	Y	Y	Y	U	Y	5	2	1	62.5
Kovalchik and Reid (2017) [60]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Lisi and Grigoletto (2021) [9]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Mackie et al. (2013) [61]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Maquirriain et al. (2016) [62]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Martínez-Gallego et al. (2013)	N	U	Y	Y	Y	Y	Y	Y	6	1	1	75.0
Martínez-Gallego et al. (2019)	N	U	Y	Y	Y	Y	Y	Y	6	1	1	75.0
McCarthy et al. (1998) [63]	N	Y	Y	Y	N	Y	Y	Y	6	2	0	75.0
Meffert et al. (2019) [64]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Mendez-Villanueva et al. (2007) [65]	N	Y	Y	N	N	Y	Y	Y	5	3	0	62.5
Morante and Brotherhood (2005) [66]	Y	N	Y	Y	Y	Y	Y	U	6	1	1	75.0
Moreno-Pérez et al. (2019) [67]	U	Y	Y	Y	Y	Y	Y	Y	7	0	1	87.5
Murphy et al. (2016) [68]	N	Y	Y	Y	Y	N	N	Y	5	3	0	62.5
Myers et al. (2016) [69]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
O'Donoghue and Ingram (2001) [70]	N	N	Y	Y	Y	Y	Y	Y	6	2	0	75.0
O'Donoghue and Liddle (1998) [71]	N	N	Y	Y	Y	Y	U	U	4	2	2	50.0
Pereira et al. (2017) [72]	N	N	Y	Y	Y	N	Y	Y	5	3	0	62.5

Table 2 (continued)

Author(s) (year)	Item number and corresponding score								Yes	No	Unclear	JBI checklist score
	1	2a	2b	4	5	6	7	8				
Pereira et al. (2016) [73]	N	U	Y	Y	Y	Y	Y	Y	6	1	1	75.0
Perri et al. (2018) [74]	U	Y	Y	Y	Y	Y	Y	U	6	0	2	75.0
Ponzano et al. (2017) [75]	N	Y	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Reid et al. (2016) [76]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Reilly and Palmer (1994) [77]	N	Y	Y	Y	N	N	Y	Y	5	3	0	62.5
Sánchez-Pay et al. (2021) [78]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Smith et al. (2018a) [79]	Y	Y	Y	Y	Y	Y	Y	Y	8	0	0	100.0
Smith et al. (2018b) [80]	Y	Y	Y	Y	Y	Y	Y	Y	8	0	0	100.0
Stare et al. (2015) [81]	N	Y	Y	Y	Y	Y	Y	N	6	2	0	75.0
Takahashi et al. (2006) [82]	N	N	Y	Y	Y	Y	Y	Y	6	2	0	75.0
Takahashi et al. (2009) [83]	N	N	Y	Y	Y	Y	Y	Y	6	2	0	75.0
Torres-Luque et al. (2011) [84]	Y	Y	Y	Y	Y	Y	Y	Y	8	0	0	100.0
Yusoff and Krasilshchikov (2021) [85]	N	N	N	Y	N	N	Y	Y	3	5	0	37.5
Whiteside and Reid (2017) [86]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Whiteside et al. (2015) [87]	Y	N	Y	Y	Y	Y	Y	Y	7	1	0	87.5
Number of studies applying the item	27	29	63	61	57	56	59	55				

Questions from the JBI Checklist: 1. Were the criteria for inclusion in the sample clearly defined? 2a. Were the study subjects described in detail? 2b. Was the setting described in detail? 4. Were objective, standard criteria used for measurement of the condition? 5. Were confounding factors identified? 6. Were strategies to deal with confounding factors stated? 7. Were the outcomes measured in a valid and reliable way? 8. Was appropriate statistical analysis used? *JBO* Joanna Briggs Institute, *N* No, *U* Unclear, *Y* Yes

average ($n = 13$) and peak running speed ($n = 12$), accelerations ($n = 7$), changes of direction ($n = 4$) and decelerations ($n = 3$). The most frequently assessed stroke counts were strokes per rally ($n = 15$), followed by strokes per match ($n = 3$), per game ($n = 3$), per minute ($n = 2$) and per second ($n = 1$). Serve speed was reported by five studies. Assessment tools used were video analysis ($n = 26$), followed by a Global Positioning System [GPS] ($n = 16$), match statistics ($n = 16$), Hawk-Eye [16] ($n = 5$), live observation ($n = 3$) and live analysis with a computerised scorebook ($n = 2$). Three studies used video analysis and GPS, and one study used match statistics and Hawk-Eye.

3.3 Methodological Quality Assessment

The overall (modified) Joanna Briggs Institute checklist scores ranged from 3/8 (37.5%) to 8/8 (100%) points (Table 2).

Among the 64 studies assessed, almost all (63/64; 98%) described the study setting in detail (item 2b), and 61/64 (95%) studies used objective and standard criteria for measuring the conditions (item 4). Most studies adequately identified the confounding variables (57/64; 89%, item 5), the strategies used to manage them (56/64; 88%, item 6) and used valid and reliable outcome measures (59/64; 92%, item 7). Only 42% of studies (27/64) reported the criteria used for inclusion in their research (item 1), and only 45% clearly

described the participants (29/64, item 2a). Most studies (55/64, 85%) used appropriate statistical analyses.

3.4 Quantitative Synthesis (Meta-analysis)

The data are presented as means or mean differences [95% CI] and for all surfaces with 95% PIs. The datasets used for analysis and the meta-analysis are available in the ESM. Brown [29], Reid [60, 76, 79, 80, 87], Stare [81], Carboch [34] and Filipcic [43] all kindly responded to our requests to provide missing data or clarification.

3.4.1 Results of Sensitivity Analyses

All analyses with different correlation parameters ($\rho = 0, 0.2, 0.4, 0.6, 1.0$) showed no impact on the results.

3.5 Duration of Play

3.5.1 Match Duration: International Level

Too few studies were available for pooling the match duration of best-of-5-set matches in male international players. The mean duration of best-of-3-set matches of international male and female players on all surfaces and on hard court are presented in Table 3. There was insufficient information for a separate meta-analysis of match duration on clay

and grass courts. The mean match duration (minutes) of international male players on all surfaces was 89.7 [79.0, 101.9] with a 95% PI of [65.7, 122.5], and on hard court 93.0 [73.1, 118.2], and of female players on all surfaces 88.0 [71.7, 107.9] with a 95% PI of [50.0, 154.7], and on hard court 98.9 [75.7, 129.1].

Three studies observed match duration of international male and female players in the same study. The match duration of best-of-3-set matches on hard court was slightly longer for male than for female players, with a mean difference of 8.8 min 95% CI [1.2, 16.4] and a 95% PI of [1.7, 16.0].

3.5.2 Match Duration: National Level

The mean match duration of junior national male and female players is presented in Table 3. There was no information for match duration on grass court of male players and on clay and grass courts of female players. The mean match duration (minutes) of national male players on hard court and clay court combined was 78.0 [69.2, 87.9], on hard court 76.6 [64.6, 90.9], on clay court 78.6 [70.3, 87.7], and of female players on hard court 72.7 [48.7, 108.7]. There were no studies on the match duration of adult national-level players. More details are provided in the ESM.

3.5.3 Rally Duration: Male Players

The mean rally duration (seconds) of international male players on all surfaces combined was 5.5 [4.9, 6.3] with a 95% PI of [3.3, 9.3]. Rally durations on hard, clay and grass courts were 5.6 [4.9, 6.5], 7.1 [6.2, 8.1], and 4.3 [3.1, 5.9] (Fig. 2).

The mean rally duration (seconds) for junior national male players for hard and clay courts combined was 8.5 [7.2, 10.0 s] with a 95% PI of [6.1, 11.9]. The mean rally duration of national male players was 12.0 s [11.4, 12.7] on clay court and 8.3 s [6.9, 10.0] on hard court, with no information available for grass court. There were no data on the rally duration of adult national male players.

3.5.4 Rally Duration: Female Players

The mean rally duration (seconds) of international female players on all surfaces was 6.4 [5.4, 7.6] with a 95% PI of [3.7, 11.0]. On hard, clay and grass courts, rally durations were 6.4 [5.0, 8.1], 8.8 [5.2, 15.0] and 5.7 [4.8, 6.7] (Fig. 3). There were too few studies for pooling rally duration for national-level female tennis players and no studies on the rally duration of adult national female players.

3.5.5 Rally Duration: Male vs Female Players

The mean difference in rally duration (seconds) of international male and female tennis players on all surfaces was -0.48 [$-2.0, 1.0$] with a 95% PI of [$-3.3, 2.3$]. On hard court, clay court and grass court, the mean differences were -0.4 [$-2.1, 1.3$], -2.0 [$-3.9, 0.0$] and -1.6 [$-5.3, 2.2$] (Fig. 4). These differences were not statistically significant. More details are provided in the ESM.

3.5.6 Effective Playing Time: International Level

The mean effective playing time (%) of international male players on all surfaces was 18.8 [15.8, 21.7], with a 95% PI of [11.7, 29.4], and on hard court, 18.2 [15.0, 22.1] (Table 4). The mean effective playing time of female players on all surfaces was 20.0 [17.2, 23.3], with a 95% PI of [13.2, 30.5], and on hard court 20.0 [15.4, 26.1]. There was insufficient information for a separate meta-analysis of male and female players' effective playing time on clay and grass courts.

When comparing the effective playing time of male and female players in the same study, we found no statistically significant differences across all surfaces or during play on hard court. The mean difference on all surfaces was -2.0 [$-6.1, 2.1$], with a 95% PI of [$-10.7, 6.6$], and on hard court -1.2 [$-4.9, 2.5$] (Table 4).

3.5.7 Effective Playing Time: National Level

The mean effective playing time (%) of junior national male players on hard court and clay was 26.2 95% CI [23.3, 29.4], with a 95% PI of [18.4, 37.2], and on hard court 25.6 [21.4, 30.7], with 95% PI [15.8, 41.6] (Table 4). Insufficient information was available for a separate meta-analysis of effective playing time on clay court, and no information was available for grass court. Based on two studies, effective playing time in national female players was 27.1%, based on two studies, with the CI not estimable. There were no data on the effective playing time of adult national-level players. More details are provided in the ESM.

3.6 On-Court Movement

3.6.1 Distance Covered: International Level

The Hawk-Eye system only measured distance covered during points and was mainly used for international players. The mean distances (metres) covered across all surfaces by international male players were 2292 [1767, 2973] with a 95% PI [1243, 4226] per best-of-5-set match, 607 [443, 831] with a

Table 3 Match duration of international and national male and female players on different court surfaces

Outcome (description)	Studies included in the meta-analysis (author(s), year)	Number of studies (<i>n</i>)	Effect size (95% CI)	Test for (sub-group) differences (<i>p</i> -value)	95% PI	<i>I</i> ² (%)	Sensitivity analysis for imputed SDs
Match duration (minutes): international level							
Men All surfaces Best-of-3	Filipcic et al. (2021) Hornery et al. (2007) Mackie (2013) Stare et al. (2015) Mendez-Villanueva et al. (2007) Moreno-Perez et al. (2019) Yusoff and Krasilshchikov (2021)	7	M: 89.7 (79.0, 101.9)	0.696	(65.7, 122.5)	61.9	M: 91.8 (79.0, 106.8)
Men Hard court Best-of-3	Filipcic et al. (2021) Hornery et al. (2007) Mackie (2013) Moreno-Perez et al. (2019) Stare et al. (2015)	5	M: 93.0 (73.1, 118.2)	NA	(52.9, 163.4)	88.1	M: 96.5 (69.8, 133.4)
Women All surfaces	Sánchez-Pay (2021) Fernandez-Fernandez et al. (2008) Fernandez-Fernandez et al. (2007) Mackie (2013) Morante and Brotherhood (2005)	5	M: 88.0 (71.7, 107.9)	0.77	(50.0, 154.7)	80.3	NA
Women Hard court	Fernandez-Fernandez et al. (2007) Mackie (2013) Morante and Brotherhood (2005) Sánchez-Pay (2021)	4	M: 98.9 (75.7, 129.1)	NA	(55.5, 176.1)	66.8	NA
Men vs women Hard court	Perri et al. (2016) Stare et al. (2015) Torres-Luque et al. (2011)	3	MD: 8.8 (1.2, 16.4)	0.039	(1.7, 16.0)	0	10.2 (−16.1, 36.5)
Match duration (minutes): national level (aged ≤ 18 years)							
Men Hard and clay court	Bergeron et al. (2007) Filipcic et al. (2021) Gallo-Salazar et al. (2019) Hoppe et al. (2014) Kilit and Arslan. (2017) Perri et al. (2016) Ponzano et al. (2017) Stare et al. (2015) Torres-Luque et al. (2011)	9	M: 78.0 (69.2, 87.9)	0.876	(52.5, 115.8)	84.75	M: 78.0 (69.2, 87.9)

Table 3 (continued)

Outcome (description)	Studies included in the meta-analysis (author(s), year)	Number of studies (<i>n</i>)	Effect size (95% CI)	Test for (sub-group) differences (<i>p</i> -value)	95% PI	<i>I</i> ² (%)	Sensitivity analysis for imputed SDs
Men Hard court	Bergeron et al. (2007) Filipic et al. (2021) Gallo-Salazar et al. (2019) Perri et al. (2016) Ponzano et al. (2017) Stare et al. (2015) Torres-Luque et al. (2011)	7	M: 76.6 (64.6, 90.9)	NA	(44.13, 133.02)	88.90	M: 76.6 (64.6, 90.9)
Men Clay court	Hoppe et al. (2014) Kilit and Arslan (2017) Ponzano et al. (2017)	3	M: 78.6 (70.3, 87.7)	NA	(61.65, 100.09)	0	NA
Women Hard court	Galé-Ansodi (2017a) Perri et al. (2016) Stare et al. (2015)	4	M: 72.7 (48.7, 108.7)	NA	(27.9, 189.8)	88.9	M: 75.9 (40.6, 141.9)

CI confidence interval, *I*² *I*-square statistic, *M* mean, *MD* mean difference, *min* minutes, *NA* not applicable, *PI* prediction interval, *SD* standard deviation

95% PI [279, 1322] per set, and 9.6 [7.6, 12.2] with a 95% PI [5.6, 16.6] per point. For female players, these numbers were 1249 [767, 2035] with a 95% PI [383, 4071] per best-of-3-set match, 573 [373, 883] with a 95% PI not estimable, and 8.3 [4.4, 15.5] with a 95% PI [1.7, 40.5] (Table 5). The distance covered per set in female players was based on only two studies. There was insufficient information available for a meta-analysis of distance covered by international male players in best-of-3-set matches.

Only the distance covered per best-of-5-set match on all surfaces and hard court could be calculated for male players. There were too few studies for female players to perform a meta-analysis on the distance per surface per match. Distances covered per set and per point were calculated for all surfaces only.

3.6.2 Distance Covered: National Level

The GPS measured the distance covered during points and moving between points, games and sets. It was primarily used for studies of national players.

The distances (metres) covered per match by junior national male players were 3313 [2870, 3826] on hard court and clay, 3200 [2322, 4411] on hard court and 3272 [3063,

3494] on clay court (Table 5). For female players, the distance covered per match was 2967 [2269, 3880] on hard court. The distances covered were reported only for hard court and clay court of male players and for hard court of female players; there was no information for distance covered on grass court.

The mean distance covered per minute on hard court was significantly higher for junior male players than for junior female players, with a mean difference of 2.4 m.min⁻¹ [1.4, 3.3].

No information was available for distance covered per set or point by junior national-level players, and no studies on the distance covered by adult national players. More details are provided in the ESM.

3.6.3 Running Speed: International Level

The average and peak running speed (m·s⁻¹) of international male tennis players on hard courts were 2.1 [0.6, 7.6] and 5.5 [4.0, 7.4] (Table 6). No or too few studies were available on average and peak running speed on clay or grass courts in male players. There was insufficient information available for a meta-analysis of the average and peak running speed of international female tennis players.




Fig. 2 Mean rally duration (seconds [s]) of international male tennis players on hard, grass and clay courts. Dashed lines on the forest plot diamonds represent the 95% prediction interval. CI confidence interval, NR not reported, RE random effects

3.6.4 Running Speed: National Level

The average and peak running speed ($m \cdot s^{-1}$) on hard court were 1.5 [0.6, 3.9] and 5.1 [4.3, 6.0] for junior national male players, and 2.9 [2.7, 3.1] and 4.2 [3.8, 4.6] for national female players (Table 6). No or insufficient data were available for running speed on grass court for male players and on clay and grass courts for female players. There were no data on the running speed of adult national players. More details are provided in the ESM.

3.7 Stroke Performance

3.7.1 Strokes Per Rally: International Level

The mean number of strokes per rally for international male players was 4.1 [3.4, 5.0] with a 95% PI of [2.5, 6.9] on all surfaces, 4.2 [3.4, 5.3] on hard court, and 4.8 [3.4, 6.7] on

clay court (Table 7). The mean number of strokes of female players was 3.9 [2.4, 6.2] with a 95% PI of [1.5, 9.8] on hard and clay courts combined. There were insufficient data available for a separate meta-analysis of strokes per rally on grass court in male players and on any surface in female players.

3.7.2 Strokes Per Rally: National Level

In junior national male players, the mean number of strokes per rally was 4.9 [3.2, 7.5] for hard and clay courts combined with a 95% PI of [1.72, 14.10], 4.6 [2.6, 8.2] for hard court, 6.5 [1.7, 25.2] for clay court and with no data for grass court (Table 7). In junior female players, the mean number of strokes per rally was 4.8 [1.1, 21.8], based on two studies. There were no data on adult national players. More details are provided in the ESM.




Fig. 3 Mean rally duration (seconds [s]) of international female tennis players on hard, grass and clay courts. *Dashed lines* on the forest plot *diagonals* represent the 95% prediction interval. *CI* confidence interval, *RE* random effects

3.7.3 Serve Speed: International Level

The mean first- and second-serve speed ($\text{km}\cdot\text{h}^{-1}$) on all surfaces were 182 [178, 187] and 149 [135, 164] for male players and 156 [151, 161] and 134 [107, 168] for female players (Table 8). Only two studies were available for the meta-analysis of the second-serve speed of female players. More details are provided in the ESM.

3.7.4 Serve Speed: National Level

There were no data on the serve speed of national male and female players.

3.8 Qualitative (Descriptive) Synthesis

Best-of-5-set match duration, games per set, points per match, work-to-rest ratio, accelerations, decelerations, changes of directions, serves per game, strokes per game and groundstroke speeds could not be included in the

meta-analysis because of missing data, or because only one or two studies were available per outcome parameter. The main results of these outcome variables are presented below (see ESM).

3.8.1 Match Duration

The median match duration (minutes) in male tennis players was 142 (75% quantile 179, mean 150) for best-of-5-set matches and 92 (75% quantile 119, mean 98) for best-of-3-set matches, based on 3712 and 16,246 matches, respectively [9]. We identified no study of similar size for female players.

3.8.2 Points, Games and Sets

At the four Grand Slams in 2016, the mean number of sets per match, games per set, points per game and points per match were 3.8, 9.8 [9.3, 10.3], 6.3 [6.1, 6.5] and 235 in male players (best-of-5), and 2.3, 9.3 [8.8, 9.8], 6.5 [6.3, 6.7] and




Fig. 4 Mean difference in rally duration (seconds [s]) between international male and female tennis players on hard, grass and clay courts. A minus sign signifies a shorter mean rally duration favour-

ing male players, and a plus sign favours female players. Dashed lines on the forest plot diamonds represent the 95% prediction intervals. *CI* confidence interval, *RE* random effects

143 in female players [30]. At Wimbledon (1992 to 1995), the mean number of sets per match, games per set, points per game and points per match were 3.7, 9.9, 6.1 and 230 in male players, and 2.3, 8.9, 6.5 and 132 in female players (95% CI not provided) [59].

Kovalchik and Reid [60] reported the median (interquartile range) number of points per match on the different court surfaces for junior and adult male and female players at the four Grand Slams in 2016, plus the 2017 Australian Open for juniors. The numbers on hard, clay and grass courts for junior male players were 130 (104–167), 131 (112–158) and 137 (118–162), and for junior female players, 122 (105–175), 122 (104–150) and 115 (100–177). For professional male players, these numbers were 216 (177–265), 213 (174–258) and 221 (180–275), and for female players, 129 (108–171), 133 (108–170) and 130 (111–169).

3.8.3 Work-to-Rest Ratio

In international male tennis players, the mean work-to-rest ratio varied from 1:3.5 to 1:5.6 on hard court [33, 43, 61], 1:2.0 to 1:3.5 on clay court [33, 65, 88] and from 1:3.7 to 1:5.0 on grass court [33, 88]. In national-level male players, the work-to-rest ratio ranged from 1:2.5 to 1:3.7 on hard court [43, 51, 58, 84] and was 1:2.1 on clay court [57, 58].

In international female players, the mean work-to-rest ratio varied from 1:3.4 to 1:4.5 on hard court [31, 61], 1:2.1 to 1:2.4 on clay court [41, 88] and 1:3.2 on grass court [88]. The work-to-rest ratio of national-level female tennis players on hard court was 1:2.2 [84].

Carboch et al. [34] reported an unusually high work-to-rest ratio that ranged from 1:5 to 1:7 in male and female players. However, after contacting the author, this turned out to be an error, and we included the correct numbers (range from 1:3.2 to 1:4.4) in the ESM.

Table 4 Effective playing time of international and national male and female players on different court surfaces

Sex court surface	Studies included in the meta-analysis (author(s), year)	Number of studies (<i>n</i>)	Effect size (95% CI)	Test for (sub-group) differences (<i>p</i> -value)	95% PI	<i>I</i> ² (%)	Sensitivity analysis for imputed SDs
Effective playing time (%): international level							
Men All surfaces	Yusoff and Krasilshchikov (2021) Filipcic et al. (2021) Mackie (2013) Mendez-Villanueva et al. (2007) Morante and Brotherhood (2005) O'Donoghue and Liddle (1998) Stare et al. (2015) Whiteside and Reid (2017)	8	M: 18.6 (15.8, 21.7)	0.563	(11.7, 29.4)	90.5	M: 18.9 (15.5, 23.0)
Men Hard court	Filipcic et al. (2021) Mackie (2013) Morante and Brotherhood (2005) Stare et al. (2015) Whiteside and Reid (2017)	5	M: 18.2 (15.0, 22.1)	NA	(10.4, 31.8)	92.5	M: 16.7 (13.2, 21.3)
Women All surfaces	Fernandez-Fernandez et al. (2007) Fernandez-Fernandez et al. (2008) Mackie (2013) Morante and Brotherhood (2005) O'Donoghue and Liddle (1998) Whiteside and Reid (2017)	6	M: 20.0 (17.2, 23.3)	0.469	(13.2, 30.5)	88.6	M: 19.6 (15.2, 25.2)
Women Hard court	Fernandez-Fernandez et al. (2007) Mackie (2013) Morante and Brotherhood (2005) Whiteside and Reid (2017)	4	M: 20.0 (15.4, 26.1)	NA	(10.4, 38.6)	92.3	M: 19.1 (12.6, 28.9)
Men vs women All surfaces	Mackie (2013) Morante and Brotherhood (2005) O'Donoghue and Liddle (1998) Whiteside and Reid (2017)	4	MD: -2.0 (-6.1, 2.1)	0.215	(-10.7, 6.6)	82.6	MD: -1.2 (-7.8, 5.5)
Men vs women Hard court	Mackie (2013) Morante and Brotherhood (2005) Whiteside and Reid (2017)	3	MD: -1.2 (-4.9, 2.5)	0.295	(-6.9, 4.5)	46.3	NA

3.8.4 Accelerations and Decelerations

The mean [95% CI] number of accelerations ($> 1.5 \text{ m}\cdot\text{s}^{-2}$) and decelerations ($< 1.5 \text{ m}\cdot\text{s}^{-2}$) in nine international

male tennis players was 18.2 [0, 40.6] and 19.1 [0, 38.9], measured using GPS and accelerometers [39]. The mean number of accelerations per minute ($> 3 \text{ m}\cdot\text{s}^{-2}$) in 12 national-level male junior tennis players measured using

Table 4 (continued)

Sex court surface	Studies included in the meta-analysis (author(s), year)	Number of studies (<i>n</i>)	Effect size (95% CI)	Test for (sub-group) differences (<i>p</i> -value)	95% PI	<i>I</i> ² (%)	Sensitivity analysis for imputed SDs
Effective playing time (%): national level (aged ≤ 18 years)							
Men Hard and clay court	Filipic et al. (2021) Gallo-Salazar et al. (2019) Kilit and Arslan (2017) Kilit and Arslan (2018) Stare et al. (2015) Torres-Luques et al. (2011)	6	M: 26.2 (23.3, 29.4)	0.86	(18.4, 37.2)	87.5	M: 25.9 (22.3, 30.1)
Men Hard court	Filipic et al. (2021) Gallo-Salazar et al. (2019) Kilit and Arslan (2018) Stare et al. (2015) Torres-Luques et al. (2011)	5	25.6 (21.4, 30.7)	NA	(15.8, 41.6)	90.7	M: 25.2 (19.5, 32.5)

CI confidence interval, *I*² I-square statistic, *M* mean, *MD* mean difference, *NA* not applicable, *PI* prediction interval, *SD* standard deviation

GPS was 0.3 [0.2, 0.4] in a morning match and 0.2 [0.1, 0.3] in the second match in the afternoon of the same day [51]. The mean number of accelerations (2 to < 4 m·s⁻² and ≥ 4 m·s⁻²) were 59 [33, 85] and 19 [0, 55] in 20 junior male players, and 62 [20, 104] and 42 [0, 106] in 20 adult regional-level male players, measured using GPS and accelerometers [54].

3.8.5 Changes of Direction

The mean [95% CI] number of changes in directions in 14 international male players was 2.5 [0.7, 4.3] on hard court and 2.4 [0.0, 5.0] on clay court [55], and 2.3 [0.0, 5.1] in eight female players on hard court [40]. At the 2012–17 Australian Open, the median (interquartile range) number of changes in direction was 6.0 (0–18) in junior male players (12 matches), 5.0 (0–22) in adult male players (21 matches), 5.0 (0–18) in junior female players (six matches) and 4.5 (0–18) in adult female players (21 matches) [60] (see ESM).

3.8.6 Stroke Counts

Whiteside and Reid reported on the number of shots per game in the first four rounds of the 2012–16 Australian Open tournaments in 18 male and 21 female players [86]. The mean number of strokes per game overall for rounds 1–4 were 15.0 [8.9, 21.2], 15.5 [9.0, 22.0], 17.4 [10.1, 24.7]

and 18.7 [9.9, 27.4] for male players, and 17.7 [11.5, 23.9], 18.2 [10.9, 25.5], 18.1 [11.9, 42.1] and 19.3 [11.8, 26.9] for female players.

The mean [95% CI] number of first serves per game in rounds 1 to 4 were 5.7 [5.1, 6.9], 5.8 [4.8, 6.9], 6.0 [4.9, 7.2] and 6.7 [4.7, 8.6] for the male players, and 6.6 [4.4, 8.7], 6.5 [5.2, 7.6], 6.5 [5.2, 7.9] and 6.9 [5.2, 8.7] for the female players. The mean number of second serves per game in rounds 1–4 were 2.1 [1.1, 3.0], 2.0 [1.1, 3.0], 2.1 [1.1, 3.1] and 2.4 [1.3, 3.6] for the male players, and 2.2 [1.3, 3.2], 2.4 [1.5, 3.3], 2.3 [1.2, 3.4] and 2.5 [1.3, 3.7] for the female players. Additional data are provided in the ESM.

3.8.7 Stroke Speeds

At the 2012–14 Australian Open, mean peak first-serve speeds were 206 km/h [183, 230] for male players and 172 km/h [151, 1993] for international female players [76]. The mean groundstrokes' speeds were 111 [100, 122] and 106 km/h [95, 118] for international male and female players [76]. Additional data are provided in the ESM.

4 Discussion

This systematic review synthesised the available evidence on the physical demands of tennis relating to the duration of play, on-court movement and stroke performance in national- and international-level male and female tennis

Table 5 Distance covered per match, set, point, and minute (m) of international and national male and female players on different court surfaces

Sex court surface	Studies included in the meta-analysis (author(s), year)	Number of studies (<i>n</i>)	Effect size (95% CI)	Test for (subgroup) differences (<i>p</i> -value)	95% PI	<i>I</i> ² (%)	Sensitivity analysis for imputed SDs
Distance covered per match (m)							
International level							
Men All surfaces Best-of-5	Cui et al. (2020a) Kovalchik and Reid (2017) Maquirriain et al. (2016) Reid et al. (2016)	4	M: 2292.3 (1767.4, 2973.1)	0.533	(1243.4, 4226.2)	85.2	M: 2382.1 (1567.5, 3620.1)
Men Hard court Best-of-5	Cui et al. (2020a) Kovalchik and Reid (2017) Reid et al. (2016)	3	M: 2164.3 (1775.4, 2638.4)	NA	(1372.68, 3412.53)	26.8	M: 2189.4 (1401.5, 3420.3)
Women All surfaces Best-of-3	Cui et al. (2018) Kovalchik and Reid (2017) Reid et al. (2016)	3	M: 1249.1 (766.6, 2035.4)	NE	(383.2, 4071.4)	87.0	M: 1304.8 (652.4, 2609.4)
National level (aged ≤ 18 years)							
Men Hard and clay court Best-of-3	Filipcic, 2021 Galé-Ansodi et al. (2017b) Gallo-Salazar et al. (2019) Hoppe et al. (2014) Hoppe et al. (2016) Kilit and Arslan (2017) Kilit and Arslan (2018) Perri et al. (2018)	8	M: 3313.6 (2870.1, 3825.7)	0.886	(2247.9, 4884.6)	85.6	NA
Men Hard court Best-of-3	Filipcic (2021) Galé-Ansodi et al. (2017b) Gallo-Salazar et al. (2019) Kilit and Arslan (2018) Perri et al. (2018)	5	M: 3200.7 (2322.3, 4411.4)	NA	(1363.1, 7515.5)	92.7	NA
Men Clay court Best-of-3	Hoppe et al. (2014) Hoppe et al. (2016) Kilit and Arslan (2017) Kilit and Arslan (2018)	4	M: 3272.3 (3063.7, 3495.0)	NA	(2940.5, 3641.5)	0	NA
Women Hard court Best-of-3	Galé-Ansodi et al. (2017a) Galé-Ansodi et al. (2017b) Perri et al. (2018)	3	M: 2966.8 (2268.6, 3879.8)	NA	(1473.1, 5975.1)	67.7	NA
Distance covered per set (m)							
International level							
Men All surfaces	Cui et al. (2020a) Pereira et al. (2017) Reid et al. (2016)	3	M: 607.0 (443.1, 831.5)	0.218	(278.6, 1322.6)	78.1	NA
Women All surfaces	Cui et al. (2020a) Reid et al. (2016)	2	M: 573.6 (372.7, 882.7)	NE	NE	84.7	NA

Table 5 (continued)

Sex court surface Best-of-3/5	Studies included in the meta-analysis (author(s), year)	Number of studies (<i>n</i>)	Effect size (95% CI)	Test for (subgroup) differences (<i>p</i> -value)	95% PI	<i>I</i> ² (%)	Sensitivity analysis for imputed SDs
Distance covered per point (m)							
International level							
Men All surfaces	Cui et al. (2020a) Filipcic (2021) Kovalchik and Reid (2017) Martínez-Gallego et al. (2019) Pereira et al. (2017) Whiteside et al. (2015)	6	M: 9.6 (7.6, 12.2)	0.925	(5.56, 16.57)	93.7	M: 9.5 (8.2, 11.0)
Men Hard court	Cui et al. (2020a) Filipcic (2021) Kovalchik and Reid (2017) Martínez-Gallego et al. (2019) Whiteside et al. (2015)	5	M: 9.7 (7.6, 12.3)	NA	(5.6, 16.8)	94.5	M: 9.5 (8.3, 11.0)
Women All surfaces	Cui et al. (2018) Kovalchik and Reid (2017) Reid et al. (2016)	3	M: 8.2 (4.4, 15.2)	NE	(1.70, 39.79)	96.7	9.4 (5.3, 16.7)
Women Hard court	Cui et al. (2018) Kovalchik and Reid (2017) Reid et al. (2016)	3	M: 8.3 (4.4, 15.5)	NE	(1.7, 40.5)	96.7	9.44 (4.78 18.63)
Distance covered per minute (m)							
National level (aged ≤ 18 years)							
Men Hard court	Fernández-Elias et al. (2020) Galé-Ansodi et al. (2017b) Galé-Ansodi et al. (2018) Gallo-Salazar et al. (2019) Perri et al. (2018)	5	M: 48.2 (45.3, 51.3)	NA	(41.5, 56.1)	66.3	NA
Women Hard court	Galé-Ansodi et al. (2017a) Galé-Ansodi et al. (2017b) Galé-Ansodi et al. (2018) Perri et al. (2018)	4	M: 45.4 (41.5, 49.7)	NA	(35.8, 57.7)	87.7	NA
Men vs women Hard court	Galé-Ansodi et al. (2017b) Galé-Ansodi et al. (2018) Perri et al. (2018)	3	MD: 2.4 (1.4, 3.3)	0.0111	(1.5, 3.2)	0	NA

CI confidence interval, *I*² I-square statistic, *m* metres, *M* mean, *MD* mean difference, *NA* not applicable, *NE* not estimable, *PI* prediction interval, *SD* standard deviation

players on hard, clay and grass courts. The main findings are summarised in Fig. 5.

4.1 Match Duration

The mean match duration (best-of-3 sets) of international tennis players was approximately 1 h and a half for male

Table 6 Average and peak running speed ($\text{m}\cdot\text{s}^{-1}$) of international and national male and female players on hard and clay courts

Outcome (description)	Studies included in the meta-analysis (author(s), year)	Number of studies (<i>n</i>)	Effect size (95% CI)	Test for (subgroup) differences (<i>p</i> -value)	95% PI	I^2 (%)	Sensitivity analysis for imputed SDs
Average running speed ($\text{m}\cdot\text{s}^{-1}$)							
International level							
Men Hard court	Martínez-Gallego et al. (2019) Martínez-Gallego et al. (2019) Reid et al. (2016) Filipcic (2021)	3	M: 2.1 (0.6, 7.6)	NA	NE	98.5	M: 2.0 (0.5, 8.4)
National level (aged ≤ 18 years)							
Men Hard and clay court	Filipcic et al. (2021) Galé-Ansodi et al. (2017b) Galé-Ansodi et al. (2018) Gallo-Salazar et al. (2019) Hoppe et al. (2014) Hoppe et al. (2016) Kilit and Arslan (2017) Kilit and Arslan (2018)	8	M: 1.1 (0.6, 2.0)	0.0473	(0.2, 7.9)	99.8	NA
Men Hard court	Filipcic et al. (2021) Galé-Ansodi et al. (2017b) Galé-Ansodi et al. (2018) Gallo-Salazar et al. (2019) Kilit and Arslan (2018)	5	M: 1.5 (0.6, 3.9)	NA	(0.16, 14.47)	99.8	NA
Men Clay court	Hoppe et al. (2014) Hoppe et al. (2016) Kilit and Arslan (2017) Kilit and Arslan (2018)	4	M: 0.7 (0.7, 0.7)	NA	(0.7, 0.7)	0	NA
Women Hard court	Galé-Ansodi et al. (2017a) Galé-Ansodi et al. (2017b) Galé-Ansodi et al. (2018)	3	M: 2.9 (2.7, 3.1)	NA	(2.44, 3.43)	30.8	NA
Peak running speed ($\text{m}\cdot\text{s}^{-1}$)							
International level							
International men, hard court	Fernández-Elias et al. (2020) Filipcic (2021) Kovalchik and Reid (2017) Whiteside et al. (2015)	4	M: 5.5 (4.0, 7.4)	NA	(3.0, 10.1)	81.8	M: 5.8 (4.9, 6.9)

Table 6 (continued)

Outcome (description)	Studies included in the meta-analysis (author(s), year)	Number of studies (<i>n</i>)	Effect size (95% CI)	Test for (subgroup) differences (<i>p</i> -value)	95% PI	<i>I</i> ² (%)	Sensitivity analysis for imputed SDs
National level (aged ≤ 18 years)							
Men Hard and clay court	Filipcic et al. (2021) Galé-Ansodi et al. (2017b) Galé-Ansodi et al. (2018) Gallo-Salazar et al. (2019) Hoppe et al. (2014) Hoppe et al. (2016)	6	M: 4.8 (4.3, 5.4)	0.11	(3.6, 6.5)	92.8	NA
Men Hard court	Filipcic et al. (2021) Galé-Ansodi et al. (2017b) Galé-Ansodi et al. (2018) Gallo-Salazar et al. (2019)	4	M: 5.1 (4.3, 6.0)	NA	(3.3 7.7)	94.5	NA

CI confidence interval, *I*² I-square statistic, *M* mean, *MD* mean difference, *NA* not applicable, *NE* not estimable, *PI* prediction interval, *SD* standard deviation

and female players across all three surfaces. These findings are comparable to the data of Lisi and Grigoletto [9], who reported a median (mean) match duration of 92 (98) minutes for best-of-3-set matches in international male players, based on 16,246 men's professional tennis matches. Unfortunately, no such large study was available for female players. The median (mean) match duration for best-of-5-set matches of international male players across all surfaces was 142 (150) minutes, based on 3712 matches. The mean match duration of junior male and female national players on hard court was around 75 min.

Lisi and Grigoletto showed that match duration was broad, with some matches lasting exceptionally long. The 95% quantile and 99.9% quantile were 154 and 200 min for best-of-3-set matches and 233 and 319 min for best-of-5-set matches. They used modelling to show how the format of the match can influence duration in best-of-5-set matches. For example, when using the no-advantage scoring system, the match duration dropped from a median of 144 to 132 min. When using one service only, the match duration dropped to 114 min, and when using both the no-advantage system and one service, the match duration dropped to 106 min.

Governing bodies in tennis have taken measures to reduce the exceptionally long match duration in men's tennis at the highest level by introducing new rules and changing the match format. In 2019, they reduced Davis Cup ties from best-of-5-set to best-of-3-set matches [89], and in 2022, the Grand Slam Tournaments jointly announced the 10-point final set tiebreak at six games-all in the fifth set [90]. The no-advantage scoring system was introduced in 2006 on the ATP and WTA tour

for doubles but is not used in singles. At the Next Gen ATP Finals, the 21-and-under-season finale, all singles matches are best-of-5-set matches to four (not six) games, with no-ad scoring [91]. In 2022, the ATP tried out new rules at this event to speed up the game even more: they shortened the time in between points to 15 s and allowed only one sit-down per set (after three games) [91].

Another important factor influencing the score, and hence the match duration, is the quality of the opposition. The more evenly matched the players are (on the day), the more points, games and sets they are likely to play, and the longer their matches will last.

4.2 Rally Duration and Rally Length (Strokes Per Rally)

Rally duration and rally length (strokes per rally) are strongly associated, and any trends in rally duration will be reflected in strokes per rally and vice versa. Important factors impacting rally duration and rally length include court surface [33, 97], playing style and tactics [94], and ball type and wear [34, 92].

4.2.1 Court Surface

Rally duration in international male and female players was longest on clay court (7.1 and 8.8 s), followed by hard court (5.6 and 6.4 s) and grass court (4.3 and 5.7 s), but only the differences in rally duration of male players on grass and clay courts reached statistical significance. The mean

Table 7 Strokes per rally of international and national male and female players on different court surfaces

Outcome (description)	Studies included in the meta-analysis (author(s), year)	Number of studies (<i>n</i>)	Effect size (95% CI)	Test for (sub-group) differences (<i>p</i> -value)	95% PI	<i>I</i> ² (%)	Sensitivity analysis for imputed SDs
Strokes per rally (#): international							
Men All surfaces	Carboch et al. (2019) Filipic et al. (2021) Hornery et al. (2007) Kovalchik and Reid (2017), Mendez-Villanueva (2007) Stare et al. (2015) Takahashi et al. (2006) Yusoff and Krasilshchikov (2021)	8	M: 4.1 (3.4, 5.0)	0.716	(2.5, 6.9)	82.7	M: 4.1 (3.2, 5.2)
Men Hard court	Carboch et al. (2019) Filipic et al. (2021) Hornery et al. (2007) Kovalchik and Reid (2017) Takahashi et al. (2006) Stare et al. (2015)	6	M: 4.2 (3.4, 5.3)	NA	(2.4, 7.5)	89.2	M: 4.3 (3.1, 5.9)
Men Clay court	Carboch et al. (2019) Hornery et al. (2007) Mendez-Villanueva (2007) Takahashi et al. (2006)	4	M: 4.8 (3.4, 6.7)	NA	(2.5, 9.2)	48.4	NA
Women Hard and clay courts	Carboch et al. (2018) Fernandez-Fernandez et al. (2007) Fernandez-Fernandez et al. (2008) Kovalchik and Reid (2017)	4	M: 3.9 (2.4, 6.2)	0.0415	(1.5, 9.8)	71.6	M: 3.3 (1.3, 8.2)
Strokes per rally (#): national (aged ≤ 18 years)							
Men Hard and clay court	Gallo-Salazar et al. (2019) Kilit and Arslan (2017) Kilit and Arslan (2018) Stare et al. (2015) Torres-Luque et al. (2011)	5	M: 4.9 (3.2, 7.50)	0.328	(1.7, 14.1)	99.0	M: 4.9 (2.7, 8.9)
Men Hard court	Gallo-Salazar et al. (2019) Kilit and Arslan (2018) Stare et al. (2015) Torres-Luque et al. (2011)	4	M: 4.6 (2.6, 8.2)	NA	(1.2, 18.0)	99.3	M: 4.4 (1.6, 12.1)

CI confidence interval, *I*² I-square statistic, *M* mean, *NA* not applicable, *PI* prediction interval, *SD* standard deviation, # number

number of strokes per rally was 4.1 [3.4, 5.0] for international male players on all surfaces and 3.9 [2.4, 6.4] for female players on hard courts and clay, with no data for female players on grass. There appeared to be a trend of more strokes per rally on clay court and fewer strokes on hard court in male players, with insufficient data for grass court. However, no statistically significant differences were

found, which may be partly related to the relatively small number of studies in this area.

Players may adjust their playing style according to the court surface, impacting rally duration. Previously, only notational methods were available to study the association between playing style and rally duration on different court surfaces. However, these methods are limited

Table 8 First- and second-serve speed ($\text{km}\cdot\text{h}^{-1}$) of international male and female players on different court surfaces

Outcome (description)	Studies included in the meta-analysis (author(s), year)	Number of studies (<i>n</i>)	Effect size (95% CI)	Test for (sub-group) differences (<i>p</i> -value)	95% PI	<i>I</i> ² (%)	Sensitivity analysis for imputed SDs
First-serve speed ($\text{km}\cdot\text{h}^{-1}$): international level							
Men All surfaces	Cui et al. (2020a) Hornery et al. (2007) Kovalchik and Reid (2017) Reid et al. (2016) Fitzpatrick et al. (2019) Brown et al. (2021)	6	M: 182.4 (178.1, 186.9)	0.099	(171.2, 194.4)	95.3	M: 183.0 (177.9, 188.3)
Men Hard court	Cui et al. (2020a) Hornery et al. (2007) Kovalchik and Reid (2017) Reid et al. (2016)	4	M: 181.9 (176.6, 187.4)	NA	(169.2, 195.6)	84.9	M: 182.7, (174.9, 191.0)
Men Clay court	Cui et al. (2020a) Fitzpatrick et al. (2019) Hornery et al. (2007)	3	M: 180.0 (168.9, 191.9)	NA	(153.8, 210.7)	84.7	NA
Women All surfaces	Kovalchik and Reid (2017) Reid et al. (2016) Fitzpatrick et al. (2019) Brown et al. (2021)	4	M: 156.1 (151.4, 161.0)	NE	(143.7, 169.6)	94.6	M: 156.9 (150.0, 164.1)
Second-serve speed ($\text{km}\cdot\text{h}^{-1}$): international level							
Men All surfaces	Hornery et al. (2007) Reid et al. (2016) Cui et al. (2020a) Brown et al. (2021)	4	M: 148.9 (135.0, 164.1)	0.268	(121.8, 182.0)	97.4	NA
Men Hard court	Cui et al. (2020a) Hornery et al. (2007) Reid et al. (2016)	3	M: 146.4 (125.3, 171.0)	NA	(99.92, 214.44)	95.8	NA
Women Hard and Grass court	Reid et al. (2016) Brown et al. (2021)	2	M: 133.7 (106.6, 167.7)	NA	NE	95.4	NA

CI confidence interval, *I*² I-square statistic, *M* mean, *NA* not applicable, *NE* not estimable, *PI* prediction interval, *SD* standard deviation

by the amount of data that can be collected and the need to define observable variables in advance [97]. Today's automated ball-tracking systems allow high-volume data collection for post-recording analysis and have provided new insights.

4.2.2 Game Characteristics

Carboch compared the game characteristics of international male and female players at the four Grand Slams [30]. He showed that the number of first serves in, points per game and games per set were reasonably similar in male and

female players across all surfaces. However, male players won more points after first and second serves (most at Wimbledon), and female players won more return games (Roland Garros, Australian Open). Men finished more points with a winner (Wimbledon, Roland Garros), and women with an unforced error (Roland Garros, Australian Open). The author concluded that the main difference between the sexes was the higher efficiency of the first and second serve in men (highest at Wimbledon) and the advantage of the second serve for the receiver in women's matches (all surfaces).

Fitzpatrick et al. [44] compared game strategy during short (0–4 shots), medium (5–8) and long (> 9) rallies during



Fig. 5 Summary of the physical demands of tennis in **a** international and **b** national players. *min* minutes, *s* seconds

men's and women's matches at Wimbledon and Roland Garros in 2016 and 2017. They showed that for male and female players, points won of 0–4 shot rally length, baseline points won, and first- and second-serve points won were most strongly associated with success. Forced errors and unforced errors were most strongly associated with losing. The authors concluded that short rallies are prevalent at Wimbledon and Roland Garros and that playing style did not drastically change for either sex when transitioning from clay to grass. They suggested that getting used to the ball-court surface interaction might be more important than focusing on a specific area of their game (such as approach shots or net play) when moving from clay to grass.

In a second study, Fitzpatrick et al. [98] analysed the short points played at Wimbledon's men's and women's matches in 2015–17. Short points were most prevalent compared to medium and long points and accounted for 72% (men) and 66% (women) of all points played. The most prevalent rally length was one shot [ace, or return error] (46.3 and 36.3%), followed by three shots (both 22.2%), two shots (return winner or error of server's second stroke, 15.7 and 20.2%), four shots (11.2 and 14.5%) and zero shots (double fault, 4.7 and 6.7%). Both points won of one shot, and two shots were associated with winning matches, and the authors concluded that the serve and serve return should be prioritised during grass court training. As short rallies are also prevalent at

Roland Garros [44], focusing on the serve and serve return seems appropriate for all court surfaces.

4.2.3 Ball Type

The ITF aimed to slow down the power and speed of serves on hard courts and speed up the games on slow courts [99]. In 2002, the ITF introduced three types of balls according to the pace of the surface. The ITF recommended using a slightly larger ball (type 3) on fast courts to reduce speed. They recommended using a somewhat harder ball (type 1) on slow courts, producing a lower bounce angle and speeding up the game. Brown and O'Donoghue [100] studied the rally duration on hard, clay and grass courts before and after introducing the different types of balls and reported that the differences in rally duration remained but were reduced in magnitude following the introduction of different balls in 2002.

4.2.4 Sex

When comparing rally duration in male and female players in the same study, we found no statistically significant difference between male and female players on hard court and clay, or across all surfaces, with insufficient studies for grass court. It is important to compare the sexes in the same study to control for confounding factors, and more studies of this type are needed.

4.2.5 Junior National Players

The mean rally duration of junior national male players was 8.3 s on hard court and 12 s on clay, slightly longer than the rally duration of international male players on the same surfaces. Hizan et al. [101] compared game characteristics of junior national and adult international players (professionals). Both juniors and professionals hit approximately 60% of their first serves in. The professionals hit more aces, fewer double faults and won a higher percentage of points when the first serves were in (73% in professional male players, 65% in professional female players, 55–60% in juniors). The juniors and professionals were comparable in the number of points won after the second serve (35–40%). The professionals won a higher percentage of points when returning second serves. Unfortunately, they did not report rally duration, and further studies on game characteristics and rally duration in junior and adult national players are warranted.

4.2.6 Time Trend

Ball types, friction coefficients of court surfaces, fitness levels and playing styles have changed. It would be therefore be interesting to determine whether there has been a time trend

in the rally duration of male and female players over the last 20 years on the different court surfaces. Unfortunately, the number of studies was too low to perform a meaningful random-effects meta-regression.

4.3 Distance Covered

The mean distance covered was 2300 m per match for international-level male players (best-of-5 sets) and 1250 m for international-level female players (best-of-3 sets). In contrast, the mean distances covered by junior national-level players were 3300 m for male players and 3000 m for female players. We recommend caution in comparing the results between national and international players because of the use of different measurement systems. The Hawk-Eye system was mainly used for international players and the GPS for junior national players. Several studies using Hawk-Eye analysis only measured the distance covered during the rally, whereas studies using GPS measured the distance covered during and between points. This may explain part of the difference in distance covered between national and international players.

Two other studies in international male players (best-of-3 sets) confirm previous findings. Filipcic et al. [43] measured the distance covered using the Sagit/tennis tracking system and reported a mean distance of 1776 ± 281 m per match in four matches. They only measured the distance covered during play. Pereira et al. [72] used the GPS and reported a much greater mean distance of 3160 ± 880 m during four matches. However, the court surface may also have influenced the outcome because Filipcic et al. studied matches on hard court and Pereira et al. matches on clay.

In contrast, GPS tends to under-report distance and speed at high-intensity efforts [102]. Therefore, distances measured using GPS are likely lower than those estimated using Hawk-Eye (if Hawk-Eye included the distance covered in between rallies). The distance covered during play (including in-between rallies and during change of ends) might be even higher than currently reported.

Reid et al. [76] reported that the mean distance covered per match by 102 male players during the 2012–14 Australian Open was 2110 ± 839 m. However, when he compared 40 top-ranked to 40 lower ranked players at these events, the match distances were 3082 ± 1075 and 2498 ± 898 m [87]. This disparity in distance covered from the same event and fundamentally the same era of players appears unusual. It seems plausible that methodological differences or the evolving measurement techniques of the tracking providers can explain this difference. Another explanation would be that higher ranked players move more continuously and anticipate and make better decisions than their lower ranked peers [103, 104]. Martínez-Gallego et al. [105] seem

to confirm this, showing that game winners covered more distance during the rally, and at a lower speed, than game losers.

4.4 Work-to-Rest Ratio

The work-to-rest ratio is determined by rally duration and the rest periods between points. Therefore, the rules on time allotted between points greatly influence the work-to-rest ratio, more so than the effective playing time. Up until 2019, the ITF [106] and WTA [107] allowed only 20 s between points, whereas the ATP [108] allowed 25 s between points. The 20-s time rule of the ITF resulted in time rule violations for 58.8% of the serves at the Australian Open men's singles event, with only 0.1% of the violations being penalised [109]. In 2018, a time clock was introduced at the US Open to speed up the game and to help the chair umpires enforce the rules regarding the time between points. Rule violations decreased (26.3%), but the average time between points did not (21.6 s) [110] compared to previous studies. In 2019, the ITF rules [111] and WTA rules [112] were aligned with the ATP, allowing 25 s between points. The differences in regulations between the organisation and rule changes over time make it difficult to compare the work-to-rest ratios of the different player groups.

The work-to-rest ratios in all studies ranged from 1:2 to 1:4.5. The ratios tended to be higher on hard and grass courts than on clay courts due to the shorter rally duration on hard and grass courts compared to clay courts. They were also higher during international-level than the junior national-level competition, again likely due to the shorter rally duration (and possibly longer rest periods) of international compared with national players.

4.5 Effective Playing Time

In addition to rally duration and rest periods between points, the effective playing time also takes into account medical time-outs (3 min for treatment after evaluation), the heat rule (10 min between the second and third set in ITF and WTA tournaments), and the resting time between change of ends (90 s) and sets (120 s). In our review, the mean effective playing time in international male and female tennis players was around 19% and slightly higher in national-level players at about 26%. This difference can partly be explained by longer rallies and shorter rest periods of national-level players, but it may also be due to fewer opportunities for medical time-outs and other breaks. For example, during televised matches of WTA players, the chair umpire may extend the change of ends and set breaks when necessary [113].

4.6 Running Speed

The peak running speed was $5 \text{ m}\cdot\text{s}^{-1}$ (18 km/h) in international male tennis players and slightly lower in female players ($4.2 \text{ m}\cdot\text{s}^{-1}$, 15.1 km/h). Players spend only a short period and cover a relatively short distance running at high speeds. International male players covered 8.7% of the distance at speeds between 4 and $5 \text{ m}\cdot\text{s}^{-1}$ and only 4.1% at speeds $> 5 \text{ m}\cdot\text{s}^{-1}$ [86]. For female players, these numbers were 5.8% and 1.8%.

In tennis, being the better player does not necessarily require the highest running speed. Whiteside et al. [87] showed that despite higher ranked players covering more distance ($3082 \pm 1075 \text{ m}$) than lower ranked players ($2498 \pm 898 \text{ m}$), their mean peak speed was lower (6.1 ± 2.5 vs $6.5 \pm 3.7 \text{ m}\cdot\text{s}^{-1}$) and their mean average speed was higher (1.3 ± 0.2 vs $1.2 \pm 0.2 \text{ m}\cdot\text{s}^{-1}$). This suggests that the higher ranked players move more continuously, whereas the lower ranked players have to accelerate more [104].

4.7 Accelerations, Decelerations and Changes of Direction

The number of accelerations and decelerations in tennis may be a better way to describe load in tennis than the mean and peak running speed because of the increased load on the legs during speed and direction changes. Male and female junior tennis players covered most of the distance during a match on hard court while accelerating or decelerating (89.7%). [47] Unfortunately, it was not possible to synthesise the data of the various studies because the cut-off points for the accelerations and decelerations varied widely (e.g. $1\text{--}2 \text{ m}\cdot\text{s}^{-2}$, $> 1.5 \text{ m}\cdot\text{s}^{-2}$, $> 2 \text{ m}\cdot\text{s}^{-2}$, $2\text{--}4 \text{ m}\cdot\text{s}^{-2}$, $> 3 \text{ m}\cdot\text{s}^{-2}$ and $> 4 \text{ m}\cdot\text{s}^{-2}$). We recommend using standardised cut-off points for accelerations and decelerations in tennis for future studies to enable comparisons.

Most studies used GPS to measure accelerations, decelerations and velocities, except one that used Hawk-Eye [86]. Different measurement techniques may explain some of the differences in the outcomes. Unfortunately, Hawk-Eye data are not publicly available, and in the study by Whiteside et al. [87], it was unclear how they computed the number of high-speed runs ($> 3 \text{ m/s}^2$) per game, making it difficult to repeat these measurements. There is low accuracy and reliability of GPS devices at high-intensity efforts, which may lead to under-reporting of distance and speed and possibly, accelerations and decelerations [102]. In particular, caution is warranted when comparing acceleration indices between different GPS brands [114].

The number of changes of direction ranged from 2 to 6 per rally, and Giles et al. [52] reported more than 400 (female player) and 600 (male player) changes of direction per match during the Australian Open (2016–18). These

direction changes, combined with the distances covered and the number of accelerations and decelerations, result in a high load on the lower extremities, for which the players must prepare.

4.8 Serve Speed

The serve speed of international male players was higher than that of female players, and the mean first-serve speed was higher than the mean second-serve speed in both male and female players. We did not find data on serve speed in national male or female players.

The serve is an important stroke in tennis, and players win more points when serving than receiving across all surfaces [29]. The success rate is higher in male than female players, and higher after the first than after the second serve. [115] The overall success rate of the serve during singles matches at Wimbledon from 2004 to 2019 was 75% and 57% for male players' first and second serve and 66% and 53% for female players [29]. Serve speed, and the chance of winning the point are correlated, both in male and female players and after the first and second serve. Other factors determining a server's success include spin and placement, but our study did not include those outcome parameters.

4.9 Training Recommendations

Training specificity is fundamental for securing optimal adaptation to ensure improved performance and enhanced recovery for the next match [116]. Therefore, understanding the competition demands of tennis is of paramount importance for coaches, strength and conditioning coaches, and tennis players to ensure they plan the appropriate training dose to maximise the fitness-fatigue response within athletes. Goossens et al. [118] studied data from the four Grand Slam Tournaments from 1992 to 2011, covering 20,320 matches. They found that the relative effort invested in winning a match negatively affected the probability of winning the next match. For men, this was the case if they had played two sets more in their previous match than their opponent had, and for women, this was one set.

Tennis players must move well to hit powerful shots with great accuracy while moving sideways, forwards or backward at variable speeds. Players need not only to be fast around the court but also to read the game well and move efficiently around the court [103]. Footwork, anticipation and tactical drills are essential components of the training programme for a developing or elite player.

Elite players (e.g. international players) have better physical fitness values and stroke performance levels than sub-elite players (e.g. regional, national and junior players) [5]. An essential part of the preparation would be to mimic the demands during competition, both in duration, frequency

and intensity, during training. Therefore, it is not enough for a tennis player to be able to run 3–5 km at an easy pace. As identified by this review, their running training should include accelerations, decelerations, changes of direction and high running speeds. Exercises to improve acceleration should be included in the training programme. The programme may consist of exercises to improve strength, power, agility for direction changes and speed.

The tennis training and strength and conditioning should mimic the work-to-rest ratios (1:2–1:4) experienced in tennis matches. Many tennis drills consist of high volume and continuous loading to acquire skills and develop consistency and economy. However, high-intensity interval exercises with adequate rest periods should be included as well, both on and off court. The serve and serve returns are essential strokes in tennis, and youth coaches should prioritise the development of an efficient serve technique and strong serve returns [101].

4.9.1 Strengths and Limitations

We provide a comprehensive overview of the literature on the physical demands of tennis. A limitation of this review is that some of the studies were conducted over a decade ago, and over time, the rules, match formats, playing surfaces and playing equipment in tennis have changed. We had to rely on the accuracy of collected data, and older data, in particular, may be limited by the quality of the GPS and video tracking. There was also a lack of standardisation in reporting, mainly related to movement variables, making it difficult to compare studies. Most studies focused on international players, with fewer on national or regional players. Most international players were adults, whereas all national-level players were juniors, which made an analysis stratified by age impossible. The relatively low number of studies per outcome variable sometimes precluded a meta-analysis or yielded wide CIs and PIs.

It would be interesting to determine whether some outcomes have a time trend. However, we could not study the influence of the covariate/variable "publication year" on different outcome variables (e.g. rally duration). The number of studies is too low (< 10–20) to perform a meaningful random-effects meta-regression.

5 Conclusions

The findings from this systematic review with meta-analysis provide a comprehensive summary of the physical demands of tennis. The results indicate that elite tennis players should have high fitness levels to handle matches of up to 3 h or more, followed by a speedy recovery. Players should focus

on footwork and movement drills and include many accelerations, decelerations and changes of directions, with appropriate work-to-rest ratios, rather than straight-line continuous running. They should prioritise the serve and serve return. The present study will assist tennis coaches, strength and conditioning trainers and tennis players in better understanding the demands of the game.

5.1 Recommendations for Future Studies

Because of inconsistent reporting methods, varying cut-off points and missing standard deviations, we could not include all data in the meta-analysis. We recommend more consistent measuring and reporting of data to enable future meta-analysts to pool meaningful data. Similarly, there is an opportunity for studies to provide access to the individual data sets to allow performing meta-analyses with age as a covariate. Given the wide variability in the data on the distance covered and the lack of sufficient data on peak and average running speed on different surfaces, more studies describing these generalised movement demands are recommended. There is also a need for improved standardisation, validity, and reliability of tracking data in tennis, especially at the professional level. This should include agreement on the treatment or filtering of raw data and expert consensus on measures even as simple as playing time. This work should then extend into the harmonisation or translation of player tracking technologies, where the performances of different systems are compared.

As most data are from professional tennis players, primarily male, more research on athletes at national and regional levels and female players is needed. Unfortunately, the tracking systems used in professional events are not publicly available and remain quite expensive, which precludes their wider use. More affordable computer vision technologies powered by artificial intelligence, and various inertial measurement units look promising in bridging this gap.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s40279-022-01807-8>.

Acknowledgements The author group would like to thank Matthew Lester for his contribution to the screening.

Declarations

Funding Tobias Saueressig received funding from the Royal Netherlands Lawn Tennis Association for the statistical analysis. The other authors received no financial support for the research, authorship and publication of this article.

Conflicts of interest BMP, MGTJ and MvR are employed by the Royal Netherlands Lawn Tennis Association. BMP is Medical and Research Consultant for the ITF and Medical Consultant for the ATP Tour. CB is employed by the Lawn Tennis Association. SW, SC, KF, NH, DCJvR,

VMP, AM, SRO, FCLO, MR, TS, LJS, JST, NW and CLA have no conflicts of interest that are directly relevant to the content of this article.

Ethics approval Not applicable.

Consent to participate Not applicable.

Consent for publication Not applicable.

Availability of data and material All data generated or analysed during this study are included in this published article and its supplementary information files.

Code availability Not applicable.

Authors' contributions BMP had the idea for the study. LJS conducted the literature search. TS performed the data analysis. BMP, MGTJ, SW, DCJvR, JST and CLA prepared the original draft. All authors critically commented on previous versions of the manuscript and read and approved the final manuscript.

References

1. International Tennis Federation (ITF). ITF global tennis report 2021. London: ITF; 2021. <http://itf.uberflip.com/i/1401406-itf-global-tennis-report-2021/9?> Accessed 21 Jun 2022.
2. Nag U. Everything you need to know about tennis courts: IOC; 2022. <https://olympics.com/en/featured-news/tennis-court-markings-dimensions-size-types-variety-surface-hard-grass-clay>. Accessed 13 Jan 2023.
3. Kovacs MS. Tennis physiology: training the competitive athlete. *Sports Med.* 2007;37(3):189–98.
4. Devine JW. Elements of excellence. *J Philos Sport.* 2022;1–17.
5. Lambrich J, Muehlbauer T. Physical fitness and stroke performance in healthy tennis players with different competition levels: a systematic review and meta-analysis. *PLoS ONE.* 2022;17(6):e0269516.
6. Xiao W, Geok SK, Bai X, Bu T, Norjali Wazir MR, Talib O, et al. Effect of exercise training on physical fitness among young tennis players: a systematic review. *Front Public Health.* 2022;10:843021.
7. Reid M, Duffield R. The development of fatigue during match-play tennis. *Br J Sports Med.* 2014;48 Suppl. 1(Suppl. 1):i7–11.
8. Barnett T, Alan B, Pollard G. Reducing the likelihood of long tennis matches. *J Sports Sci Med.* 2006;5(4):567–74.
9. Lisi F, Grigoletto M. Modeling and simulating durations of men's professional tennis matches by resampling match features. *J Sports Analyt.* 2021;7:57–75.
10. Haake SJ, Chadwick SG, Dignall RJ, Goodwill S, Rose P. Engineering tennis: slowing the game down. *Sports Eng.* 2000;3(2):131–43.
11. Baodong Y. Hawkeye technology using tennis match. *Comput Model New Technol.* 2014;18(12c):400–2.
12. Page MJ, Moher D, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. PRISMA 2020 explanation and elaboration: updated guidance and exemplars for reporting systematic reviews. *BMJ.* 2021;372.
13. Otten R, de Vries R, Schoonmade L. Amsterdam Efficient Deduplication (AED) method (Version 1). Amsterdam: Zenodo; 2019.

















14. Bramer WM, Giustini D, de Jonge GB, Holland L, Bekhuis T. De-duplication of database search results for systematic reviews in EndNote. *J Med Libr Assoc.* 2016;104(3):240–3.
15. Ouzzani M, Hammady H, Fedorowicz Z, Elmagarmid A. Rayyan: a web and mobile app for systematic reviews. *Syst Rev.* 2016;5(1):1–10.
16. Hawk-Eye Innovations. Pioneering & inspiring change in sport 2022. <https://www.hawkeyeinnovations.com/>. Accessed 13 Jan 2023.
17. JBI. Critical appraisal tools Adelaide: JBI Collaboration; 2022. <https://jbi.global/critical-appraisal-tools>. Accessed 13 Jan 2023.
18. Tanner-Smith EE, Tipton E, Polanin JR. Handling complex meta-analytic data structures using robust variance estimates: a tutorial in R. *J Dev Life-Course Criminol.* 2016;2(1):85–112.
19. Fisher Z, Tipton E. robmeta: an R-package for robust variance estimation in meta-analysis. arXiv:150302220. 2015.
20. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials.* 1986;7(3):177–88.
21. Harrer M, Cuijpers P, Furukawa TA, Ebert DD. Doing meta-analysis with R: a hands-on guide. New York: Chapman and Hall/CRC; 2021.
22. Friedrich JO, Adhikari NK, Beyene J. The ratio of means method as an alternative to mean differences for analyzing continuous outcome variables in meta-analysis: a simulation study. *BMC Med Res Methodol.* 2008;8(1):1–15.
23. IntHout J, Ioannidis JP, Rovers MM, Goeman JJ. Plea for routinely presenting prediction intervals in meta-analysis. *BMJ Open.* 2016;6(7): e010247.
24. R Core Team. R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing; 2020. <https://www.r-project.org/>. Accessed 13 Jan 2023.
25. Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw.* 2010;36(3):1–48.
26. Higgins JP, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. *Cochrane handbook for systematic reviews of interventions.* Chichester: Wiley; 2019.
27. Bergeron MF, McLeod KS, Coyle JF. Core body temperature during competition in the heat: National Boys' 14s Junior Championships. *Br J Sports Med.* 2007;41(11):779–83.
28. Brown E, O'Donoghue P. Gender and surface effect on elite tennis strategy. *ITF Coach Sport Science Rev.* 2008;15(46):9–11.
29. Brown EG. A faster serve has more impact on success for female elite tennis players than males. *Int J Perform Anal Sport.* 2021;21(4):600–10.
30. Carboch J. Comparison of game characteristics of male and female tennis players at grand-slam tournaments in 2016. *Trends Sport Sci.* 2017;4(24):151–5.
31. Carboch J, Plachá K. Development of rally pace and other match characteristics in women's matches in the Australian Open 2017. *J Phys Educ Sport.* 2018;18(2):1079–83.
32. Carboch J, Placha K, Sklenarik M. Rally pace and match characteristics of male and female tennis matches at the Australian Open 2017. *J Hum Sport Exerc.* 2018;13(4):743–51.
33. Carboch J, Siman J, Sklenarik M, Blau M. Match characteristics and rally pace of male tennis matches in three Grand Slam tournaments. *Phys Activity Rev.* 2019;7:49–56.
34. Carboch J, Blau M, Sklenarik M, Siman J, Placha K. Ball change in tennis: How does it affect match characteristics and rally pace in Grand Slam tournaments? *J Hum Sports Exerc.* 2020;15(1):153–62.
35. Cui Y, Gómez M-Á, Gonçalves B, Liu H, Sampaio J. Effects of experience and relative quality in tennis match performance during four Grand Slams. *Int J Perform Anal Sport.* 2017;17(5):783–801.
36. Cui Y, Gómez M, Gonçalves B, Sampaio J. Performance profiles of professional female tennis players in grand slams. *PLoS ONE.* 2018;13(7): e0200591.
37. Cui Y, Zhao Y, Liu H, Gómez M, Wei R, Liu Y. Effect of a seeding system on competitive performance of elite players during major tennis tournaments. *Front Psychol.* 2020;11:1294.
38. Cui Y, Liu H, Gómez M, Liu H, Gonçalves B. Set-to-set performance variation in tennis Grand Slams: play with consistency and risks. *J Hum Kinet.* 2020;73:153–63.
39. Fernández-Elías V, Courel-Ibáñez J, Pérez-López A, Jodra P, Moreno-Pérez V, Coso JD, et al. Acute beetroot juice supplementation does not improve match-play activity in professional tennis players. *J Am Nutr Assoc.* 2022;41(1):30–7.
40. Fernandez-Fernandez J, Mendez-Villanueva A, Fernandez-Garcia B, Terrados N. Match activity and physiological responses during a junior female singles tennis tournament. *Br J Sports Med.* 2007;41(11):711–6.
41. Fernandez-Fernandez J, Sanz-Rivas D, Fernandez-Garcia B, Mendez-Villanueva A. Match activity and physiological load during a clay-court tennis tournament in elite female players. *J Sports Sci.* 2008;26(14):1589–95.
42. Filipcic A, Zecic M, Reid M, Crespo M, Panjan A, Nejc S. Differences in performance indicators of elite tennis players in the period 1991–2010. *J Phys Educ Sport.* 2015;15(4):671–7.
43. Filipcic A, Leskosek B, Crespo M, Filipcic T. Matchplay characteristics and performance indicators of male junior and entry professional tennis players. *Int J Sports Sci Coach.* 2021;16(3):768–76.
44. Fitzpatrick A, Stone JA, Choppin S, Kelley J. Important performance characteristics in elite clay and grass court tennis match-play. *Int J Perform Anal Sport.* 2019;19(6):942–52.
45. Galé AC. Youths tennis players' velocity and acceleration in match play. *Revista Internacional de Deportes Colectivos.* 2014;18:50–65.
46. Galé-Ansodi C, Castellano J, Usabiaga O. Effects of different surfaces in time-motion characteristics in youth elite tennis players. *Int J Perform Anal Sport.* 2016;16(3):860–70.
47. Galé-Ansodi C, Castellano J, Usabiaga O. More acceleration and less speed to assess physical demands in female young tennis players. *Int J Perform Anal Sport.* 2017;17(6):872–84.
48. Galé-Ansodi C, Castellano J, Usabiaga O. Physical profile of young tennis players in the tennis match-play using global positioning systems. *J Phys Educ Sport.* 2017;17(2):826–32.
49. Galé-Ansodi C, Castellano J, Usabiaga O. Differences between running activity in tennis training and match-play. *Int J Perform Anal Sport.* 2018;18(5):855–67.
50. Gallo-Salazar C, Areces F, Abián-Vicén J, Lara B, Salinero JJ, Gonzalez-Millán C, et al. Enhancing physical performance in elite junior tennis players with a caffeinated energy drink. *Int J Sports Physiol Perform.* 2015;10(3):305–10.
51. Gallo-Salazar C, Del Coso J, Sanz-Rivas D, Fernandez-Fernandez J. Game activity and physiological responses of young tennis players in a competition with two consecutive matches in a day. *Int J Sports Physiol Perform.* 2019;14(7):887–93.
52. Giles B, Kovalchik S, Reid M. A machine learning approach for automatic detection and classification of changes of direction from player tracking data in professional tennis. *J Sports Sci.* 2020;38(1):106–13.
53. Hoppe MW, Baumgart C, Bornefeld J, Sperlich B, Freiwald J, Holmberg HC. Running activity profile of adolescent tennis players during match play. *Pediatr Exerc Sci.* 2014;26(3):281–90.
54. Hoppe MW, Baumgart C, Freiwald J. Do running activities of adolescent and adult tennis players differ during play? *Int J Sports Physiol Perform.* 2016;11(6):793–801.

55. Hornery DJ, Farrow D, Mujika I, Young W. An integrated physiological and performance profile of professional tennis. *Br J Sports Med.* 2007;41(8):531–6 (**discussion 6**).
56. Johnson CD, McHugh MP. Performance demands of professional male tennis players. *Br J Sports Med.* 2006;40(8):696–9.
57. Kilit B, Arslan E. Physiological responses and time-motion characteristics of young tennis players: comparison of serve vs. return games and winners vs. losers matches. *Int J Perform Anal Sport.* 2017;17(5):684–94.
58. Kilit B, Arslan E. Playing tennis matches on clay court surfaces are associated with more perceived enjoyment response but less perceived exertion compared to hard courts. *Acta Gymnica.* 2018;48(4):147–52.
59. SSRN. On the independence and identical distribution of points in tennis. 1998. <https://ssrn.com/abstract=138096>. Accessed 13 Apr 2022.
60. Kovalchik SA, Reid M. Comparing matchplay characteristics and physical demands of junior and professional tennis athletes in the era of big data. *J Sports Sci Med.* 2017;16(4):489–97.
61. Mackie D. The effects of gender on the work to rest ratio of elite level tennis players. Cardiff: Cardiff School of Sport; 2013.
62. Maquirriain J, Baglione R, Cardey M. Male professional tennis players maintain constant serve speed and accuracy over long matches on grass courts. *Eur J Sport Sci.* 2016;16(7):845–9.
63. McCarthy PR, Thorpe RDW, C. Body fluid loss during competitive tennis match-play. In: Lees A, Maynard J, Hughes MR, Reilly T, editors. *Science and racket sports II*. London: E & FN Spon; 1998. pp. 52–5.
64. Meffert D, O'Shannessy C, Born P, Grambow R, Vogt T. Tennis at tiebreaks: addressing elite players' performance for tomorrow's coaching. *German J Exerc Sport Res.* 2019;49(3):339–44.
65. Mendez-Villanueva A, Fernandez-Fernandez J, Bishop D, Fernandez-Garcia B, Terrados N. Activity patterns, blood lactate concentrations and ratings of perceived exertion during a professional singles tennis tournament. *Br J Sports Med.* 2007;41(5):296–300 (**discussion**).
66. Morante SB, John. Match characteristics of professional singles tennis. *J Med Sci Tennis.* 2005;10(3):12–3.
67. Moreno-Pérez V, López-Samanes Á, Domínguez R, Fernández-Elías VE, González-Frutos P, Fernández-Ruiz V, et al. Acute effects of a single tennis match on passive shoulder rotation range of motion, isometric strength and serve speed in professional tennis players. *PLoS ONE.* 2019;14(4): e0215015.
68. Murphy AP, Duffield R, Kellett A, Reid M. A comparison of the perceptual and technical demands of tennis training, simulated match play, and competitive tournaments. *Int J Sports Physiol Perform.* 2016;11(1):40–7.
69. Myers NL, Sciascia AD, Kibler WB, Uhl TL. Volume-based interval training program for elite tennis players. *Sports Health.* 2016;8(6):536–40.
70. O'Donoghue P, Ingram B. A notational analysis of elite tennis strategy. *J Sports Sci.* 2001;19(2):107–15.
71. O'Donoghue P, Liddle D. A notational analysis of time factors of elite men's and ladies' singles tennis on clay and grass surfaces. In: Lees A, Maynard J, Hughes M, Reilly T, editors. *Science and racket sports II*. London: E & FN Spon; 1998. p. 241–6.
72. Pereira TJC, Nakamura FY, de Jesus MT, Vieira CL, Misuta MS, de Barros RM, et al. Analysis of the distances covered and technical actions performed by professional tennis players during official matches. *J Sports Sci.* 2017;35(4):361–8.
73. Pereira LA, Freitas V, Arruda Moura F, Saldanha Aoki M, Loturco I, Yuzo NF. The activity profile of young tennis athletes playing on clay and hard courts: preliminary data. *J Hum Kinet.* 2016;50:211–8.
74. Perri T, Norton KI, Bellenger CR, Murphy AP. Training loads in typical junior-elite tennis training and competition: implications for transition periods in a high-performance pathway. *Int J Perform Anal Sport.* 2018;18(2):327–38.
75. Ponzano M, Gollin M. Movement analysis and metabolic profile of tennis match play: comparison between hard courts and clay courts. *Int J Perform Anal Sport.* 2017;17(3):220–31.
76. Reid M, Morgan S, Whiteside D. Matchplay characteristics of Grand Slam tennis: implications for training and conditioning. *J Sports Sci.* 2016;34(19):1791–8.
77. Reilly TP, J. Investigation of exercise intensity in male singles lawn tennis. In: Reilly T, Hughes ML, Lees A, editors. *Science and racket sports*. London: E & FN Spon; 1994. pp. 10–3.
78. Sánchez-Pay A, Ortega-Soto JA, Sánchez-Alcaraz BJ. Notational analysis in female Grand Slam tennis competitions. *Kinesiology.* 2021;53(1):154–61.
79. Smith MT, Reid M, Kovalchik S, Wood T, Duffield R. Heat stress incidence and matchplay characteristics in women's Grand Slam tennis. *J Sci Med Sport.* 2018;21(7):666–70.
80. Smith MT, Reid M, Kovalchik S, Woods TO, Duffield R. Heat stress incident prevalence and tennis matchplay performance at the Australian Open. *J Sci Med Sport.* 2018;21(5):467–72.
81. Stare M, Žibrat U, Filipic A. Stroke effectiveness in professional and junior tennis. *Kinesiology Slovenica.* 2015;21(2):39–50.
82. Takahashi H, Wada T, Maeda A, Kodama M, Nishizono H, Kurata H. The relationship between court surface and tactics in tennis using a computerized scorebook. *Int J Perform Anal Sport.* 2006;6(2):15–25.
83. Takahashi H, Wada T, Maeda A, Kodama M, Nishizono H. An analysis of time factors in elite male tennis players using the computerised scorebook for tennis. *Int J Perform Anal Sport.* 2009;9(3):314–9.
84. Torres-Luque G, Cabello-Manrique D, Hernández-García R, Garatachea N. An analysis of competition in young tennis players. *Eur J Sport Sci.* 2011;11(1):39–43.
85. Yusoff SSB, Krasilshchikov O. Match and game performance variables in elite and junior men singles tennis players. *Sport Mont.* 2021;19:189–93.
86. Whiteside D, Reid M. External match workloads during the first week of Australian Open tennis competition. *Int J Sports Physiol Perform.* 2017;12(6):756–63.
87. Whiteside D, Bane MK, Reid M, editors. Differentiating top-ranked male tennis players from lower-ranked players using Hawk-Eye data: an investigation of the 2018 2014 Australian Open tournaments. In: 33rd international society of biomechanics in sports conference; 29 June–3 July 2015; Poitiers, France.
88. O'Donoghue P, Liddle D. A notational analysis of time factors of elite men's and ladies' singles tennis on clay and grass surfaces. In: Lees A, Maynard J, Reilly T, editors. *Science and racket sports II*. London: E & FN Spon; 1998. pp. 241–6.
89. ITF. Historic Davis Cup reforms approved at AGM London: ITF; 2018. <https://www.daviscup.com/en/news/290057.aspx>. Accessed 13 Jan 2023.
90. Hrdlicka J, Moreton G, Hewitt I, McNulty M. Grand Slam Tournaments jointly announce 10-point final set tie-break at six games all: AELTC; 2022. https://www.wimbledon.com/en_GB/news/articles/2022-03-16/grand_slam_tournaments_jointly_announce_10point_final_set_tiebreak_at_six_games_all.html. Accessed 13 Jan 2023.
91. ATP. Rules and innovations Florida: ATP 2022. <https://www.nextgenatpfinals.com/en/event/rules-and-innovations>. Accessed 13 Jan 2023.
92. Miller S. Modern tennis rackets, balls, and surfaces. *Br J Sports Med.* 2006;40(5):401–5.

93. Doğan İ, Revan S, Arıkan Ş. Analysis of tennis competitions on different court surfaces. *Turk J Sport Exerc.* 2021;23(1):60–6.
94. Kovalchik S, Reid M. A shot taxonomy in the era of tracking data in professional tennis. *J Sports Sci.* 2018;36(18):2096–104.
95. Périard JD, Racinais S, Knez WL, Herrera CP, Christian RJ, Girard O. Thermal, physiological and perceptual strain mediate alterations in match-play tennis under heat stress. *Br J Sports Med.* 2014;48(Suppl. 1):i32–8.
96. Prieto-Lage I, Paramés-González A, Argibay-González JC, Reguera-López-de-la-Osa X, Ordóñez-Álvarez S, Gutiérrez-Santiago A. Match analysis in women's tennis on clay, grass and hard courts. *Int J Environ Res Public Health.* 2022;19(13):7955.
97. Martin C, Thevenet D, Zouhal H, Mornet Y, Delès R, Crestel T, et al. Effects of playing surface (hard and clay courts) on heart rate and blood lactate during tennis matches played by high-level players. *J Strength Cond Res.* 2011;25(1):163–70.
98. Fitzpatrick A, Stone JA, Choppin S, Kelley J. Investigating the most important aspect of elite grass court tennis: short points. *Int J Sports Sc Coach.* 2021;16(5):1178–86.
99. Rudzki K. ITF introduces three types of balls to counter power game. Independent. 2001. <https://www.independent.co.uk/sport/tennis/itf-introduces-three-types-of-balls-to-counter-power-game-9206610.html>. Accessed 21 June 2022.
100. Brown E. Gender and surface effects on elite tennis strategy. *ITF Coach Sport Sci Rev.* 2007;15(46):9–11.
101. Hizan H, Whipp P, Reid M. Comparison of serve and serve return statistics of high performance male and female tennis players from different age-groups. *Int J Perform Anal Sport.* 2011;11(2):365–75.
102. Vickery WM, Dascombe BJ, Baker JD, Higham DG, Spratford WA, Duffield R. Accuracy and reliability of GPS devices for measurement of sports-specific movement patterns related to cricket, tennis, and field-based team sports. *J Strength Cond Res.* 2014;28(6):1697–705.
103. Giles B, Peeling P, Dawson B, Reid M. How do professional tennis players move? The perceptions of coaches and strength and conditioning experts. *J Sports Sci.* 2019;37(7):726–34.
104. Kolman NS, Kramer T, Elferink-Gemser MT, Huijgen BCH, Visscher C. Technical and tactical skills related to performance levels in tennis: a systematic review. *J Sports Sci.* 2019;37(1):108–21.
105. Martínez-Gallego R, J FG, James N, Pers J, Ramón-Llin J, Vuckovic G. Movement characteristics of elite tennis players on hard courts with respect to the direction of ground strokes. *J Sports Sci Med.* 2013;12(2):275–81.
106. ITF. ITF 2018 Rules of tennis London: ITF Ltd; 2018. <http://itf.uberflip.com/i/920624-2018-rules-of-tennis/15?%29>. Accessed 15 Mar 2022.
107. WTA. Women's Tennis Association 2018 Official Rulebook. USA: WTA Tour, Inc; 2018. <http://wtafiles.wtatennis.com/pdf/publications/2018WTARulebook.pdf>. Accessed 13 Jan 2023.
108. ATP. The 2018 ATP official rulebook. USA: ATP Tour; 2018.
109. Kolbinger O, Großmann S, Lames M. A closer look at the prevalence of time rule violations and the inter-point time in men's Grand Slam tennis. *J Sports Analytics.* 2019;5:75–84.
110. Mühlberger A, Kolbinger O. The serve clock reduced rule violations, but did not speed up the game: a closer look at the inter-point time at the 2018 US Open. *J Hum Sport Exerc.* 2021;16(3):528–40.
111. ITF WorldTennisTour. 2019 Men's & Women's ITF World Tennis Tour Regulations. London: ITF Ltd; 2019. <https://aut.uy/wp/wp-content/uploads/2019/06/2019-ITF-Mens-Womens-Regulations.pdf>. Accessed 13 Jan 2023.
112. WTA. 2019 WTA Official Rulebook. USA: WTA; 2019. <http://wtafiles.wtatennis.com/pdf/publications/2019WTARulebook.pdf>. Accessed 13 Jan 2023.
113. WTA. WTA Tour 2022 Official Rulebook. USA: WTA; 2022. <https://photoresources.wtatennis.com/wta/document/2022/01/26/125189f7-fe9f-4aaf-8ff4-88973e54bd9a/2022-WTA-Rulebook-1-26-2022-.pdf>. Accessed 13 Jan 2023.
114. Brosnan RJ, Watson G, Stuart W, Twentyman C, Kitic CM, Schmidt M. The validity, reliability, and agreement of global positioning system units-can we compare research and applied data? *J Strength Cond Res.* 2021;36(12):3330–8.
115. Mecheri S, Rioult F, Mantel B, Kauffmann F, Benguigui N. The serve impact in tennis: first large-scale study of big Hawk-Eye data. *Statistical analysis and data mining. ASA Data Sci J.* 2016;9(5):310–25.
116. Reilly T, Morris T, Whyte G. The specificity of training prescription and physiological assessment: a review. *J Sports Sci.* 2009;27(6):575–89.
117. Hughes DC, Ellefsen S, Baar K. Adaptations to endurance and strength training. *Cold Spring Harb Perspect Med.* 2018;8(6):a029769.
118. Goossens RD, Kempeneers J, Koning HR, Spijksma FCR. Winning in straight sets helps in Grand Slam tennis. *Int J Perform Anal Sport.* 2015;15(3):1007–21.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Authors and Affiliations

Babette M. Pluim^{1,2,3,29,30}  · Marleen G. T. Jansen^{4,5}  · Samuel Williamson⁶ · Cain Berry⁷ · Silvia Camporesi⁸  · Kristina Fagher⁹  · Neil Heron^{10,11}  · Dina C. Janse van Rensburg^{1,12}  · Víctor Moreno-Pérez¹³  · Andrew Murray^{14,15}  · Seán R. O'Connor¹⁶  · Fábio C. L. de Oliveira¹⁷  · Machar Reid¹⁸  · Miriam van Reijen^{4,19} · Tobias Saueressig²⁰  · Linda J. Schoonmade²¹  · Jane S. Thornton^{22,23}  · Nick Webborn^{24,25,26}  · Clare L. Ardern^{27,28} 

✉ Babette M. Pluim
b.pluim@knltnb.nl

¹ Section Sports Medicine, Faculty of Health Sciences, University of Pretoria, Pretoria, South Africa

² Amsterdam Collaboration on Health and Safety in Sports (ACHSS), IOC Research Center of Excellence, Amsterdam, The Netherlands

³ Medical Department, Royal Netherlands Lawn Tennis Association (KNLTB), Amstelveen, The Netherlands

- 4 Toptennis Department, Royal Netherlands Lawn Tennis Association (KNLTB), Amstelveen, The Netherlands
- 5 Centre for Human Movement Sciences, University Medical Centre Groningen, University of Groningen, Groningen, The Netherlands
- 6 English Institute of Sport, London, UK
- 7 Lawn Tennis Association, London, UK
- 8 Department of Political Science, University of Vienna, Vienna, Austria
- 9 Rehabilitation Medicine Research Group, Department of Health Sciences, Lund University, Lund, Sweden
- 10 Centre for Public Health, Queen's University Belfast, Belfast, Northern Ireland
- 11 School of Medicine, Keele University, Staffordshire, UK
- 12 Medical Committee, World Netball, Manchester, UK
- 13 Center for Translational Research in Physiotherapy, Department of Pathology and Surgery, Miguel Hernández University of Elche, San Joan, Spain
- 14 Sport and Exercise, University of Edinburgh. Pleasance, Edinburgh, UK
- 15 European Tour Health and Performance Institute, European Tour Golf, Virginia Water, UK
- 16 School of Psychology, Queen's University Belfast, Belfast, UK
- 17 Research Unit in Sport and Physical Activity (CIDAF), Faculty of Sport Sciences and Physical Education, University of Coimbra, Coimbra, Portugal
- 18 Tennis Australia, Melbourne, VIC, Australia
- 19 National Olympic Committee TeamNL, Arnhem, The Netherlands
- 20 Physio Meets Science GmbH, Leimen, Germany
- 21 University Library, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands
- 22 Return to Health and Performance Lab, Western Centre for Public Health and Family Medicine, Schulich School of Medicine and Dentistry, Western University, London, ON, Canada
- 23 Fowler Kennedy Sport Medicine Clinic, Western University, London, ON, Canada
- 24 IPC Medical Committee, Bonn, Germany
- 25 Centre for Sport and Exercise Science and Medicine (SESAME), School of Sport and Service Management, University of Brighton, Brighton, UK
- 26 School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, UK
- 27 Department of Family Practice, University of British Columbia, Vancouver, BC, Canada
- 28 Sport and Exercise Medicine Research Centre, La Trobe University, Melbourne, VIC, Australia
- 29 Department of Orthopaedic Surgery and Sports Medicine, Amsterdam UMC location University of Amsterdam, Amsterdam, The Netherlands
- 30 Academic Center for Evidence-based Sports Medicine, ACES, Amsterdam, The Netherlands