

IX INTERNATIONAL SYMPOSIUM ON AGRICULTURAL SCIENCES

24th September 2020
Banja Luka
Bosnia and Herzegovina

BOOK OF ABSTRACTS



AGRORES 2020



BOOK OF ABSTRACTS

IX International Symposium on Agricultural Sciences "AgroReS 2020"
24 September, 2020; Banja Luka, Bosnia and Herzegovina

Publisher

University of Banja Luka
Faculty of Agriculture
University City
Bulevar vojvode Petra Bojovića 1A
78000 Banja Luka, Republic of Srpska, B&H

Editor in Chief

Željko Vaško

Technical Editors

Biljana Rogić

Circulation

online on the website <https://agrores.net/zbornici/>

CIP - Каталогизacija у публикацији

Народна и универзитетска библиотека

Републике Српске, Бања Лука

631(048.3)

INTERNATIONAL Symposium on Agricultural
Sciences (9 ; Banja Luka ; 2020)

Book of Abstracts [Elektronski izvor] / 9th
International Symposium on Agricultural Sciences
"AgroReS 2020", 24 September, 2020, Banja Luka,
Bosnia and Herzegovina ; [organizer University of
Banjaluka, Faculty of Agriculture ; editor in chief Željko
Vaško]. - Banja Luka : Faculty of Agriculture =
Poljoprivredni fakultet, 2020

Način pristupa (URL): <https://agrores.net/zbornici/>

ISBN 978-99938-93-63-9

COBISS.RS-ID 129416961

IX INTERNATIONAL SYMPOSIUM ON
AGRICULTURAL SCIENCES



BOOK OF ABSTRACTS

24 September, 2020
Banja Luka
Bosnia and Herzegovina

P1_10

Determination of the fumonisins content in different small grains

Iva Savić¹, Milica Nikolić¹, Vesna Kandić¹, Dejan Dodig¹, Ana Obradović¹,
Danijela Ristić¹, Slavica Stanković¹

¹ *Maize Research Institute "Zemun Polje", Serbia*

Corresponding author: Iva Savić, isavic@mrizp.rs

Abstract

Fungi of the *Fusarium* genus belonging to the Liseola section can synthesise fumonisins of greater or smaller concentrations. Maize is a primary host of these species. In recent years their presence has been also observed in small grains. The aim of this study was to observe the concentrations of fumonisins synthesised by small grains after artificial inoculation. Twelve isolates of the following species were selected from the collection of fungal cultures of the Maize Research Institute, Zemun Polje for artificial inoculation: *Fusarium verticillioides* (6), *Fusarium subglutinans* (3) and *Fusarium proliferatum* (3). The concentrations of synthesised fumonisins were analysed in four small grains: wheat (Aurelia), barely (Nektar), triticale (Zenit) and durum (Cosmostar). The artificial inoculation was performed with the hand sprayer when more than a half of tested plants were in the full-blossom stage. Inoculation of plants was done in four replications. The amount of inoculum (spore concentration was 1×10^{-6} per 1 ml) was 20 ml per a group of 20 spikes. The isolate of *Fusarium graminearum* species was used for spike inoculation in the positive control, while sterile distilled water was used in the negative control. Inoculated spikes were covered with wet PVC bags that were removed after 48h. After harvest, fumonisins were analysed by the ELISA test (Tecna, Italy). According to obtained results not a single isolate of the observed species synthesised fumonisins in the barley crop. In the remaining crops, isolates of *F. subglutinans* species synthesised fumonisins in low concentrations (0.793-24.949 ppm), while the corresponding values of isolates of *F. proliferatum* species were high and ranged from 35.886 to 60.000 ppm. Isolates of *F. verticillioides* species had low values in wheat (2.162-7.925 ppm), while these values in durum were high (29.610-47.174 ppm). The mean values of synthesised fumonisins in the triticale crop were low (1.357-32.587 ppm).

Key words: cereals, *Fusarium*, fumonisins