



## Priority management actions for alien freshwater fish species in Australia

John D. Koehn & Rachel F. MacKenzie

To cite this article: John D. Koehn & Rachel F. MacKenzie (2004) Priority management actions for alien freshwater fish species in Australia, *New Zealand Journal of Marine and Freshwater Research*, 38:3, 457-472, DOI: [10.1080/00288330.2004.9517253](https://doi.org/10.1080/00288330.2004.9517253)

To link to this article: <https://doi.org/10.1080/00288330.2004.9517253>



Published online: 30 Mar 2010.



Submit your article to this journal [↗](#)



Article views: 1492



View related articles [↗](#)



Citing articles: 39 View citing articles [↗](#)

## Priority management actions for alien freshwater fish species in Australia

JOHN D. KOEHN

Arthur Rylah Institute for Environmental Research  
Department of Sustainability and Environment  
123 Brown Street, Heidelberg  
VIC 3084, Australia  
email: John.Koehn@dse.vic.gov.au

RACHEL F. MACKENZIE

Queensland Fisheries Service  
GPO Box 46, Brisbane  
Q 4001, Australia  
email: Rachel.Mackenzie@nrm.qld.gov.au

**Abstract** In Australia, alien freshwater fish are continuing to steadily increase in number of species (reported in this paper to be 43), abundance, and distribution. In general however, their impacts are not well quantified in either environmental or economic terms and current management to reduce their impacts is limited and lacking direction. Although carp *Cyprinus carpio* have received some attention, very little is known about the impacts and even the distribution of most species. There is a lack of recognition of the problem, inconsistency in legislation, policy, and approaches across jurisdictions, and no nationally coordinated on-the-ground management actions. Where legislation and policy is available it is not always used to good effect. This paper provides a synthesis of existing knowledge of alien fishes in Australia, suggests a new management approach, and recommends priority management actions.

**Keywords** invasive fishes; control; impacts; management; benefit:cost; carp

## INTRODUCTION

Alien fish species have received considerable attention internationally with the integrity of aquatic ecosystems being challenged worldwide by species' invasions (Moyle & Light 1996). Biological exchange is considered a relatively more important threat to the biodiversity of freshwater ecosystems compared to other ecosystems because of the intentional and unintentional release of organisms (Sala et al. 2000). The resultant loss of biodiversity caused by alien species is generally severe and well documented (Lodge & Shrader-Frechette 2003), although such losses have not necessarily been well documented in Australia. Some attention has been focused on assessment of potential invaders, however the most likely short-term impacts of alien species will come from those already present in the country, region, or catchment. Concern has been expressed at the lack of recognition, commitment, consistency of approach, coordination, and on-ground actions in relation to alien freshwater fish species in a deteriorating Australian situation. Although many reviews and management plans have been undertaken, few have been fully implemented or widely published and many only consider individual species or particular areas.

The purpose of this paper is to synthesise the Australian literature (including "grey" literature), identify current management frameworks for invasive freshwater fish species in Australia, assess the current status in terms of species knowledge and management, and suggest a new management approach and a list of priority management actions.

## METHODS


Information in this paper has been compiled from existing literature, including unpublished reports, and discussions with State and Commonwealth agencies. However, it does not attempt to provide a comprehensive review of all literature relating to alien fish species. Several general reviews of

established aquatic species have been conducted, covering a range of aspects of their biology and management (Arthington & Mitchell 1986; Arthington 1991; Arthington & Bluhdorn 1995; Arthington & McKenzie 1997; Arthington et al. 1999; Clarke et al. 2000). Recently a range of more detailed reviews for some species and regions have also been produced that include management plans and recommendations and these are listed in Table 3. Unfortunately, most of these publications are not widely available but their recommendations have been collated and have assisted in forming the list of priority actions set out in this paper.

The terminology relating to invasive species is large and many terms are used interchangeably. In this paper, alien species refer to those species intentionally or accidentally dispersed by human agency outside their historically known native range (Department of Primary Industries Queensland (DPIQ) 2001; [www.iucn.org/themes/ssc/pubs/policy/invasives](http://www.iucn.org/themes/ssc/pubs/policy/invasives)). Introduced species are those alien species found in the wild, but not yet breeding successfully, whereas established species refer to those species that have bred successfully and formed a self-sustaining population (Williamson & Fitter 1996). Translocation is the movement of living organisms from one area, with free release into another ([www.iucn.org/themes/ssc/pubs/policy/transe.htm](http://www.iucn.org/themes/ssc/pubs/policy/transe.htm)). Quite clearly, by definition, the term alien species does include favoured angling species such as brown trout and rainbow trout and Australian species translocated outside their natural range. There has been a traditional reluctance by some fisheries agencies, which actively advocate and stock such species, to adequately address their negative environmental impacts (Jackson et al. 2004). Some alien species also meet the criteria to be considered as pest species, a pest being any species that has a negative economic or ecological impact (Olsen 1998; DPIQ 2001).

### Current species status

The total number of alien fish species (31) (Lintermans 2004) established on mainland Australia, is greater than the number of established alien mammals (25) and birds (20) and far more than amphibians (1) or reptiles (4) (Bomford 2001). A total of 43 alien fish species have now been recorded in the wild in Australia (this paper) and McNee (2002) identified 1181 alien freshwater fish species that have been present in Australia, mostly in aquaria, over the past 40 years. Only 481 of these are species on the current permitted import list and



**Fig. 1** Numbers of alien fish species reported in popular fish books from 1967 to 2002. Data from: Weatherley & Lake (1967); McDowall (1980, 1996); and Allen et al. (2002).

other species may be present as the result of illegal imports (Kailola 2000), which are estimated to comprise 5–10% of fish imported into Australia (Australian Quarantine Inspection Service (AQIS) 1999). Australia has five of the world's eight "worst" invasive fish taxa as detailed by the International Union for the Conservation of Nature (Lowe et al. 2000): brown trout (*Salmo trutta* L.), carp (*Cyprinus carpio* L.), Mozambique tilapia (*Oreochromis mossambicus* Peters), rainbow trout (*Oncorhynchus mykiss* (Walbaum)), and gambusia (*Gambusia affinis/holbrooki* Baird & Girard).

The number of alien species included in popular fish books may be one measure of public recognition of alien fish species becoming established in Australia. This number of species has grown steadily from eight in 1967 (Weatherley & Lake 1967) to 21 in 2002 (Allen et al. 2002) (Fig. 1). In more specialised publications, McKay (1989) reported the presence of 20 alien freshwater fish, Arthington et al. (1999) reported 37, and this paper reports 43.

Australia has a relatively depauperate freshwater fish fauna by world standards with slightly over 200 species (Allen et al. 2002). More than 16% of these are considered to be under serious conservation threat nationally (Crook 2001) and detrimental interactions with alien species are considered a threat to 77% of these species (Jackson et al. 1993).

The invasion of carp in Australia illustrates how quickly an introduced fish species can spread and dominate fish communities (in numbers and biomass) in the absence of early intervention and an effective management strategy. Following the introduction of the "Boolarra" strain in Victoria in the early 1960s, carp have become the most abundant large freshwater fish in south-east Australia, now

**Table 1** Summary of distribution and reason for introduction for established alien freshwater fish species in Australia (Arthington & Bluhdorn 1995; Arthington et al. 1999; Clarke et al. 2000; Clunie et al. 2002). Reason for introduction: P, pet and aquarium; O, ornamental; B, biological control; R, recreational angling; A, aquaculture; and S, ship ballast. (VIC, Victoria; NSW, New South Wales; ACT, Australian Capital Territory; QLD, Queensland; NT, Northern Territory; WA, Western Australia; SA, South Australia; TAS, Tasmania.)

Species		Reason for introduction	Drainage division	Distribution (States)
Family and scientific name	Common name			
<b>SALMONIDAE</b>				
<i>Oncorhynchus mykiss</i>	rainbow trout	R	2, 3, 4, 5, 6	VIC, NSW, ACT, WA, SA, TAS
<i>Salmo trutta</i>	brown trout	R	1, 2, 3, 4, 5, 6	VIC, NSW, ACT, QLD, WA, SA, TAS
<i>Salmo salar</i>	Atlantic salmon	R	2, 3, 4	VIC, NSW, SA, TAS
<i>Salvelinus fontinalis</i>	brook trout	R	2, 3, 4	NSW, TAS
<b>CYPRINIDAE</b>				
<i>Carassius auratus</i>	goldfish	O	1, 2, 3, 4, 5, 6,	VIC, NSW, ACT, QLD, WA, SA, TAS
<i>Cyprinus carpio</i>	carp	O, A	1, 2, 3, 4, 5, 6	VIC, NSW, ACT, QLD, WA, SA, TAS
<i>Rutilus rutilus</i>	roach	R	2, 4	VIC, NSW
<i>Tinca tinca</i>	tench	R	2, 3, 4	VIC, NSW, TAS, SA
<b>COBITIDAE</b>				
<i>Misgurnus anguillicaudatus</i>	oriental weatherloach	O	1, 2, 4, 5	VIC, NSW, ACT, QLD, SA
<b>POECILIIDAE</b>				
<i>Gambusia holbrooki</i>	eastern gambusia	B	1, 2, 3, 4, 5, 6, 7, 8, 10	VIC, NSW, ACT, QLD, NT, WA, SA, TAS
<i>Phalloceros caudimaculatus</i>	caudo/one-spot livebearer	O, B(?)	2, 6	NSW, WA
<i>Poecilia latipinna</i>	sailfin molly	O, B	1	QLD
<i>Poecilia reticulata</i>	guppy	O	1, 2, 6, 8	NSW, QLD, WA, NT
<i>Xiphophorus helleri</i>	green swordtail	O	1, 2, 6, 8, 10	NSW, QLD, WA, NT
<i>Xiphophorus maculatus</i>	platy	O	1, 2, 8, 10	NSW, QLD, NT
<b>PERCIDAE</b>				
<i>Perca fluviatilis</i>	redfin perch	R	2, 3, 4, 5, 6	VIC, NSW, ACT, WA, SA, TAS
<b>CICHLIDAE</b>				
<i>Archocentrus nigrofasciatus</i>	convict cichlid	O	1, 2	VIC, QLD
<i>Astronotus ocellatus</i>	oscar	O	1	QLD
<i>Oreochromis mossambicus</i>	tilapia	O	1, 7	QLD, WA
<i>Tilapia mariae</i>	black mangrove cichlid	O	1, 2	VIC, QLD
<b>GOBIIDAE</b>				
<i>Acanthogobius flavimanus</i>	yellowfin goby	S	2	VIC, NSW

distributed over more than 1 million km<sup>2</sup> (KoeHN et al. 2000). The spread of tilapia in northern Australia is another example of a species becoming established as a result of a lack of action in the early stages of invasion and continued human movement of fish. Tilapia were released into ornamental ponds in the 1970s and have formed large populations around the cities of Townsville and Cairns (Fig. 1) (Arthington et al. 1984). Tilapia have formed separate, established populations around Brisbane and recently expanded into the Burnett River catchment (DPIQ 2001). Similarly, in Western Australia, tilapia, which were previously restricted to the Gascoyne river in the Pilbara drainage (basin 7), have "spread" to rivers immediately to the north and south (Morgan et al. 2003). Although the distribution of carp and tilapia is reasonably well documented, this is not the

situation for most other species. Literature reviews and contact with State agencies have shown that although invasions into new areas are common, few surveys have been conducted, and data have not been collated and are often of poor quality. Reports of many invasions come from the public and research work rather than dedicated alien fish surveys. As such, records of alien species are not reported in a consistent way and even the generalised distributions given in Tables 1 and 2 may be incomplete. In particular, whether or not established populations occur is uncertain for many species.

In addition to alien species, it is believed that at least 51 native fish species have been translocated outside their natural range in Australia (Lintermans 2004). Banded grunter (*Amniataba percooides* Gunther), has been declared a noxious species in

**Table 2** Summary of distribution and reason for introduction for introduced alien freshwater fish species in Australia (Allen 1989; Arthington et al. 1999; Raadik 2001, 2003a,b; Webb 1994, cited in Raadik 2003b; Raadik unpubl. data). Reason for introduction: O, ornamental; R, recreational angling. (VIC, Victoria; NSW, New South Wales; ACT, Australian Capital Territory; QLD, Queensland; NT, Northern Territory; WA, Western Australia; SA, South Australia; TAS, Tasmania.)

Family and scientific name	Common name	Reason for introduction	Distribution	
			Basins	States
<b>SALMONIDAE</b>				
<i>Oncorhynchus tshawytscha</i>	Chinook salmon	R	2	VIC
<b>CYPRINIDAE</b>				
<i>Puntius conchonius</i>	rosy barb	O	1, 2	QLD, NSW
<i>Puntius tetrazona</i>	Sumatra barb	O	1	QLD
<i>Tanichthys albonubes</i>	white-cloud mountain minnow	O	2	NSW
<b>CYPRINODONTIDAE</b>				
<i>Jordanella floridae</i>	American flagfish	O	1	QLD
<b>CICHLIDAE</b>				
<i>Aequidens pulchrus</i>	blue acara	O	2	VIC, QLD
<i>Aequidens rivulatus</i>	green terror	O	1	QLD
<i>Amphilophus citrinellus</i>	midas cichlid	O	1	QLD
<i>Amphilophus labiatus</i>	red devil	O	2	VIC
<i>Cichlasoma octofasciatum</i>	Jack Dempsey	O	1	QLD
<i>Cichlasoma severum</i>	banded cichlid	O	1	QLD
<i>Cichlasoma synspilum</i>	redhead cichlid	O	1	QLD
<i>Cichlasoma trimaculatum</i>	three spot cichlid	O	1	QLD
<i>Cichlasoma brasiliensis</i>	pearl cichlid	O	1	QLD
<i>Haplochromis burtoni</i>	Burton's haplochromis	O	1	QLD
<i>Hemichromis guttatus</i>	jewel cichlid	O	1	QLD
<i>Heros severus</i>	green severum	O	1	QLD
<i>Labeotrophus/Pseudotrophus</i>	Hybrid?	O	2	VIC
<i>Tilapia zillii</i>	redbelly tilapia	O	1	QLD
<i>Thorichthys meeki</i>	firemouth	O	1	QLD
<b>BELONTIIDAE</b>				
<i>Trichogaster trichopterus</i>	three-spot gourami	O	1	QLD
<b>CYPRINIDAE</b>				
Unidentified cyprinid	?	R, O?	1	QLD

**Fig. 2** Map of Australian drainage division boundaries (dotted lines) and State borders (solid lines, bold text).



NSW (NSW Fisheries 2002; Table 1) and concern has been expressed at the potential introduction of Murray cod (*Maccullochella peelii peelii* Mitchell) and golden perch (*Macquaria ambigua* Richardson) into Western Australia for aquaculture and recreational fishing (Morgan et al. 2002). Murray cod have illegally been translocated into waterholes in the south of the Northern Territory, raising concerns that they may find their way onto the Finke River system, a system naturally devoid of a large piscivorous fish (H. Larsen pers. comm.). Native and exotic species may also escape from aquaculture facilities with examples including brown and rainbow trout, the native Australian bass (*Macquaria novemaculeata* Steindachner), trout cod (*Maccullochella macquariensis* Cuvier), and short-finned eels (*Anguilla australis* Richardson). The recent arrival into Victoria of the Barcoo grunter (*Scortum barcoo* McCulloch & Waite), native to Lake Eyre and catchments in Queensland and the Northern Territory, gives cause for concern (Koehn 2002). Records of translocations are erratic, need to be formalised, and should be included as a component of alien species management plans.

### Current legislation and management

Australia's 11 main river basin boundaries do not generally coincide with jurisdictional boundaries, meaning that alien fish species are managed, at least

in part, by nine different governments (Fig. 2) operating under different legislation, with different management approaches (Arthington & McKenzie 1997). Not surprisingly, there is inconsistency in approach across these jurisdictions. For example, until recently, carp have had a different legal status in States within the Murray-Darling Basin. In Queensland they are declared noxious, which means it is unlawful to possess them live or dead. They are also declared "noxious" in Victoria meaning they cannot be returned to the water alive. However, although they were only declared as a "class 1" noxious species in New South Wales in December 2002, this listing still allows for their continued legal possession, sale, and distribution, including ornamental "Koi" carp (NSW Fisheries 2002). This inconsistency has significant implications for carp dispersal in eastern Australia and it has been recommended that their legal status be changed to enable national coordination to control their spread (Georges & Cottingham 2002). Some States use the terms "controlled", "undesirable", or "pest" in their legislation when referring to alien fish species. The range of terminology used, and inconsistent definitions, make it difficult to compare the legislative status of alien fish between states. The number of taxa listed as "noxious" varies across jurisdictions, from none in the Australian Capital Territory (ACT) to over 80 in Victoria, with many

alien fish species in many States having no legal status at all. It could also be said that where legislation exists, it is generally poorly adhered to, enforced, or publicised. Fisheries compliance is sparsely resourced, is rarely focused on alien fish, and must be accompanied by education aimed to change community attitudes.

It is fair to say that alien fish species management in Australia has been overly focused on carp, with many other damaging species having received little attention (e.g., *Gambusia*, salmonids). Queensland is the only State to develop a multi-species pest management strategy (DPIQ 2001). This document sits under the "Queensland Pest Animal Strategy" (Queensland Department of Natural Resources and Mines 2002) and although a National Carp Strategy was developed and endorsed by the Murray-Darling Basin Ministerial Council, the recommended actions are not necessarily implemented by State agencies. Often, where policy and legislation instruments are available, they have not been used to assist on-ground alien fish species management, and fish are generally not included in State pest management policies. In Victoria, a Potentially Threatening Process encompassing alien fish introductions was listed under the Flora and Fauna Guarantee Act 1992 (Scientific Advisory Committee 1992), but no Action Statement (management plan) detailing an appropriate response to ameliorate this threat has yet been produced. "The Victorian Pest Management—A Framework for Action" (Natural Resources & Environment (NRE) 2001) is focused upon weeds and more traditional terrestrial vertebrates such as rabbits and foxes, rather than freshwater species. The Victorian River Health Strategy (NRE 2002) pays scant regard to alien species, with fish not included as taxa for the monitoring of river health. However, changes in the geographic distribution of indigenous and exotic fish species has been recommended as an environmental health indicator (Office for the Commissioner for the Environment 1988) and fish and alien fish species have been considered in other measures of river health (Harris 1995; Whittington et al. 2001). A recent nomination for listing of "The introduction of live fish to waters outside their natural range within a river catchment after 1770" as a Key Threatening Process under the Commonwealth Environmental Protection and Biodiversity Conservation Act 1999 was unsuccessful. The ability of this legislation to list widespread threatening processes is yet to be proven, although predation by *Gambusia holbrooki* has been listed as a Key

Threatening Process under the New South Wales Threatened Species Conservation Act 1995.

Management of vertebrate pests in Australia continues to focus on terrestrial animals despite several alien fish species meeting pest species criteria because "they threaten the environmental and personal resources valued by humans" (Olsen 1998). The inclusion of carp in the Bureau of Rural Sciences Vertebrate pest series (Koehn et al. 2000) is recognition of their potential as a pest species, but formal steps need to be taken for them to be included in the development of vertebrate pest plans and associated funding. Carp are the largest and most visible of introduced fish species, have received the most public attention, and now rate in the public eye along with rabbits and foxes as a vertebrate pest species in south-eastern Australia (Koehn 2001). Fish have only recently been included in papers presented at an Australian Vertebrate Pests Conference (Koehn 2001; Mackenzie & Bryant 2001; Stuart et al. 2001), and changes to the traditional view of vertebrate pests is urgently needed to consider the environmental damage caused by alien fish species.

### Current knowledge

Although our understanding of the impacts of alien fish is poor, and there is a lack of coordination, a review of the literature shows there is a range of information available that could form the basis of improved management of alien freshwater fish species in Australia. This information is of three types: (1) general strategic documents; (2) area based assessments; and (3) reviews of individual species. However, a coordinated approach such as that outlined for marine pests (National Taskforce on the Prevention and Management of Marine Pest Incursions 1999) is needed.

#### *General strategic documents*

The major issue is not whether we understand what management actions need to be undertaken for alien fish, but whether there is the will to implement these actions in a consistent and coordinated way. Carp have received the most publicity of alien fish species, have been widely reviewed (Hume et al. 1983; Roberts et al. 1997; Koehn et al. 2000), and been the subject of a National Strategy (Braysher & Barrett 2000; Carp Control Coordinating Group 2000a,b). Although this strategy relates only to this species, it can easily be adapted to include other alien species, as has been done in Queensland (DPIQ 2001), forming the basis of a national alien species

management plan. The Action Plan for the Murray-Darling Basin, developed for alien species by the National Carp and Pest Fish Taskforce, a community focused organisation (Murray-Darling Association Inc. 2003), could also be expanded and incorporated into a national strategy. The “National Policy for the Translocation of Live Aquatic Organisms” (Ministerial Council on Forestry, Fisheries and Aquaculture 1999) is seen as a valuable document, whose recommendations and processes unfortunately are not always being followed across the country. This document could be updated from the more detailed 43 recommendations made at the recent Murray-Darling Basin workshop on managing fish translocation and stocking (Phillips 2002). Although concentrating on the Murray-Darling Basin, this document has wider relevance. Issues of alien species relating to interbasin water transfers have been addressed by Todd et al. (2002).

#### *Area based assessments*

A risk assessment has been undertaken for alien species within the Murray-Darling Basin (Clunie et al. 2000). A pre-emptive assessment for the management of tilapia, should they enter the Murray-Darling Basin (Braysher 2001), also provides a useful example that could be followed for other species. Carp have been studied comprehensively in the Barmah-Millewa area (Stuart & Jones 2002), which has been identified to be an important area for their recruitment (Stuart et al. 2000). These documents together with regional carp management plans could be adapted more widely to similar areas and other species.

#### *Reviews of individual species*

Of the 43 alien species considered in Tables 1 and 2, brief reviews have been conducted for 21 established species (Arthington & Bluhdorn 1995). More comprehensive reviews have also been conducted for several of these species (Table 3) providing important information regarding biology, impacts, management recommendations, and knowledge gaps. However, little information appears to be available on introduced or establishing species (Table 2), and no studies provide detailed predictions for future spread or impacts. There is currently no red list of the worst potential aquatic invaders and, consequently, no additional procedures to stop them before they arrive. However, there is potential under the Environmental Protection and Biodiversity Conservation Act for the establishment and maintenance of a list of non-native species that pose

a threat to Australian biodiversity. Kailola (2000) provides initial steps toward such a list. Risk assessment has been considered for some of these species, but no detailed predictions have been made for future spread. It is these more recent introductions that have the greatest potential for spread and further impacts. Clear examples of these include the rapid spread of oriental weatherloach (*Misgurnus anguillicaudatus* Cantor) (Koster et al. 2002a,b) and tilapia (DPIQ 2001) in many regions.

#### **Knowledge gaps**

There are few detailed species-specific risk assessments (e.g., Barlow & Lisle 1987; Townsend & Winterbourn 1992) predicting the future impacts of introduced fish species that are already established (Arthington & Bluhdorn 1995; Coates & Ulaiwi 1995; Clunie et al. 2001). Such predictive assessments need to be undertaken for other species in Australia, giving consideration to dispersal pathways (e.g., Lintermans 2004) and climatic limitations (Sutherst et al. 1996, 1999). Our knowledge of the biology and ecology of many of these species, especially under Australia conditions, is limited. Our understanding of the impacts of most alien species is still rudimentary, even for direct impacts such as predation, let alone impacts at the community or ecosystem levels (Townsend 2003). An understanding of impacts is important to build the case for the importance of adequate alien species management and to provide some basis for benefit:cost analyses of management options (Choquenot et al. 2004). Understanding impacts is an essential component of best practice vertebrate pest management, which is based on the concept of managing impacts rather than numbers (Braysher 1993). Evaluation of the environmental and economic impacts of the world's best known biotic invasion, the zebra mussel (*Dreissena polymorpha* Pallas) into the North American Great Lakes region, has underpinned public support and management actions (Lodge 1993; Nalepa & Schloesser 1993; Strayer 1999; Pimentel et al. 2000). There is the need for similar evaluations in Australia.

Potential control techniques for most aquatic pest species are limited. Options for carp are explored in detail in Roberts & Tilzey (1997) and Koehn et al. (2000) and include: removal (commercial and recreational); environmental rehabilitation; environmental manipulation; biomanipulation (e.g., adding predators); exclusion; poisoning; and future biological controls (e.g., Thresher unpubl. data). As many of these “biocontrol” options have yet to be



developed and tested, they are not available at present and usually involve long time frames for development, testing, approval, and deployment. Other more specific options for carp (are currently under development) (e.g., separation cages, Stuart et al. 2003). Despite the public favouring fishing and removal, these options are only applicable to some larger species in certain areas, and are unlikely to make a significant contribution to the control of invasive fish species in Australia. It can however have significant value in increasing public awareness of the pest species. This lack of realistic control options highlights the need to support actions of the "National Carp Management Strategy" (Carp Control Coordinating Group 2000a) which utilises a range of management options, highlights the importance of preventing future spread, and recognises the importance of gaining community support. As eradication after establishment is usually impossible, management should concentrate on preventing new introductions and limiting future spread (Lodge et al. 1998). This strategic approach should be adopted for alien fish species other than carp, as has been done in Queensland (DPIQ 2001).

### Future management actions

The goal of management for alien species is not a reduction in numbers *per se*, but a reduction in the impacts caused by each species (Lodge & Shrader-Frechette 2003) (e.g., predation, competition, ecosystem changes). This together with the use of pest management principles (Braysher 1993; Bomford & Tilzey 1997) and adoption of integrated approaches will give the best results. Clearly defining the problem, objectives, and measurables is important before action is taken. This will require the development of new skills and organisational links that go beyond the traditional roles of fisheries agencies and current management expertise.

Range expansions within catchments can occur by natural movement, especially for mobile species, but most transfer between catchments is human assisted (Lintermans 2004). Range expansions should be managed on the basis of drainage basins, with a hierarchy of containment to river reach, river basin, then major drainage division (Fig. 1). This requires coordinated efforts across State and jurisdictional boundaries. Transfer of fish as bait by anglers, either through accident or ignorance, or

**Table 3** Summary of knowledge for established alien freshwater fish species in Australia. (C, comprehensive; M, mostly; P, partial; N, not at all. Impacts: Pr, predation on native fish; H, habitat utilisation; HD, habitat destruction.)

Common name	Impacts		Management plan /recommendations	Knowledge gaps	
	Reviewed	Known		/research plan	References*
Rainbow trout	C	M (Pr)	Yes	Yes	1; 2; 3; 5; 11; 14
Atlantic salmon	C	P (Pr)	Yes	Yes	1; 2; 11
Brown trout	C	M(Pr)	Yes	Yes	1; 2; 3; 5; 11; 14
Brook trout	C	P(Pr)	Yes	Yes	1; 3; 11
Goldfish	P	N	No	No	1; 3
Carp	C	P (H,HD)	Yes	Yes	1; 3; 5; 6; 8; 11; 15; 16
Roach	P	N	Yes	No	1; 3; 11
Tench	P	N	No	No	1; 3; 11
Oriental weatherloach	C	P	Yes	Yes	1; 3; 11; 12
Eastern gambusia	C	M (Pr)	Yes	Yes	1; 4; 5; 8; 9; 11
Caudo/one-spot livebearer	P	M	No	No	1; 3; 11
Sailfin molly	P	M	No	No	1; 3
Guppy	P	M	No	No	1; 3; 5; 11
Green swordtail	P	M	No	No	1; 3; 13
Platy	P	M	No	No	1; 3
Redfin perch	P	M (Pr)	No	No	1; 3; 10
Convict cichlid	P	N	No	No	1; 3; 8
Tilapia	C	N	Yes	Yes	1; 3; 5; 7; 8; 11
Black mangrove cichlid	N	N	No	No	1; 3; 8
Yellowfin goby	N	N	No	No	1; 3

\*1, Arthington & Bluhdorn (1995); 2, Cadwallader (1996); 3, Arthington & McKenzie (1997); 4, Gill et al. (1999); 5, Clarke et al. (2000); 6, Koehn et al. (2000); 7, Braysher (2001); 8, DPIQ (2001); 9, McKay et al. (2001); 11, Morgan & Gill (2001); 11, Clunie et al. (2002); 12, Koster et al. (2002a,b); 13, Morgan et al. (2002); 14, Crowl et al. (1992); 15, Hume et al. (1983); 16, Roberts et al. (1997).

deliberately by “coarse” fishers wanting to establish new fishing grounds, has been recognised as a major source of invasion into new catchments both in Australia and New Zealand (McDowall 1997; Koehn et al. 2000). Social research and appropriate education, management, and enforcement options are required for this issue. The inappropriate disposal of fish available in the aquarium trade remains an increasing problem (McDowall 2004), with 12 of the 21 established species and 21 of the 22 introduced species (77% of alien fish overall) (Tables 1 and 2) being introduced to Australia as ornamental fish. Federal legislation needs to be reviewed to be more inclusive of the species present within the aquarium trade, then adequately enforced. States also need to ensure they have a consistent approach to managing introductions from the aquarium trade. Several species have also invaded as a result of interbasin water transfers (Waters et al. 2000; Todd et al. 2002) and the environmental consequences of these transfers need to be factored into the cost of water, a topical issue in Australia at the moment. Screens installed on dams in Queensland to prevent these transfers have cost more than AU\$1 million with ongoing arguments about the responsibility for payment. These issues need to be addressed at political and social levels with the support and assistance of aquarium and water industries, recreational anglers, and other stakeholders.


Management of priority species has mostly concentrated on species that are already well-established (Arthington & Bluhdorn 1995). While prevention of further spread remains a priority in the management of alien fish, concentrating on newly introduced and establishing species may often provide greater benefits. Most alien fish species occur in south-eastern Australia, particularly the Murray-Darling Basin (Clarke et al. 2000). The alien fish problem in the Murray-Darling Basin provides an important perspective on the future of many other river basins if continued invasion is not prevented. Areas such as the Lake Eyre drainage basin (with a catchment of more than 1.3 million km<sup>2</sup>), the Kimberley, Pilbara, and Northern Territory need priority protection from invasive species such as carp and *Gambusia*.

Dedicated monitoring is essential to alien species management and must relate to clear objectives. However, current resources are insufficient to enable research and management agencies to carry out this monitoring without support from the community. To date, despite significant public interest, there has been no organised community monitoring and

management programme for alien fish. Groups such as “Carpbusters” in Queensland have huge community support and enthusiasm but this needs to be harnessed to provide information suitable for management. Waterwatch is a national community-based monitoring network with 2000 groups monitoring more than 6000 sites in over 200 catchments across Australia. The extent of their monitoring means there are opportunities for Waterwatch to provide data on the occurrence of alien fish species in waterways. With the appropriate training and resources, this existing network could provide an extensive monitoring network (J. McCoy pers. comm.). Involving the community in monitoring activities also provides them with a sense of ownership of the pest fish issue and an improved understanding of the complexities of managing alien fish species.

Community education is a vital component of pest fish management for a number of reasons. First, by raising community awareness of the issue there is less potential for humans to spread alien species. Second, the public can assist with reporting infestations and finally, funding for on-ground natural resource management activities in Australia is largely directed through regional Natural Resource Management (NRM) bodies. These bodies determine NRM priorities for their region and funding is allocated according to those priorities. They also co-ordinate and support other community based activities such as Landcare and Integrated Catchment Management. Therefore, to access funding and support for on-ground alien species management, regional bodies need sufficient information about the significance of the alien species threat and how the community can assist. A broad range of information needs to be developed regarding the threat alien species pose, methods of transferral, and potential management actions. There also needs to be an improved approach to reporting of pest fish sightings by the public with well-publicised and staffed telephone “hotlines”. Queensland has developed a pest fish reporting kit that contains detailed instructions on the information required, how to identify fish, describe their location and other useful information, with a more detailed version provided to Queensland Boating and Fisheries Patrol officers.

Encompassing four States, the ACT and Commonwealth Government jurisdictions, the Murray-Darling Basin (basin 4) can be seen as a microcosm of the rest of Australia in terms of natural resource management. The multi-jurisdictional



**Fig. 3** Potential management framework for coordinated alien freshwater fish species management in Australia.

management currently provided by the Murray-Darling Basin Commission (MDBC) for fish ([www.mdbc.gov.au](http://www.mdbc.gov.au)) could potentially provide a model framework for the management of alien species across Australia. A suggested national management framework involves a lead coordinating agency (probably Environment & Heritage or Agriculture, Fisheries & Forestry Australia), that reports to relevant Ministerial Councils and oversees the formulation and implementation of a National Alien Freshwater Fish Management Plan through the combined efforts of a Management and Science Committee (comprising State and technical representation) and a Community Stakeholder Committee (Fig. 3). The Community Stakeholder Committee could be derived from a restructure of the membership of the existing National Carp and Pest Fish Taskforce (MDA Inc. 2003). The management and science committee may subsequently need to form specialist working groups for particular areas of concern (e.g., angling, water transfer, aquariums). Links to organisations such as the Vertebrate Pests Committee, MDBC, Cooperative Research Centre for Pest Animal Control, AQIS, and research institutions also need to be considered. In recognising that there is a clear need for representation from State agencies, management of pest fish undertaken in Australia must also be supported and coordinated at a Commonwealth level.

This paper deals with only alien fish species, but it is logical that other aquatic invaders (e.g., plants and invertebrates) can be linked into the suggested national strategy. Benchmarking of native and alien fish populations and other environmental parameters are needed before the implementation of management actions to provide baseline data against which the success of actions can be measured.

### Recommended priority actions

The priority actions listed may be conducted over both short and long time frames and should be subjected to a rigorous ranking process undertaken by the managing agencies. Importance will vary between species, areas, and actions already undertaken. The difficulties of comparing the effects of smaller scale interventions to wider actions such as habitat improvement (Harris 1997) and assessing the impacts of alien fish species with those of flow regulation (Gehrke et al. 1995) and other human impacts (Driver et al. 1997) will always be problematic. The integration of science (e.g., alien species biology) with applied management actions however, will improve environmental outcomes (see: Department of Conservation 2003).

#### *Nationally coordinated management*

- (1) Engage an appropriate national agency to coordinate alien freshwater fish species management in Australia.
- (2) Adopt a suitable management framework for the development and implementation of a National Alien Fish Management Strategy, recognising roles, responsibilities and cost sharing arrangements.
- (3) Ensure national consistency in jurisdictional legislation and management approaches including compliance.
- (4) Stringently review and revise existing national permitted and "noxious" fish species lists. Organise a national summit to ensure consistency of lists and compliance across jurisdictions.
- (5) Develop an appropriate strategic process for the non-proliferation and disposal of noxious or unwanted species.

- (6) List deleterious actions of alien fish (e.g., predation) as a Key Threatening Process under relevant State and Commonwealth legislation.
- (7) Recognise fish as potential vertebrate pests in national and State vertebrate pest policies, programmes, and legislation.
- (8) Recognise translocation and stocking of native fish outside their natural range as a component of alien species management.
- (9) Update the National Translocation Policy (see: Phillips 2003) and ensure compliance.
- (10) Develop a national distributional database for alien species and fish translocations.
- (11) Develop criteria to prioritise alien fish species management actions.
- (26) Initiate dedicated alien species monitoring at a community and scientific level.
- (27) Determine appropriate resources necessary for alien species management. Undertake and report bioeconomic assessments (benefit:costs) and evaluation of potential and existing management actions.
- (28) Initiate a research strategy—with priority areas being: alien species knowledge, control and management techniques (including ecosystem rehabilitation), predictive capabilities, quantification of impacts, risk assessment.

#### *Use of existing information*

- (12) Include alien fish management in existing natural resource management frameworks.
- (13) Undertake recommended actions in existing management reports on alien species.

#### *On-ground management*

- (14) Quantify impacts and threats posed by alien fish species within the broader ecosystem context.
- (15) Use pest management principles as a basis for management of alien species.
- (16) Minimise future spread of established and establishing species.
- (17) Engage non-fisheries agencies in alien fish management such as regional NRM bodies.
- (18) Integrate alien fish species management into other natural resource management plans and place in context with other rehabilitation measures for aquatic ecosystems.
- (19) Determine priority native species/habitats for protection e.g., Lake Eyre Basin.
- (20) Develop a consistent strategy for the control of trade and movement of fishes.
- (21) Include alien species distribution and abundance as a measure of river health.
- (22) Establish and resource rapid response plans, processes, and teams to eradicate new infestations and facilitate reporting processes for new invasions.
- (23) Rank sites for control actions.

#### *Knowledge*

- (24) Undertake distributional surveys and collate distributional data on alien fishes.
- (25) Undertake risk assessments for the potential spread and impacts of alien fish species.

#### *Community*

- (29) Restructure and fund National Carp and Pest Fish Taskforce to provide the role of engaging the community in alien fish management.
- (30) Engage key stakeholder groups such as anglers, aquarium and water industries.
- (31) Develop and implement an education and awareness programme to improve community education about alien fish species at a range of levels—political, departmental, enforcement, waterwatch, fisheries, aquarium trade, general public.
- (32) Engage the community in alien species management and regional planning and through “demonstration” alien fish sites. Utilise existing community networks to incorporate alien species awareness e.g., Landcare, conservation groups, Invasive Species Council Inc.
- (33) Undertake social research to minimise deliberate transfers (especially “coarse” anglers).
- (34) Incorporate alien fish into Waterwatch and other community monitoring schemes.

## **CONCLUSION**

A review of the current state of knowledge in Australia indicates that there is much information on many alien freshwater fish species, and we propose a management approach that can be used to formulate a national framework for managing and reducing the impacts of alien species in Australia. It is not lack of knowledge that is impeding our progress managing alien fish but a lack of coordination and perhaps a lack of will at all levels to acknowledge that alien fish really are a problem.

## ACKNOWLEDGMENTS

We thank Jim Barrett, Tarmo Raadik, Richard Tilzey, Alex McNee, and Ivor Stuart for their views and reviews of this paper; inputs from David Morgan, Helen Larsen, and Andria Marshall as representatives of the Australian Society for Fish Biology exotic fishes committee; and John McCoy for his advice regarding Waterwatch monitoring.

## REFERENCES

- Allen, G. R. 1989: Freshwater fishes of Australia. United States, THF Publications Inc.
- Allen, G. R.; Midgley, S. H.; Allen, M. 2002: Field guide to the freshwater fishes of Australia. Australia, Western Australian Museum and CSIRO Publishing.
- Arthington, A. H. 1991: Ecological and genetic impacts of introduced and translocated freshwater fishes in Australia. *Canadian Journal of Fisheries and Aquatic Sciences* 48 (Suppl. 1): 33–43.
- Arthington, A. H.; Bluhdorn, D. R. 1995: Improved management of exotic aquatic fauna: R&D for Australian rivers. Canberra, Land and Water Resources and Development Corporation. *Occasional Paper 04/95*.
- Arthington, A.; McKenzie, F. 1997: Review of impacts of displaced/introduced fauna associated with inland waters. Australia: State of the Environment Technical Paper Series (Inland Waters). Canberra, Department of the Environment.
- Arthington, A. H.; Mitchell, D. S. 1986: Aquatic invading species. In: Groves, E. H.; Burdon, J. J. ed. Ecology of biological invasions: an Australian perspective. Canberra, Australian Academy of Sciences. Pp. 34–52.
- Arthington, A. H.; McKay, R. J.; Russell, D. J.; Milton, D. A. 1984: Occurrence of the introduced cichlid *Oreochromis mossambicus* in Queensland. *Australian Journal of Marine and Freshwater Research* 35: 267–262.
- Arthington, A. H.; Kailola, P. J.; Woodland, D. J.; Zaluki, J. M. 1999: Baseline environmental data relevant to an evaluation of quarantine risk potentially associated with the importation to Australia of ornamental finfish. Report to the Australian Quarantine and Inspection Service. Canberra, ACT, Department of Agriculture, Fisheries and Forestry.
- Australian Quarantine Inspection Service 1999: Exotic fish trade. Report to the National Fisheries Compliance Committee. Canberra, AQIS.
- Barlow, C. J.; Lisle, A. 1987: Biology of the Nile perch (*Lates niloticus* (Pisces: Centropomidae) with reference to its proposed role as a sport fish in Australia. *Biological Conservation* 39: 269–289.
- Bishop, K. A., 1999: Threatened fish species surveying and habitat assessment in the emigrant creek catchment, Richmond River system, New South Wales. Report prepared in relation to the proposed Pacific Highway upgrading: The Ballina Bypass, for Connell Wagner Pty Ltd on behalf of the NSW Roads and Traffic Authority.
- Bomford, M. 2001: Can we quantify the risk that imported vertebrates could establish wild pest populations in Australia? In: Proceedings of the 12th Australian vertebrate pest conference, Melbourne, Victoria, 21–25 May 2001. Victoria, Natural Resources and Environment. Pp. 178–183.
- Bomford, M.; Tilzey, R. 1997: Pest management principles for European carp. In: Roberts, J.; Tilzey R. ed. Controlling carp: exploring the options for Australia. Proceedings of a workshop 22–24 October 1996, Albury. Canberra, CSIRO and Murray-Darling Basin Commission. Pp. 9–20.
- Braysher, M. 1993: Managing vertebrate pests: principles and strategies. Canberra, Bureau of Resource Sciences, Australian Government Publishing Service.
- Braysher, M. 2001: Report on the workshop for the management of Tilapia in the northern Murray-Darling Basin, Toowoomba, 2–3 April, 2001. Canberra, Applied Ecology Research Group, University of Canberra.
- Braysher, M.; Barrett, J. 2000: Ranking areas for action: a guide for carp management groups. Canberra, Murray-Darling Basin Commission.
- Cadwallader, P. L. 1996: Overview of the impacts of introduced salmonids on Australian native fauna. Canberra, Australian Nature Conservation Agency.
- Carp Control Coordinating Group 2000a: National management strategy for carp control 2000–2005. Canberra, Murray-Darling Basin Commission.
- Carp Control Coordinating Group 2000b: Future directions for research into carp. Canberra, Murray-Darling Basin Commission.
- Choquenot, D.; Nicol, S.; Koehn, J. 2004: Bioeconomic modelling in the development of invasive species policy. *New Zealand Journal of Marine and Freshwater Research* 38: 419–428.
- Clarke, G. M.; Grosse, S.; Matthews, M.; Catling, P. C.; Baker, B.; Hewitt, C. L.; Crowther, D.; Saddler, S. 2000: State of the environment indicators for exotic environmental pest species. Australia: State of the Environment Technical Report Series. Canberra, Environment Australia.

- Clunie, P.; Stuart, I.; Jones, M.; Schreiber, S.; McKay, S.; O'Connor, J.; McLaren, D.; Weiss, J.; Gunasekera, L.; Roberts, J. 2002: A risk assessment of the impacts of pest species in the riverine environment in the Murray-Darling Basin. Report produced for the Murray-Darling Basin Commission, Strategic investigations and Riverine Program, Project R2006. Victoria, Freshwater Ecology, Arthur Rylah Institute for Environmental Research.
- Coates, D.; Ulaiwi, W. K. 1995: A simple model for predicting ecological impacts of introduced aquatic organisms: a case study of common carp *Cyprinus carpio* L., in the Sepik-Ramu river basin, Papua New Guinea. *Fisheries Management and Ecology* 2: 227–242.
- Crook, D. 2001: Threatened species supplement. *Australian Society for Fish Biology Newsletter* 31(2): 37–41.
- Department of Conservation 2003: Managing invasive freshwater fish in New Zealand. Proceedings of a workshop hosted by the Department of Conservation, 10–12 May 2001, Hamilton. 174 p.
- Department of Natural Resources and Environment 2002: Healthy rivers, healthy communities and regional growth. Victorian river health strategy. Victoria, Department of Natural Resources and Environment.
- Department of Primary Industries, Queensland 2001: Control of exotic pest fishes. An operational strategy for Queensland waters 2000–2005. Queensland, Department of Primary Industries.
- Driver, P. D.; Harris, J. H.; Norris, R. H.; Closs, G. P. 1997: The role of the natural environment and human impacts in determining biomass densities of common carp in New South Wales rivers. *In*: Harris, J. H.; Gehrke, P. C. *ed.* Fish and rivers in stress. The New South Wales rivers survey. Cronulla, Canberra, NSW Office of Conservation and Cooperative Research Centre for Freshwater Ecology. Pp. 225–250.
- Gehrke, P. C.; Brown, P.; Sciller, C. B.; Mofatt, D. B.; Bruce, A. M. 1995: River regulation and fish communities in the Murray-Darling River system, Australia. *Regulated Rivers: Research and Management* 11: 363–375.
- Georges, A.; Cottingham, P. 2002: Biodiversity in inland waters—priorities for its protection and management. Recommendations from the 2001 Fenner Conference on the Environment. Cooperative Research Centre for Freshwater Ecology Technical Report 1/2002. Canberra, CRCFE.
- Gill, H. S.; Hambleton, S. J.; Morgan, D. L. 1999: Is *Gambusia holbrooki* a major threat to the native freshwater fishes of south-western Australia? *In*: Seret, B.; Sire, J.-Y. *ed.* Proceedings of the 5th Indo-Pacific Fish Conference (Noumea, 3–8 November 1997). Paris, Societe Francaise d'Ichtyologie & Institut de Recherche pour le Development. Pp. 79–87.
- Harris, J. H. 1995: The use of fish in ecological assessments. *Australian Journal of Ecology* 20: 65–80.
- Harris, J. H. 1997: Environmental rehabilitation for carp control. *In*: Roberts, J.; Tilzey, R. *ed.* Controlling carp: exploring the options for Australia. Proceedings of a workshop 22–24 October 1996, Albury. Canberra, CSIRO; Murray-Darling Basin Commission. Pp. 21–36.
- Ivanstovff, W.; Aarn, 1999: Detection of predation on Australian native fishes by *Gambusia holbrooki*. *Marine and Freshwater Research* 50: 467–468.
- Jackson, J. E.; Raadik, T. A.; Lintermans, M.; Hammer, M. 2004: Alien salmonids in Australia: impediments to effective management, and future directions. *New Zealand Journal of Marine and Freshwater Research* 38: 447–455.
- Jackson, P. D.; Koehn, J. D.; Wager, R. 1993: Appendix 1: Australia's threatened fishes 1992 listing—Australian Society for Fish Biology. *In*: Hancock, D. A. *ed.* Sustainable fisheries through sustaining fish habitats. Proceedings of the Australian Society for Fish Biology Workshop, Victor Harbour, South Australia, 12–13 August. Bureau of Resource Sciences. Canberra, AGPS. Pp. 213–227.
- Kailola, P. J. 1990: Translocated and exotic fishes: towards a cooperative role for industry. *In*: Pollard, D. A. *ed.* Introduced and translocated fishes and their ecological effects. *Australian Society for Fish Biology Workshop Proceedings No. 8*. Magnetic Island, Townsville, Queensland 24–25 August 1989. Canberra, Australian Government Publishing Service. Pp. 31–38.
- Kailola, P. J. 2000: Development of an alert list for non-native freshwater fishes. Final report to Environment Australia, Canberra.
- Koehn, J. 2001: Carp—a major vertebrate pest in Australia. *In*: Proceedings of the 12th Australian vertebrate pest conference, Melbourne, Victoria, 21–25 May 2001. Victoria, Natural Resources and Environment. Pp. 173–178.
- Koehn, J. 2002: Freshwater invaders in Victoria. *Under Control—Pest Plant and Animal News* 22 (December 2002): 12–14.
- Koehn, J. D.; Brumley, A. R.; Gehrke, P. C. 2000: Managing the impacts of carp. Canberra, Bureau of Resource Sciences.

- Koster, W. M.; Raadik, T. A.; Clunie, P. 2002a: Scoping study of the potential spread and impact of the exotic fish Oriental weatherloach in the Murray-Darling Basin, Australia: a resource document. Report to Agriculture, Fisheries, Forestry, Australia under Murray-Darling 2001 FishRehab Program. Victoria, Freshwater Ecology, Arthur Rylah Institute for Environmental Research.
- Koster, W. M.; Raadik, T. A.; Clunie, P. 2002b: Scoping study of the potential spread and impact of the exotic fish Oriental weatherloach in the Murray-Darling Basin, Australia: A draft management strategy. Report to Agriculture, Fisheries, Forestry, Australia under Murray-Darling 2001 FishRehab Program. Victoria, Freshwater Ecology, Arthur Rylah Institute for Environmental Research.
- Lintermans, M. 2004: Human-assisted dispersal of alien freshwater fish species in Australia. *New Zealand Journal of Marine and Freshwater Research* 38: 481–501.
- Lodge, D. M.; Shrader-Frechette, K. 2003: Nonindigenous species: ecological explanation, environmental ethics, and public policy. *Biological Conservation* 17(1): 31–37.
- Lodge, D. M.; Steen, R. A.; Brown, K. M.; Covich, A. P.; Bronmark, C.; Gavey, J. E.; Klosiewski, S. P. 1998: Predicting impact of freshwater exotic snails on biodiversity: challenges in spatial scaling. *Australian Journal of Ecology* 23: 53–67.
- Lowe, S. J.; Browne, M.; Boudjelas, S. 2000: 100 of the world's worst invasive alien species. Auckland, New Zealand, IUCN/SSC Invasive Species Specialist Group (ISSG).
- Mackenzie, R.; Bryant, M. 2001: Controlling exotic pest fishes in Queensland: a strategic approach. In: Proceedings of the 12th Australian vertebrate pest conference, Melbourne, Victoria, 21–25 May 2001. Victoria, Natural Resources and Environment. Pp. 178–183.
- McDowall, R. M. ed. 1980: Freshwater fishes in south-eastern Australia. Sydney, Reed.
- McDowall, R. M. ed. 1996: Freshwater fishes in south-eastern Australia. Sydney, Reed.
- McDowall, R. M. 1997: A brief history of carp in New Zealand. In: Roberts, J.; Tiltz, R. ed. Controlling carp: exploring the options for Australia. Proceedings of a workshop, 22–24 October 1996, Albury. Canberra, CSIRO & Murray-Darling Basin Commission. Pp. 1–9
- McDowall, R. M. 2004: Shoot first, and then ask questions: a look at aquarium fish imports and invasiveness in New Zealand. *New Zealand Journal of Marine and Freshwater Research* 38: 503–510.
- McKay, R. J. 1989: Exotic and translocated freshwater fishes in Australia. In: De Silva, S. S. ed. Exotic aquatic organisms in Asia. Proceedings of the workshop on introduction of aquatic organisms in Asia. *Asian Fisheries Society Special Publication* 3. Pp. 21–34.
- McKay, S.; Clunie, P.; Gillespie, G.; Raadik, T.; Saddler, S.; O'Brien, T.; Ryan, T.; Aland, G. 2001: Predation by *Gambusia holbrooki*, a review of the literature. A report to the New South Wales National Parks and Wildlife Service. Victoria, Freshwater Ecology, Arthur Rylah Institute for Environmental Research.
- McNee, A. 2002: A national approach to the management of exotic fish species in the aquarium trade: An inventory of exotic freshwater fish species. Report for Fisheries Resources Research Fund. Canberra, Department of Agriculture, Fisheries and Forestry, Australia.
- Ministerial Council on Forestry, Fisheries and Aquaculture 1999: National policy for the translocation of live aquatic organisms—issues, principles and guidelines for implementation. Canberra, Bureau of Rural Sciences.
- Morgan, D. L.; Gill, H. S. 2001: The green swordtail *Xiphophorus helleri* Heckel (Poeciliidae): another aquarium fish established in the wild in Western Australia. *Records of the Western Australian Museum* 20: 349–352.
- Morgan, D.; Gill, H.; Allen, M.; Maddern, M. 2003: Distribution and biology of fish in inland waters of the Pilbara (Indian Ocean) Drainage Division. Report to the Natural Heritage Trust, March 2003. Project Number 003026.
- Morgan, D. L.; Hambleton, S. J.; Gill, H. S.; Beatty, S. J. 2002: Distribution, biology and likely impacts of the introduced redfin perch (*Perca fluviatilis*) (Percidae) in Western Australia. *Marine and Freshwater Research* 53: 1211–1221.
- Moyle, P. B.; Light, T. 1996: Biological invasions of fresh water: empirical rules and assembly theory. *Biological Conservation* 78: 149–161.
- Murray-Darling Association Inc. 2003: Community and local government involvement in the management of alien and pest fish in the Murray-Darling Basin. An Action Plan. Albury, Murray-Darling Association.
- Nalepa, T. F.; Schloesser, D. W. ed. 1993: Zebra mussels: biology, impact and control. Michigan, Lewis, Ann Arbor.
- National Taskforce on the Prevention and Management of Marine Pest Incursions 1999: Report of the taskforce. Canberra, Joint SCC/SCFA National Taskforce on the Prevention and Management of Marine Pest Incursions.

- New South Wales Fisheries 2003: Aquatic pests in NSW. *Fishfact NSW1122*.
- NSW National Parks and Wildlife Service 2002: Predation by *Gambusia holbrooki*—the plague minnow. Draft threat abatement plan. Hurstville, NSW, NPWS.
- Office for the Commissioner for the Environment 1988: State for the environment report 1988. Victoria's inland waters. Victoria, Office for the Commissioner for the Environment.
- Olsen, P. 1998: Australia's pest animals. New solutions to old problems. Australia, Bureau of Rural Sciences and Kangaroo Press.
- Phillips, B. *comp.* 2003: Managing fish translocation and stocking in the Murray-Darling Basin. Workshop held in Canberra 25–26 September 2002: Statement, recommendations and supporting papers. Report 03/02. Australia, World Wildlife Fund.
- Pimentel, D.; Lach, L.; Zuniga, R.; Morrison, D. 2000: Environmental and economic costs on nonindigenous species in the United States. *BioScience* 50(1): 53–65.
- Pollard, D. A. 1990a: Recommendations of the workshop. In: Pollard, D. A. *ed.* Introduced and translocated fishes and their ecological effects. *Australian Society for Fish Biology Workshop Proceedings No. 8*. Magnetic Island, Townsville, Queensland 24–25 August 1989. Canberra, Australian Government Publishing Service. Pp. 180–181.
- Pollard, D. A. 1990b: *ed.* Introduced and translocated fishes and their ecological effects. *Australian Society for Fish Biology Workshop Proceedings No. 8*. Magnetic Island, Townsville, Queensland 24–25 August 1989. Canberra, Australian Government Publishing Service.
- Queensland Department of Natural Resources and Mines 2002: Queensland Pest Animal Strategy. Queensland, Department of Natural Resources and Mines.
- Raadik, T. 2001: Alien zone. Report of the exotic fishes sub-committee—to May 2001. *Australian Society for Fish Biology Newsletter* 31(1): 26–32.
- Raadik, T. 2003a: Alien zone (II). Report of the exotic fishes sub-committee—to May 2001. *Australian Society for Fish Biology Newsletter* 32(2): 60–67.
- Raadik, T. 2003b: Alien zone (III). Exotic fishes committee report—to June 2003. *Australian Society for Fish Biology Newsletter* 33(1): 45–49.
- Roberts, J.; Tilzey, R. *ed.* 1997: Controlling carp: exploring the options for Australia. Proceedings of a workshop 22–24 October 1996, Albury. Canberra, CSIRO and Murray-Darling Basin Commission.
- Rowland, S. J. 2001: Record of the banded grunter *Amniataba percoides* (Teraponidae) from the Clarence River, New South Wales. *Australian Zoologist* 31(4): 603–607.
- Sala, O. E.; Chapin, F. S. III; Armesto, J. J.; Berlow, E.; Bloomfield, J.; Dirzo, R.; Huber-Sanwald, E.; Huenneke, L. F.; Jackson, R. B.; Kinzig, A.; Leemans, R.; Lodge, D. M.; Mooney, H. A.; Oesterheld, M.; Poff, N. L.; Sykes, M. T.; Walker, B. H.; Walker, M.; Wall, D. H. 2000: Global biodiversity scenarios for the year 2010. *Science* 287: 1770–1774.
- Scientific Advisory Committee 1992: Flora and Fauna Guarantee Act final recommendation on a nomination for listing. Deliberate or accidental introduction of live fish into public or private waters within a Victorian river catchment in which the taxon to which the fish belongs cannot reliably be inferred to have been present prior to the year 1770 AD. Victoria, Department of Conservation, Forests and Lands.
- Strayer, D. L. 1999: Effects of alien species on freshwater mollusks in North America. *Journal of the North American Benthological Society* 18: 74–98.
- Stuart, I.; Jones, M. 2002: Ecology and management of common carp in the Barmah-Millewa forest. Final report of the point source management of carp project to Agriculture, Fisheries and Forestry Australia. Victoria, Arthur Rylah Institute for Environmental Research.
- Stuart, I.; Jones, M.; Koehn, J. 2001: Targeting spawning habitats to control carp populations. In: Proceedings of the 12th Australian vertebrate pest conference, Melbourne, Victoria, 21–25 May 2001. Victoria, Natural Resources and Environment. Pp. 178–183.
- Stuart, I.; McKenzie, J.; Williams, A.; Holt, T. 2003: Separation cages for removal of carp from Murray-Darling Basin fishways. Final Report on preliminary trials of the William's carp separation cage. Murray-Darling Basin Commission Project R2104. February 2003. Canberra, Murray-Darling Basin Commission.
- Sutherst, R. W.; Floyd, R. B.; Maywald, G. F. 1996: The potential geographical distribution of the cane toad, *Bufo marinus* L. in Australia. *Conservation Biology* 10: 294–299.
- Sutherst, R. W.; Maywald, G. F.; Yonow, T.; Stevens, P. M. 1999: CLIMEX: Predicting the effects of climate on plants and animals. Collingwood, CSIRO Publishing.



- The Ecology Lab. 2002: Fish fauna study—Oxleyan pygmy perch, Yelgun to Chinderah—Contract No. F/99/01. Progress Report for Sampling Times 1–6. Report to NSW Road and Traffic Authority, Pacific Highway Office by The Ecology Lab Pty Ltd, Brookvale, NSW.
- Todd, C. R.; McKay, S. F.; Conallin, A. T.; Close, P. C.; Raadik, T. A. 2002: Aquatic biota introduction from inter-basin water transfer; Murray-Darling Basin, Australia. Report to Department of Agriculture, Fisheries and Forestry, Australia. Victoria, Freshwater Ecology, Department of Natural Resources and Environment.
- Townsend, C. R. 2003: Individual, population, community and ecosystem consequences of a fish invader in New Zealand streams. *Conservation Biology* 17(1): 38–47.
- Townsend, C. R.; Winterbourn, M. J. 1992: Assessment of the environmental risk posed by an exotic fish: the proposed introduction of channel catfish (*Ictalurus punctatus*) to New Zealand. *Conservation Biology* 6: 273–282.
- Wager, R.; Unmack, P. J. 2000: Fishes of the Lake Eyre catchment of central Australia. Queensland, Department of Primary Industries.
- Waters, J. M.; Shirley, M.; Closs, G. P. 2002: Hydroelectric development and translocation of *Galaxias brevipinnis*: a cloud at the end of the tunnel? *Canadian Journal of Fisheries and Aquatic Sciences* 59(1): 49–56.
- Weatherley, A. H.; Lake, J. S. 1967: Introduced fish species in Australian inland waters. In: Weatherley, A. H. ed. Australian inland waters and their fauna. Canberra, Australian National University Press. Pp. 217–239.
- Webb, A. C. 1994: Ecological aspects of the Mozambique mouthbrooder, *Oreochromis mossambicus*, and other introduced cichlids in northern Queensland. Unpublished MSc thesis, Faculty of Biological Sciences, James Cook University, Townsville.
- Whittington, J.; Coysh, J.; Davies, P.; Dyer, F.; Gawne, B.; Lawrence, I.; Liston, P.; Norris, R.; Robinson, W.; Thoms, M. 2001: Development of a framework for the sustainable river audit. Draft Final Report Project R2004. Canberra, Cooperative Research Centre for Freshwater Ecology.
- Williamson, M. H.; Fitter, A. 1996: The characteristics of successful invaders. *Biological Conservation* 78: 163–170.