

## Microbiome enrichment from contaminated marine sediments unveils novel bacterial strains for petroleum hydrocarbon and heavy metal bioremediation

Dell'Anno, Filippo; van Zyl, Leonardo Joaquim; Trindade, Marla; Buschi, Emanuela; Cannavacciuolo, Antonio; Pepi, Milva; Sansone, Clementia; Brunet, Christophe; Ianora, Adrianna; de Pascale, Donatella; Golyshin, Peter; Dell'Anno, Antonio; Rastelli, Eugene

### Environmental Pollution

DOI:  
[10.1016/j.envpol.2022.120772](https://doi.org/10.1016/j.envpol.2022.120772)

Published: 15/01/2023

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

*Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):*  
Dell'Anno, F., van Zyl, L. J., Trindade, M., Buschi, E., Cannavacciuolo, A., Pepi, M., Sansone, C., Brunet, C., Ianora, A., de Pascale, D., Golyshin, P., Dell'Anno, A., & Rastelli, E. (2023). Microbiome enrichment from contaminated marine sediments unveils novel bacterial strains for petroleum hydrocarbon and heavy metal bioremediation. *Environmental Pollution*, 317, Article 120772. Advance online publication. <https://doi.org/10.1016/j.envpol.2022.120772>

#### Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

1 **Microbiome enrichment from contaminated marine sediments unveils novel**  
2 **bacterial strains for petroleum hydrocarbon and heavy metal bioremediation**

3 Filippo Dell'Anno<sup>1§</sup>, Leonardo Joaquim van Zyl<sup>2</sup>, Marla Trindade<sup>2</sup>, Emanuela Buschi<sup>3</sup>, Antonio  
4 Cannavacciuolo<sup>4</sup>, Milva Pepi<sup>4</sup>, Clementina Sansone<sup>1</sup>, Christophe Brunet<sup>1</sup>, Adrianna Ianora<sup>1</sup>,  
5 Donatella de Pascale<sup>1</sup>, Peter N. Golyshin<sup>5</sup>, Antonio Dell'Anno<sup>6§</sup>, Eugenio Rastelli<sup>3\*§</sup>

6 <sup>1</sup>*Department of Marine Biotechnology, Stazione Zoologica "Anton Dohrn", Villa Comunale, 80121 Naples,*  
7 *Italy; [filippo.dellanno@szn.it](mailto:filippo.dellanno@szn.it) (FD), [clementina.sansone@szn.it](mailto:clementina.sansone@szn.it) (CS), [brunet@szn.it](mailto:brunet@szn.it) (CB), [ianora@szn.it](mailto:ianora@szn.it)*  
8 *(A.I.), [donatella.depascale@szn.it](mailto:donatella.depascale@szn.it) (DdP);*

9 <sup>2</sup>*Department of Biotechnology, Institute for Microbial Biotechnology and Metagenomics, University of the*  
10 *Western Cape, Bellville 7535, Cape Town, South Africa; [lvanzyl@uwc.ac.za](mailto:lvanzyl@uwc.ac.za) (LJZ), [ituffin@uwc.ac.za](mailto:ituffin@uwc.ac.za) (MT);*

11 <sup>3</sup>*Department of Marine Biotechnology, Stazione Zoologica "Anton Dohrn", Fano Marine Centre, Viale*  
12 *Adriatico I-N, 61032 Fano, Italy; [emanuela.buschi@szn.it](mailto:emanuela.buschi@szn.it) (EB), [eugenio.rastelli@szn.it](mailto:eugenio.rastelli@szn.it) (ER);*

13 <sup>4</sup>*Department of Integrative Marine Ecology, Stazione Zoologica "Anton Dohrn", Fano Marine Centre, Viale*  
14 *Adriatico I-N, 61032 Fano, Italy; [antonio.cannavacciuolo@szn.it](mailto:antonio.cannavacciuolo@szn.it) (AC), [milva.pepi@szn.it](mailto:milva.pepi@szn.it) (MP);*

15 <sup>5</sup>*Centre for Environmental Biotechnology, School of Natural Sciences, Bangor University, Gwynedd LL57*  
16 *2UW, UK; [p.golyshin@bangor.ac.uk](mailto:p.golyshin@bangor.ac.uk) (PNG);*

17 <sup>6</sup>*Department of Life and Environmental Sciences, Università Politecnica delle Marche, Via Brecce Bianche,*  
18 *60131 Ancona, Italy; [a.dellanno@univpm.it](mailto:a.dellanno@univpm.it) (AD).*

19


20 \*Correspondence: [eugenio.rastelli@szn.it](mailto:eugenio.rastelli@szn.it) (ER).

21 § These authors contributed equally.



Sampling of contaminated marine sediments @Sarno River Mouth

Microbiome enrichment




- **Halomonas (26.7%)**
- **Alcanivorax (23.7%)**
- Chromohalobacter (7.3%)
- Alkaliphilus (5.7%)
- Marinobacter (5.6%)
- Pseudomonas (3.9%)
- Ruegeria (3.8%)
- Pseudoalteromonas (2.8%)
- Burkholderia (1.5%)
- Marinomonas (1.1%)
- Others (<1%)


Bacterial isolation



Bioremediation experiments



Genomic basis of bioremediation and screening for novel useful enzymes



## 22 **Abstract**

23 Petroleum hydrocarbons and heavy metals are some of the most widespread contaminants affecting  
24 marine ecosystems, urgently needing effective and sustainable remediation solutions. Microbial-  
25 based bioremediation is gaining increasing interest as an effective, economically and environmentally  
26 sustainable strategy. Here, we hypothesized that the heavily polluted coastal area facing the Sarno  
27 River mouth, which discharges >3 tons of polycyclic aromatic hydrocarbons (PAHs) and ~15 tons of  
28 heavy metals (HMs) into the sea annually, hosts unique microbiomes including marine bacteria useful  
29 for PAHs and HMs bioremediation. We thus enriched the microbiome of marine sediments,  
30 contextually selecting for HM-resistant bacteria. The enriched mixed bacterial culture was subjected  
31 to whole-DNA sequencing, metagenome-assembled-genomes (MAGs) annotation, and further sub-  
32 culturing to obtain the major bacterial species as pure strains. We obtained two novel isolates  
33 corresponding to the two most abundant MAGs (*Alcanivorax xenomutans* strain-SRM1 and  
34 *Halomonas alkaliantarctica* strain-SRM2), and tested their ability to degrade PAHs and remove  
35 HMs. Both strains exhibited high PAHs degradation (60-100%) and HMs removal (21-100%) yield,  
36 and we described in detail >60 genes in their MAGs to unveil the possible genetic basis for such  
37 abilities. Most promising yields (~100%) were obtained towards naphthalene, pyrene and lead. We  
38 propose these novel bacterial strains and related genetic repertoire to be further exploited for effective  
39 bioremediation of marine environments contaminated with both PAHs and HMs.

40 **Keywords:** polycyclic aromatic hydrocarbons; heavy metals; bioremediation; next-generation  
41 sequencing; marine biotechnology.

## 42 **1. Introduction**

43 Pollution of coastal environments due to organic (e.g., polycyclic aromatic hydrocarbons, PAHs)  
44 and/or inorganic (e.g., heavy metals, HMs) contaminants can determine major detrimental effects on  
45 the marine food web and human health (Tashla et al., 2018; Fuentes-Gandara et al., 2018; Loflen et  
46 al., 2018; Buah-Kwofie et al., 2018). PAHs, HMs and metalloids are known to strongly affect  
47 biological systems such as cell membranes, organelles and enzymes, causing cell cycle alteration,  
48 carcinogenesis or apoptosis (Tchounwou et al., 2001; Sutton et al., 2002; Yedjou and Tchounwou,  
49 2007a,b; Beyersmann and Hartwig, 2008; Patlolla et al., 2009; Kim et al., 2015; Costa et al., 2022).  
50 Such contaminants can persist in the environment for a long time and can be subjected to  
51 bioaccumulation and/or biomagnification processes, which increase their potential to cause harm  
52 (Oyetibo et al., 2017). Conventional methods for the removal of PAHs and HMs include chemical  
53 treatments (e.g., precipitation, oxidation and reduction), electrochemical techniques or physical  
54 adsorption (Fenyvesi et al., 2019). Unfortunately, such methods typically produce special wastes  
55 (e.g., toxic sludge and by-products) and are expensive, ineffective at low pollutant concentrations and  
56 highly energy-demanding (Joshi, 2017; Priyadarshane and Das, 2021). The use of microorganisms  
57 (especially, bacteria, fungi and microalgae) may be a promising alternative or complementary  
58 strategy to such conventional tools, due to several microbial characteristics including high  
59 biodegradation/detoxification efficiency towards several contaminants, high surface area-to-volume  
60 ratio and the ability to grow at high concentrations of toxic pollutants (Zouboulis et al., 2004;  
61 Kordialik-Bogacka and Diowksz, 2014). Microbial-based remediation strategies are also considered  
62 to be one of the most sustainable approaches due to low carbon footprint of the overall  
63 decontamination process (Kuppusamy et al., 2017; Dell'Anno et al., 2020; Jain et al., 2022).  
64 Microbes effectively degrade PAHs by several mechanisms including assimilation, intracellular  
65 detoxification, and/or co-metabolism mediated by the activity of specific enzymes such as oxygenase,  
66 dehydrogenase and ligninolytic enzymes (Johnsen et al., 2005; Ladino-Orjuela et al., 2016; Saravanan  
67 et al., 2021). Microbial-mediated HMs' sorption, leaching or transformation are well-known as

68 inexpensive and highly efficient HM bioremediation processes (Leung et al., 2000; Aryal et al., 2010;  
69 Sahmoune, 2018; Quiton et al., 2018; Cai et al., 2019). Co-contamination by PAHs and HMs is  
70 typically encountered in most marine polluted sites (El-Mufleh et al., 2014; Ali et al., 2021) and  
71 several microbes can exert a combined action towards such contaminants by bioleaching, biosorption  
72 and biodegradation, which increase their overall removal and/or detoxification capacity (Liu et al.,  
73 2017; Ali et al., 2021). Examples of this synergistic approach include microbial secretion - under  
74 mixed PAHs and HMs contamination - of enzymes with useful bioremediation activities (Chen et al.,  
75 2020) and of extracellular polysaccharides (EPSs). EPSs are particularly relevant targets for  
76 bioremediation studies, as they consist of a wide array of polymers and functional groups, that can  
77 simultaneously bind metals (thus enhancing HM extraction/removal from contaminated matrices)  
78 (Amoozegar et al., 2012; Ates, 2015; Little et al., 2014; Mohite et al., 2017; Gupta and Diwan, 2017;  
79 Cao et al., 2022; Cheng et al., 2022), and increase PAH solubilization and enzymatic degradation  
80 efficiency (Gutierrez et al., 2013; Alaba et al., 2018).

81 Environmental matrices that are naturally or experimentally enriched with organic and/or inorganic  
82 contaminants are an underexploited source of novel microbes resistant to contaminants and involved  
83 in their biodegradation/detoxification (Beolchini et al., 2009; Adams et al., 2015; Fodelianakis et al.,  
84 2015; Dell'Anno et al., 2021; Wang et al., 2021). Currently, high-throughput sequencing technologies  
85 help to uncover such microbial bioremediation potential, providing novel insights on the diversity of  
86 useful environmental microbes and their repertoire of genes involved in PAH and HM bioremediation  
87 (Czaplicki and Gunsch, 2016; Dell'Anno et al., 2021; Meng et al., 2022; Hassan et al., 2022; Sharma  
88 et al., 2022).

89 In this study, we selected the highly anthropically-impacted and severely contaminated coastal area  
90 at the mouth of the Sarno River (Gulf of Naples, Mediterranean Sea), one the most polluted rivers in  
91 Europe discharging large amounts of PAHs and HMs into the sea (Montuori and Triassi, 2012;  
92 Montuori et al., 2013). Based on such high inputs of contaminants at this site, we postulated that here,  
93 marine sediments could host microbiomes enriched in bacterial taxa that typically characterize marine

94 areas affected by oil-spills and/or industrial activities, and possibly useful for PAHs and HMs  
95 bioremediation. We collected marine sediments from one of the most polluted sites in this area, and  
96 we adopted a laboratory approach based on the enrichment of the sediment microbiome, subsequent  
97 selection and culturing of bacteria resistant to HMs, and next-generation sequencing coupled with  
98 laboratory experiments to unveil their potential for PAHs and HMs bioremediation.

99

## 100 **2. Materials and Methods**

### 101 **2.1. Sediment sampling, microbiome enrichment and selection of bacteria useful for PAH and** 102 **HM bioremediation**

103 Sampling was performed at the mouth of the Sarno River (Fig. 1). Surface sediments were collected  
104 by a Van Veen grab, placed into sterile Whirl-Pak bags (Nasco), and then stored at 4°C in the dark  
105 until processing. One gram of sediment was added to a 1000 ml flask containing 200 ml of Marine  
106 Broth (Difco, Marine Broth 2216). Inoculated flasks were mixed and incubated at 28°C in the dark.  
107 After 2 weeks of incubation, the enriched microbiome was plated by streaking onto marine agar  
108 (Difco, Marine Agar) added with a mix of  $Pb^{2+}$  (500 ppm),  $As^{3+}$  (500 ppm), and  $Cd^{2+}$  (10 ppm) and  
109 incubated at 28°C for 48 hours. Above HMs were selected as the most relevant in the study area, and  
110 the applied concentrations were significantly higher than those determined in the sediments  
111 (Montuori et al., 2013), to isolate HM-resistant bacteria with possible PAH and HM bioremediation  
112 ability. PAHs were not added at this stage, as we hypothesized that bacterial PAH degraders were  
113 already abundant in the original contaminated sediments, and our rationale was to select those able  
114 to also tolerate high HM concentrations. As the diversity of potential PAH-degrading bacteria is  
115 potentially high in contaminated sediments, we acknowledge that alternative enrichment strategies  
116 (e.g., contextual addition of PAHs at this stage) may have led to different/additional bacterial isolates  
117 than those obtained in our study.

118 Following incubation, high microbial growth was observed, which was confirmed to be a multi-  
119 species bacterial culture by preliminary Sanger sequencing of the 16S rRNA genes, hence analyzed  
120 by whole DNA shotgun sequencing, as described below.

121

## 122 **2.2. Whole DNA shotgun sequencing and analysis of bacterial metagenome-assembled genomes** 123 **(MAGs)**

124 The total genomic DNA (gDNA) obtained from the enriched mixed culture was extracted with the  
125 DNeasy Blood & Tissue kit, according to the manufacturer's instructions. The DNA concentration  
126 was determined using the Qubit™ dsDNA HS assay kit with a Qubit fluorometer (Thermo Fisher,  
127 Waltan, US). Sequence library preparation of gDNA was performed using the Nextera DNA Flex kit  
128 (Illumina, Hayward, USA) with 1 ng input DNA according to the manufacturer's instructions. The  
129 resultant libraries were sequenced on an Illumina MiSeq instrument using a MiSeq Reagent kit V2  
130 (500 cycles) with a 10% phiX v3 spike, generating  $2 \times 250$  bp reads. Preliminary metagenome  
131 processing and taxonomic and functional annotation were performed in MG-RAST under default  
132 settings (Meyer et al., 2008). Read assembly was performed using CLC Genomics Workbench  
133 version 11. Briefly, the raw reads were trimmed and demultiplexed, and contigs  $\leq 500$  bp were  
134 removed from the final assembly. Binning of metagenomic contigs was performed using MyCC (Lin  
135 and Liao, 2016) while completeness and contamination of MAGs and genome quality were  
136 determined using CheckM with the lineage-specific workflow and default parameters (Parks et al.,  
137 2015).

138

## 139 **2.3. Isolation and identification of single bacterial strains**

140 To obtain single colonies of pure bacterial strains, the enriched mixed culture was re-plated on marine  
141 agar (Difco, Marine Agar) added with HMs as described above, through serial dilution. After several  
142 re-streaking cycles of 48-hours incubation at 28°C, two main colony morphologies were observed,  
143 and each re-streaked until confirmed to be pure by PCR analysis and Sanger sequencing targeting



144 16S rRNA gene (by universal bacterial primers E9F-5'-GAGTTTGATCCTGGCTCAG-3' and  
145 U1510R-5'-GGTTACCTTGTTACGACTT-3'; Rodriguez-Caballero et al., 2012). All polymerase  
146 chain reactions (PCR) were carried out in a Perkin Elmer Thermocycler (Gene Amp PCR system  
147 6700) in a 50 µl reaction volume containing 1× PCR buffer, 200 µM of each dNTP, 0.5 µM of each  
148 primer, 0.2 U of Taq Gold polymerase (Applied Biosystems, Waltham, MA, US) and 1 ng of template  
149 DNA. Thermal cycling conditions were 5 min denaturation at 94 °C; 30 cycles of 94 °C for 30 s, 55  
150 °C for 30 s and 72°C for 90 s; final elongation step at 72°C for 5 min. The PCR products were  
151 analyzed by agarose gel (1.2% w/v) electrophoresis in TAE buffer solution (40 mM Tris-acetate, 1  
152 mM ethylenediaminetetraacetic acid, EDTA) containing 0.5 µg ml<sup>-1</sup> (w/v) ethidium bromide. The  
153 amplicons were purified and sequenced using an ABI PRISM 377 automated sequencer (Applied  
154 Biosystems). The sequencing data were processed using Chromas Pro v. 1.5a software  
155 (Technelysium, South Brisbane, QLD, Australia) for alignment and manual editing of sequences. The  
156 consensus sequences of the isolates were compared with those deposited in GenBank using BLAST.  
157 The two bacterial strains were matched with the two corresponding MAGs obtained by whole-DNA  
158 shotgun sequencing by comparing their 16S rRNA gene sequences following nucleotide alignment  
159 performed through a local blast in the annotation system. The pure bacterial strains were then tested  
160 in experiments to assess their bioremediation ability to degrade PAHs and remove HMs, as described  
161 below.

162

#### 163 **2.4. Laboratory tests to assess the PAH degradation and HM removal ability of the bacterial** 164 **isolates**

165 Each bacterial isolate was incubated in flasks (T175, TPP tissue culture flasks, final volume 250 ml  
166 Marine Broth; starting inoculum of 8 x 10<sup>7</sup> cells ml<sup>-1</sup>), and subjected to two treatments. These  
167 included: i) addition of naphthalene, pyrene and phenanthrene (ratio of 1:1:1 with a total  
168 concentration of 242 ppm); ii) addition of arsenic (As<sup>3+</sup>; 14 ppm), lead (Pb<sup>2+</sup>; 331 ppm), cadmium  
169 (Cd<sup>2+</sup>; 1 ppm), copper (Cu<sup>2+</sup>; 74 ppm), and zinc (Zn<sup>2+</sup>; 899 ppm). Controls were included for each of

170 the two treatments, following the same procedure but without bacterial inoculum. All experimental  
171 microcosms were set up in triplicate. The concentration of the toxic compounds used for  
172 bioremediation experiments was selected based on the average values found in the surrounding  
173 marine area (Montuori and Triassi, 2012; Montuori et al., 2013). The flasks were incubated for 27  
174 days at 28°C, and bacterial growth was monitored by OD<sub>600</sub> at days 0, 9 and 27. The quantification  
175 of PAHs and HMs was conducted, respectively, by gas chromatography-mass spectrometry (GC-MS;  
176 EPA8270) (Casillo et al., 2018), and by inductively coupled plasma atomic emission spectroscopy  
177 (ICP-OES; EPA6010) (EPA, 2014) on aliquots from each experimental treatment. The PAHs  
178 degradation yield was calculated for each experimental treatment by comparing the concentrations of  
179 each contaminant at the beginning and at the end of the incubations. For HMs, aliquots of each  
180 experimental treatment at the end of incubations were first centrifuged (1000 x g, 5 min), and the HM  
181 concentration in the pellet was used to calculate the % of HMs removal, as the % of HM mass in the  
182 pellet compared to the HM mass added at the beginning of incubations.

183

## 184 **2.5. MAGs functional annotation and comparative genomics**

185 The genome taxonomy database (GTDB) (<https://gtdb.ecogenomic.org/>) implemented through K-  
186 Base ([www.kbase.us](http://www.kbase.us)) was used to perform the whole-genome based classification of the two MAGs  
187 obtained following whole-DNA shotgun sequencing and corresponding to the two bacterial strains  
188 used in the PAHs degradation and HM removal experiments in this study. Species relatedness was  
189 evaluated through the average nucleotide identity (ANI) analysis (Rodriguez and Konstantinidis,  
190 2016; Han et al., 2016) by comparing the de novo genomes with the genomes selected following the  
191 GTDB output. The obtained genomes were annotated by RAST (Overbeek et al., 2014) providing an  
192 automated functional annotation or hypothetical protein annotation for each open reading frame  
193 (ORF) identified on the genome. KEGG was used for metabolic prediction (Kanehisa et al., 2017).  
194 Following automated identification of genes involved in hydrocarbon degradation or metal  
195 resistance/detoxification/removal, manual verification of the annotated ORFs was conducted against

196 the SwissProt database. In addition, the sequences flanking the genes of interest were manually  
197 annotated to better understand their genomic context, accurately delineate the regions involved in  
198 these functions and for synteny comparison, and visualized using Easyfig (Sullivan et al., 2011). The  
199 superimposition analysis of dioxygenases belonging to the two MAGs have been performed using  
200 the web portal for protein modelling, prediction and analysis Phyre2 (Kelley et al., 2015), and enzyme  
201 similarity was assessed according to Zhang and Skolnick, 2004.

202

## 203 **2.6. Statistical analyses**

204 To test for differences in the experimental results, Student T and Fisher-Snedecor tests were carried  
205 out using PAST3 software (Hammer et al., 2001).

206

## 207 **3. Results and Discussion**

### 208 **3.1 Characterization of the enriched microbiome and of the bacterial isolates tested for PAHs** 209 **and HMs bioremediation**

210 The shotgun sequencing of the total DNA, extracted from the selectively enriched mixed bacterial  
211 culture obtained in this study from the contaminated marine sediments of the Sarno River mouth,  
212 resulted in >3.1 million high-quality reads (average sequence length 201±59 bp; Supplementary Table  
213 S1). Taxonomic annotation of this enriched metagenome showed a dominance of two  
214 Oceanospirillales (Gammaproteobacteria) genera: *Alcanivorax* (~24% of total reads) and *Halomonas*  
215 (~27% of total reads) (Fig. 2A). Several other bacterial taxa were detected in the metagenome, though  
216 at a much lower relative abundance. These included other Gammaproteobacteria (*Chromohalobacter*,  
217 *Marinobacter*, *Pseudomonas*, *Pseudoalteromonas* and *Marinomonas*), Clostridia (*Alkaliphilus*),  
218 Alphaproteobacteria (*Ruegeria*), and Betaproteobacteria (*Burkholderia*) (Fig. 2A). All of these  
219 bacterial taxa have previously been reported to typically increase rapidly in abundance during oil  
220 spills, to degrade hydrocarbons and to tolerate/detoxify heavy metals (Gutierrez et al., 2013;  
221 Dubinsky et al., 2013; Kumar et al., 2019; Dell'Anno et al., 2021; Huo et al., 2014; Liu et al., 2019;

222 Ramasamy et al., 2020; Ghosh et al., 2022). This suggests that our microbiome enrichment strategy  
223 was successful in selectively boosting autochthonous bacterial taxa with promising potential for  
224 petroleum hydrocarbon and HM remediation. This was also supported by the preliminary functional  
225 annotation of the metagenome (Fig. 2B), which highlighted that the functions putatively related to  
226 hydrocarbon degradation and to resistance/interaction with heavy metals represented an important  
227 portion of the overall reads count (>6%). The major role of such functional features was further  
228 suggested by the fact that their representation was quantitatively similar compared to fundamental  
229 cell processes such as cell respiration, DNA metabolism or membrane transport (Fig. 2B). It should  
230 be noted that our enrichment approach, resulting in the virtual absence of Archaea in the enriched  
231 metagenome, might have overlooked possible syntrophic relationships among bacteria and archaea  
232 in the original sediments, whose relevance in petroleum hydrocarbons degradation has been  
233 highlighted by recent independent studies (Liu et al., 2018, 2021; Harindintwali et al., 2022).

234 The reads assembly and MAGs reconstruction and annotation resulted in two dominant MAGs with  
235 genome completeness between 99-100%, which were classified as *Alcanivorax xenomutans* and  
236 *Halomonas alkaliantarctica* based on GTDB-Tk whole-genome based classification, and supported  
237 by 16S rRNA gene sequence identities (Supplementary Tables S2-S4). The ANI analysis indicated  
238 *Alcanivorax xenomutans* strain KS-293 (Barbato et al., 2015; ANI score 99.10%) and *Halomonas*  
239 *alkaliantarctica* strain CRSS (Poli et al., 2007; ANI score 97.13%) as their respective closest  
240 relatives. Two additional partial MAGs (4-32% completeness) were recovered from the metagenome  
241 and classified in the genera *Pseudoalteromonas* and *Alkalphilus*, but these were not analysed further.

242 The completeness and coverage of the four reconstructed MAGs reflected the relative contribution of  
243 their reads to the overall sequence count (Fig. 2A), further suggesting that the obtained enriched  
244 mixed bacterial culture was dominated by *A. xenomutans* and *H. alkaliantarctica*, with minor  
245 contribution by other bacterial taxa.

246 The sequential and selective sub-culturing from the enriched mixed bacterial culture allowed us to  
247 obtain the two strains that matched the two full-reconstructed MAGs based on 16S rRNA gene

248 sequence identity (100%). We hence refer to the two novel strains and related genomes obtained in  
249 this study as *A. xenomutans* strain SRM1 and *H. alkaliantarctica* strain SRM2 (Supplementary Fig.  
250 S1, S2).

251 The laboratory tests conducted to assess their ability for PAHs and HMs remediation showed high  
252 PAHs degradation (ranging for both strains from 60% for phenanthrene to 100% for both naphthalene  
253 and pyrene; Fig. 3A), as well as high HMs removal yields (34-91% for As, 79-94% for Cd, 21-70%  
254 for Cu, 50-89% for Zn, and 94-100% for Pb), with highest values observed with *H. alkaliantarctica*  
255 strain SRM2 for As, Cu, and Zn; Fig. 3B). Notably, we observed that the culturing of *A. xenomutans*  
256 strain SRM1 and *H. alkaliantarctica* strain SRM2 with addition of PAHs resulted in growth rates  
257 almost double to those of control conditions (Supplementary Fig. S3), which agrees with previous  
258 independent evidence that these bacterial genera include taxa able to exploit hydrocarbons to produce  
259 cell biomass (Mnif et al., 2009; Rahul et al., 2014). Both strains exhibited resistance to the high  
260 concentrations of mixed HMs tested in our study (>1300 ppm, considering the sum of As, Cd, Cu,  
261 Zn, and Pb) (Fig. 3B, Supplementary Fig. S3), indicating that these strains can be particularly useful  
262 for PAH biodegradation of marine matrices that are simultaneously highly polluted with different  
263 HMs. Indeed, the use of bacterial strains able to both degrade organic contaminants and tolerate toxic  
264 inorganic compounds present in the target matrix can help to reduce failure risk in bioremediation of  
265 environments that display high loads of mixed toxic contaminants (Thompson et al., 2005; Nwuche  
266 and Ugoji, 2008; Alisi et al., 2009; Tyagi et al., 2011; Dueholm et al., 2015).

267 Overall, these results confirm several previous reports showing that *Halomonas* spp. and *Alcanivorax*  
268 spp. include members that display high resistance towards toxic organic and inorganic contaminants  
269 (Rahul et al., 2014; Fu et al., 2018, Catania et al., 2018; Dell'Anno et al., 2020) and can degrade  
270 PAHs (Budiyanto et al., 2018; Kadri et al., 2018).

271 The observed higher degradation rates of naphthalene and pyrene than phenanthrene (Fig. 3A) may  
272 be counterintuitive, as it is generally assumed that bacterial biodegradation of low-molecular-weight  
273 PAHs (such as naphthalene and phenanthrene, with  $\leq 3$  aromatic rings) occurs faster than for high-

274 molecular-weight PAHs (with  $\geq 4$  aromatic rings, like pyrene). Despite previous investigations  
275 support the expected pattern of a higher PAH recalcitrance with increasing molecular weight (Yu et  
276 al., 2005; Haritash and Kaushik, 2009; Thavamani et al., 2012; Nzila et al., 2021), other studies have  
277 reported the opposite trend, similar to that observed in the present study (Sohn et al., 2004;  
278 Wongwongsee et al., 2013; Vaidya et al., 2017, 2018). This can be explained by different enzymatic  
279 and metabolic pathways among microbes, by differences in the specific chemical-physical conditions  
280 applied in the laboratory (which can differentially influence the degradability of different PAHs;  
281 Leahy and Colwell, 1990; Bagby et al., 2017), as well as by possible interactions among different  
282 PAHs, such as inhibition or enhancement of the biodegradation of specific high-molecular-weight  
283 PAHs in the presence of specific low-molecular-weight PAHs (Guha et al., 1999; Vaidya et al., 2018).  
284 As PAHs-contaminated sites typically display mixtures of high- and low-molecular-weight PAHs  
285 (Bezza and Chirwa, 2017), we can conclude that our two bacterial strains, able to enhance  
286 biodegradation rates of high-molecular-weight PAHs in the presence of low-molecular-weight PAHs,  
287 can represent a significant advantage for environmental bioremediation applications.

288 Regarding the ability of the two tested strains to tolerate and remove HMs from contaminated  
289 matrices, our results confirm previous findings obtained using other *Halomonas* and *Alcanivorax*  
290 species. *Halomonas elongate*, *Halomonas halophila* and others displayed 50-94% removal yields  
291 towards Pb, Cd and/or Cr (Amoozegar et al., 2012; Murugavelh and Mohanty, 2012; Asksonthong et  
292 al., 2018; Abdel-Razik et al., 2020), while Pb removal ability has already been documented for some  
293 *Alcanivorax* sp. (da Costa Waite et al., 2016; Dell'Anno et al., 2020; Ramasamy et al., 2020).  
294 Notwithstanding, the current knowledge on the promising application of *Halomonas* sp. and  
295 *Alcanivorax* sp. for the bioremediation of HMs is still in its infancy and should be investigated further,  
296 also including tests for larger sets of HMs (Pennafirme et al., 2015; Verma and Kuila, 2019; Cecchi  
297 et al., 2021; Dell'Anno et al., 2020).

298

### 299 **3.2. Genetic basis for PAHs biodegradation**

300 Understanding the genetic basis and metabolic processes involved in microbial removal of petroleum  
301 hydrocarbons is fundamental to optimize bioremediation strategies, enabling tailored amendments to  
302 favor specific bacterial metabolism, genetic bioengineering, or discovery of useful  
303 enzymes/compounds (Schneiker et al., 2006; Dell'Anno et al., 2021; Sharma et al., 2022). Our results  
304 from high throughput sequencing allowed us to identify the genes and metabolic pathways potentially  
305 involved in the bioremediation ability of the tested *A. xenomutans* strain SRM1 and *H.*  
306 *alkaliantarctica* strain SRM2. In particular, several genes involved in the metabolism of aromatic  
307 compounds were identified through the automatic functional annotation of their MAGs  
308 (Supplementary Fig. S4). Even though the two strains possessed a different number of enzymes  
309 involved in these metabolic pathways (36 in *A. xenomutans* strain SRM1, 61 in *H. alkaliantarctica*  
310 strain SRM2; Supplementary Fig. S4), they performed similarly in the laboratory tests for PAHs  
311 degradation (Fig. 3A,B). We cannot exclude that additional genes for enzymes known to be involved  
312 in petroleum hydrocarbon degradation and apparently missing in the two MAGs were not identified  
313 through automatic annotation, due to divergence of DNA sequences from those available in current  
314 public databases. To gain a more complete picture of the main pathways for hydrocarbon degradation  
315 in the two novel strains, we manually checked the organization of the genomic regions that contain  
316 the genes of major interest, including those for hydrocarbon degradation peripheral pathways (*cis*-  
317 hydroxylation and *trans*-hydroxylation pathways) and central degradation routes (catechuate,  
318 protocatechuate, homoprotocatechuate, homogentisate and phenylacetic pathways), as detailed  
319 below.

### 320 *3.2.1 Peripheral pathways for hydrocarbon degradation*

321 The *cis*-hydroxylation pathway usually starts following the activity of ring hydroxylating  
322 dioxygenase enzymes (RHDs) (Peng et al., 2019), which have large ( $\alpha$ ) and small ( $\beta$ ) subunits  
323 (Kauppi et al., 1998). The alpha subunit (RHD $\alpha$ ) contains two conserved regions (the [Fe<sub>2</sub>-S<sub>2</sub>] Rieske  
324 center and the mononuclear iron-containing catalytic domain), which promote the incorporation of  
325 molecular oxygen into the aromatic ring forming a *cis*-dihydrodiol. We found RHDs in both *A.*

326 *xenomutans* strain SRM1 and *H. alkaliantarctica* strain SRM2 MAGs (Fig. 4A-D). The sequence of  
327 *A. xenomutans* strain SRM1 showed ~100% identity with that of *A. xenomutans* P40 (Fu et al. 2018)  
328 (Fig. 4A). Comparison with *A. dieselolei* B5 (Lai et al., 2012) highlighted a lower correlation but still  
329 high overall similarity (homology of  $\geq 70\%$ ) with 11 genes (including the RHD gene) (Fig. 4A).  
330 Other proteins involved in hydrocarbon detoxification and degradation processes were found in *A.*  
331 *xenomutans* strain SRM1 when analyzing the flanking region of RHD, including glutathione s-  
332 transferase, linear amide C-N hydrolase, aldo-keto reductase and nitrite reductase (Lloyd-Jones and  
333 Lau, 1997; Al-Turki, 2009; Cao et al., 2015; Imperato et al., 2019; Salam and Ishaq, 2019). The  
334 comparison of the sequences flanking the RHD in *H. alkaliantarctica* strain SRM2 MAG, with those  
335 of close relatives (Fig. 4B) highlighted that in all genomes the genes encoding the RHD and  
336 ferredoxin (both directly involved in the degradation of hydrocarbons), as well as serin  
337 hydroxymethyl transferase, sarcosine oxidase  $\alpha,\beta,\gamma$ -subunits and formyltetrahydrofolate  
338 deformylase, were all located in the same genomic region. Such genomic organization supports  
339 previous evidence that genes associated with glycine and serine metabolism are involved in  
340 hydrocarbon degradation (Yan and Wu, 2017). The 3D superimposition analysis revealed a high  
341 similarity for both RHDs with naphthalene 1,2 dioxygenases (Fig. 4C,D), which are enzymes  
342 involved in the first ring hydroxylation of multiple PAHs, including naphthalene, phenanthrene,  
343 anthracene, dibenzothiophene and fluorene (Park and Crowley, 2006). Although it will be necessary  
344 to confirm the specific function of the RHDs identified, the results of 3D modeling suggest a similar  
345 ability in degrading PAHs (based on TM scores of 0.89 and 0.73 respectively for *A. xenomutans* strain  
346 SRM1 and *H. alkaliantarctica* strain SRM2). Surprisingly, such high similarity between the two  
347 enzymes' structures corresponded to only 30% identity in their secondary sequences. This further  
348 suggests that the lack of matches of our MAGs DNA sequences for some of the other major genes  
349 involved in hydrocarbon degradation may actually be due to evolutionary genetic divergence from  
350 currently known bacterial genomes deposited in public databases.



351 An alternative to the *cis*-hydroxylation pathway is represented by the *trans*-hydroxylation pathway,  
352 in which the cytochrome P450 system (CYP450) catalyzes a *trans*-dihydrodiols formation by the  
353 epoxidation of the aromatic ring by epoxide hydrolase (Moody et al., 2005). For the *A. xenomutans*  
354 strain SRM1 MAG, we found highly conserved homologs of the CYP450 within the 3 closest  
355 relatives retrieved from NCBI (100% identity with *Alcanivorax xenomutans* sp 40, and >70% identity  
356 with *Alcanivorax* sp N3-2A and *Alcanivorax dieselolei* B5) (Fig. 5A). Within the same genome region  
357 containing CYP450, we also found two genes involved in the detoxification and biodegradation of  
358 xenobiotics: glutathione-disulfide reductase (Moron et al., 1979) and a Rieske domain non-heme  
359 oxygenase (Barry and Challis, 2013). For *H. alkaliantarctica* strain SRM2, the BLASTp analysis of  
360 the CYP450 sequence found within its MAG showed no homologs within the 3 most similar reference  
361 sequences retrieved from NCBI (of *H. axialiensis* Althf1, *H. olivaria* TYRC17 and *H. aestuari* Hb3),  
362 despite the flanking region included other conserved genes (e.g., cytochrome C, nitrogen metabolism  
363 and membrane transporters genes) (Fig. 5B). As the abovementioned three closest relatives were not  
364 isolated from matrices heavily polluted by petroleum hydrocarbons and do not show genetic bases  
365 for PAHs degradation (Tsurumaki et al., 2019; Nagata et al., 2019; Kim et al., 2018), we can expect  
366 that this CYP450 system acquired by this novel *H. alkaliantarctica* strain SRM2 may have conferred  
367 specific PAH-degradation abilities to this strain. As CYP450s are broadly distributed across the tree  
368 of life and are considered the most versatile biocatalysts in nature because of the wide variety of  
369 substrate structures they can react with (Nelson, 2018; Yeom et al., 2021; Haas et al., 2022), further  
370 studies are needed to test the possible degradation/detoxification activity of the enzymes we identified  
371 towards other contaminants besides petroleum hydrocarbons.

### 372 3.2.2. Central pathways for hydrocarbon degradation

373 The activity of RHD generates salicylate that enters the catechol pathway, a classic bacterial central  
374 hydrocarbon degradation route mainly found in proteobacteria and actinobacteria (Nešvera et al.,  
375 2015). In both MAGs, we identified the salicylate hydroxylase, which catalyzes the formation of  
376 catechol acid (a substrate of the catechol 1-2 dioxygenase enzyme; Nešvera et al., 2015), as well as

377 the terminal enzymes ( $\beta$ -keto adipate succinyl-CoA transferase or  $\beta$ -keto adypyl thiolase), which  
378 convert  $\beta$ -keto adipic acid in acetyl-CoA and succinyl-CoA (Peng et al., 2008). Additionally, we found  
379 a hydroxiquinol 1,2 dioxygenase (BLASTp e-value 0, score 1461, id. 95.5%) suggested to promote  
380 the formation of  $\beta$ -keto adipic acid (Ferraroni et al., 2005). These findings suggest that the catechol  
381 pathway could contribute to the observed PAH degradation ability of both strains.

382 Following CYP450-mediated trans-hydroxylation, the degradation of hydrocarbons proceeds *via* the  
383 protocatechuate metabolic pathway, which starts with an hydroxybenzoate hydroxylase that  
384 transforms 4-hydroxybenzoate into 3,4 hydroxybenzoate (Fuchs et al., 2011). This enzyme has been  
385 identified only in *H. alkali antarctica* strain SRM2 MAG (Supplementary Fig. S5), which also  
386 possessed a protocatechuate 3,4-dioxygenase, 3-carboxy-*cis,cis*-muconolactone cycloisomerase and  
387 4-carboxymuconolactone decarboxylase. Again, we cannot exclude that genes with similar function  
388 but low sequence-similarity may be present also in *A. xenomutans* strain SRM1 MAG. Conversely,  
389 both MAGs possessed the terminal enzymes of the protocatechuate pathway (including  $\beta$ -keto adipate  
390 enol-lactonase, 3-oxo adipate CoA transferase,  $\beta$ -keto adypil-CoA thiolase) capable of catalyzing the  
391 production of succinyl-CoA. The comparison of the *H. alkali antarctica* strain SRM2 gene sequences  
392 involved in this pathway with their homologs, generally showed high conservation (>70%)  
393 (Supplementary Fig. S5). To the best of our knowledge, this is the first evidence for genes associated  
394 with the protocatechuate degradation pathway organized as an operon in a *Halomonas* sp. (Corti  
395 Monzón et al., 2018). Further analyses are needed to assess if this feature is peculiar to the strains  
396 considered here, or common across *Halomonas* spp. and/or other bacterial taxa.

397 Several other genes of the homoprotocatechuate pathway (Méndez et al., 2011) were also identified  
398 in both MAGs (Fig. 6A,B). Notably, the genome region containing the genes for the  
399 homoprotocatechuate pathway in *A. xenomutans* strain SRM1 was highly conserved with that of close  
400 *Alcanivorax* sp. relatives (Fig. 6A), while that of *H. alkali antarctica* SRM2 showed no apparent  
401 homologs in currently known *Halomonas* sp. genomes (Fig. 6B). We thus suggest that deeper  
402 investigation of the homoprotocatechuate pathway of *H. alkali antarctica* strain SRM2 through

403 mutation, heterologous expression or proteomics may unveil novel mechanisms for PAH  
404 biodegradation.

405 Finally, *H. alkaliantarctica* strain SRM2 also displayed a complete set of genes for the phenylacetic  
406 pathway within a specific genomic region, highly conserved with closest *Halomonas* sp. genomes  
407 available for comparison (Supplementary Fig. S6), indicating that this strain possesses a particularly  
408 wide array of genes for PAHs degradation. Additional enzymes involved in the homogentisate  
409 pathway (Arias-Barrau et al., 2004; Guazzaroni et al., 2013), were detected in both MAGs, including  
410 maleylacetoacetate isomerase, fumarylacetoacetate hydrolase, and 4-hydroxyphenilpyruvate  
411 dioxygenase (data not shown).

412

### 413 **3.3. Genetic basis for HMs removal/detoxification**

414 The blastp search against the SwissProt database identified in both MAGs the genetic basis for  
415 exopolysaccharide (EPS) biosynthesis, which may not only facilitate PAH degradation by enhancing  
416 the efficiency of all abovementioned PAH-degrading enzymes (Gutierrez et al., 2013; Alaba et al.,  
417 2018), but also confer metal-binding properties contributing to explain the observed ability of these  
418 strains to resist to and to sequester HMs (Amoozegar et al., 2012; Gupta and Diwan, 2017; Cao et al.,  
419 2022; Cheng et al., 2022). Specifically, a UDP-glucose 4-epimerase (e-value 0, score 1709, 97%  
420 identity), a UDP-glucose pyrophosphorylase (e-value  $7.5 \times 10^{-180}$ , score 1300, 90% identity), and a poly-  
421 beta-1,6-N-acetyl-D-glucosamine synthase (e-value 0, score 2011, 81% identity) were identified in  
422 *A. xenomutans* strain SRM1, with the latter two also present in *H. alkaliantarctica* strain SRM2  
423 (respectively, e-value  $2.6 \times 10^{-165}$ , score 1206, identity 77.3%, and e-value 0, score 1977, identity 86.1%).

424 A deeper insight on the production of EPS by the tested strains would likely allow optimizing of their  
425 use for simultaneous bioremediation of PAHs and HMs in different environmental conditions.

426 The annotation of the two MAGs by the automated RAST pipeline, followed by blastp check against  
427 the SwissProt database, shed light on further mechanisms possibly involved in the ability of both  
428 strains to resist to and to remove/detoxify HMs (Supplementary Fig. S7). In particular, several genes

429 coded for efflux pumps that selectively and non-selectively regulate the transport of multiple metals,  
430 including *czcA*, *czcB*, *czcC*, *czcD* able to transport cobalt, zinc and cadmium, as well as *corC*, specific  
431 for the transport of cobalt and magnesium. In addition, several genes were identified encoding for  
432 proteins involved in the detoxification of copper, including a copper chaperone, copper homeostasis  
433 protein CutE and CutF, copper resistance protein B, C, D, copper ATP ase, Cu-sensing two-  
434 component system response regulator and Cu-responsive transcriptional regulator. Notably, the  
435 automated RAST pipeline also identified multicopper and blue multicopper oxidases (Supplementary  
436 Fig. S7), which may be synergistically involved both in Cu-detoxification/sequestration and act as  
437 laccase-like multicopper oxidases for the degradation of PAHs and other organic contaminants  
438 (Cooksey, 1994; Arregui et al., 2019; Ramasamy et al., 2020; Zhang et al., 2020).

439 Several genes coding for proteins known to confer resistance to and/or detoxify arsenic were found  
440 in both MAGs (Supplementary Fig. S7), and our manual annotation of flanking gene sequences  
441 highlighted operon-like structures (Figure 7A-B) similar to “arsenic islands” previously described  
442 (Wu et al., 2018). In *A. xenomutans* strain SRM1, this MAG region (Figure 7A) included an arsenic  
443 resistance protein ArsH, arsenic transporter ArsB, arsenate reductase ArsC and ArsR regulator (Rosen  
444 and Liu, 2009; Chang et al., 2018). This genome region of *A. xenomutans* strain SRM1 showed on  
445 average  $\geq 70\%$  similarity with its closest *Alcanivorax* sp. relatives (Figure 7A), indicating high  
446 conservation of this As operon and flanking regions across different *Alcanivorax* species. Similarly,  
447 in *H. alkaliantarctica* strain SRM2, the region comprising the arsenic operon displayed high  
448 conservation with the homologous sequences of close *Halomonas* sp. relatives (Figure 7B). Notably,  
449 all the ORFs of the arsenic resistance operon in *H. alkaliantarctica* strain SRM2 MAG are contiguous,  
450 whereas additional ORFs (of un-identified function) are present in the three reference genomes, that  
451 separate ORF 7 from ORFs 8-9. As the three reference *Halomonas* genomes were obtained from  
452 matrices not contaminated by As (Nagata et al., 2019; Williamson et al., 2016) we can argue that this  
453 observed simplification of the arsenic operon may have enhanced the ability of *H. alkaliantarctica*  
454 strain SRM2 to detoxify/remove As. The functional implications of the observed differences in the

455 structure of the arsenic resistance operon across different *Halomonas* species remains to be further  
456 investigated, to understand which gene asset may perform better for different bioremediation  
457 purposes.

458 A complete mercury-resistance operon (Boyd and Barkay, 2012) was also identified in *H.*  
459 *alkaliantarctica* strain SRM2 (Fig. 7C), whose coding sequence was highly conserved within the  
460 closest relatives retrieved from NCBI (*H. axialiensis* Althf1 and *H. sp.* ZM 3, which were isolated  
461 from HM-rich hydrothermal vents in the Pacific Ocean and from a mineral waste repository,  
462 respectively; Dziewit et al., 2013; Tsurumaki et al., 2019). The structure of the identified mercury  
463 operon showed typical features identified in other *Halomonas* species (Boyd and Barkay, 2012),  
464 including two transcriptional regulators MerR, a mercuric transport protein MerT (able to transport  
465 Hg(II) to the cytoplasm), a periplasmic Hg-binding protein MerP, a mercuric reductase MerA, and an  
466 organomercurial lyase MerB. Based on the presence of this Mer operon, and especially of MerP, we  
467 can expect that, even if not directly assessed in our bioremediation tests, *H. alkaliantarctica* strain  
468 SRM2 may also perform Hg<sup>2+</sup> biosorption (Huang et al., 2003).

469 Notably, several of the genes of the arsenic and mercury operons we identified have been documented  
470 to cross-react with other toxic metals. For instance, ArsH has been reported to also detoxify and  
471 enhance the precipitation of chromium by reducing Cr(VI) to Cr(III) (Xue et al., 2014), ArsB to also  
472 detoxify the hazardous metalloid antimony (Meng et al., 2004), and MerP to be involved in the  
473 biosorption of other HMs such as nickel, chromium, copper and zinc (Kao et al., 2008; Hsueh et al.,  
474 2017). This suggests that similar cross-reactivity with multiple HMs may contribute to explain the  
475 overall high removal efficiency towards the different HMs observed in our bioremediation  
476 experiments, possibly extending our findings to additional HMs not directly investigated here.

477 Nevertheless, we acknowledge that other processes such as extracellular electron transfer and  
478 electrocatalysis, not assessed in the present study, may be involved in HMs bioremediation (Liu et  
479 al., 2018) and as such deserve further investigations.

480

481 **4. Conclusions**

482 In summary, our multidisciplinary study based on an integrated approach that combines experimental  
483 microbiome enrichment, next-generation sequencing and selective culturing, allowed us to obtain two  
484 novel *Alcanivorax* and *Halomonas* strains with promising bioremediation potential. Both strains were  
485 shown to be resistant to, and to detoxify or remove multiple PAHs and HMs, and hence represent  
486 promising candidates for developing bioremediation applications (e.g., bioaugmentation or *ex situ*  
487 treatments) in environments contaminated by combinations of toxic pollutants. Finally, the contextual  
488 analysis of their genomic repertoire highlights the presence of genes and/or operons that are proposed  
489 as possible bioengineering targets, to further enhance the observed ability of these or other bacterial  
490 strains to serve for environmental bioremediation purposes.

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507 **CRedit authorship contribution statement**

508 AD, ER, and FD conceived the study. FD, MP, ER, and CS conducted the field work. FD, ER, LJZ,  
509 and MT conducted the laboratory analyses. FD, ER, LJZ, MT, AC, EB, PNG and AD contributed to  
510 data elaboration. ER and FD wrote the draft of the manuscript. All authors critically revised the article  
511 and contributed to its finalization.

512

513 **Declaration of competing interest**

514 The authors declare that they have no known competing financial interests or personal relationships  
515 that could have appeared to influence the work reported in this paper.

516

517 **Acknowledgements**

518 This study was supported by the projects ABbaCo funded by the Italian Ministry for Education,  
519 University and Research (grant number C62F16000170001), MERCES (H2020-SC5-2015, grant  
520 number 689518), and SZN-Polytechnic University of Marche OPEN University PhD grant to FD.  
521 PNG acknowledges the Natural Environment Research Council (NERC) for funding ‘Plastic Vectors’  
522 project NE/S004548/1. We would like to thank Marco Cannavacciuolo, Francesco Terlizzi and  
523 Gianluca Zazo for field support, as well as Dr. Trevor Willis for useful comments on our work.

524

525 **Appendix A. Supplementary data**

526 Supplementary data to this article can be found online at XXX

527 **Data Availability Statement**

528 The sequencing data of the two bacterial strains described in this work have been deposited in NCBI  
529 (accession number: PRJNA899357). The bacterial strains obtained in our study are maintained at the  
530 Stazione Zoologica Anton Dohrn, National Institute for Marine Biology, Ecology and Biotechnology,  
531 and are available upon request of collaboration.

532

533 **References**

- 534 Abdel-Razik, M. A., Azmy, A. F., Khairalla, A. S., & AbdelGhani, S. (2020). Metal bioremediation  
535 potential of the halophilic bacterium, *Halomonas* sp. strain WQL9 isolated from Lake Qarun,  
536 Egypt. *The Egyptian Journal of Aquatic Research*, 46, 19-25. doi:10.1016/j.ejar.2019.11.009.
- 537 Adams, G. O., Fufeyin, P. T., Okoro, S. E., & Ehinomen, I. (2015). Bioremediation, biostimulation  
538 and bioaugmentation: a review. *International Journal of Environmental Bioremediation &*  
539 *Biodegradation*, 3, 28-39. doi:10.12691/ijebb-3-1-5.
- 540 Alaba, P. A., Oladoja, N. A., Sani, Y. M., Ayodele, O. B., Mohammed, I. Y., Olupinla, S. F., & Daud,  
541 W. M. W. (2018). Insight into wastewater decontamination using polymeric adsorbents. *Journal*  
542 *of Environmental Chemical Engineering*, 6, 1651-1672. doi:10.1016/j.jece.2018.02.019.
- 543 Ali, M., Song, X., Ding, D., Wang, Q., Zhang, Z., & Tang, Z. (2021). Bioremediation of PAHs and  
544 heavy metals co-contaminated soils: Challenges and enhancement strategies. *Environmental*  
545 *Pollution*, 118686. doi:10.1016/j.envpol.2021.118686
- 546 Alisi, C., Musella, R., Tasso, F., Ubaldi, C., Manzo, S., Cremisini, C., & Sprocati, A. R. (2009).  
547 Bioremediation of diesel oil in a co-contaminated soil by bioaugmentation with a microbial  
548 formula tailored with native strains selected for heavy metals resistance. *Science of the Total*  
549 *Environment*, 407(8), 3024-3032. doi:10.1016/j.scitotenv.2009.01.011.
- 550 Al-Turki, A. I. (2009). Microbial Polycyclic Aromatic Hydrocarbons degradation in soil. *Research*  
551 *Journal of Environmental Toxicology*, 3(1), 1-8. doi:10.3923/rjet.2009.1.8.
- 552 Amoozegar, M. A., Ghazanfari, N., & Didari, M. (2012). Lead and cadmium bioremoval by  
553 *Halomonas* sp., an exopolysaccharide-producing halophilic bacterium. *Progress in Biological*  
554 *Sciences*, 2(1), 1-11. doi:10.22059/PBS.2012.24820
- 555 Arias-Barrau, E., Olivera, E. R., Luengo, J. M., Fernández, C., Galán, B., García, J. L., ... &  
556 Minambres, B. (2004). The homogentisate pathway: a central catabolic pathway involved in the  
557 degradation of L-phenylalanine, L-tyrosine, and 3-hydroxyphenylacetate in *Pseudomonas putida*.  
558 *Journal of Bacteriology*, 186(15), 5062-5077. doi:10.1128/JB.186.15.5062-5077.2004
- 559 Arregui, L., Ayala, M., Gómez-Gil, X., Gutiérrez-Soto, G., Hernández-Luna, C. E., Herrera de Los  
560 Santos, M., ... & Valdez-Cruz, N. A. (2019). Laccases: structure, function, and potential  
561 application in water bioremediation. *Microbial Cell Factories*, 18(1), 1-33.



- 562 Aryal, M., Ziagova, M., & Liakopoulou-Kyriakides, M. (2010). Study on arsenic biosorption using  
563 Fe (III)-treated biomass of *Staphylococcus xylosus*. *Chemical Engineering Journal*, 162(1), 178-  
564 185. doi:10.1016/j.cej.2010.05.026
- 565 Asksonthong, R., Siripongvutikorn, S., & Usawakesmanee, W. (2018). Heavy metal removal ability  
566 of *Halomonas elongata* and *Tetragenococcus halophilus* in a media model system as affected by  
567 pH and incubation time. *International Food Research Journal*, 25, 234-240.  
568 www.ifrj.upm.edu.my/25%20(01)%202018/(30).pdf
- 569 Ates, O. (2015). Systems biology of microbial exopolysaccharides production. *Frontiers in*  
570 *bioengineering and biotechnology*, 3, 200. doi:10.3389/fbioe.2015.00200
- 571 Bagby, S. C., Reddy, C. M., Aeppli, C., Fisher, G. B., & Valentine, D. L. (2017). Persistence and  
572 biodegradation of oil at the ocean floor following Deepwater Horizon. *Proceedings of the National*  
573 *Academy of Sciences*, 114(1), E9-E18. doi:10.1073/pnas.161011011
- 574 Barbato, M., Mapelli, F., Chouaia, B., Crotti, E., Daffonchio, D., & Borin, S. (2015). Draft genome  
575 sequence of the hydrocarbon-degrading bacterium *Alcanivorax dieselolei* KS-293 isolated from  
576 surface seawater in the Eastern Mediterranean Sea. *Genome announcements*, 3(6), e01424-15.  
577 doi:10.1128/genomeA.01424-15
- 578 Barry, S. M., & Challis, G. L. (2013). Mechanism and catalytic diversity of Rieske non-heme iron-  
579 dependent oxygenases. *ACS catalysis*, 3(10), 2362-2370. doi:10.1021/cs400087p
- 580 Beolchini, F., Dell'Anno, A., De Propriis, L., Ubaldini, S., Cerrone, F., & Danovaro, R. (2009). Auto-  
581 and heterotrophic acidophilic bacteria enhance the bioremediation efficiency of sediments  
582 contaminated by heavy metals. *Chemosphere*, 74(10), 1321-1326.  
583 doi:10.1016/j.chemosphere.2008.11.057
- 584 Beyersmann, D., & Hartwig, A. (2008). Carcinogenic metal compounds: recent insight into molecular  
585 and cellular mechanisms. *Archives of toxicology*, 82(8), 493-512. doi:10.1007/s00204-008-0313-  
586 y
- 587 Bezza, F. A., & Chirwa, E. M. N. (2017). The role of lipopeptide biosurfactant on microbial  
588 remediation of aged polycyclic aromatic hydrocarbons (PAHs)-contaminated soil. *Chemical*  
589 *Engineering Journal*, 309, 563-576. doi:10.1016/j.cej.2016.10.055.
- 590 Boyd, E. S., & Barkay, T. (2012). The mercury resistance operon: from an origin in a geothermal  
591 environment to an efficient detoxification machine. *Frontiers in microbiology*, 3, 349.  
592 doi:10.3389/fmicb.2012.00349.

- 593 Buah-Kwofie, A., Humphries, M. S., & Pillay, L. (2018). Bioaccumulation and risk assessment of  
594 organochlorine pesticides in fish from a global biodiversity hotspot: iSimangaliso Wetland Park,  
595 South Africa. *Science of the total environment*, 621, 273-281.  
596 doi:10.1016/j.scitotenv.2017.11.212.
- 597 Budiyanto, F., Thukair, A., Al-Momani, M., Musa, M. M., & Nzila, A. (2018). Characterization of  
598 halophilic bacteria capable of efficiently biodegrading the high-molecular-weight polycyclic  
599 aromatic hydrocarbon pyrene. *Environmental Engineering Science*, 35(6), 616-626.  
600 doi:10.1089/ees.2017.0244.
- 601 Cai, X., Zheng, X., Zhang, D., Iqbal, W., Liu, C., Yang, B., ... & Mao, Y. (2019). Microbial  
602 characterization of heavy metal resistant bacterial strains isolated from an electroplating  
603 wastewater treatment plant. *Ecotoxicology and environmental safety*, 181, 472-480.  
604 doi:10.1016/j.ecoenv.2019.06.036.
- 605 Cao, J., Lai, Q., Yuan, J., & Shao, Z. (2015). Genomic and metabolic analysis of fluoranthene  
606 degradation pathway in *Celeribacter indicus* P73T. *Scientific reports*, 5(1), 1-12.  
607 doi:10.1038/srep07741.
- 608 Cao, X., Xu, L., Chen, Y. P., Decho, A. W., Cui, Z., & Lead, J. R. (2022). Contribution, composition,  
609 and structure of EPS by in vivo exposure to elucidate the mechanisms of nanoparticle-enhanced  
610 bioremediation to metals. *Environmental Science & Technology*, 56(2), 896-906.
- 611 Casillo, A., Lanzetta, R., Parrilli, M., & Corsaro, M. M. (2018). Exopolysaccharides from marine and  
612 marine extremophilic bacteria: structures, properties, ecological roles and applications. *Marine*  
613 *drugs*, 16(2), 69. doi:10.3390/md16020069.
- 614 Catania, V., Cappello, S., Di Giorgi, V., Santisi, S., Di Maria, R., Mazzola, A., ... & Quatrini, P.  
615 (2018). Microbial communities of polluted sub-surface marine sediments. *Marine Pollution*  
616 *Bulletin*, 131, 396-406. doi:10.1016/j.marpolbul.2018.04.015.
- 617 Cecchi, G., Cutroneo, L., Di Piazza, S., Besio, G., Capello, M., & Zotti, M. (2021). Port Sediments:  
618 Problem or Resource? A Review Concerning the Treatment and Decontamination of Port  
619 Sediments by Fungi and Bacteria. *Microorganisms*, 9(6), 1279.  
620 doi:10.3390/microorganisms9061279
- 621 Chang, J. S., Yoon, I. H., & Kim, K. W. (2018). Arsenic biotransformation potential of microbial  
622 arsH responses in the biogeochemical cycling of arsenic-contaminated groundwater.  
623 *Chemosphere*, 191, 729-737. doi:10.1016/j.chemosphere.2017.10.044.

- 624 Chen, Z., Zhang, X., Ma, Y., Song, Y., Li, Y., Geng, G., & Huang, Y. (2020). Anammox biofilm  
625 system under the stress of Hg (II): Nitrogen removal performance, microbial community dynamic  
626 and resistance genes expression. *Journal of hazardous materials*, 395, 122665.  
627 doi:10.1016/j.jhazmat.2020.122665.
- 628 Cheng, H., Mai, Z., Wang, Y., Liu, D., & Sun, Y. (2022). Role of extracellular polymeric substances  
629 in metal sequestration during mangrove restoration. *Chemosphere*, 135550.
- 630 Cooksey, D. A. (1994). Molecular mechanisms of copper resistance and accumulation in bacteria.  
631 *FEMS microbiology reviews*, 14(4), 381-386.
- 632 Corti Monzón, G., Nisenbaum, M., Herrera Seitz, M. K., & Murialdo, S. E. (2018). New findings on  
633 aromatic compounds' degradation and their metabolic pathways, the biosurfactant production and  
634 motility of the halophilic bacterium *Halomonas* sp. KHS3. *Current microbiology*, 75(8), 1108-  
635 1118. doi:10.1007/s00284-018-1497-x
- 636 Costa, P. G., Marube, L. C., Artifon, V., Escarrone, A. L., Hernandes, J. C., Zebral, Y. D., &  
637 Bianchini, A. (2022). Temporal and spatial variations in metals and arsenic contamination in  
638 water, sediment and biota of freshwater, marine and coastal environments after the Fundão dam  
639 failure. *Science of The Total Environment*, 806, 151340. doi:10.1016/j.scitotenv.2021.151340
- 640 Czaplicki, L. M., & Gunsch, C. K. (2016). Reflection on molecular approaches influencing state-of-  
641 the-art bioremediation design: culturing to microbial community fingerprinting to omics. *Journal*  
642 *of environmental engineering (New York, NY)*, 142(10). doi:10.1061/(ASCE)EE.1943-  
643 7870.0001141
- 644 da Costa Waite, C. C., da Silva, G. O. A., Bitencourt, J. A. P., Sabadini-Santos, E., & Crapez, M. A.  
645 C. (2016). Copper and lead removal from aqueous solutions by bacterial consortia acting as  
646 biosorbents. *Marine pollution bulletin*, 109(1), 386-392. doi:10.1016/j.marpolbul.2016.05.044
- 647 Dell'Anno, F., Rastelli, E., Sansone, C., Brunet, C., Ianora, A., & Dell'Anno, A. (2021). Bacteria,  
648 fungi and microalgae for the bioremediation of marine sediments contaminated by petroleum  
649 hydrocarbons in the omics era. *Microorganisms*, 9(8), 1695.  
650 doi:10.3390/microorganisms9081695.
- 651 Dell'Anno, F., Brunet, C., van Zyl, L. J., Trindade, M., Golyshin, P. N., Dell'Anno, A., ... & Sansone,  
652 C. (2020). Degradation of hydrocarbons and heavy metal reduction by marine bacteria in highly  
653 contaminated sediments. *Microorganisms*, 8(9), 1402. doi:10.3390/microorganisms8091402.

- 654 Dell'Anno, F., Rastelli, E., Tangherlini, M., Corinaldesi, C., Sansone, C., Brunet, C., ... & Dell'Anno,  
655 A. (2021). Highly contaminated marine sediments can host rare bacterial taxa potentially useful  
656 for bioremediation. *Frontiers in microbiology*, 12, 584850. doi:10.3389/fmicb.2021.584850.
- 657 Dell'Anno, F., van Zyl, L. J., Trindade, M., Brunet, C., Dell'Anno, A., Ianora, A., & Sansone, C.  
658 (2021). Metagenome-assembled genome (MAG) of *Oceancaulis alexandrii* NP7 isolated from  
659 Mediterranean Sea polluted marine sediments and its bioremediation potential. *G3*, 11(9), jkab210.  
660 doi:10.1093/g3journal/jkab210.
- 661 Dubinsky, E. A., Conrad, M. E., Chakraborty, R., Bill, M., Borglin, S. E., Hollibaugh, J. T., ... &  
662 Andersen, G. L. (2013). Succession of hydrocarbon-degrading bacteria in the aftermath of the  
663 Deepwater Horizon oil spill in the Gulf of Mexico. *Environmental science & technology*, 47(19),  
664 10860-10867. doi:10.1021/es401676y.
- 665 Dueholm, M. S., Marques, I. G., Karst, S. M., D'Imperio, S., Tale, V. P., Lewis, D., ... & Nielsen, J.  
666 L. (2015). Survival and activity of individual bioaugmentation strains. *Bioresource Technology*,  
667 186, 192-199. doi:10.1016/j.biortech.2015.02.111.
- 668 Dziewit, L., Pyzik, A., Matlakowska, R., Baj, J., Szuplewska, M., & Bartosik, D. (2013).  
669 Characterization of *Halomonas* sp. ZM3 isolated from the Zelazny Most post-flotation waste  
670 reservoir, with a special focus on its mobile DNA. *BMC microbiology*, 13(1), 1-13.  
671 doi:10.1186/1471-2180-13-59.
- 672 El-Mufleh, A., Béchet, B., Basile-Doelsch, I., Geffroy-Rodier, C., Gaudin, A., & Ruban, V. (2014).  
673 Distribution of PAHs and trace metals in urban stormwater sediments: combination of density  
674 fractionation, mineralogy and microanalysis. *Environmental Science and Pollution Research*,  
675 21(16), 9764-9776. doi:10.1007/s11356-014-2850-7
- 676 EPA, U.S. 2014. "Method 6020B Inductively Coupled Plasma-Mass Spectrometry," 1–33.
- 677 Fenyvesi, É., Gruiz, K., Morillo, E., & Villaverde, J. (2019). Traditional and innovative methods for  
678 physical and chemical remediation of soil contaminated with organic contaminants. *Engineering*  
679 *Tools for Environmental Risk Management*, 4, 285-362. doi:10.1201/b20405-9.
- 680 Ferraroni, M., Seifert, J., Travkin, V. M., Thiel, M., Kaschabek, S., Scozzafava, A., ... & Briganti, F.  
681 (2005). Crystal structure of the hydroxyquinol 1, 2-dioxygenase from *Nocardioides simplex* 3E, a  
682 key enzyme involved in polychlorinated aromatics biodegradation. *Journal of Biological*  
683 *Chemistry*, 280(22), 21144-21154. doi:10.1074/jbc.M500666200.

- 684 Fodelianakis, S., Antoniou, E., Mapelli, F., Magagnini, M., Nikolopoulou, M., Marasco, R., ... &  
685 Kalogerakis, N. (2015). Allochthonous bioaugmentation in ex situ treatment of crude oil-polluted  
686 sediments in the presence of an effective degrading indigenous microbiome. *Journal of hazardous*  
687 *materials*, 287, 78-86. doi:10.1016/j.jhazmat.2015.01.038.
- 688 Fu, X., Lai, Q., Dong, C., Wang, W., & Shao, Z. (2018). Complete genome sequence of *Alcanivorax*  
689 *xenomutans* P40, an alkane-degrading bacterium isolated from deep seawater. *Marine Genomics*,  
690 38, 1-4. doi:10.1016/j.margen.2017.05.010.
- 691 Fuchs, G., Boll, M., & Heider, J. (2011). Microbial degradation of aromatic compounds—from one  
692 strategy to four. *Nature Reviews Microbiology*, 9(11), 803-816. doi:10.1038/nrmicro2652.
- 693 Fuentes-Gandara, F., Pinedo-Hernández, J., Marrugo-Negrete, J., & Díez, S. (2018). Human health  
694 impacts of exposure to metals through extreme consumption of fish from the Colombian Caribbean  
695 Sea. *Environmental geochemistry and health*, 40(1), 229-242. doi:10.1007/s10653-016-9896-z.
- 696 Ghosh, S., Bhattacharya, J., Nitnavare, R., & Webster, T. J. Microbial Remediation of Metals by  
697 Marine Bacteria. In *Development in Wastewater Treatment Research and Processes*, Eds.  
698 Rodriguez-Couto S. & Shah, M.P. pp. 131–158. Elsevier. doi:10.1016/B978-0-323-85839-  
699 7.00011-6.
- 700 Guazzaroni, M. E., Herbst, F. A., Lores, I., Tamames, J., Peláez, A. I., Lopez-Cortés, N., ... & Ferrer,  
701 M. (2013). Metaproteogenomic insights beyond bacterial response to naphthalene exposure and  
702 bio-stimulation. *The ISME journal*, 7(1), 122-136. doi:10.1038/ismej.2012.82
- 703 Guha, S., Peters, C. A., & Jaffé, P. R. (1999). Multisubstrate biodegradation kinetics of naphthalene,  
704 phenanthrene, and pyrene mixtures. *Biotechnology and bioengineering*, 65(5), 491-499.  
705 doi:10.1002/(SICI)1097-0290(19991205)65:5<491::AID-BIT1>3.0.CO;2-H
- 706 Gupta, P., & Diwan, B. (2017). Bacterial exopolysaccharide mediated heavy metal removal: a review  
707 on biosynthesis, mechanism and remediation strategies. *Biotechnology Reports*, 13, 58-71.  
708 doi:10.1016/j.btre.2016.12.006.
- 709 Gutierrez, T., Singleton, D. R., Berry, D., Yang, T., Aitken, M. D., & Teske, A. (2013). Hydrocarbon-  
710 degrading bacteria enriched by the Deepwater Horizon oil spill identified by cultivation and DNA-  
711 SIP. *The ISME journal*, 7(11), 2091-2104. doi:10.1038/ismej.2013.98
- 712 Haas, J., Hayward, A., Buer, B., Maiwald, F., Nebelsiek, B., Glaubitz, J., et al. (2022). Phylogenomic  
713 and functional characterization of an evolutionary conserved cytochrome P450-based insecticide

714 detoxification mechanism in bees. *Proceedings of the National Academy of Sciences*, 119(26),  
715 e2205850119. doi:10.1073/pnas.2205850119

716 Hammer, Ø., Harper, D. A., & Ryan, P. D. (2001). PAST: Paleontological statistics software package  
717 for education and data analysis. *Palaeontologia electronica*, 4(1), 9.

718 Han, N., Qiang, Y., & Zhang, W. (2016). ANIttools web: a web tool for fast genome comparison  
719 within multiple bacterial strains. *Database*, 2016. 1-5. doi:10.1093/database/baw084.

720 Haritash, A. K., & Kaushik, C. P. (2009). Biodegradation aspects of polycyclic aromatic  
721 hydrocarbons (PAHs): a review. *Journal of hazardous materials*, 169(1-3), 1-15.  
722 doi:10.1016/j.jhazmat.2009.03.137.

723 Hassan, S., Khurshid, Z., Bhat, S. A., Kumar, V., Ameen, F., & Ganai, B. A. (2022). Marine bacteria  
724 and omic approaches: A novel and potential repository for bioremediation assessment. *Journal of*  
725 *Applied Microbiology*, 133(4), 2299-2313. Doi.org/10.1111/jam.15711.

726 Harindintwali, J. D., Xiang, L., Wang, F., Chang, S. X., Zhao, Z., Mei, Z., et al. (2022). Syntrophy  
727 of bacteria and archaea in the anaerobic catabolism of hydrocarbon contaminants. *Critical Reviews*  
728 *in Environmental Science and Technology*, 1-27. doi:10.1080/10643389.2022.2134702

729 Hsueh, Y. H., Lin, K. S., Wang, Y. T., & Chiang, C. L. (2017). Copper, nickel, and zinc cations  
730 biosorption properties of Gram-positive and Gram-negative MerP mercury-resistance proteins.  
731 *Journal of the Taiwan Institute of Chemical Engineers*, 80, 168-175.

732 Huang, C. C., Su, C. C., Hsieh, J. L., Tseng, C. P., Lin, P. J., & Chang, J. S. (2003). Polypeptides for  
733 heavy-metal biosorption: capacity and specificity of two heterogeneous MerP proteins. *Enzyme*  
734 *and Microbial Technology*, 33(4), 379-385.

735 Huo, Y. Y., Li, Z. Y., Cheng, H., Wang, C. S., & Xu, X. W. (2014). High quality draft genome  
736 sequence of the heavy metal resistant bacterium *Halomonas zincidurans* type strain B6T.  
737 *Standards in genomic sciences*, 9(1), 1-9. doi:10.1186/1944-3277-9-30.

738 Imperato, V., Portillo-Estrada, M., McAmmond, B. M., Douwen, Y., Van Hamme, J. D., Gawronski,  
739 S. W., ... & Thijs, S. (2019). Genomic diversity of two hydrocarbon-degrading and plant growth-  
740 promoting *Pseudomonas* species isolated from the oil field of Bóbrka (Poland). *Genes*, 10(6), 443.  
741 doi:10.3390/genes10060443.

742 Jain, M., Khan, S. A., Sharma, K., Jadhao, P. R., Pant, K. K., Ziora, Z. M., & Blaskovich, M. A.  
743 (2022). Current perspective of innovative strategies for bioremediation of organic pollutants from  
744 wastewater. *Bioresource Technology*, 344, 126305. doi:10.1016/j.biortech.2021.126305

745 Johnsen, A. R., Wick, L. Y., & Harms, H. (2005). Principles of microbial PAH-degradation in soil.  
746 *Environmental pollution*, 133(1), 71-84. doi:10.1016/j.envpol.2004.04.015

747 Joshi, N. C. (2017). Heavy metals, conventional methods for heavy metal removal, biosorption and  
748 the development of low cost adsorbent. *Eur. J. Pharm. Med. Res*, 4, 388-393.

749 Kadri, T., Magdouli, S., Rouissi, T., & Brar, S. K. (2018). Ex-situ biodegradation of petroleum  
750 hydrocarbons using *Alcanivorax borkumensis* enzymes. *Biochemical Engineering Journal*, 132,  
751 279-287. doi:10.1016/j.bej.2018.01.014.

752 Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y., & Morishima, K. (2017). KEGG: new  
753 perspectives on genomes, pathways, diseases and drugs. *Nucleic acids research*, 45(D1), D353-  
754 D361. doi:10.1093/nar/gkw1092.

755 Kao, W. C., Huang, C. C., & Chang, J. S. (2008). Biosorption of nickel, chromium and zinc by MerP-  
756 expressing recombinant *Escherichia coli*. *Journal of hazardous materials*, 158(1), 100-106.

757 Kauppi, B., Lee, K., Carredano, E., Parales, R. E., Gibson, D. T., Eklund, H., & Ramaswamy, S.  
758 (1998). Structure of an aromatic-ring-hydroxylating dioxygenase–naphthalene 1, 2-dioxygenase.  
759 *Structure*, 6(5), 571-586. doi:10.1016/S0969-2126(98)00059-8.

760 Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N., & Sternberg, M. J. (2015). The Phyre2 web  
761 portal for protein modeling, prediction and analysis. *Nature protocols*, 10(6), 845-858.  
762 doi:10.1038/nprot.2015.053

763 Kim, H. S., Kim, Y. J., & Seo, Y. R. (2015). An overview of carcinogenic heavy metal: molecular  
764 toxicity mechanism and prevention. *Journal of cancer prevention*, 20(4), 232.  
765 doi:10.15430/jcp.2015.20.4.232.

766 Kim, S. J., Do, K. T., & Park, S. J. (2018). Complete genome of *Halomonas aestuarii* Hb3, isolated  
767 from tidal flat. *Marine genomics*, 37, 43-45. doi:10.1016/j.margen.2017.11.007

768 Kordialik-Bogacka, E., & Diowksz, A. (2014). Metal uptake capacity of modified *Saccharomyces*  
769 *pastorianus* biomass from different types of solution. *Environmental Science and Pollution*  
770 *Research*, 21(3), 2223-2229. doi:10.1007/s11356-013-2144-5

- 771 Kumar, A. G., Rajan, N. N., Kirubakaran, R., & Dharani, G. (2019). Biodegradation of crude oil using  
772 self-immobilized hydrocarbonoclastic deep sea bacterial consortium. *Marine pollution bulletin*,  
773 146, 741-750. doi:10.1016/j.marpolbul.2019.07.006.
- 774 Kuppusamy, S., Thavamani, P., Venkateswarlu, K., Lee, Y. B., Naidu, R., & Megharaj, M. (2017).  
775 Remediation approaches for polycyclic aromatic hydrocarbons (PAHs) contaminated soils:  
776 Technological constraints, emerging trends and future directions. *Chemosphere*, 168, 944-968.  
777 doi:10.1016/j.chemosphere.2016.10.115
- 778 Ladino-Orjuela, G., Gomes, E., Silva, R. D., Salt, C., & Parsons, J. R. (2016). Metabolic pathways  
779 for degradation of aromatic hydrocarbons by bacteria. *Reviews of Environmental Contamination*  
780 *and Toxicology* 237, 105-121. doi:10.1007/978-3-319-23573-8\_5
- 781 Lai, Q., Li, W., & Shao, Z. (2012). Complete Genome Sequence of *Alcanivorax dieselolei* Type  
782 Strain B5. *Journal of Bacteriology*, 194(23), 6674–6674. doi:10.1128/JB.01813-12.
- 783 Leahy, J. G., & Colwell, R. R. (1990). Microbial degradation of hydrocarbons in the environment.  
784 *Microbiological reviews*, 54(3), 305-315. doi:10.1128/mr.54.3.305-315.1990
- 785 Leung, W. C., Wong, M. F., Chua, H., Lo, W., Yu, P. H. F., & Leung, C. K. (2000). Removal and  
786 recovery of heavy metals by bacteria isolated from activated sludge treating industrial effluents  
787 and municipal wastewater. *Water science and technology*, 41(12), 233-240.  
788 doi:10.2166/wst.2000.0277
- 789 Lin, H. H., & Liao, Y. C. (2016). Accurate binning of metagenomic contigs via automated clustering  
790 sequences using information of genomic signatures and marker genes. *Scientific reports*, 6, 24175.  
791 doi:10.1038/srep24175.
- 792 Little, D. J., Li, G., Ing, C., DiFrancesco, B. R., Bamford, N. C., Robinson, H., ... & Howell, P. L.  
793 (2014). Modification and periplasmic translocation of the biofilm exopolysaccharide poly- $\beta$ -1, 6-  
794 N-acetyl-d-glucosamine. *Proceedings of the National Academy of Sciences*, 111(30), 11013-  
795 11018. doi:10.1073/pnas.1406388111.
- 796 Liu, S. H., Zeng, G. M., Niu, Q. Y., Liu, Y., Zhou, L., Jiang, L. H., ... & Cheng, M. (2017).  
797 Bioremediation mechanisms of combined pollution of PAHs and heavy metals by bacteria and  
798 fungi: A mini review. *Bioresour. Technol.*, 224, 25-33. doi:10.1016/j.biortech.2016.11.095.
- 799 Liu, X., Li, M., Castelle, C. J., Probst, A. J., Zhou, Z., Pan, J., ... & Gu, J. D. (2018a). Insights into  
800 the ecology, evolution, and metabolism of the widespread Woese archaeal lineages. *Microbiome*,  
801 6(1), 1-16. doi: 10.1186/s40168-018-0488-2.



- 802 Liu, X., Shi, L., & Gu J-D. (2018b). Microbial electrocatalysis: Redox mediators responsible for  
803 extracellular electron transfer. *Biotechnology Advances*, 36, 1815-1827. doi:  
804 10.1016/j.biotechadv.2018.07.001.
- 805 Liu, X., Wang, Y., & Gu, J. D. (2021). Ecological distribution and potential roles of Woesearchaeota  
806 in anaerobic biogeochemical cycling unveiled by genomic analysis. *Computational and Structural*  
807 *Biotechnology Journal*, 19, 794-800. doi: 10.1016/j.csbj.2021.01.013.
- 808 Liu, X. X., Hu, X., Cao, Y., Pang, W. J., Huang, J. Y., Guo, P., & Huang, L. (2019). Biodegradation  
809 of phenanthrene and heavy metal removal by acid-tolerant *Burkholderia fungorum* FM-2.  
810 *Frontiers in microbiology*, 10, 408. doi:10.3389/fmicb.2019.00408.
- 811 Lloyd-Jones, G., & Lau, P. C. (1997). Glutathione S-transferase-encoding gene as a potential probe  
812 for environmental bacterial isolates capable of degrading polycyclic aromatic hydrocarbons.  
813 *Applied and Environmental Microbiology*, 63(8), 3286-3290. doi:10.1128/aem.63.8.3286-  
814 3290.1997
- 815 Loflen, C. L., Buck, T., Bonnema, A., & Heim, W. A. (2018). Pollutant bioaccumulation in the  
816 California spiny lobster (*Panulirus interruptus*) in San Diego Bay, California, and potential human  
817 health implications. *Marine pollution bulletin*, 128, 585-592.  
818 doi:10.1016/j.marpolbul.2018.02.001.
- 819 Mendez, V., Agullo, L., Gonzalez, M., & Seeger, M. (2011). The homogentisate and  
820 homoprotocatechuate central pathways are involved in 3- and 4-hydroxyphenylacetate degradation  
821 by *Burkholderia xenovorans* LB400. *PLoS One*, 6(3), e17583. doi:10.1371/journal.pone.0017583
- 822 Meng, Y. L., Liu, Z., & Rosen, B. P. (2004). As (III) and Sb (III) uptake by GlpF and efflux by ArsB  
823 in *Escherichia coli*. *Journal of Biological Chemistry*, 279(18), 18334-18341.
- 824 Meng, S., Peng, T., Liu, X., Wang, H., Huang, T., Gu, J. D., & Hu, Z. (2022). Ecological role of  
825 bacteria involved in the biogeochemical cycles of mangroves based on functional genes detected  
826 through GeoChip 5.0. *Msphere*, 7(1), e00936-21. doi: <https://doi.org/10.1128/msphere.00936-21>.
- 827 Meyer, F., Paarmann, D., D'Souza, M., Olson, R., Glass, E. M., Kubal, M., ... & Edwards, R. (2008).  
828 The metagenomics RAST server—a public resource for the automatic phylogenetic and functional  
829 analysis of metagenomes. *BMC bioinformatics*, 9(1), 1-8. doi:10.1186/1471-2105-9-386

- 830 Mnif, S., Chamkha, M., & Sayadi, S. (2009). Isolation and characterization of *Halomonas* sp. strain  
831 C2SS100, a hydrocarbon - degrading bacterium under hypersaline conditions. *Journal of applied*  
832 *microbiology*, 107(3), 785-794. doi:10.1111/j.1365-2672.2009.04251.x
- 833 Mohite, B. V., Koli, S. H., Narkhede, C. P., Patil, S. N., & Patil, S. V. (2017). Prospective of microbial  
834 exopolysaccharide for heavy metal exclusion. *Applied biochemistry and biotechnology*, 183(2),  
835 582-600. doi:10.1007/s12010-017-2591-4.
- 836 Montuori, P., & Triassi, M. (2012). Polycyclic aromatic hydrocarbons loads into the Mediterranean  
837 Sea: Estimate of Sarno River inputs. *Marine pollution bulletin*, 64(3), 512-520.  
838 doi:10.1016/j.marpolbul.2012.01.003.
- 839 Montuori, P., Lama, P., Aurino, S., Naviglio, D., & Triassi, M. (2013). Metals loads into the  
840 Mediterranean Sea: estimate of Sarno River inputs and ecological risk. *Ecotoxicology*, 22(2), 295-  
841 307. doi:10.1007/s10646-012-1026-9.
- 842 Moody, J. D., Freeman, J. P., & Cerniglia, C. E. (2005). Degradation of benz [a] anthracene by  
843 *Mycobacterium vanbaalenii* strain PYR-1. *Biodegradation*, 16(6), 513-526. doi:10.1007/s10532-  
844 004-7217-1.
- 845 Moron, M. S., Depierre, J. W., & Mannervik, B. (1979). Levels of glutathione, glutathione reductase  
846 and glutathione S-transferase activities in rat lung and liver. *Biochimica et biophysica acta (BBA)-*  
847 *general subjects*, 582(1), 67-78. doi:10.1016/0304-4165(79)90289-7
- 848 Murugavelh, S., & Mohanty, K. (2012). Bioreduction of hexavalent chromium by free cells and cell  
849 free extracts of *Halomonas* sp. *Chemical engineering journal*, 203, 415-422.  
850 doi:10.1016/j.cej.2012.07.069
- 851 Nagata, S., Ii, K. M., Tsukimi, T., Miura, M. C., Galipon, J., & Arakawa, K. (2019). Complete  
852 genome sequence of *Halomonas olivaria*, a moderately halophilic bacterium isolated from olive  
853 processing effluents, obtained by nanopore sequencing. *Microbiology Resource Announcements*,  
854 8(18), e00144-19. doi:10.1128/mra.00144-19.
- 855 Nelson, D. R. (2018). Cytochrome P450 diversity in the tree of life. *Biochimica et Biophysica Acta*  
856 *(BBA)-Proteins and Proteomics*, 1866(1), 141-154. doi:10.1016/j.bbapap.2017.05.003.
- 857 Nešvera, J., Rucká, L., & Pátek, M. (2015). Catabolism of phenol and its derivatives in bacteria:  
858 genes, their regulation, and use in the biodegradation of toxic pollutants. *Advances in applied*  
859 *microbiology*, 93, 107-160. doi:10.1016/bs.aambs.2015.06.002.

- 860 Nwuche, C. O., & Ugoji, E. O. (2008). Effects of heavy metal pollution on the soil microbial activity.  
861 International Journal of Environmental Science & Technology, 5(3), 409-414.  
862 doi:10.1007/BF03326036.
- 863 Nzila, A., Musa, M. M., Sankara, S., Al-Momani, M., Xiang, L., & Li, Q. X. (2021). Degradation of  
864 benzo [a] pyrene by halophilic bacterial strain *Staphylococcus haemolyticus* strain 10SBZ1A. PloS  
865 one, 16(2), e0247723. doi:10.1371/journal.pone.0247723.
- 866 Overbeek, R., Olson, R., Pusch, G. D., Olsen, G. J., Davis, J. J., Disz, T., ... & Stevens, R. (2014).  
867 The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology  
868 (RAST). Nucleic acids research, 42(D1), D206-D214. doi:10.1093/nar/gkt1226.
- 869 Oyetibo, G. O., Miyauchi, K., Huang, Y., Chien, M. F., Ilori, M. O., Amund, O. O., & Endo, G.  
870 (2017). Biotechnological remedies for the estuarine environment polluted with heavy metals and  
871 persistent organic pollutants. International Biodeterioration & Biodegradation, 119, 614-625.  
872 doi:10.1016/j.ibiod.2016.10.005.
- 873 Park, J. W., & Crowley, D. E. (2006). Dynamic changes in *nahAc* gene copy numbers during  
874 degradation of naphthalene in PAH-contaminated soils. Applied microbiology and biotechnology,  
875 72(6), 1322-1329. doi:10.1007/s00253-006-0423-5.
- 876 Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P., & Tyson, G. W. (2015). CheckM:  
877 assessing the quality of microbial genomes recovered from isolates, single cells, and  
878 metagenomes. Genome research, 25(7), 1043-1055. doi:10.1101/gr.186072.114.
- 879 Patlolla, A. K., Barnes, C., Yedjou, C., Velma, V. R., & Tchounwou, P. B. (2009). Oxidative stress,  
880 DNA damage, and antioxidant enzyme activity induced by hexavalent chromium in Sprague -  
881 Dawley rats. Environmental Toxicology: An International Journal, 24(1), 66-73.  
882 doi:10.1002/tox.20395.
- 883 Peng, R. H., Xiong, A. S., Xue, Y., Fu, X. Y., Gao, F., Zhao, W., ... & Yao, Q. H. (2008). Microbial  
884 biodegradation of polyaromatic hydrocarbons. FEMS microbiology reviews, 32(6), 927-955.  
885 doi:10.1111/j.1574-6976.2008.00127.x.
- 886 Peng, T., Luo, A., Kan, J., Liang, L., Huang, T., & Hu, Z. (2018). Identification of a ring-  
887 hydroxylating dioxygenases capable of anthracene and benz [a] anthracene oxidization from  
888 *Rhodococcus* sp. P14. Microbial Physiology, 28(4), 183-189. doi:10.1159/000494384.

- 889 Pennafirme, S., Lima, I., Bitencourt, J. A., Crapez, M. A. C., & Lopes, R. T. (2015). Microbial biofilm  
890 study by synchrotron X-ray microscopy. *Radiation Physics and Chemistry*, 116, 116-119.  
891 doi:10.1016/j.radphyschem.2015.05.040
- 892 Poli, A., Esposito, E., Orlando, P., Lama, L., Giordano, A., de Appolonia, F., ... & Gambacorta, A.  
893 (2007). *Halomonas alkaliantarctica* sp. nov., isolated from saline lake Cape Russell in Antarctica,  
894 an alkalophilic moderately halophilic, exopolysaccharide-producing bacterium. *Systematic and*  
895 *applied microbiology*, 30(1), 31-38. doi:10.1016/j.syapm.2006.03.003
- 896 Priyadarshane, M., & Das, S. (2021). Biosorption and removal of toxic heavy metals by metal  
897 tolerating bacteria for bioremediation of metal contamination: A comprehensive review. *Journal*  
898 *of Environmental Chemical Engineering*, 9(1), 104686. doi:10.1016/j.jece.2020.104686
- 899 Quiton, K. G., Doma Jr, B., Futralan, C. M., & Wan, M. W. (2018). Removal of chromium (VI) and  
900 zinc (II) from aqueous solution using kaolin-supported bacterial biofilms of Gram-negative *E. coli*  
901 and Gram-positive *Staphylococcus epidermidis*. *Sustainable Environment Research*, 28(5), 206-  
902 213. doi:10.1016/j.serj.2018.04.002.
- 903 Rahul, K., Sasikala, C., Tushar, L., Debadrita, R., & Ramana, C. V. (2014). *Alcanivorax xenomutans*  
904 sp. nov., a hydrocarbonoclastic bacterium isolated from a shrimp cultivation pond. *International*  
905 *journal of systematic and evolutionary microbiology*, 64, 3553-3558. doi:10.1099/ijs.0.061168-0.
- 906 Ramasamy, K. P., Rajasabapathy, R., Lips, I., Mohandass, C., & James, R. A. (2020). Genomic  
907 features and copper biosorption potential of a new *Alcanivorax* sp. VBW004 isolated from the  
908 shallow hydrothermal vent (Azores, Portugal). *Genomics*, 112(5), 3268-3273.  
909 doi:10.1016/j.ygeno.2020.06.015
- 910 Rodriguez-Caballero, A., Ramond, J. B., Welz, P. J., Cowan, D. A., Odlare, M., & Burton, S. G.  
911 (2012). Treatment of high ethanol concentration wastewater by biological sand filters: Enhanced  
912 COD removal and bacterial community dynamics. *Journal of environmental management*, 109,  
913 54-60. doi:10.1016/j.jenvman.2012.05.005.
- 914 Rodriguez-R, L. M., & Konstantinidis, K. T. (2016). The enveomics collection: a toolbox for  
915 specialized analyses of microbial genomes and metagenomes (No. e1900v1). *PeerJ Preprints*.  
916 doi:10.7287/peerj.preprints.1900.
- 917 Rosen, B. P., & Liu, Z. (2009). Transport pathways for arsenic and selenium: a minireview.  
918 *Environment international*, 35(3), 512-515. doi:10.1016/j.envint.2008.07.023

- 919 Sahmoune, M. N. (2018). Performance of *Streptomyces rimosus* biomass in biosorption of heavy  
920 metals from aqueous solutions. *Microchemical Journal*, 141, 87-95.  
921 doi:10.1016/j.microc.2018.05.009
- 922 Salam, L. B., & Ishaq, A. (2019). Biostimulation potentials of corn steep liquor in enhanced  
923 hydrocarbon degradation in chronically polluted soil. *3 Biotech*, 9(2), 1-20. doi:10.1007/s13205-  
924 019-1580-4.
- 925 Saravanan, A., Kumar, P. S., Vo, D. V. N., Jeevanantham, S., Karishma, S., & Yaashikaa, P. R.  
926 (2021). A review on catalytic-enzyme degradation of toxic environmental pollutants: Microbial  
927 enzymes. *Journal of Hazardous Materials*, 419, 126451. doi:10.1016/j.jhazmat.2021.126451
- 928 Schneiker, S., Dos Santos, V. A., Bartels, D., Bekel, T., Brecht, M., Buhrmester, J., ... & Golyshin,  
929 P. N. (2006). Genome sequence of the ubiquitous hydrocarbon-degrading marine bacterium  
930 *Alcanivorax borkumensis*. *Nature biotechnology*, 24(8), 997-1004. doi:10.1038/nbt1232
- 931 Sharma, P., Singh, S. P., Iqbal, H. M., & Tong, Y. W. (2022). Omics approaches in bioremediation  
932 of environmental contaminants: An integrated approach for environmental safety and  
933 sustainability. *Environmental Research*, 211, 113102. doi:10.1016/j.envres.2022.113102
- 934 Sohn, J. H., Kwon, K. K., Kang, J. H., Jung, H. B., & Kim, S. J. (2004). *Novosphingobium*  
935 *pentaromativorans* sp. nov., a high-molecular-mass polycyclic aromatic hydrocarbon-degrading  
936 bacterium isolated from estuarine sediment. *International journal of systematic and evolutionary*  
937 *microbiology*, 54(5), 1483-1487. doi:10.1099/ijms.0.02945-0.
- 938 Sullivan, M. J., Petty, N. K., & Beatson, S. A. (2011). Easyfig: a genome comparison visualizer.  
939 *Bioinformatics*, 27(7), 1009-1010. doi:10.1093/bioinformatics/btr039.
- 940 Sutton, D. J., Tchounwou, P. B., Ninashvili, N., & Shen, E. (2002). Mercury Induces Cytotoxicity  
941 Transcriptionally Activates Stress Genes in Human Liver Carcinoma (HepG2) Cells. *International*  
942 *Journal of Molecular Sciences*, 3(9), 965-984. doi:10.3390/i3090965.
- 943 Tashla, T., Prodanovic, R., Boskovic, J., Zuza, M., Solesa, D., Ljubojevic, D., & Puvaca, N. (2018).  
944 Persistent organic pollutants and heavy metals and the importance of fish as a bio-indicator of  
945 environmental pollution. *Concepts of Dairy & Veterinary Sciences*, 2(2), 168-70.  
946 doi:10.32474/cdvs.2018.02.000131.
- 947 Tchounwou, P. B., Ishaque, A. B., & Schneider, J. (2001). Cytotoxicity and transcriptional activation  
948 of stress genes in human liver carcinoma cells (HepG2) exposed to cadmium chloride. *Molecular*  
949 *and Cellular Biochemistry*, 222(1), 21-28. doi:10.1023/A:1017922114201.

- 950 Thavamani, P., Megharaj, M., & Naidu, R. (2012). Bioremediation of high molecular weight  
951 polyaromatic hydrocarbons co-contaminated with metals in liquid and soil slurries by metal  
952 tolerant PAHs degrading bacterial consortium. *Biodegradation*, 23(6), 823-835.  
953 doi:10.1007/s10532-012-9572-7.
- 954 Thompson, I. P., Van Der Gast, C. J., Ciric, L., & Singer, A. C. (2005). Bioaugmentation for  
955 bioremediation: the challenge of strain selection. *Environmental Microbiology*, 7(7), 909-915.  
956 doi:10.1111/j.1462-2920.2005.00804.x.
- 957 Tsurumaki, M., Deno, S., Galipon, J., & Arakawa, K. (2019). Complete genome sequence of  
958 halophilic deep-sea bacterium *Halomonas axialensis* strain Althf1. *Microbiology Resource*  
959 *Announcements*, 8(31), e00839-19. doi:10.1128/MRA.00839-19.
- 960 Tyagi, M., da Fonseca, M. M. R., & de Carvalho, C. C. (2011). Bioaugmentation and biostimulation  
961 strategies to improve the effectiveness of bioremediation processes. *Biodegradation*, 22(2), 231-  
962 241. doi:10.1007/s10532-010-9394-4.
- 963 Vaidya, S., Jain, K., & Madamwar, D. (2017). Metabolism of pyrene through phthalic acid pathway  
964 by enriched bacterial consortium composed of *Pseudomonas*, *Burkholderia*, and *Rhodococcus*  
965 (PBR). *3 Biotech*, 7(1), 1-15. doi:10.1007/s13205-017-0598-8.
- 966 Vaidya, S., Devpura, N., Jain, K., & Madamwar, D. (2018). Degradation of chrysene by enriched  
967 bacterial consortium. *Frontiers in microbiology*, 9, 1333. doi:10.3389/fmicb.2018.01333.
- 968 Verma, S., & Kuila, A. (2019). Bioremediation of heavy metals by microbial process. *Environmental*  
969 *Technology & Innovation*, 14, 100369. doi:10.1016/j.eti.2019.100369.
- 970 Wang, D., Qin, L., Liu, E., Chai, G., Su, Z., Shan, J., ... & Lin, Y. (2021). Biodegradation performance  
971 and diversity of enriched bacterial consortia capable of degrading high-molecular-weight  
972 polycyclic aromatic hydrocarbons. *Environmental Technology*, 1-12.  
973 doi:10.1080/09593330.2021.1946163.
- 974 Williamson, A., De Santi, C., Altermark, B., Karlsen, C., & Hjerde, E. (2016). Complete genome  
975 sequence of *Halomonas* sp. R5-57. *Standards in Genomic Sciences*, 11(1), 1-9.  
976 doi:10.1186/s40793-016-0192-4.
- 977 Wongwongsee, W., Chareanpat, P., & Pinyakong, O. (2013). Abilities and genes for PAH  
978 biodegradation of bacteria isolated from mangrove sediments from the central of Thailand. *Marine*  
979 *pollution bulletin*, 74(1), 95-104. doi:10.1016/j.marpolbul.2013.07.025.

- 980 Yan, S., & Wu, G. (2017). Reorganization of gene network for degradation of polycyclic aromatic  
981 hydrocarbons (PAHs) in *Pseudomonas aeruginosa* PAO1 under several conditions. *Journal of*  
982 *Applied Genetics*, 58(4), 545-563. doi:10.1007/s13353-017-0402-9.
- 983 Yedjou, C. G., & Tchounwou, P. B. (2007). N-acetyl-l-cysteine affords protection against lead-  
984 induced cytotoxicity and oxidative stress in human liver carcinoma (HepG2) cells. *International*  
985 *journal of environmental research and public health*, 4(2), 132-137.  
986 doi:10.3390/ijerph2007040007.
- 987 Yedjou, C. G., & Tchounwou, P. B. (2007). In-vitro cytotoxic and genotoxic effects of arsenic  
988 trioxide on human leukemia (HL-60) cells using the MTT and alkaline single cell gel  
989 electrophoresis (Comet) assays. *Molecular and cellular biochemistry*, 301(1), 123-130.  
990 doi:10.1007/s11010-006-9403-4.
- 991 Yeom, S. J., Le, T. K., & Yun, C. H. (2021). P450-driven plastic-degrading synthetic bacteria. *Trends*  
992 *in Biotechnology*. doi:10.1016/j.tibtech.2021.06.003.
- 993 Yu, S. H., Ke, L., Wong, Y. S., & Tam, N. F. Y. (2005). Degradation of polycyclic aromatic  
994 hydrocarbons by a bacterial consortium enriched from mangrove sediments. *Environment*  
995 *International*, 31(2), 149-154. doi:10.1016/j.envint.2004.09.008
- 996 Zhang, Y., & Skolnick, J. (2004). Scoring function for automated assessment of protein structure  
997 template quality. *Proteins: Structure, Function, and Bioinformatics*, 57(4), 702-710.  
998 doi:10.1002/prot.20264.
- 999 Zhang, Y., Lin, D. F., Hao, J., Zhao, Z. H., & Zhang, Y. J. (2020). The crucial role of bacterial  
1000 laccases in the bioremediation of petroleum hydrocarbons. *World Journal of Microbiology and*  
1001 *Biotechnology*, 36(8), 1-10.
- 1002 Zouboulis, A. I., Loukidou, M. X., & Matis, K. A. (2004). Biosorption of toxic metals from aqueous  
1003 solutions by bacteria strains isolated from metal-polluted soils. *Process biochemistry*, 39(8), 909-  
1004 916. doi:10.1016/S0032-9592(03)00200-0.

1005 **Figure Legends**

1006 **Figure 1.** Map of the study area with the position of the sampling site at Sarno River Mouth  
1007 (Tyrrhenian Sea; 40°43'42.01"N, 14°28'0.45"E).

1008

1009 **Figure 2A-B.** Taxonomic and functional annotation of the enriched metagenome. A) taxonomic  
1010 classification at the genus level based on M5NR database (standard cutoff: alignment length 15bp; e-  
1011 value  $e^{-5}$ ; id. 60%). B) functional classification based on Subsystems ontology (level 1; standard  
1012 cutoff: alignment length 15bp; e-value  $e^{-5}$ ; id. 60%), with functions putatively related to hydrocarbon  
1013 degradation and to resistance/interaction with heavy metals contained in the subsystems “stress  
1014 response” and “metabolism of aromatic compounds” subsystems.

1015

1016 **Figure 3A-B.** Removal of PAHs and HMs by the two novel bacterial strains isolated from the  
1017 enriched metagenome. A) Degradation of PAHs (naphthalene, phenanthrene and pyren) under control  
1018 (no bacteria) and treated conditions (i.e., addition of *A. xenomutans* strain SRM1 or *H.*  
1019 *alkaliantarctica* strain SRM2). B) Removal of HMs (As, Cd, Cu, Zn, Pb) under control (no bacteria)  
1020 and treated conditions (i.e., addition of *A. xenomutans* strain SRM1 or *H. alkaliantarctica* strain  
1021 SRM2). Reported are average values and standard deviations.

1022

1023 **Figure 4A-D.** Comparative genomics of genetic regions for peripheral pathways for hydrocarbon  
1024 degradation with ring hydroxylating dioxygenases enzymes (RHDs). A) Comparison of contig 9  
1025 from *A. xenomutans* strain SRM1 MAG (containing the ORF 4 encoding for RHD) with three closest  
1026 sequences from the NCBI database. In the right, the complete list of genes encoded within contig 9  
1027 (ORFs 1-10) and those encoded by the 3 other reference genomes (ORFs 11-29). B) Comparison of  
1028 contig 29 from *H. alkaliantarctica* strain SRM2 MAG (containing the ORF 2 encoding for RHD)  
1029 with three closest sequences from the NCBI database. In the right, the complete list of genes encoded  
1030 within contig 29 (ORFs 1-17) and those encoded by the 3 other reference genomes (ORFs 18-30). C)  
1031 Superimposition of RHD of *H. alkaliantarctica* strain SRM2 (in red) with naphthalene 1,2-  
1032 dioxygenase crystal structure from *Pseudomonas* sp. strain C18 (in blue)  
1033 [[www.rcsb.org/structure/2hmm](http://www.rcsb.org/structure/2hmm)]. D) Superimposition of RHD of *A. xenomutans* strain SRM1 (in red)  
1034 with naphthalene 1,2-dioxygenase crystal structure from *Rhodococcus* sp. NCIMB 12038 (in blue)  
1035 [[www.rcsb.org/structure/2b1x](http://www.rcsb.org/structure/2b1x)].

1036



1037 **Figure 5A-B.** Comparative genomics of genetic regions for peripheral pathways for hydrocarbon  
1038 degradation with cytochrome P450 (CYP450). A) Comparison of contig 76 from *A. xenomutans* strain  
1039 SRM1 MAG (containing the ORF 12 encoding for CYP450) with three closest sequences from the  
1040 NCBI database. In the right, the complete list of genes encoded within contig 76 (ORFs 1-20) and  
1041 those encoded by the 3 other reference genomes (ORFs 21-29). B) Comparison of contig 15 from *H.*  
1042 *alkaliantarctica* strain SRM2 MAG (containing the ORF 6 encoding for CYP450) with three closest  
1043 sequences from the NCBI database. In the right, the complete list of genes encoded within contig 15  
1044 (ORFs 1-14) and those encoded by the 3 other reference genomes (ORFs 15-27).

1045

1046 **Figure 6A-B.** Comparative genomics of genetic regions for central pathways for hydrocarbon  
1047 degradation with genes for the homoprotocatechuate pathway. A) Comparison of contig 65 from *A.*  
1048 *xenomutans* strain SRM1 MAG (containing the ORF 7-17 encoding for the genes of the  
1049 homoprotocatechuate pathway) with three closest sequences from the NCBI database. In the right,  
1050 the complete list of genes encoded within contig 65 (ORFs 1-20) and those encoded by the 3 other  
1051 reference genomes (ORFs 21-38). B) Comparison of contig 29 from *H. alkaliantarctica* strain SRM2  
1052 MAG (containing the ORF 4-14 encoding for the genes of the homoprotocatechuate pathway) with  
1053 three closest sequences from the NCBI database. In the right, the complete list of genes encoded  
1054 within contig 29 (ORFs 1-17) and those encoded by the 3 other reference genomes (ORFs 18-47).

1055

1056 **Figure 7A-C.** Comparative genomics of genetic regions for heavy metal removal/detoxification. A)  
1057 Comparison of contig 20 from *A. xenomutans* strain SRM1 MAG (containing the ORFs 11-18  
1058 encoding the arsenic resistance operon-like genomic region) with three closest sequences from the  
1059 NCBI database. In the right, the complete list of genes encoded within contig 20 (ORFs 1-18) and  
1060 those encoded by the 3 other reference genomes (ORFs 19-24). B) Comparison of contig 68 from *H.*  
1061 *alkaliantarctica* strain SRM2 MAG (containing the ORFs 2-12 encoding the arsenic resistance  
1062 operon-like genomic region) with three closest sequences from the NCBI database. In the right, the  
1063 complete list of genes encoded within contig 68 (ORFs 1-15) and those encoded by the 3 other  
1064 reference genomes (ORFs 16-29). C) Comparison of contig 30 from *H. alkaliantarctica* strain SRM2  
1065 MAG (containing the ORFs 6-11 encoding the mercury-resistance operon) with three closest  
1066 sequences from the NCBI database. In the right, the complete list of genes encoded within contig 30  
1067 (ORFs 1-15) and those encoded by the 3 other reference genomes (ORFs 16-23).



