

## Core stability and low-back pain: a causal fallacy

Augeard, Nathan; Carroll, Sean Paul

*Published in:*  
Journal of Exercise Rehabilitation

*DOI:*  
[10.12965/jer.1938198.099](https://doi.org/10.12965/jer.1938198.099)

*Publication date:*  
2019

*Document Version*  
Publisher's PDF, also known as Version of record

[Link to publication in ResearchOnline](#)

*Citation for published version (Harvard):*  
Augeard, N & Carroll, SP 2019, 'Core stability and low-back pain: a causal fallacy', *Journal of Exercise Rehabilitation*, vol. 15, no. 3, pp. 493-495. <https://doi.org/10.12965/jer.1938198.099>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

## Core stability and low-back pain: a causal fallacy

Nathan Augeard\*, Sean Paul Carroll

School of Health and Life Sciences, Glasgow Caledonian University, Glasgow, UK


Dear Editor,

We would firstly like to congratulate Seo and Kim on the publication of their research article “*The effects of Gyrotonic expansion system exercise and trunk stability exercise on muscle activity and lumbar stability for the subjects with chronic low-back pain.*” However, we find some of the claims made by the authors to be unsubstantiated and reductionist of the complex biopsychosocial phenomenon which is chronic low-back pain. We feel that the methods utilised and the discussion that followed largely did not see the forest for the trees they were looking at.

Firstly, the authors state that back pain refers to “pain around the lumbar vertebrae” and do not use appropriate evidence to support their claim that chronic low-back pain is caused by posture or “imbalanced and weak muscles.” Evidence clearly suggests that the majority of the low-back pain presentation is nonspecific, meaning that it is not possible to identify a single origin of the patient’s symptoms (Deyo and Weinstein, 2001; Hartvigsen et al., 2018; Koes et al., 2006; Maher et al., 2017). Therefore, stating that posture is the cause of chronic low-back pain is, at best, reductionist and not evidence-based. At worse, it promotes a narrative potentially harmful to patients (Darlow, 2016; Darlow et al., 2013). High-quality evidence suggests a clear lack of association between incidence of low-back pain and awkward postures (Roffey et al., 2010), physical features such as facet joint degeneration (Hartvigsen et al., 2018; Maas et al., 2017) or magnetic resonance imaging changes (Brinjikji et al., 2015; Steffens et al., 2014; Tonosu et al., 2017). It could be argued that the increased rates of structural failure observed *in vitro* models within the “neutral” zone (Gooyers and Callaghan, 2015) support a biomedical

approach to low-back pain. However, these changes have been deemed largely unavoidable (Arjmand and Shirazi-Adl, 2005). Furthermore, the validity of cadaveric *in vitro* models to humans is a debate in itself, largely opposed by the high variability and the adaptability of the body. While high loads applied in end-range could potentially lead to structural changes, these changes are rarely directly related to pain (Brinjikji et al., 2015; Steffens et al., 2014; Tonosu et al., 2017). It is more scientifically accurate to think of posture change as a symptom modification (Lehman, 2018) rather than a direct treatment of a disease, of which low back pain is rarely (Deyo and Weinstein, 2001; Waddell, 1987).

The literature supports the premise of the intervention so far as that exercise is beneficial for low back pain (Gordon et al., 2016). However, the claim that core stability exercise is one of the most effective options for low-back pain is without merit and clearly lacks supporting evidence. It has been demonstrated thoroughly that in those with low-back pain there is no long term difference between different exercise modalities including different modes of “core stability” (Michaelson et al., 2016; Saragiotto et al., 2016; Shnayderman and Katz-Leurer, 2013; Smith et al., 2014; Vasseljen et al., 2012; Wang et al., 2012). Perhaps their utilisation may be more effective acutely. However, acute response to physiotherapy treatment is a complex phenomenon, which could be influenced by many nonspecific factors (e.g., therapeutic alliance or patient expectations) and specific factors (e.g., symptom modification or neurophysiological changes) (Bialosky et al., 2009; Bjorbaekmo and Mengshoel, 2016; Kinney et al., 2018; O’Keeffe et al., 2016). Moreover, core activation may not even vary between symptomatic and healthy asymptomatic people (Gorbet et al.,

\*Corresponding author: Nathan Augeard  <https://orcid.org/0000-0002-9154-7233>  
School of Health and Life Sciences, Glasgow Caledonian University,  
Cowcaddens Road, Glasgow G4 0BA, UK  
E-mail: [naugea200@caledonian.ac.uk](mailto:naugea200@caledonian.ac.uk)  
Received: April 11, 2019 / Accepted: April 19, 2019

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

2010), nor can core stability changes predict improvements in disability following specific exercise (Mannion et al., 2012). Strength training can be beneficial but the larger effect sizes are noted only for full body programmes (Searle et al., 2015), and given that strength training is a complex intervention that has multiple biological and psychosocial benefits, further work is required before we can make any assumptions about cause and effect (Jay et al., 2015; Pedersen and Febbraio, 2008). Therefore, it would appear that the common denominators at present for improvements in the literature were seeking or receiving care for low-back pain, and an increase in activity (Alzahrani et al., 2019). This is hardly surprising when we consider the overall literature suggesting that systemic markers (Klyne et al., 2018), sleep, physical activity, diet (Yang and Haldeman, 2018), depression and mental health (Jarvik et al., 2005; Klyne et al., 2018; Pinheiro et al., 2015) are all partial mediators of development and prognosis. This is confounded further by the comparison of Gyrotonic exercise to another treatment and not including an appropriate control group. As highlighted earlier, there is a wide range of good-quality literature to support the complex and biopsychosocial nature of chronic low-back pain beyond “neutral” posture. Relying on a biomedical paradigm to explain a complex biopsychosocial problem is inadequate (Engel, 1977; Waddell, 1987).

Finally, several of the claims made with regards to the Gyrotonic exercise are unsubstantiated. For example, the statements “these natural movements improve muscle strength and flexibility along with breathing and collaborative muscle contraction” and “the arch and curl motion has been considered to create a connection between the erector spinae and the external oblique” are used to develop a discussion regarding the importance of these exercises for chronic low-back pain. However, the authors failed to provide any credible evidence to support their claims. Only one reference was used to support the first statement (Campbell and Miles, 2006). Upon reading this article, it rapidly becomes apparent that the authors only offer a description of the method, with no evidence or data to support their claims (Campbell and Miles, 2006). As per Hitchen’s razor: what can be asserted without evidence can be dismissed without evidence.

In conclusion, we believe that the authors have failed to consider the wider literature and make bold claims without considering the evidence-based biopsychosocial model within their study. As humans, we are all prone to confirmation bias; however, as scientists we have a unique responsibility to display clinical equipoise and follow systematic processes in methodology and reporting to progress our field forward on stable foundations.

## CONFLICT OF INTEREST

No potential conflict of interest relevant to this article was reported.

## REFERENCES

- Alzahrani H, Shirley D, Cheng SW, Mackey M, Stamatakis E. Physical activity and chronic back conditions: A population-based pooled study of 60,134 adults. *J Sport Heal Sci* 2019 January 11 [Epub]. <https://doi.org/10.1016/j.jsjshs.2019.01.003>
- Arjmand N, Shirazi-Adl A. Biomechanics of changes in lumbar posture in static lifting. *Spine (Phila Pa 1976)* 2005;30:2637-2648.
- Bialosky JE, Bishop MD, Price DD, Robinson ME, George SZ. The mechanisms of manual therapy in the treatment of musculoskeletal pain: a comprehensive model. *Man Ther* 2009;14:531-538.
- Bjorbaekmo WS, Mengshoel AM. “A touch of physiotherapy” - the significance and meaning of touch in the practice of physiotherapy. *Physiother Theory Pract* 2016;32:10-19.
- Brinjikji W, Luetmer PH, Comstock B, Bresnahan BW, Chen LE, Deyo RA, Halabi S, Turner JA, Avins AL, James K, Wald JT, Kallmes DF, Jarvik JG. Systematic literature review of imaging features of spinal degeneration in asymptomatic populations. *AJNR Am J Neuroradiol* 2015;36:811-816.
- Campbell J, Miles W. Analyzing the Gyrotonic® arch and curl. *J Bodyw Mov Ther* 2006;10:147-153.
- Darlow B. Beliefs about back pain: the confluence of client, clinician and community. *Int J Osteopath Med* 2016;20:53-61.
- Darlow B, Dowell A, Baxter GD, Mathieson F, Perry M, Dean S. The enduring impact of what clinicians say to people with low back pain. *Ann Fam Med* 2013;11:527-534.
- Deyo RA, Weinstein JN. Low back pain. *N Engl J Med* 2001;344:363-370.
- Engel GL. The need for a new medical model: a challenge for biomedicine. *Science* 1977;196:129-136.
- Gooyers CE, Callaghan JP. Exploring interactions between force, repetition and posture on intervertebral disc height loss and bulging in isolated porcine cervical functional spinal units from sub-acute-failure magnitudes of cyclic compressive loading. *J Biomech* 2015;48:3701-3708.
- Gorbet N, Selkow NM, Hart JM, Saliba S. No Difference in Transverse Abdominis Activation Ratio between Healthy and Asymptomatic Low Back Pain Patients during Therapeutic Exercise. *Rehabil Res Pract* 2010;2010:459738.
- Gordon R, Bloxham S. A systematic review of the effects of exercise and physical activity on non-specific chronic low back pain. *Healthcare*

- (Basel) 2016;4(2):22.
- Hartvigsen J, Hancock MJ, Kongsted A, Louw Q, Ferreira ML, Genevay S, Hoy D, Karppinen J, Pransky G, Sieper J, Smeets RJ, Underwood M; Lancet Low Back Pain Series Working Group. What low back pain is and why we need to pay attention. *Lancet* 2018;391:2356-2367.
- Jarvik JG, Hollingworth W, Heagerty PJ, Haynor DR, Boyko EJ, Deyo RA. Three-year incidence of low back pain in an initially asymptomatic cohort: clinical and imaging risk factors. *Spine (Phila Pa 1976)* 2005;30:1541-1548.
- Jay K, Brandt M, Hansen K, Sundstrup E, Jakobsen MD, Schraefel MC, Sjøgaard G, Andersen LL. Effect of individually tailored biopsychosocial workplace interventions on chronic musculoskeletal pain and stress among laboratory technicians: Randomized controlled trial. *Pain Physician* 2015;18:459-471.
- Kinney M, Seider J, Beaty AF, Coughlin K, Dyal M, Clewley D. The impact of therapeutic alliance in physical therapy for chronic musculoskeletal pain: A systematic review of the literature. *Physiother Theory Pract* 2018 Sep 28:1-13 [Epub]. <https://doi.org/10.1080/09593985.2018.1516015>.
- Klyne DM, Barbe MF, van den Hoorn W, Hodges PW. ISSLS PRIZE IN CLINICAL SCIENCE 2018: longitudinal analysis of inflammatory, psychological, and sleep-related factors following an acute low back pain episode-the good, the bad, and the ugly. *Eur Spine J* 2018;27:763-777.
- Koes BW, van Tulder MW, Thomas S. Diagnosis and treatment of low back pain. *BMJ* 2006;332:1430-1434.
- Lehman GJ. The Role and Value of Symptom-Modification Approaches in Musculoskeletal Practice. *J Orthop Sports Phys Ther* 2018;48:430-435.
- Maas ET, Juch JN, Ostelo RW, Groeneweg JG, Kallewaard JW, Koes BW, Verhagen AP, Huygen FJ, van Tulder MW. Systematic review of patient history and physical examination to diagnose chronic low back pain originating from the facet joints. *Eur J Pain* 2017;21:403-414.
- Maher C, Underwood M, Buchbinder R. Non-specific low back pain. *Lancet* 2017;389:736-747.
- Mannion AF, Caporaso F, Pulkovski N, Sprott H. Spine stabilisation exercises in the treatment of chronic low back pain: a good clinical outcome is not associated with improved abdominal muscle function. *Eur Spine J* 2012;21:1301-1310.
- Michaelson P, Holmberg D, Aasa B, Aasa U. High load lifting exercise and low load motor control exercises as interventions for patients with mechanical low back pain: A randomized controlled trial with 24-month follow-up. *J Rehabil Med* 2016;48:456-463.
- O'Keefe M, Cullinane P, Hurley J, Leahy I, Bunzli S, O'Sullivan PB, O'Sullivan K. What influences patient-therapist interactions in musculoskeletal physical therapy? Qualitative systematic review and meta-synthesis. *Phys Ther* 2016;96:609-622.
- Pedersen BK, Febbraio MA. Muscle as an endocrine organ: focus on muscle-derived interleukin-6. *Physiol Rev* 2008;88:1379-1406.
- Pinheiro MB, Ferreira ML, Refshauge K, Ordoñana JR, Machado GC, Prado LR, Maher CG, Ferreira PH. Symptoms of depression and risk of new episodes of low back pain: A systematic review and meta-analysis. *Arthritis Care Res (Hoboken)* 2015;67:1591-1603.
- Roffey DM, Wai EK, Bishop P, Kwon BK, Dagenais S. Causal assessment of awkward occupational postures and low back pain: results of a systematic review. *Spine J* 2010;10:89-99.
- Saragiotto BT, Maher CG, Yamato TP, Costa LO, Costa LC, Ostelo RW, Macedo LG. Motor control exercise for nonspecific low back pain: A cochrane review. *Spine (Phila Pa 1976)* 2016;41:1284-1295.
- Searle A, Spink M, Ho A, Chuter V. Exercise interventions for the treatment of chronic low back pain: a systematic review and meta-analysis of randomised controlled trials. *Clin Rehabil* 2015;29:1155-1167.
- Shnayderman I, Katz-Leurer M. An aerobic walking programme versus muscle strengthening programme for chronic low back pain: a randomized controlled trial. *Clin Rehabil* 2013;27:207-214.
- Smith BE, Littlewood C, May S. An update of stabilisation exercises for low back pain: a systematic review with meta-analysis. *BMC Musculoskelet Disord* 2014;15:416.
- Steffens D, Hancock MJ, Maher CG, Williams C, Jensen TS, Latimer J. Does magnetic resonance imaging predict future low back pain? A systematic review. *Eur J Pain* 2014;18:755-765.
- Tonosu J, Oka H, Higashikawa A, Okazaki H, Tanaka S, Matsudaira K. The associations between magnetic resonance imaging findings and low back pain: A 10-year longitudinal analysis. *PLoS One* 2017;12:e0188057.
- Vasseljen O, Unsgaard-Tøndel M, Westad C, Mork PJ. Effect of core stability exercises on feed-forward activation of deep abdominal muscles in chronic low back pain: a randomized controlled trial. *Spine (Phila Pa 1976)* 2012;37:1101-1108.
- Waddell G. 1987 Volvo award in clinical sciences. A new clinical model for the treatment of low-back pain. *Spine (Phila Pa 1976)* 1987;12:632-644.
- Wang XQ, Zheng JJ, Yu ZW, Bi X, Lou SJ, Liu J, Cai B, Hua YH, Wu M, Wei ML, Shen HM, Chen Y, Pan YJ, Xu GH, Chen PJ. A meta-analysis of core stability exercise versus general exercise for chronic low back pain. *PLoS One* 2012;7:e52082.
- Yang H, Haldeman S. Behavior-related factors associated with low back pain in the US adult population. *Spine (Phila Pa 1976)* 2018;43: 28-34.