

## Illinois Wesleyan University Digital Commons (a) IWU

John Wesley Powell Student Research Conference

1994, 5th Annual JWP Conference

Apr 23rd, 9:00 AM - 4:00 PM

Toward an Understanding of Alzheimer's Disease VI: A Comparison of the Effects of Bilateral Injections of  $\beta A$  (1-42) and  $\beta A$  (25-35) into Hippocampus on the Acquisition of a Spatial Task in the Male Rat

Steven M. Bond *Illinois Wesleyan University* 

Wayne A. Dornan, Faculty Advisor *Illinois Wesleyan University* 

Follow this and additional works at: http://digitalcommons.iwu.edu/jwprc

Steven M. Bond and Wayne A. Dornan, Faculty Advisor, "Toward an Understanding of Alzheimer's Disease VI: A Comparison of the Effects of Bilateral Injections of  $\beta A$  (1-42) and  $\beta A$  (25-35) into Hippocampus on the Acquisition of a Spatial Task in the Male Rat" (April 23, 1994). *John Wesley Powell Student Research Conference*. Paper 9. http://digitalcommons.iwu.edu/jwprc/1994/posters/9

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

## TOWARD AN UNDERSTANDING OF ALZHEIMER'S DISEASE VI: A COMPARISON OF THE EFFECTS OF BILATERAL INJECTIONS OF BA (1-42) AND BA (25-35) INTO HIPPOCAMPUS ON THE ACQUISITION OF A SPATIAL TASK IN THE MALE RAT

Steven M. Bond and Wayne A. Dornan\*, Department of Psychology, IWU

Alzheimer's disease (AD) is a neurodegenerative condition which presently affects more than 4 million Americans. Cognitively, AD manifests itself through gradual memory loss, learning disruption, and dementia. The neuropathology of AD is not completely understood though it is generally agreed that the neurons of the hippocampus are among the neuroanatomical structures selectively targeted in the disease. Extracellular neuritic plaques and intracellular neurofibrillary tangles have been identified as characteristic neurohistological features that may play a role in AD etiology. Accumulating evidence suggests that the aggregation of the 39-43 amino acid protein, beta amyloid (BA), may exhibit neurotoxicity either individually or synergistically with excitotoxins in the brain. Dornan, Kang, McCampbell, and Kang (Neuroreport, 1993) reported that bilateral injections of BA (25-35) in combination with a subthreshold dose of ibotenic acid (IBO) produced a dramatic disruption in acquisition of spatial learning in the rat. This deficit was accompanied by cell death in the hippocampus that mimicked the degeneration observed in AD. In that study, however, injections of BA (25-35) alone had no effect on the acquisition of spatial learning. At present, whether fragments of BA or the full protein produce neurotoxicity in neurons remains somewhat controversial. Therefore, in this study the effects of bilateral injections into the hippocampus of BA (1-42) on the acquisition of spatial learning in the rat were compared with BA (25-35), saline, and a subthreshold dose of ibotenic acid. Comparisons were made using a partially baited 8-arm radial arm maze. To assess the effects of these injections on spatial learning, the following parameters were recorded: session latency, latency to first choice, total number of choices, total correct choices, reference memory errors, working memory correct and incorrect errors, and choice accuracy. The results of this study will be presented at the conference.