



Illinois Wesleyan University Digital Commons @ IWU

John Wesley Powell Student Research
Conference

2004, 15th Annual JWP Conference

Apr 17th, 11:00 AM - 12:00 PM

Protection from the Effects of Ultraviolet Radiation by the Encapsulating Structures of Embryos of *Physa* SP., a Freshwater, Pulmonate Snail

Stacy L. Nielsen

Illinois Wesleyan University

Elizabeth Balsler, Faculty Advisor

Illinois Wesleyan University

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

Nielsen, Stacy L. and Balsler, Faculty Advisor, Elizabeth, "Protection from the Effects of Ultraviolet Radiation by the Encapsulating Structures of Embryos of *Physa* SP., a Freshwater, Pulmonate Snail" (2004). *John Wesley Powell Student Research Conference*. 3.

<http://digitalcommons.iwu.edu/jwprc/2004/oralpres3/3>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact digitalcommons@iwu.edu.

©Copyright is owned by the author of this document.

Oral Presentation O3.3

PROTECTION FROM THE EFFECTS OF ULTRAVIOLET RADIATION
BY THE ENCAPSULATING STRUCTURES OF EMBRYOS OF
PHYSA SP., A FRESHWATER, PULMONATE SNAIL

Stacy L. Nielsen and Elizabeth Balser*
Department of Biology, Illinois Wesleyan University

The genus *Physa* includes freshwater pulmonate snails that inhabit shallow environments well within depths penetrated by ultraviolet (UV) radiation. The distribution patterns of egg masses of *Physa* sp. indicate that the snails lay egg masses in sunlit areas and the masses are attached to rocks and debris such as leaves and twigs. Each mass consists of a viscous jelly covering that encases a variable number of embryos. Each of the embryos is individually surrounded by an egg capsule. Field-collected masses had debris and epifaunal organisms attached to the jelly. Preliminary studies of the effect of UV radiation on the developing embryos of these snails indicated that embryos surrounded by the capsule and jelly coat experienced no detriment from UV exposure. However, removal of the jelly coat from encapsulated embryos and subsequent exposure to UV radiation resulted in a >90% mortality. The UV absorption of the jelly coat was measured in both laboratory-raised and field-collected egg masses using a UV-Vis spectrophotometer. Absorption of energy by the jelly covering was between the wavelengths of 275-300 nm. This is consistent with absorption of UV-B radiation. The fact that no difference was detected in absorbance between field and laboratory egg masses suggests that the jelly is providing protection, rather than any organism or material adhered to the egg jelly.