



## Illinois Wesleyan University Digital Commons @ IWU

---

John Wesley Powell Student Research  
Conference

1998, 9th Annual JWP Conference

---

Apr 18th, 9:00 AM - 10:30 AM

# Synthesis of a New Family of Diamines: Components for Supramolecular Architectures

Jonathan T. Brockman  
*Illinois Wesleyan University*

Rebecca Roesner, Faculty Advisor  
*Illinois Wesleyan University*

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

---

Jonathan T. Brockman and Rebecca Roesner, Faculty Advisor, "Synthesis of a New Family of Diamines: Components for Supramolecular Architectures" (April 18, 1998). *John Wesley Powell Student Research Conference*. Paper 2.  
<http://digitalcommons.iwu.edu/jwprc/1998/posters/2>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.

## Poster Presentation 2

SYNTHESIS OF A NEW FAMILY OF DIAMINES:  
COMPONENTS FOR SUPRAMOLECULAR ARCHITECTURESJonathan T. Brockman and Rebecca Roesner\*  
Department of Chemistry, Illinois Wesleyan University

The practical limits of manufacturing smaller and smaller electronic components is fast approaching, and scientists have been exploring the use of molecular and macromolecular electronic devices. Supramolecular systems have been designed and demonstrated to function as molecular switches, molecular wires or photoelectric devices. One supramolecular structure is the rotaxane which is composed of a linear molecular string (linker) threaded through a macrocyclic molecular ring with bulky blocking groups attached to the ends of the linker to prevent dethreading. The current goal is the synthesis of a family of diamine linkers of various lengths that can be attached to the bulky blocking group. Synthesis is being achieved through the reaction of *p*-hydroxyacetanilide with dibromoalkanes in basic media, followed by basic hydrolysis.<sup>1</sup> An example with dibromobutane is shown in Figure 1.




Figure 1

<sup>1</sup>Bartulin, J.; Ramos, M.L.; Rivas, B.L. *Polymer Bulletin* 15, 405-409 (1986).